

**A FIRST COURSE
IN
LINEAR ALGEBRA**

A FIRST COURSE
IN
LINEAR ALGEBRA
MAT2040 Notebook

Prof. Tom Luo

The Chinese University of Hong Kong, Shenzhen

Prof. Ruoyu Sun

University of Illinois Urbana-Champaign



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen

Copyright ©2004 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 646-8600, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services please contact our Customer Care Department with the U.S. at 877-762-2974, outside the U.S. at 317-572-3993 or fax 317-572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print, however, may not be available in electronic format.

Library of Congress Cataloging-in-Publication Data:

Survey Methodology / Robert M. Groves . . . [et al.].

p. cm.—(Wiley series in survey methodology)

“Wiley-Interscience.”

Includes bibliographical references and index.

ISBN 0-471-48348-6 (pbk.)

1. Surveys—Methodology. 2. Social

sciences—Research—Statistical methods. I. Groves, Robert M. II. Series.

HA31.2.S873 2004

001.4'33—dc22

2004044064

Printed in the United States of America.

10 9 8 7 6 5 4 3 2 1

Contents

Contributors	v
Foreword	vii
Preface	ix
Acknowledgments	xi
Acronyms	xiii
1 Week1	1
1.1 Tuesday	1
1.1.1 Introduction	1
1.1.2 Gaussian Elimination	3
1.1.3 Complexity Analysis	11
1.1.4 Brief Summary	12
1.2 Thursday	14
1.2.1 Row-Echelon Form	14
1.2.2 Matrix Multiplication	16
1.2.3 Special Matrices	19
1.3 Friday	21
1.3.1 Matrix Multiplication	21
1.3.2 Elementary Matrix	22
1.3.3 Properties of Matrix	24
1.3.4 Permutation Matrix	26
1.3.5 LU decomposition	29
1.3.6 LDU decomposition	33
1.3.7 LU Decomposition with row exchanges	35
1.4 Assignment One	36

2	Week2	39
2.1	Tuesday	39
2.1.1	Review	39
2.1.2	Special matrix multiplication case	41
2.1.3	Inverse	44
2.2	Wednesday	49
2.2.1	Remarks on Gaussian Elimination	49
2.2.2	Properties of matrix	50
2.2.3	matrix transpose	53
2.3	Assignment Two	55
2.4	Friday	56
2.4.1	symmetric matrix	56
2.4.2	Interaction of inverse and transpose	57
2.4.3	Vector Space	58
2.5	Assignment Three	68
3	Week3	71
3.1	Tuesday	71
3.1.1	Introduction	71
3.1.2	Review of 2 weeks	73
3.1.3	Examples of solving equations	74
3.1.4	How to solve a general rectangular	79
3.2	Thursday	85
3.2.1	Review	85
3.2.2	Remarks on solving linear system equations	88
3.2.3	Linearly dependence	90
3.2.4	Basis and dimension	94
3.3	Friday	99
3.3.1	Review	99

3.3.2	More on basis and dimension	100
3.3.3	What is rank?	102
3.4	Assignment Four	110
4	Midterm	113
4.1	Sample Exam	113
4.2	Midterm Exam	120
5	Week4	127
5.1	Friday	127
5.1.1	Linear Transformation	127
5.1.2	Example: differentiation	130
5.1.3	Basis Change	135
5.1.4	Determinant	137
5.2	Assignment Five	144
6	Week5	147
6.1	Tuesday	147
6.1.1	Formulas for Determinant	147
6.1.2	Determinant by Cofactors	152
6.1.3	Determinant Applications	153
6.1.4	Orthogonality and Projection	156
6.2	Thursday	160
6.2.1	Orthogonality and Projection	160
6.2.2	Least Squares Approximations	165
6.2.3	Projections	168
6.3	Friday	171
6.3.1	Orthonormal basis	171
6.3.2	Gram-Schmidt Process	174

6.3.3	The Factorization $A = QR$.	180
6.3.4	Function Space	183
6.3.5	Fourier Series	184
6.4	Assignment Six	186
7	Week6	187
7.1	Tuesday	187
7.1.1	Summary of last two weeks	187
7.1.2	Eigenvalues and eigenvectors	191
7.1.3	Products and Sums of Eigenvalue	196
7.1.4	Application: Page Rank and Web Search	197
7.2	Thursday	200
7.2.1	Review	200
7.2.2	Similarity and eigenvalues	200
7.2.3	Diagonalization	203
7.2.4	Powers of A	208
7.2.5	Nondiagonalizable Matrices	209
7.3	Friday	210
7.3.1	Review	210
7.3.2	Fibonacci Numbers	210
7.3.3	Imaginary Eigenvalues	212
7.3.4	Complex Numbers	214
7.3.5	Complex Vectors	214
7.3.6	Spectral Theorem	220
7.3.7	Hermitian matrix	221
7.4	Assignment Seven	223
8	Week7	227
8.1	Tuesday	227
8.1.1	Quadratic form	227

8.1.2	Positive Definite Matrices	232
8.2	Thursday	241
8.2.1	SVD: Singular Value Decomposition	241
8.2.2	Remark on SVD decomposition	245
8.2.3	Best Low-Rank Approximation	253
8.3	Assignment Eight	255
9	Final Exam	257
9.1	Sample Exam	257
9.2	Final Exam	264
10	Solution	271
10.1	Assignment Solutions	271
10.1.1	Solution to Assignment One	271
10.1.2	Solution to Assignment Two	277
10.1.3	Solution to Assignment Three	280
10.1.4	Solution to Assignment Four	286
10.1.5	Solution to Assignment Five	297
10.1.6	Solution to Assignment Six	303
10.1.7	Solution to Assignment Seven	311
10.1.8	Solution to Assignment Eight	321
10.2	Midterm Exam Solutions	328
10.2.1	Sample Exam Solution	328
10.2.2	Midterm Exam Solution	338
10.3	Final Exam Solutions	346
10.3.1	Sample Exam Solution	346
10.3.2	Final Exam Solution	357

A	This is Appendix Title	371
A.1	This is First Level Heading	371
A.1.1	This is Second Level Heading	372

Contributors

ZHI-QUAN LUO, Shenzhen Research Institute of Big Data, Lecturer

RUOYU SUN, Industrial and Enterprise Systems Engineering, Lecturer

JIE WANG, The Chinese University of Hongkong, Shenzhen, Typer

Foreword

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Preface

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

place

date

Acknowledgments

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

I. R. S.

Acronyms

ASTA	Arrivals See Time Averages
BHCA	Busy Hour Call Attempts
BR	Bandwidth Reservation
b.u.	bandwidth unit(s)
CAC	Call / Connection Admission Control
CBP	Call Blocking Probability(-ies)
CCS	Centum Call Seconds
CDTM	Connection Dependent Threshold Model
CS	Complete Sharing
DiffServ	Differentiated Services
EMLM	Erlang Multirate Loss Model
erl	The Erlang unit of traffic-load
FIFO	First in - First out
GB	Global balance
GoS	Grade of Service
ICT	Information and Communication Technology
IntServ	Integrated Services
IP	Internet Protocol
ITU-T	International Telecommunication Unit – Standardization sector
LB	Local balance
LHS	Left hand side

LIFO	Last in - First out
MMPP	Markov Modulated Poisson Process
MPLS	Multiple Protocol Labeling Switching
MRM	Multi-Retry Model
MTM	Multi-Threshold Model
PASTA	Poisson Arrivals See Time Averages
PDF	Probability Distribution Function
pdf	probability density function
PFS	Product Form Solution
QoS	Quality of Service
r.v.	random variable(s)
RED	random early detection
RHS	Right hand side
RLA	Reduced Load Approximation
SIRO	service in random order
SRM	Single-Retry Model
STM	Single-Threshold Model
TCP	Transport Control Protocol
TH	Threshold(s)
UDP	User Datagram Protocol

2.4. Friday

2.4.1. symmetric matrix

Definition 2.5 [symmetric matrix] A $n \times n$ matrix \mathbf{A} is a **symmetric matrix** if we have $\mathbf{A}^T = \mathbf{A}$, which means $a_{ij} = a_{ji}$ for all i, j . ■

For example, the matrix \mathbf{A} shown below is a symmetric matrix:

$$\text{symmetric matrix } \mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} = \mathbf{A}^T$$

Definition 2.6 [skew-symmetric matrix] A $n \times n$ matrix \mathbf{A} is a **skew-symmetric matrix** or say, **anti-symmetric matrix** if we have $\mathbf{A} = -\mathbf{A}^T$. ■

For example, matrix \mathbf{B} shown below is a skew-symmetric matrix:

$$\text{skew-symmetric matrix } \mathbf{B} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = -\mathbf{B}^T$$

Theorem 2.3 Any $n \times n$ matrix can be decomposed as the sum of a *symmetric* and a *skew-symmetric* matrix.

Proofoutline. Given any $n \times n$ matrix \mathbf{A} , we can write \mathbf{A} as:

$$\mathbf{A} = \underbrace{\frac{\mathbf{A} + \mathbf{A}^T}{2}}_{\text{symmetric}} + \underbrace{\frac{\mathbf{A} - \mathbf{A}^T}{2}}_{\text{skew-symmetric}}$$

■

2.4.2. Interaction of inverse and transpose

Proposition 2.5 If \mathbf{A} exists, then \mathbf{A}^T also exists, and $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.

Proof.

$$(\mathbf{A}^{-1}\mathbf{A})^T = \mathbf{A}^T(\mathbf{A}^{-1})^T = \mathbf{I} \implies (\mathbf{A}^{-1})^T = (\mathbf{A}^T)^{-1}$$

■

Corollary 2.1 If matrix \mathbf{A} is symmetric and invertible, then \mathbf{A}^{-1} remains symmetric.

Proof.

$$(\mathbf{A}^{-1})^T = (\mathbf{A}^T)^{-1} = \mathbf{A}^{-1} \implies \mathbf{A}^{-1} \text{ is symmetric.}$$

■

Proposition 2.6 If $\mathbf{M} = \begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}$, then $\mathbf{M}^T = \begin{bmatrix} \mathbf{A}^T & \mathbf{C}^T \\ \mathbf{B}^T & \mathbf{D}^T \end{bmatrix}$.

Corollary 2.2 Given matrix $\mathbf{M} = \begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}$, matrix \mathbf{M} is symmetric if and only if

$$\mathbf{A} = \mathbf{A}^T, \mathbf{D} = \mathbf{D}^T, \mathbf{B}^T = \mathbf{C}.$$

Proposition 2.7 Suppose \mathbf{A} is invertible and symmetric. When we do LDU decomposition such that $\mathbf{A} = \mathbf{LDU}$, \mathbf{U} is exactly \mathbf{L}^T .

Proofoutline. Note that

$$\mathbf{A}^T = (\mathbf{LDU})^T = \mathbf{U}^T \mathbf{D}^T \mathbf{L}^T = \mathbf{A} = \mathbf{LDU}.$$

Since \mathbf{D} is diagonal matrix, we have $\mathbf{D} = \mathbf{D}^T$. It follows that

$$\mathbf{U}^T \mathbf{D} \mathbf{L}^T = \mathbf{LDU} = \mathbf{A}.$$

Since \mathbf{U}^T is also a lower triangular matrix, \mathbf{L}^T is also an upper triangular matrix, $\mathbf{U}^T \mathbf{D} \mathbf{L}^T$ is also the LDU decomposition of \mathbf{A} .

Due to the uniqueness of LDU decomposition, we obtain $\mathbf{U}^T = \mathbf{L}, \mathbf{L}^T = \mathbf{U}$. ■

2.4.3. Vector Space

We move to a new topic: vector spaces.

From Numbers to Vectors. We know matrix calculation (such as $\mathbf{Ax} = \mathbf{b}$) involves many numbers, but they are just linear combinations of n vectors.

Third Level Understanding. This topic moves from numbers and vectors to a third level of understanding (the highest level). Instead of individual column vectors, we look at "spaces" of vectors. And this topic will end with the "Fundamental Theorem of Linear Algebra".

Matrix Calculation: Numbers \implies Vectors \implies Spaces

We begin with the typical vector space, which is denoted as \mathbb{R}^n .

Definition 2.7 [Real Space] The space \mathbb{R}^n contains all column vectors v such that v has n real number entries. ■

Notation. We denote vectors as *a column between brackets*, or *along a line using commas and parentheses*:

$$\begin{bmatrix} 4 \\ \pi \end{bmatrix} \text{ is in } \mathbb{R}^2 \quad (1, 1, 1) \text{ is in } \mathbb{R}^3.$$

Definition 2.8 [vector space] A **vector space** \mathbf{V} is a set of vectors such that these vectors satisfy *vector addition* and *scalar multiplication*:

- **vector addition:** If vector v and w is in \mathbf{V} , then $v + w \in \mathbf{V}$.
- **scalar multiplication:** If vector $v \in \mathbf{V}$, then $cv \in \mathbf{V}$ for any real numbers c .

In other words, the set of vectors is **closed** under *addition* $v + w$ and *multiplication* cv .

In other words,

any linear combination is closed under vector space.

Proposition 2.8 Every vector space must contain the zero vector.

Proof. Given $v \in V \implies -v \in V \implies v + (-v) = \mathbf{0} \in V$. ■

■ **Example 2.3**

$$V = \left\{ \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \\ \vdots \end{pmatrix} \mid \{a_n\} \text{ is infinite length sequences.} \right\}$$

is a vector space.

This is because for any vector $v = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \\ \vdots \end{pmatrix}$, $w = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \\ \vdots \end{pmatrix}$, we can define vector addition

and scalar multiplication as follows:

$$v + w = \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ \vdots \\ a_n + b_n \\ \vdots \end{pmatrix} \quad cv = \begin{pmatrix} ca_1 \\ ca_2 \\ \vdots \\ ca_n \\ \vdots \end{pmatrix} \quad \text{for any } c \in \mathbb{R}.$$

$$\mathbf{V} = \text{span} \left\{ v_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{4} \\ \vdots \\ \frac{1}{2^n} \\ \vdots \end{pmatrix}, v_2 = \begin{pmatrix} \frac{1}{3} \\ \frac{1}{9} \\ \vdots \\ \frac{1}{3^n} \\ \vdots \end{pmatrix}, v_3 = \begin{pmatrix} \frac{1}{4} \\ \frac{1}{16} \\ \vdots \\ \frac{1}{4^n} \\ \vdots \end{pmatrix} \right\}$$

$$= \{ \alpha_1 v_1 + \alpha_2 v_2 + \alpha_3 v_3 \mid \alpha_1, \alpha_2, \alpha_3 \in \mathbb{R} \}$$

is also vector space.

Definition 2.9 [Span] The **span** of a collection of vectors $\mathbf{a}_1, \dots, \mathbf{a}_n \in \mathbb{R}^m$ is defined as:

$$\text{span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\} = \left\{ \mathbf{y} \in \mathbb{R}^m \mid \mathbf{y} = \sum_{i=1}^n \alpha_i \mathbf{a}_i, \alpha_i \in \mathbb{R} \right\},$$

i.e., it is the set of all linear combinations of $\mathbf{a}_1, \dots, \mathbf{a}_n$. ■

How to check \mathbf{V} is a vector space?

Given any two vectors u, w in \mathbf{V} , suppose

$$u = \alpha_1 v_1 + \alpha_2 v_2 + \alpha_3 v_3, \quad w = \beta_1 v_1 + \beta_2 v_2 + \beta_3 v_3,$$

then we obtain:

$$\begin{aligned} \gamma_1 u + \gamma_2 w &= \gamma_1 (\alpha_1 v_1 + \alpha_2 v_2 + \alpha_3 v_3) + \gamma_2 (\beta_1 v_1 + \beta_2 v_2 + \beta_3 v_3) \\ &= (\gamma_1 \alpha_1 + \gamma_2 \beta_1) v_1 + (\gamma_1 \alpha_2 + \gamma_2 \beta_2) v_2 + (\gamma_1 \alpha_3 + \gamma_2 \beta_3) v_3 \end{aligned}$$

where $\gamma_1, \gamma_2 \in \mathbb{R}$. Hence any linear combination of u and w are also in \mathbf{V} . Hence \mathbf{V} is a vector space. ■

■ **Example 2.4** $\mathbf{F} = \{f(x) \mid f: [0,1] \mapsto \mathbb{R}\}$ is also a vector space. (verify it by yourself.)

This vector space \mathbf{F} contains all real functions defined on $[0,1]$, and it is infinite dimensional.

Given two functions f and g in \mathbf{F} , the inner product of f and g is defined as:

$$\langle f, g \rangle := \int_0^1 f(x)g(x) dx$$

Also, we can use the span to form a vector space:

$$\mathbf{F} = \text{span}\{\sin x, x^3, e^x\} = \{\alpha_1 \sin x + \alpha_2 x^3 + \alpha_3 e^x \mid \alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}\}$$

This set \mathbf{F} is also a vector space. ■

■ Example 2.5

$$\mathbf{V} = \left\{ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix} \middle| a_{ij} \in \mathbb{R} \text{ for } i = 1, 2; j = 1, 2, 3. \right\}$$

is a vector space. Moreover, it is equivalent to the span of six basic vectors:

$$\mathbf{V} = \text{span} \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$$

We say that \mathbf{V} is 6-dimensional without introducing the definition of dimension formally. ■

■ Example 2.6

$$\mathbf{V} = \left\{ \begin{bmatrix} a_{ij} \end{bmatrix}_{3 \times 3} \middle| \text{any } 3 \times 3 \text{ matrices} \right\}$$

is also a vector space.

Obviously, it is 9-dimensional. We usually denote it as $\dim(\mathbf{V}) = 9$.

$$\mathbf{V}_1 = \left\{ \begin{bmatrix} a_{ij} \end{bmatrix}_{3 \times 3} \middle| \text{any } 3 \times 3 \text{ symmetric matrices} \right\}$$

is a special vector space.

Notice that $\mathbf{V}_1 \subset \mathbf{V}$, so we say \mathbf{V}_1 is a *subspace* of \mathbf{V} . In the future we will know $\dim(\mathbf{V}_1) = 6 < 9$. ■

2.4.3.1. The solution to $\mathbf{Ax} = \mathbf{0}$

We can use vector space to discuss the solution to system of equation. Firstly, let's introduce some definitions:

Definition 2.10 [homogeneous equations] A system of linear equations is said to be **homogeneous** if the constants on the righthand side are all zero. In other words, $\mathbf{Ax} = \mathbf{0}$ is said to be **homogeneous**. ■

Definition 2.11 [column space] The column space consists of all linear combinations of the columns of matrix \mathbf{A} . In other words, for the matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$ given by $\mathbf{A} = \begin{bmatrix} a_1 & a_2 & \dots & a_n \end{bmatrix}$, its column space is denoted as

$$\mathbf{C}(\mathbf{A}) := \text{span}(a_1, a_2, \dots, a_n) \subset \mathbb{R}^m.$$

Definition 2.12 [null space] The null space of a matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$ consists of all solutions to $\mathbf{Ax} = \mathbf{0}$, which can be denoted as

$$\mathbf{N}(\mathbf{A}) = \{\mathbf{x} \mid \mathbf{Ax} = \mathbf{0}\} \subset \mathbb{R}^n.$$

Proposition 2.9 The null space $\mathbf{N}(\mathbf{A})$ is a vector space.

Proofoutline. For any two vectors $\mathbf{x}, \mathbf{y} \in \mathbf{N}(\mathbf{A})$, we have $\mathbf{Ax} = \mathbf{0}, \mathbf{Ay} = \mathbf{0}$.

$$\implies \mathbf{A}(\alpha\mathbf{x} + \beta\mathbf{y}) = \alpha(\mathbf{Ax}) + \beta(\mathbf{Ay}) = \alpha\mathbf{0} + \beta\mathbf{0} = \mathbf{0} \quad \alpha, \beta \in \mathbb{R}.$$

Since the linear combination of \mathbf{x} and \mathbf{y} is also in $\mathbf{N}(\mathbf{A})$, $\mathbf{N}(\mathbf{A})$ is a vector space. ■

■ **Example 2.7** Describe the null space of $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 5 & 0 \\ 2 & 3 \end{bmatrix}$.

Obviously, converting matrix into linear system of equation we obtain:

$$\begin{cases} x_1 + 0x_2 = 0 \\ 5x_1 + 0x_2 = 0 \\ 2x_1 + 3x_2 = 0 \end{cases}$$

We can easily obtain the solution $\begin{cases} x_1 = 0 \\ x_2 = 0 \end{cases}$. Hence the null space is $\mathbf{N}(\mathbf{A}) = \mathbf{0}$. ■

■ **Example 2.8** Describe the null space of $\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 5 & 4 & 9 \\ 2 & 3 & 5 \end{bmatrix}$.

In the next lecture we will know its null space is a line.

We find that $\mathbf{A} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = \mathbf{0}$, so $\begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$ is a special solution.

Note that *the null space contains all linear combinations of special solutions*. Hence

the null space is $\mathbf{N}(\mathbf{A}) = \left\{ c \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} \mid c \in \mathbb{R} \right\}$. ■

2.4.3.2. The complete solution to $\mathbf{Ax} = \mathbf{b}$

In order to find all solutions of $\mathbf{Ax} = \mathbf{b}$, (\mathbf{A} may not be square matrix), let's introduce two kinds of solutions:

Definition 2.13 [Particular & Special Solution] For the system of equations $\mathbf{Ax} = \mathbf{b}$,

there are two kinds of solutions:

$\mathbf{x}_{\text{particular}}$ The particular solution that solves $\mathbf{Ax} = \mathbf{b}$

$\mathbf{x}_{\text{nullspace}}$ The special solutions that solves $\mathbf{Ax} = \mathbf{0}$

There is a theorem that helps us to obtain the complete solution to $\mathbf{Ax} = \mathbf{b}$.

Theorem 2.4 Any solution to $\mathbf{Ax} = \mathbf{b}$ can be represented as $\mathbf{x}_{\text{complete}} = \mathbf{x}_p + \mathbf{x}_n$.

Proof. Sufficiency. Given $\mathbf{x}_{\text{complete}} = \mathbf{x}_p + \mathbf{x}_n$, it suffices to show $\mathbf{x}_{\text{complete}}$ is the solution to $\mathbf{Ax} = \mathbf{b}$.

Note that

$$\mathbf{Ax}_{\text{complete}} = \mathbf{A}(\mathbf{x}_p + \mathbf{x}_n) = \mathbf{Ax}_p + \mathbf{Ax}_n = \mathbf{b} + \mathbf{0} = \mathbf{b}.$$

Hence $\mathbf{x}_{\text{complete}}$ is the solution to $\mathbf{Ax} = \mathbf{b}$. ■

Necessity. Suppose \mathbf{x}^* is the solution to $\mathbf{Ax} = \mathbf{b}$, it suffices to show \mathbf{x}^* could be represented as $\mathbf{x}_p + \mathbf{x}_n$.

It suffices to show $\mathbf{x}^* - \mathbf{x}_p \in N(\mathbf{A})$.

Notice that $\mathbf{A}(\mathbf{x}^* - \mathbf{x}_p) = \mathbf{Ax}^* - \mathbf{Ax}_p = \mathbf{b} - \mathbf{b} = \mathbf{0} \implies \mathbf{x}^* - \mathbf{x}_p \in N(\mathbf{A})$. ■

■ **Example 2.9** Let's study a system that has $n = 2$ unknowns but only $m = 1$ equation:

$$x_1 + x_2 = 2.$$

It's easy to check that the particular solution is $\mathbf{x}_p = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$, the special solutions are

$\mathbf{x}_n = c \begin{pmatrix} 1 \\ -1 \end{pmatrix}$, c can be taken arbitrarily.

Hence the complete solution for the equations could be written as

$$\mathbf{x}_{complete} = \mathbf{x}_p + \mathbf{x}_n = \begin{pmatrix} c+1 \\ -c+1 \end{pmatrix}.$$

So we summarize that if there are n unknowns and m equations such that $m < n$, then $\mathbf{Ax} = \mathbf{b}$ is **underdetermined** (It may have infinitely many solutions since the special solutions could be infinite).




Figure 2.1: Complete solution = one particular solution + all nullspace solutions

2.4.3.3. Row-Echelon Matrices

Given $m \times n$ rectangular matrix \mathbf{A} , we can still do Gaussian Elimination to convert \mathbf{A} into \mathbf{U} , where \mathbf{U} is of **Row Echelon form**. The whole process could be expressed as:

$$\mathbf{PA} = \mathbf{LDU}.$$

where \mathbf{L} is $m \times m$ **lower triangular** matrix, \mathbf{U} is $m \times n$ matrix that is of **row echelon form**.

■ **Example 2.10** Here is a 4×7 row echelon matrix with the three pivots **1** highlighted

in blue:

$$U = \begin{bmatrix} 1 & \times & \times & \times & \times & \times & \times \\ 0 & 1 & \times & \times & \times & \times & \times \\ 0 & 0 & 0 & 0 & 0 & 1 & \times \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- Columns 3,4,5,7 have no pivots, and we say the free variables are x_3, x_4, x_5, x_7 .
- Columns 1,2,6 have pivots, and we say the pivot variables are x_1, x_2, x_6 .

Moreover, we can continue Gaussian Elimination to convert U into R that is of **reduced row echelon form**:

$$R = \begin{bmatrix} 1 & 0 & \times & \times & \times & 0 & \times \\ 0 & 1 & \times & \times & \times & 0 & \times \\ 0 & 0 & 0 & 0 & 0 & 1 & \times \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The reduced row echelon matrix R has zeros above the pivots as well as below. Zeros above the pivots come from upward elimination. ■

R Remember the two steps (forward and back elimination) in solving $Ax = b$:

1. **Forward Elimination** takes A to U . (or its reduced form R)
2. **Back Elimination** in $Ux = c$ or $Rx = d$ produces x .

2.4.3.4. Problem Size Analysis

When faced with $m \times n$ matrix A , notice that m refers to the **number of equations**, n refers to the **number of variables**. Assume r denotes **number of pivots**, then we know r is also the **number of pivot variables**, $n - r$ is the **number of free variables**. Finally we have $m - r$ **redundant equations** and r **irredundant equations**. In next lecture, we will introduce the definition for r formally (rank).

