

**A FIRST COURSE  
IN  
LINEAR ALGEBRA**



---

**A FIRST COURSE**  
**IN**  
**LINEAR ALGEBRA**  
**MAT2040 Notebook**

---

**Prof. Tom Luo**

*The Chinese University of Hong Kong, Shenzhen*

**Prof. Ruoyu Sun**

*University of Illinois Urbana-Champaign*



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen

Copyright ©2004 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 646-8600, or on the web at [www.copyright.com](http://www.copyright.com). Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008.

**Limit of Liability/Disclaimer of Warranty:** While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services please contact our Customer Care Department with the U.S. at 877-762-2974, outside the U.S. at 317-572-3993 or fax 317-572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print, however, may not be available in electronic format.

**Library of Congress Cataloging-in-Publication Data:**

Survey Methodology / Robert M. Groves . . . [et al.].

p. cm.—(Wiley series in survey methodology)

“Wiley-Interscience.”

Includes bibliographical references and index.

ISBN 0-471-48348-6 (pbk.)

1. Surveys—Methodology. 2. Social

sciences—Research—Statistical methods. I. Groves, Robert M. II. Series.

HA31.2.S873 2004

001.4'33—dc22

2004044064

Printed in the United States of America.

10 9 8 7 6 5 4 3 2 1

# Contents

<b>Contributors</b>	<b>v</b>
<b>Foreword</b>	<b>vii</b>
<b>Preface</b>	<b>ix</b>
<b>Acknowledgments</b>	<b>xi</b>
<b>Acronyms</b>	<b>xiii</b>
<b>1 Week1</b>	<b>1</b>
<b>1.1 Tuesday</b>	<b>1</b>
1.1.1 Introduction	1
1.1.2 Gaussian Elimination	3
1.1.3 Complexity Analysis	11
1.1.4 Brief Summary	12
<b>1.2 Thursday</b>	<b>14</b>
1.2.1 Row-Echelon Form	14
1.2.2 Matrix Multiplication	16
1.2.3 Special Matrices	19
<b>1.3 Friday</b>	<b>21</b>
1.3.1 Matrix Multiplication	21
1.3.2 Elementary Matrix	22
1.3.3 Properties of Matrix	24
1.3.4 Permutation Matrix	26
1.3.5 LU decomposition	29
1.3.6 LDU decomposition	33
1.3.7 LU Decomposition with row exchanges	35
<b>1.4 Assignment One</b>	<b>36</b>

<b>2</b>	<b>Week2</b>	<b>39</b>
<b>2.1</b>	<b>Tuesday</b>	<b>39</b>
2.1.1	Review	39
2.1.2	Special matrix multiplication case	41
2.1.3	Inverse	44
<b>2.2</b>	<b>Wednesday</b>	<b>49</b>
2.2.1	Remarks on Gaussian Elimination	49
2.2.2	Properties of matrix	50
2.2.3	matrix transpose	53
<b>2.3</b>	<b>Assignment Two</b>	<b>55</b>
<b>2.4</b>	<b>Friday</b>	<b>56</b>
2.4.1	symmetric matrix	56
2.4.2	Interaction of inverse and transpose	57
2.4.3	Vector Space	58
<b>2.5</b>	<b>Assignment Three</b>	<b>68</b>
<b>3</b>	<b>Week3</b>	<b>71</b>
<b>3.1</b>	<b>Tuesday</b>	<b>71</b>
3.1.1	Introduction	71
3.1.2	Review of 2 weeks	72
3.1.3	Examples of solving equations	73
3.1.4	How to solve a general rectangular	78
<b>3.2</b>	<b>Thursday</b>	<b>83</b>
3.2.1	Review	83
3.2.2	Remarks on solving linear system equations	86
3.2.3	Linear dependence	88
3.2.4	Basis and dimension	90
<b>3.3</b>	<b>Friday</b>	<b>96</b>
3.3.1	Review	96

3.3.2	More on basis and dimension	97
3.3.3	What is rank?	99
<b>3.4</b>	<b>Assignment Four</b>	<b>110</b>
<b>4</b>	<b>Midterm</b>	<b>113</b>
<b>4.1</b>	<b>Sample Exam</b>	<b>113</b>
<b>4.2</b>	<b>Midterm Exam</b>	<b>120</b>
<b>5</b>	<b>Week4</b>	<b>127</b>
<b>5.1</b>	<b>Friday</b>	<b>127</b>
5.1.1	Linear Transformation	127
5.1.2	Example: differentiation	130
5.1.3	Basis Change	135
5.1.4	Determinant	137
<b>5.2</b>	<b>Assignment Five</b>	<b>144</b>
<b>6</b>	<b>Week5</b>	<b>147</b>
<b>6.1</b>	<b>Tuesday</b>	<b>147</b>
6.1.1	Formulas for Determinant	147
6.1.2	Determinant by Cofactors	152
6.1.3	Determinant Applications	153
6.1.4	Orthogonality and Projection	156
<b>6.2</b>	<b>Thursday</b>	<b>160</b>
6.2.1	Orthogonality and Projection	160
6.2.2	Least Squares Approximations	165
6.2.3	Projections	168
<b>6.3</b>	<b>Friday</b>	<b>171</b>
6.3.1	Orthonormal basis	171
6.3.2	Gram-Schmidt Process	174

6.3.3	The Factorization $A = QR$ .	180
6.3.4	Function Space	183
6.3.5	Fourier Series	184
<b>6.4</b>	<b>Assignment Six</b>	<b>186</b>
<b>7</b>	<b>Week6</b>	<b>187</b>
<b>7.1</b>	<b>Tuesday</b>	<b>187</b>
7.1.1	Summary of last two weeks	187
7.1.2	Eigenvalues and eigenvectors	191
7.1.3	Products and Sums of Eigenvalue	196
7.1.4	Application: Page Rank and Web Search	197
<b>7.2</b>	<b>Thursday</b>	<b>200</b>
7.2.1	Review	200
7.2.2	Similarity and eigenvalues	200
7.2.3	Diagonalization	203
7.2.4	Powers of $A$	208
7.2.5	Nondiagonalizable Matrices	209
<b>7.3</b>	<b>Friday</b>	<b>210</b>
7.3.1	Review	210
7.3.2	Fibonacci Numbers	210
7.3.3	Imaginary Eigenvalues	212
7.3.4	Complex Numbers	214
7.3.5	Complex Vectors	214
7.3.6	Spectral Theorem	220
7.3.7	Hermitian matrix	221
<b>7.4</b>	<b>Assignment Seven</b>	<b>223</b>
<b>8</b>	<b>Week7</b>	<b>227</b>
<b>8.1</b>	<b>Tuesday</b>	<b>227</b>
8.1.1	Quadratic form	227



8.1.2	Positive Definite Matrices . . . . .	232
<b>8.2</b>	<b>Thursday</b>	<b>241</b>
8.2.1	SVD: Singular Value Decomposition . . . . .	241
8.2.2	Remark on SVD decomposition . . . . .	245
8.2.3	Best Low-Rank Approximation . . . . .	253
<b>8.3</b>	<b>Assignment Eight</b>	<b>255</b>
<b>9</b>	<b>Final Exam . . . . .</b>	<b>257</b>
<b>9.1</b>	<b>Sample Exam</b>	<b>257</b>
<b>9.2</b>	<b>Final Exam</b>	<b>264</b>
<b>10</b>	<b>Solution . . . . .</b>	<b>271</b>
<b>10.1</b>	<b>Assignment Solutions</b>	<b>271</b>
10.1.1	Solution to Assignment One . . . . .	271
10.1.2	Solution to Assignment Two . . . . .	277
10.1.3	Solution to Assignment Three . . . . .	280
10.1.4	Solution to Assignment Four . . . . .	286
10.1.5	Solution to Assignment Five . . . . .	297
10.1.6	Solution to Assignment Six . . . . .	303
10.1.7	Solution to Assignment Seven . . . . .	311
10.1.8	Solution to Assignment Eight . . . . .	321
<b>10.2</b>	<b>Midterm Exam Solutions</b>	<b>328</b>
10.2.1	Sample Exam Solution . . . . .	328
10.2.2	Midterm Exam Solution . . . . .	338
<b>10.3</b>	<b>Final Exam Solutions</b>	<b>346</b>
10.3.1	Sample Exam Solution . . . . .	346
10.3.2	Final Exam Solution . . . . .	357

<b>A</b>	<b>This is Appendix Title .....</b>	<b>371</b>
<b>A.1</b>	<b>This is First Level Heading</b>	<b>371</b>
A.1.1	This is Second Level Heading .....	372

# Contributors

ZHI-QUAN LUO, Shenzhen Research Institute of Big Data, Lecturer

RUOYU SUN, Industrial and Enterprise Systems Engineering, Lecturer

JIE WANG, The Chinese University of Hongkong, Shenzhen, Typer



# Foreword

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.



# Preface

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

place

date





# Acknowledgments

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

I. R. S.



# Acronyms

ASTA	Arrivals See Time Averages
BHCA	Busy Hour Call Attempts
BR	Bandwidth Reservation
b.u.	bandwidth unit(s)
CAC	Call / Connection Admission Control
CBP	Call Blocking Probability(-ies)
CCS	Centum Call Seconds
CDTM	Connection Dependent Threshold Model
CS	Complete Sharing
DiffServ	Differentiated Services
EMLM	Erlang Multirate Loss Model
erl	The Erlang unit of traffic-load
FIFO	First in - First out
GB	Global balance
GoS	Grade of Service
ICT	Information and Communication Technology
IntServ	Integrated Services
IP	Internet Protocol
ITU-T	International Telecommunication Unit – Standardization sector
LB	Local balance
LHS	Left hand side

LIFO	Last in - First out
MMPP	Markov Modulated Poisson Process
MPLS	Multiple Protocol Labeling Switching
MRM	Multi-Retry Model
MTM	Multi-Threshold Model
PASTA	Poisson Arrivals See Time Averages
PDF	Probability Distribution Function
pdf	probability density function
PFS	Product Form Solution
QoS	Quality of Service
r.v.	random variable(s)
RED	random early detection
RHS	Right hand side
RLA	Reduced Load Approximation
SIRO	service in random order
SRM	Single-Retry Model
STM	Single-Threshold Model
TCP	Transport Control Protocol
TH	Threshold(s)
UDP	User Datagram Protocol

## 3.3. Friday

### 3.3.1. Review

**Proposition 3.2** Undetermined system  $\mathbf{Ax} = \mathbf{b}$  with  $m < n$ , i.e., number of equations  $<$  number of unknowns, has **no solution** or **infinitely many solutions**.

We want to understand the meaning of rank: number of "real" equations.

Then we introduce definition of *linearly independence* and *linearly dependence*.

The linear dependence has relation with the system:

**Proposition 3.3**  $\mathbf{Ax} = \mathbf{0}$  has nonzero solutions if and only if the column vectors of  $\mathbf{A}$  are dep.

Combining proposition (3.3) with (3.2), we derive the corollary:

**Corollary 3.3** Any  $(n + 1)$  vectors in  $\mathbb{R}^n$  are dep.

**Proposition 3.4** Undetermined system  $\mathbf{Ax} = \mathbf{b}$  with  $m \geq n$ , i.e., number of equations  $\geq$  number of unknowns may have **no solution** or **unique solution** or **infinitely many solutions**.

From this proposition we derive the corollary immediately:

**Corollary 3.4** Any  $(n - 1)$  vectors in  $\mathbb{R}^n$  cannot span the whole space.

Then we introduce the definition of basis:

**Definition 3.6** [Basis] A set of ind. vectors that span this space is called the **basis** of this space. ■

Then we introduce a theorem saying that **All basis of a given vector space have the same size**.

Thus we introduce **dimension** to denote the *number of vectors in a basis*.

### 3.3.2. More on basis and dimension

The basis of a given vector space has to satisfy two conditions:

$$\underbrace{\text{linear independence}}_{\text{not too many}} + \underbrace{\text{span the space}}_{\text{not too few}}$$

The **ind.** constraint let the size of basis not too many. For example, if given 1000 vectors of  $\mathbb{R}^3$ , they are very likely to be dep.

**Spanning the space** let the size of basis not too few. For example, given only 3 vectors of  $\mathbb{R}^{100}$ , they cannot span the whole space obviously.

We claim that:

$$\begin{aligned} \text{A basis} &= \text{maximal ind. set} \\ &= \text{minimal spanning set} \end{aligned}$$

**Definition 3.7** [spanning set]  $v_1, v_2, \dots, v_n$  is said to be the spanning set of  $V$  if

$$V = \text{span}\{v_1, v_2, \dots, v_n\}.$$

■ **Example 3.14**  $v_1 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$  is not a basis of  $\mathbb{R}^3$ .

We can add  $v_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ , which is ind. of  $v_1$ . But  $v_1, v_2$  still don't form a basis.

If we add one more vector  $v_3 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ , then  $v_1, v_2, v_3$  form a basis of  $\mathbb{R}^3$ . ■

**Theorem 3.3** Let  $V$  be a space of dimension  $n > 0$ , then

1. Any set of  $n$  ind. vectors span  $V$ .
2. Any  $n$  vectors that span  $V$  are ind.

Here is the proof outline, but you should complete the proof in detail.

*proofoutline.* 1. Suppose  $v_1, v_2, \dots, v_n$  are ind. and  $v$  is an arbitrary vector in  $V$ .

Firstly, show that  $v_1, v_2, \dots, v_n, v$  is dep., thus derive the equation  $c_1v_1 + c_2v_2 + \dots + c_nv_n + c_{n+1}v = \mathbf{0}$ . Argue that the scalar  $c_{n+1} \neq 0$ . Then we can express  $v$  in form of  $v_1, v_2, \dots, v_n$ , i.e.,  $v_1, v_2, \dots, v_n$  span  $V$ .

2. Suppose  $v_1, v_2, \dots, v_n$  span  $V$ . Assume  $v_1, v_2, \dots, v_n$  are dep. Then show that  $v_n$  could be written as form of other  $(n-1)$  vectors, it follows that  $v_1, v_2, \dots, v_{n-1}$  still span  $V$ . If  $v_1, v_2, \dots, v_{n-1}$  are also dep, we can continue eliminating one vector. We continue this way until we get an ind. spanning set with  $k < n$  elements, which contradicts  $\dim(V) = n$ . Therefore,  $v_1, v_2, \dots, v_n$  must be ind. ■

■ **Example 3.15**  $\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$  are ind.  $\implies$  they span  $\mathbb{R}^3$ . ■

### 3.3.2.1. Clarification of dimension

Firstly, we need to understand “set”:

1.  $P \triangleq \{\text{All polynomials}\} = \text{span}\{1, x, x^2, \dots\} \implies \dim(P) = \infty$ .
2.  $P_3 \triangleq \{\text{All polynomials with degree} \leq 3\} = \text{span}\{1, x, x^2, x^3\} \implies \dim(P) = 4$ .
3.  $Q \triangleq \text{span}\{x^2, 1 + x^3 + x^{10}, x^{300}\} \implies \dim(Q) = 3$ .

- Ⓡ dim of space  $\neq$  dim of the space it lives in.  
 For example, the line in  $\mathbb{R}^{100}$  has dim 1.

### 3.3.3. What is rank?

**Definition 3.8** [Rank] The rank of matrix  $\mathbf{A}$  is defined as the **number of nonzero pivots of rref of  $\mathbf{A}$** .

■ **Example 3.16**

$$\mathbf{A} = \begin{bmatrix} 1 & 3 & 3 & 4 \\ 2 & 6 & 9 & 7 \\ -1 & -3 & 3 & 4 \end{bmatrix} \xrightarrow{\text{row transform}} \mathbf{U} = \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\mathbf{U}$  has two pivots, hence  $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{U}) = 2$ .

However, the definition for rank is too complicated, can we define rank of  $\mathbf{A}$  directly?

**Key question: What quantity is not changed under row transformation?**

*Answer:* Dimension of row space.

**Definition 3.9** [column space] The **column space** of a matrix is the subspace of  $\mathbb{R}^n$  spanned by the columns.

In other words, suppose  $\mathbf{A} = \left[ \begin{array}{c|c|c} a_1 & \dots & a_n \end{array} \right]$ , the column space of  $\mathbf{A}$  is given by

$$\mathcal{C}(\mathbf{A}) = \text{span}\{a_1, a_2, \dots, a_n\}.$$

**Definition 3.10** [row space] The **row space** of a matrix is the subspace of  $\mathbb{R}^n$  spanned by the rows.



Suppose  $\mathbf{A} = \begin{bmatrix} a_1 \\ \dots \\ a_n \end{bmatrix}$ , the row space of  $\mathbf{A}$  is given by

$$\mathcal{R}(\mathbf{A}) = \text{span}\{a_1, a_2, \dots, a_n\}.$$

The **row space** of  $\mathbf{A}$  is essentially  $\mathcal{R}(\mathbf{A}) := \mathcal{C}(\mathbf{A}^T)$ , i.e., the column space of  $\mathbf{A}^T$ . ■

**Proposition 3.5** Row transformation doesn't change the row space

*Proof.* After row transformation, **new rows are linear combinations of old rows**.

Hence we have  $\mathcal{R}(\text{new rows}) \subset \mathcal{R}(\text{old rows})$ .

More specifically, assuming  $\mathbf{A} \xrightarrow{\text{Row Transform}} \mathbf{B}$ , then we have  $\mathcal{R}(\mathbf{B}) \subset \mathcal{R}(\mathbf{A})$ .

Since row transformations are invertible, we also have  $\mathbf{B} \xrightarrow{\text{Row Transform}} \mathbf{A}$ , thus we have  $\mathcal{R}(\mathbf{A}) \subset \mathcal{R}(\mathbf{B})$ .

In conclusion, we obtain  $\mathcal{R}(\mathbf{B}) = \mathcal{R}(\mathbf{A})$ . ■

Hence  $\text{rank}(\mathbf{A}) = \text{pivots of } \mathbf{U} = \dim(\text{row}(\mathbf{U})) = \dim(\text{row}(\mathbf{A}))$ .

Hence we have a much simpler definition for rank:

**Definition 3.11** [rank] The **dimension of the row space** is the **rank of a matrix**, i.e.,

$$\text{rank}(\mathbf{A}) = \dim(\mathcal{R}(\mathbf{A})).$$

In the example (3.15), we find  $\dim(\text{row}(\mathbf{A})) = \dim(\text{col}(\mathbf{A})) = 2$ , is this a coincidence? *The fundamental theorem of linear algebra* gives this answer:

**Theorem 3.4** The row space and column space both have the **same** dimension  $r$ .

We call  $\dim(\mathcal{C}(\mathbf{A}))$  as *column rank*;  $\dim(\mathcal{R}(\mathbf{A}))$  as *row rank*.

In brevity, **column rank=row rank= rank**, i.e.,

$$\dim(\mathcal{C}(\mathbf{A})) = \dim(\mathcal{R}(\mathbf{A})) = \text{rank}(\mathbf{A}), \text{ for matrix } \mathbf{A}$$

Let's discuss an example to have an idea of proving it.

■ **Example 3.17**

$$\mathbf{A} = \begin{bmatrix} 1 & 3 & 3 & 4 \\ 2 & 6 & 9 & 7 \\ -1 & -3 & 3 & 4 \end{bmatrix} \xrightarrow{\text{row transform}} \mathbf{U} = \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We notice that **column rank of  $\mathbf{A} = 2$**  and **column rank of  $\mathbf{U} = 2$** .

Why do they have the same **column space dimension**?

**Wrong reason:**  $\mathbf{A}$  and  $\mathbf{U}$  has the same column space. This is false. For

example, the first column of  $\mathbf{A}$  is  $\begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \notin \text{col}(\mathbf{U})$ . The column spaces of  $\mathbf{A}$  and  $\mathbf{U}$  are **different**, but the dimension of them are **equal**.

**Right reason:**  $\mathbf{A}\mathbf{x} = \mathbf{0}$  iff.  $\mathbf{U}\mathbf{x} = \mathbf{0}$ . The same combinations of the columns are zero (or nonzero) for  $\mathbf{A}$  and  $\mathbf{U}$ .

In other words, the  $r$  pivot columns (for both  $\mathbf{A}$  and  $\mathbf{U}$ ) are independent; the  $(n - r)$  free columns (for both  $\mathbf{A}$  and  $\mathbf{U}$ ) are dependent.

For example, for  $\mathbf{U}$ , column 1 and 3 are ind.(pivot columns); column 2 and 4 are dep.(free columns).

For  $\mathbf{A}$ , column 1 and 3 are also ind.(pivot columns); column 2 and 4 are also dep.(free columns). ■

This example shows that **Row transformation doesn't change independence relations of columns**. We give a formal proof below:

**Proposition 3.6** Suppose matrix  $\mathbf{A}$  is converted into  $\mathbf{B}$  by row transformation. If a set of columns of  $\mathbf{A}$  are ind. then so are the corresponding columns of  $\mathbf{B}$ .

*Proof.* Assume  $\mathbf{A} = \left[ \begin{array}{c|c|c} a_1 & \dots & a_n \end{array} \right], \mathbf{B} = \left[ \begin{array}{c|c|c} b_1 & \dots & b_n \end{array} \right]$ .

Without loss of generality (We often denote it as "WLOG"), we assume  $a_1, a_2, \dots, a_k$  are ind.(We can achieve it by switching columns.)

We define the sub-matrices  $\hat{\mathbf{A}} = \left[ \begin{array}{c|c|c} a_1 & \dots & a_k \end{array} \right]$  and  $\hat{\mathbf{B}} = \left[ \begin{array}{c|c|c} b_1 & \dots & b_k \end{array} \right]$ .

1. Notice that  $\hat{\mathbf{A}}$  could be converted into  $\hat{\mathbf{B}}$  by row transformation.

Hence  $\hat{\mathbf{A}}\mathbf{x} = \mathbf{0}$  and  $\hat{\mathbf{B}}\mathbf{x} = \mathbf{0}$  has the same solutions.

2. On the other hand,  $a_1, a_2, \dots, a_k$  are ind. columns.

Hence  $\hat{\mathbf{A}}\mathbf{x} = \mathbf{0}$  has the only zero solution.

Combining (1) and (2),  $\hat{\mathbf{B}}\mathbf{x} = \mathbf{0}$  has the only zero solution. Hence  $b_1, b_2, \dots, b_k$  are ind. ■

We can answer why the coincidence shown in the example, i.e.,  $\mathbf{A}$  and  $\mathbf{U}$  has the same column space dimension:

**Proposition 3.7** Row transformation doesn't change the column rank.

*Proof.* Assume  $\mathbf{A} \xrightarrow{\text{row transform}} \mathbf{B}$ .

Suppose  $\dim(\mathcal{C}(\mathbf{A})) = r$ , then we pick  $r$  ind. columns of  $\mathbf{A}$ . After row transformation, they are still ind. Hence  $\dim(\mathcal{C}(\mathbf{B})) \geq r = \dim(\mathcal{C}(\mathbf{A}))$ .

Since row transformations are invertible, we get  $\mathbf{B} \xrightarrow{\text{row transform}} \mathbf{A}$ . Similarly,  $\dim(\mathcal{C}(\mathbf{A})) \geq \dim(\mathcal{C}(\mathbf{B}))$ .

Hence  $\dim(\mathcal{C}(\mathbf{A})) = \dim(\mathcal{C}(\mathbf{B}))$ . ■

Combining proposition (3.5) and (3.7), we can proof theorem (3.4):

*Proof for theorem 3.4.* Assume  $\mathbf{A} \xrightarrow{\text{row transform}} \mathbf{U}(\text{rref})$ .

- Proposition (3.5)  $\implies \dim(\mathcal{R}(\mathbf{A})) = \dim(\mathcal{R}(\mathbf{U}))$ .
- Proposition (3.7)  $\implies \dim(\mathcal{C}(\mathbf{A})) = \dim(\mathcal{C}(\mathbf{U}))$ .
- Notice that  $\dim(\mathcal{R}(\mathbf{U}))$  denotes the number of pivots,  $\dim(\mathcal{C}(\mathbf{U}))$  denotes the number of pivot columns. Obviously,  $\dim(\mathcal{R}(\mathbf{U})) = \dim(\mathcal{C}(\mathbf{U}))$ .

Hence  $\dim(\mathcal{R}(\mathbf{A})) = \dim(\mathcal{C}(\mathbf{A}))$ . ■

R  $\dim(\mathcal{R}(\mathbf{U}))$  essentially denotes the number of “real” equations.  $\dim(\mathcal{C}(\mathbf{U}))$  denotes the number of “real” variables.

So Theorem 3.4 implies that the number of “real” equations should equal to the number of “real” variables.

### 3.3.3.1. What is the null space dimension?

Assume the system  $\mathbf{Ax} = \mathbf{b}$  has  $n$  variables.

**Proposition 3.8** For matrix  $\mathbf{A}$ ,

$$\text{rank}(\mathbf{A}) + \text{rank}(N(\mathbf{A})) = n.$$

*Proof.* **Number of pivot variables + Number of free variables =  $n$ .** ■

Note that  $\mathbf{b} \in \text{col}(\mathbf{A})$  iff.  $\mathbf{Ax} = \mathbf{b}$  for some  $\mathbf{x}$ .

Hence  $\mathcal{C}(\mathbf{A})$  denotes all possible vectors in the form  $\mathbf{Ax}$ . Hence we call  $\mathcal{C}(\mathbf{A})$  as “range space” of  $\mathbf{A}$ , which is denoted as  $\text{range}(\mathbf{A})$ .

Equivalently, we have  $\dim(\text{range}(\mathbf{A})) + \dim(N(\mathbf{A})) = n$ .

**Proposition 3.9** If  $\mathbf{Ax} = \mathbf{b}$  has at least one solution, then  $\text{rank}(\mathbf{A}) = \text{rank}\left(\begin{bmatrix} \mathbf{A} & \mathbf{b} \end{bmatrix}\right)$ .

■ **Example 3.18** Suppose  $\mathbf{A} = \begin{bmatrix} a_1 & a_2 & a_3 \end{bmatrix}$ . If  $\mathbf{Ax} = \mathbf{b}$  has at least one solution, then  $\text{rank}\left(\begin{bmatrix} a_1 & a_2 & a_3 \end{bmatrix}\right) = \text{rank}\left(\begin{bmatrix} a_1 & a_2 & a_3 & b \end{bmatrix}\right)$ . ■

*Proofoutline.*

$$\mathbf{Ax} = \mathbf{b} \iff \mathbf{b} \in \mathcal{C}(\mathbf{A})$$

Hence  $\mathbf{b}$  is the linear combination of columns of  $\mathbf{A}$ . Adding one more column  $\mathbf{b}$  into  $\mathbf{A}$  doesn't change the dimension of  $\mathcal{C}(\mathbf{A})$ . Hence  $\text{rank}(\mathbf{A}) = \text{rank}\left(\begin{bmatrix} \mathbf{A} & \mathbf{b} \end{bmatrix}\right)$ . ■

**Proposition 3.10** If  $\text{rank}(\mathbf{A}) \leq n - 1$  for  $m \times n$  matrix  $\mathbf{A}$ , then  $\mathbf{Ax} = \mathbf{b}$  has **no solution** or **infinitely many solutions**.

*Proofoutline.*

$$\dim(\mathcal{C}(\mathbf{A})) + \dim(N(\mathbf{A})) = n \implies \dim(N(\mathbf{A})) \geq 1$$

So we have special solutions for  $\mathbf{Ax} = \mathbf{b}$ . For the particular solution, if doesn't exist, then we have no solution, otherwise we have infinitely many solutions. ■

**Definition 3.12** [Full Rank] For  $m \times n$  matrix  $\mathbf{A}$ , if  $\text{rank}(\mathbf{A}) = \min(m, n)$ , then we say  $\mathbf{A}$  is full rank. ■

**Theorem 3.5** For  $n \times n$  matrix  $\mathbf{A}$ , it is invertible iff.  $\text{rank}(\mathbf{A}) = n$ .

*Proof. Sufficiency.* Assume  $\text{rank}(\mathbf{A}) = r < n$ , then by row transformation, we can convert  $\mathbf{A}$  into  $\mathbf{U} := \begin{bmatrix} \mathbf{I}_r & \mathbf{B} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}$  (rref), where  $\mathbf{B} \in \mathbb{R}^{r \times (n-r)}$ . We can represent this process in matrix notation:

$$\mathbf{PA} = \mathbf{U} := \begin{bmatrix} \mathbf{I}_r & \mathbf{B} \\ \mathbf{0} & \mathbf{0} \end{bmatrix},$$

where  $\mathbf{P}$  is the product of row transformation matrices, which is obviously invertible.

Since  $\mathbf{A}$  is invertible, we let  $\mathbf{A}^{-1} = \begin{bmatrix} \mathbf{C}_1 \\ \mathbf{C}_2 \end{bmatrix}_{(r+(n-r)) \times n}$ . It follows that

$$\mathbf{P} = \mathbf{PI}_n = \mathbf{P}(\mathbf{AA}^{-1}) = (\mathbf{PA})\mathbf{A}^{-1} = \mathbf{UA}^{-1} = \begin{bmatrix} \mathbf{I}_r & \mathbf{B} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C}_1 \\ \mathbf{C}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{C}_1 + \mathbf{BC}_2 \\ \mathbf{0} \end{bmatrix}.$$

Since  $\mathbf{P}$  has  $(n - r)$  zero rows as shown above, it is not invertible, which is a contradiction.

*Necessity.* If  $\mathbf{A}$  is full rank, then it has  $n$  pivots, then by row transformation we can convert it into  $\mathbf{I}$ (rref). We can represent this process in matrix notation:

$$\mathbf{PA} = \mathbf{I}$$

where  $\mathbf{P}$  is the product of row transformation matrix. Hence  $\mathbf{P}$  is the left inverse of  $\mathbf{A}$ ,  $\mathbf{A}$  is invertible. ■

### 3.3.3.2. Matrices of rank 1

#### ■ Example 3.19

$$\mathbf{A} = \begin{bmatrix} 2 & 1 & 1 \\ 4 & 2 & 2 \\ 8 & 4 & 4 \\ -2 & -1 & -1 \end{bmatrix} \xrightarrow{\mathbf{v}^T = \begin{bmatrix} 2 & 1 & 1 \end{bmatrix}} \begin{bmatrix} \mathbf{v}^T \\ 2\mathbf{v}^T \\ 4\mathbf{v}^T \\ -\mathbf{v}^T \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 4 \\ -1 \end{bmatrix} \mathbf{v}^T \xrightarrow{\mathbf{u} = \begin{bmatrix} 1 & 2 & 4 & -1 \end{bmatrix}^T} \mathbf{u} \mathbf{v}^T$$

Here  $\text{rank}(\mathbf{A}) = 1$ . ■

**Proposition 3.11** Every rank 1 matrix  $\mathbf{A}$  has the form  $\mathbf{A} = \mathbf{u} \mathbf{v}^T$  = column vector  $\times$  row vector.

You may prove it directly by SVD decomposition (we will learn it later, but note that most theorems or propositions could be proved by SVD). Alternatively, we have another proof:

*Proof.* We set

$$\mathbf{A} = \begin{bmatrix} \mathbf{c}_1 \\ \mathbf{c}_2 \\ \vdots \\ \mathbf{c}_n \end{bmatrix},$$

where  $\mathbf{c}_i$  is row vector. WLOG, we set  $\mathbf{c}_1 \neq \mathbf{0}$  and  $\mathbf{c}_1 = \begin{pmatrix} a_1 b_1 & a_1 b_2 & \dots & a_1 b_n \end{pmatrix}$ , where  $a_1 \neq 0$ , and  $b_i (i = 1, \dots, n)$  are not all zero.

Since  $\text{rank}(\mathbf{A}) = 1$ , we have  $\dim(\mathcal{R}(\mathbf{A})) = 1$ . Hence other  $\mathbf{c}_i$  are dep. with  $\mathbf{c}_1$ . So we set

$$b_i = \frac{a_i}{a_1} \text{ for } i = 1, 2, \dots, n.$$

Thus we construct the form of  $\mathbf{A}$ :

$$\mathbf{A} = \begin{bmatrix} a_1b_1 & a_1b_2 & \dots & a_1b_n \\ a_2b_1 & a_2b_2 & \dots & a_2b_n \\ \vdots & \vdots & & \vdots \\ a_nb_1 & a_nb_2 & \dots & a_nb_n \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \begin{bmatrix} b_1 & b_2 & \dots & b_n \end{bmatrix}$$

■

Question: What about the form of rank 2?

Answer: By SVD, it has the form  $\mathbf{u}_1\mathbf{v}_1^T + \mathbf{u}_2\mathbf{v}_2^T$ .

Enjoy Your Midterm!



