

**A FIRST COURSE
IN
LINEAR ALGEBRA**

A FIRST COURSE
IN
LINEAR ALGEBRA
MAT2040 Notebook

Prof. Tom Luo

The Chinese University of Hong Kong, Shenzhen

Prof. Ruoyu Sun

University of Illinois Urbana-Champaign



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen

Copyright ©2004 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 646-8600, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services please contact our Customer Care Department with the U.S. at 877-762-2974, outside the U.S. at 317-572-3993 or fax 317-572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print, however, may not be available in electronic format.

Library of Congress Cataloging-in-Publication Data:

Survey Methodology / Robert M. Groves . . . [et al.].

p. cm.—(Wiley series in survey methodology)

“Wiley-Interscience.”

Includes bibliographical references and index.

ISBN 0-471-48348-6 (pbk.)

1. Surveys—Methodology. 2. Social

sciences—Research—Statistical methods. I. Groves, Robert M. II. Series.

HA31.2.S873 2004

001.4'33—dc22

2004044064

Printed in the United States of America.

10 9 8 7 6 5 4 3 2 1

Contents

Contributors	v
Foreword	vii
Preface	ix
Acknowledgments	xi
Acronyms	xiii
1 Week1	1
1.1 Tuesday	1
1.1.1 Introduction	1
1.1.2 Gaussian Elimination	3
1.1.3 Complexity Analysis	11
1.1.4 Brief Summary	12
1.2 Thursday	14
1.2.1 Row-Echelon Form	14
1.2.2 Matrix Multiplication	16
1.2.3 Special Matrices	19
1.3 Friday	21
1.3.1 Matrix Multiplication	21
1.3.2 Elementary Matrix	22
1.3.3 Properties of Matrix	24
1.3.4 Permutation Matrix	26
1.3.5 LU decomposition	29
1.3.6 LDU decomposition	33
1.3.7 LU Decomposition with row exchanges	35
1.4 Assignment One	36

2	Week2	39
2.1	Tuesday	39
2.1.1	Review	39
2.1.2	Special matrix multiplication case	41
2.1.3	Inverse	44
2.2	Wednesday	48
2.2.1	Remarks on Gaussian Elimination	48
2.2.2	Properties of matrix	49
2.2.3	matrix transpose	51
2.3	Assignment Two	55
2.4	Friday	56
2.4.1	symmetric matrix	56
2.4.2	Interaction of inverse and transpose	57
2.4.3	Vector Space	58
2.5	Assignment Three	68
3	Week3	71
3.1	Tuesday	71
3.1.1	Introduction	71
3.1.2	Review of 2 weeks	73
3.1.3	Examples of solving equations	74
3.1.4	How to solve a general rectangular	79
3.2	Thursday	85
3.2.1	Review	85
3.2.2	Remarks on solving linear system equations	88
3.2.3	Linearly dependence	90
3.2.4	Basis and dimension	94
3.3	Friday	99
3.3.1	Review	99

3.3.2	More on basis and dimension	100
3.3.3	What is rank?	102
3.4	Assignment Four	110
4	Midterm	113
4.1	Sample Exam	113
4.2	Midterm Exam	120
5	Week4	127
5.1	Friday	127
5.1.1	Linear Transformation	127
5.1.2	Example: differentiation	130
5.1.3	Basis Change	135
5.1.4	Determinant	137
5.2	Assignment Five	144
6	Week5	147
6.1	Tuesday	147
6.1.1	Formulas for Determinant	147
6.1.2	Determinant by Cofactors	152
6.1.3	Determinant Applications	153
6.1.4	Orthogonality and Projection	156
6.2	Thursday	160
6.2.1	Orthogonality and Projection	160
6.2.2	Least Squares Approximations	165
6.2.3	Projections	168
6.3	Friday	171
6.3.1	Orthonormal basis	171
6.3.2	Gram-Schmidt Process	174

6.3.3	The Factorization $A = QR$.	180
6.3.4	Function Space	183
6.3.5	Fourier Series	184
6.4	Assignment Six	186
7	Week6	187
7.1	Tuesday	187
7.1.1	Summary of last two weeks	187
7.1.2	Eigenvalues and eigenvectors	191
7.1.3	Products and Sums of Eigenvalue	196
7.1.4	Application: Page Rank and Web Search	197
7.2	Thursday	200
7.2.1	Review	200
7.2.2	Similarity and eigenvalues	200
7.2.3	Diagonalization	203
7.2.4	Powers of A	208
7.2.5	Nondiagonalizable Matrices	209
7.3	Friday	210
7.3.1	Review	210
7.3.2	Fibonacci Numbers	210
7.3.3	Imaginary Eigenvalues	212
7.3.4	Complex Numbers	214
7.3.5	Complex Vectors	214
7.3.6	Spectral Theorem	220
7.3.7	Hermitian matrix	221
7.4	Assignment Seven	223
8	Week7	227
8.1	Tuesday	227
8.1.1	Quadratic form	227

8.1.2	Positive Definite Matrices	232
8.2	Thursday	241
8.2.1	SVD: Singular Value Decomposition	241
8.2.2	Remark on SVD decomposition	245
8.2.3	Best Low-Rank Approximation	253
8.3	Assignment Eight	255
9	Final Exam	257
9.1	Sample Exam	257
9.2	Final Exam	264
10	Solution	271
10.1	Assignment Solutions	271
10.1.1	Solution to Assignment One	271
10.1.2	Solution to Assignment Two	277
10.1.3	Solution to Assignment Three	280
10.1.4	Solution to Assignment Four	286
10.1.5	Solution to Assignment Five	297
10.1.6	Solution to Assignment Six	303
10.1.7	Solution to Assignment Seven	311
10.1.8	Solution to Assignment Eight	321
10.2	Midterm Exam Solutions	328
10.2.1	Sample Exam Solution	328
10.2.2	Midterm Exam Solution	338
10.3	Final Exam Solutions	346
10.3.1	Sample Exam Solution	346
10.3.2	Final Exam Solution	357

A	This is Appendix Title	371
A.1	This is First Level Heading	371
A.1.1	This is Second Level Heading	372

Contributors

ZHI-QUAN LUO, Shenzhen Research Institute of Big Data, Lecturer

RUOYU SUN, Industrial and Enterprise Systems Engineering, Lecturer

JIE WANG, The Chinese University of Hongkong, Shenzhen, Typer

Foreword

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Preface

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

place

date

Acknowledgments

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

I. R. S.

Acronyms

ASTA	Arrivals See Time Averages
BHCA	Busy Hour Call Attempts
BR	Bandwidth Reservation
b.u.	bandwidth unit(s)
CAC	Call / Connection Admission Control
CBP	Call Blocking Probability(-ies)
CCS	Centum Call Seconds
CDTM	Connection Dependent Threshold Model
CS	Complete Sharing
DiffServ	Differentiated Services
EMLM	Erlang Multirate Loss Model
erl	The Erlang unit of traffic-load
FIFO	First in - First out
GB	Global balance
GoS	Grade of Service
ICT	Information and Communication Technology
IntServ	Integrated Services
IP	Internet Protocol
ITU-T	International Telecommunication Unit – Standardization sector
LB	Local balance
LHS	Left hand side

LIFO	Last in - First out
MMPP	Markov Modulated Poisson Process
MPLS	Multiple Protocol Labeling Switching
MRM	Multi-Retry Model
MTM	Multi-Threshold Model
PASTA	Poisson Arrivals See Time Averages
PDF	Probability Distribution Function
pdf	probability density function
PFS	Product Form Solution
QoS	Quality of Service
r.v.	random variable(s)
RED	random early detection
RHS	Right hand side
RLA	Reduced Load Approximation
SIRO	service in random order
SRM	Single-Retry Model
STM	Single-Threshold Model
TCP	Transport Control Protocol
TH	Threshold(s)
UDP	User Datagram Protocol

Chapter 2

Week2

2.1. Tuesday

2.1.1. Review

2.1.1.1. Solving a system of linear Equations

Gaussian Elimination. For the system of equations $\mathbf{Ax} = \mathbf{b}$, it has three cases for its solutions:

$$\mathbf{Ax} = \mathbf{b} \left\{ \begin{array}{l} \text{unique solution} \\ \text{no solution} \\ \text{infinitely many solutions} \end{array} \right.$$

We claim that

if for this system of equation it has **infinitely** many solutions, then *its columns(or rows) could be linearly combined to zero nontrivially.*

Let's raise an example to explain this statement. Let's use an augmented matrix to represent $\mathbf{Ax} = \mathbf{b}$ (Assume \mathbf{A} is a 3×3 matrix):

$$\mathbf{Ax} = \mathbf{b} \iff \left[\begin{array}{ccc|c} a_{11} & a_{12} & a_{13} & b_1 \\ a_{21} & a_{22} & a_{23} & b_2 \\ a_{31} & a_{32} & a_{33} & b_3 \end{array} \right]$$

When focusing on the columns, we may have the question: in which case does its columns could be linearly combined to zero? That means we need to choose the

coefficients c_1, c_2, c_3 such that

$$c_1 \begin{pmatrix} a_{11} \\ a_{21} \\ a_{31} \end{pmatrix} + c_2 \begin{pmatrix} a_{12} \\ a_{22} \\ a_{32} \end{pmatrix} + c_3 \begin{pmatrix} a_{13} \\ a_{23} \\ a_{33} \end{pmatrix} = 0$$

- It's obvious that when $c_1 = c_2 = c_3 = 0$ we can linearly combine the columns. So $c_1 = c_2 = c_3 = 0$ is the *trivial* solution.
- But is there any **nontrivial** solution? We claim that if this system of equation has *infinitely* many solutions, we could linearly combine the columns *nontrivially*. We will prove this statement in the end of this lecture.

If we focus on the rows, we may have the similar question and conclusion.

Matrix to describe Gaussian Elimination.

1. Firstly let's consider the nonsingular matrix \mathbf{A} without row exchange case. We find that postmultiplying elementary matrix has the same effect as doing gaussian elimination. If we finally convert \mathbf{A} into *upper triangular matrix* \mathbf{U} , we can write this process in matrix notation:

$$\mathbf{E}_n \dots \mathbf{E}_1 \mathbf{A} = \mathbf{U} \implies \mathbf{A} = (\mathbf{E}_n \dots \mathbf{E}_1)^{-1} \mathbf{U} \implies \mathbf{A} = \mathbf{E}_1^{-1} \dots \mathbf{E}_n^{-1} \mathbf{U}$$

- (a) If we define $\mathbf{L} := \mathbf{E}_1^{-1} \dots \mathbf{E}_n^{-1}$, which is a lower triangular matrix, then we finally decompose \mathbf{A} into the product of two triangular matrix:

$$\mathbf{A} = \mathbf{L}\mathbf{U}$$

- (b) We can further decompose \mathbf{A} into product of three matrices to make the diagonal entries of \mathbf{U} and \mathbf{L} to be **one**:

$$\mathbf{A} = \mathbf{L}\mathbf{D}\mathbf{U}$$

Recall that the LDU decomposition is unique for any matrix.

2. If we have to do row exchange, the process for converting \mathbf{A} into \mathbf{U} may be like the form:

$$\mathbf{E} \cdots \mathbf{EPE} \cdots \mathbf{EPE} \cdots \mathbf{EA} = \mathbf{U},$$

but we can always do row exchange first to combine all elementary matrix together, which means we can convert this process into:

$$\mathbf{E} \cdots \mathbf{EPA} = \mathbf{U} \implies \mathbf{PA} = \mathbf{LU}$$

Also, we can do LDU decomposition to get $\mathbf{PA} = \mathbf{LDU}$.

2.1.2. Special matrix multiplication case

Firstly let's introduce a new type of vector named unit vector:

Definition 2.1 [unit vector] An i th unit vector is given by:

$$e_i = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Only in i th row its entry is 1, other entries of e_i are all 0. ■

Then let's discuss some interesting matrix multiplication cases:

1. (a) Given $m \times n$ matrix $\mathbf{A} = [a_{ij}]_{m \times n}$, the product $\mathbf{A}e_i$ is given by:

$$\mathbf{A}e_i = [a_{.i}]$$

where $\begin{bmatrix} a_{:i} \end{bmatrix}$ denotes the i th column of \mathbf{A} . (It is from the MATLAB or Julia language.)

(b) Also, given a row vector $e_j^T := \begin{bmatrix} 0 & 0 & \dots & 1 & \dots & 0 \end{bmatrix}$, the product $e_j^T \mathbf{A}$ is given by:

$$e_j^T \mathbf{A} = \begin{bmatrix} a_{j:} \end{bmatrix},$$

where $\begin{bmatrix} a_{j:} \end{bmatrix}$ denotes the j th row of \mathbf{A} .

2. Secondly, we want to compute the product $\mathbf{1}^T \mathbf{A} \mathbf{1}$, where $\mathbf{1}$ denotes a column vector that all entries of $\mathbf{1}$ are 1 and $\mathbf{1}^T$ denotes the corresponding row vector.

Let's first compute $\mathbf{A} \times \mathbf{1}$, where $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\mathbf{1} \in \mathbb{R}^n$:

$$\mathbf{A} \times \mathbf{1} = \begin{pmatrix} \sum_{j=1}^n a_{1j} \\ \sum_{j=1}^n a_{2j} \\ \vdots \\ \sum_{j=1}^n a_{mj} \end{pmatrix}$$

It follows that

$$\mathbf{1}^T \mathbf{A} \mathbf{1} = \mathbf{1}^T (\mathbf{A} \mathbf{1}) = \mathbf{1}^T \begin{pmatrix} \sum_{j=1}^n a_{1j} \\ \sum_{j=1}^n a_{2j} \\ \vdots \\ \sum_{j=1}^n a_{mj} \end{pmatrix} = \langle \mathbf{1}, \mathbf{A} \mathbf{1} \rangle = \sum_{i=1}^m \sum_{j=1}^n a_{ij},$$

3. For vectors $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, we can compute $x^T \mathbf{A} y$:

$$x^T \mathbf{A} y = x^T \begin{pmatrix} \sum_{j=1}^n a_{1j} y_j \\ \sum_{j=1}^n a_{2j} y_j \\ \vdots \\ \sum_{j=1}^n a_{mj} y_j \end{pmatrix} = \sum_{i=1}^m x_i \left(\sum_{j=1}^n a_{ij} y_j \right) = \sum_{i,j} a_{ij} x_i y_j$$

4. For vectors $x \in \mathbb{R}^n$, $y \in \mathbb{R}^n$, you should distinguish $x^T y$ and xy^T :

$$x^T y = \langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

$$xy^T = \begin{bmatrix} x_1 y_1 & x_1 y_2 & \dots & x_1 y_n \\ x_2 y_1 & x_2 y_2 & \dots & x_2 y_n \\ \vdots & & \ddots & \\ x_n y_1 & x_n y_2 & \dots & x_n y_n \end{bmatrix} = \left[x_i y_j \right]_{n \times n}$$

5. For vectors $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, we can compute $x^T A y$ by using block matrix:

Firstly, We partition A into four parts:

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}_{(m_1+m_2) \times (n_1+n_2)}.$$

Then we partition vector x and y respectively:

$$x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}_{m_1+m_2}, \quad y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}_{n_1+n_2},$$

where x_1 has m_1 rows, x_2 has m_2 rows, y_1 has n_1 rows, y_2 has n_2 rows.

Then we can compute $x^T A y$:

$$x^T A y = \begin{bmatrix} x_1^T & x_2^T \end{bmatrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \sum_{i=1}^2 \sum_{j=1}^2 x_i^T A_{ij} y_j.$$

6.

Proposition 2.1 Postmultiplying Q for the vector $v = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ has the same effect of rotating v in the plane *anticlockwise* by the angle θ , where

$$Q = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}.$$

Proof. We convert vector v into the form $v = \begin{bmatrix} \rho \cos \varphi \\ \rho \sin \varphi \end{bmatrix}$, where $\rho = \sqrt{x_1^2 + x_2^2}$, and $\varphi = \arctan(\frac{x_2}{x_1})$. Hence we obtain the product of \mathbf{Q} and v :

$$\mathbf{Q}v = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \rho \cos \varphi \\ \rho \sin \varphi \end{bmatrix} = \begin{bmatrix} \rho \cos \theta \cos \varphi - \rho \sin \theta \sin \varphi \\ \rho \cos \theta \sin \varphi + \rho \sin \theta \cos \varphi \end{bmatrix} = \begin{bmatrix} \rho \cos(\theta + \varphi) \\ \rho \sin(\theta + \varphi) \end{bmatrix}$$

This is the form that this vector has been rotated anticlockwise by the angle θ . ■

7. Given $m \times n$ matrix $\mathbf{A} = [a_{ij}]$, how to flip this matrix vertically? We just need to postmultiply a special matrix:

$$\begin{bmatrix} 0 & & & 1 \\ & & 1 & \\ & \ddots & & \\ 1 & & & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} a_{m1} & a_{m2} & \cdots & a_{mn} \\ a_{(m-1)1} & a_{(m-1)2} & \cdots & a_{(m-1)n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{11} & a_{12} & \cdots & a_{1n} \end{bmatrix}$$

If we aftermultiply this matrix for the matrix \mathbf{A} , we can flip \mathbf{A} horizontally:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} 0 & & & 1 \\ & & 1 & \\ & \ddots & & \\ 1 & & & 0 \end{bmatrix} = \begin{bmatrix} a_{1n} & a_{1(n-1)} & \cdots & a_{11} \\ a_{2n} & a_{2(n-1)} & \cdots & a_{21} \\ \vdots & \vdots & \ddots & \vdots \\ a_{mn} & a_{m(n-1)} & \cdots & a_{m1} \end{bmatrix}$$

2.1.3. Inverse

Let's introduce the definition for inverse matrix:

Definition 2.2 [Inverse matrix] For $n \times n$ matrix \mathbf{A} , the matrix \mathbf{B} is said to be the **inverse** of \mathbf{A} if we have $\mathbf{AB} = \mathbf{BA} = \mathbf{I}$. If such \mathbf{B} exists, we say matrix \mathbf{A} is **invertible** or **nonsingular**. ■

And inverse matrix has some interesting properties:

Proposition 2.2 **Matrix inverse is Unique.** In other words, if we have $\mathbf{AB}_1 = \mathbf{B}_1\mathbf{A} = \mathbf{I}$

and $AB_2 = B_2A = I$, then we obtain $B_1 = B_2$.

Proof.

$$\begin{aligned} AB_1 = I &\implies B_2AB_1 = B_2I \implies B_2AB_1 = B_2 \\ &\implies (B_2A)B_1 = IB_1 = B_1 = B_2. \end{aligned}$$

■

Proposition 2.3 If we have both $AB = I$ and $CA = I$, then we have $C = B$.

Proof. On the one hand, we have

$$CAB = C(AB) = CI = C$$

On the other hand, we obtain:

$$CAB = (CA)B = IB = B$$

Hence we have $C = B$.

■

2.1.3.1. How to compute inverse? When does it exist?

Assuming the inverse of $n \times n$ matrix A exists, and we define it to be

$$A^{-1} := X = \left[x_1 \mid x_2 \mid \dots \mid x_n \right] = \left[x_{ij} \right]$$

By definition, we have $AX = I$. We write it into block columns:

$$AX = A \left[x_1 \mid x_2 \mid \dots \mid x_n \right] = I = \left[e_1 \mid e_2 \mid \dots \mid e_n \right],$$

where e_1, e_2, \dots, e_n are all unit vectors.

Hence we obtain

$$A \left[x_1 \mid x_2 \mid \dots \mid x_n \right] = \left[Ax_1 \mid Ax_2 \mid \dots \mid Ax_n \right] = \left[e_1 \mid e_2 \mid \dots \mid e_n \right].$$

Thus we only need to compute n system of equations $\mathbf{A}x_i = e_i, i = 1, \dots, n$ to get the columns of the inverse matrix \mathbf{X} . Or equivalently, we need to do Gaussian Elimination to convert the augmented matrix $\left[\mathbf{A} \mid \mathbf{I}\right]$ into the form $\left[\mathbf{I} \mid \mathbf{X}\right]$. Once we have done that, we get the inverse of \mathbf{A} immediately. Let's discuss an example to show how to achieve it:

■ **Example 2.1** Assuming we have only 3 systems of equations to solve. And we put them altogether into one Augmented matrix. And the right side of augmented matrix is an identity matrix

$$\begin{aligned}
 \left[\mathbf{A} \mid e_1 \mid e_2 \mid e_3\right] &= \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{array}\right] \xrightarrow[\mathbf{E}_{21}=\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}]{\mathbf{E}_{31}=\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}} \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 8 & 3 & 1 & 0 & 1 \end{array}\right] \\
 &\xrightarrow[\mathbf{E}_{13}=\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}]{\mathbf{E}_{32}=\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}} \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}\right] \xrightarrow[\mathbf{E}_{13}=\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}]{\mathbf{E}_{23}=\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}\right] \\
 &\Rightarrow \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & 1 & 0 & \frac{1}{2} & -\frac{3}{8} & -\frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}\right] \xrightarrow[\mathbf{E}_{12}=\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}]{\mathbf{E}_{12}=\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}} \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & -\frac{5}{8} & -\frac{6}{8} \\ 0 & 1 & 0 & \frac{1}{2} & -\frac{3}{8} & -\frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}\right]
 \end{aligned}$$

$$\Rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & -\frac{5}{16} & -\frac{6}{16} \\ 0 & 1 & 0 & \frac{1}{2} & -\frac{3}{8} & -\frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The final augmented matrix is equivalent to the system $IX = \begin{bmatrix} \frac{12}{16} & -\frac{5}{16} & -\frac{6}{16} \\ \frac{1}{2} & -\frac{3}{8} & -\frac{1}{4} \\ -1 & 1 & 1 \end{bmatrix}$.

Hence we obtain the inverse: $A^{-1} = X = \begin{bmatrix} \frac{12}{16} & -\frac{5}{16} & -\frac{6}{16} \\ \frac{1}{2} & -\frac{3}{8} & -\frac{1}{4} \\ -1 & 1 & 1 \end{bmatrix}$. ■

Then let's study in which case does the inverse exist:

Theorem 2.1 The inverse of $n \times n$ matrix A exists if and only if $Ax = b$ has a unique solution.

Proofoutline. The inverse of $n \times n$ matrix A exists

\Leftrightarrow none pivot values of A is zero. $\Leftrightarrow Ax = b$ has a unique solution $x = A^{-1}b$. ■

At the end, let's prove the claim at the beginning of the lecture:

Theorem 2.2 Let A be $n \times n$ matrix, the following statements are equivalent:

1. Columns of A can be linearly combined to zero nontrivially.
2. $Ax = 0$ has infinitely many solutions.
3. Row vectors of A can be linearly combined to zero nontrivially.

Proofoutline. The following statements are equivalent:

- Columns of A can be linearly combined to zero nontrivially.
- Given $A = \begin{bmatrix} a_1 & a_2 & \dots & a_n \end{bmatrix}$, then there exists x_i 's that are not all zero such that

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 0.$$

- $Ax = 0$ has a nonzero solution \bar{x} .

- $2\bar{x}, 3\bar{x}, \dots$ are also solutions to $A\mathbf{x} = \mathbf{0}$.
- $A\mathbf{x} = \mathbf{0}$ has infinitely many solutions.
- A^{-1} does not exist. (otherwise we will only have unique solution $A^{-1} \times \mathbf{0} = \mathbf{0}$.)
- Gaussian Elimination breaks down, i.e., there exists zero row in the row echelon form.
- Row vectors of A can be linearly combined to zero nontrivially.

■

