

Motivic Homotopy Theory

Chow.



Contents

1	Unstable Homotopy Category	1
1.1	Simplicial Methods for Algebraic Geometry	2
1.1.1	Simplicial Objects	3
1.1.2	Simplicial Homotopy Theory for Presheaves and Sheafification . .	4
1.1.3	Hypercovers and Bousfield localization	5
1.2	Motivic Homotopy Category	6
1.2.1	Realization to Motives	7
1.2.2	Homotopy Purity	8
1.3	Properties of Motivic Homotopy Category	8
2	Stable Motivic Homotopy Theory	9
2.1	T-Spectra	9
2.1.1	Motivic Spheres	9
2.1.2	Stable Weak Equivalence	10
2.1.3	Bigraded Cohomology	11
2.1.4	Motivic Eilenberg-MacLane Spectrum	11

1 Unstable Homotopy Category

In order to refine the triangulated category of Voevodsky's motives and apply more tools from algebraic topology, we need well-behaved homotopy theory for schemes. Homotopy theory for schemes also allow us to construct motivic cohomology and algebraic K-theory which are looked as generalized cohomology theory defined with Brown representativity theorem.

1.1 Simplicial Methods for Algebraic Geometry

Let us firstly recall some simplicial homotopy theory. Δ is category consists of objects such as

$$[n] = 0 \rightarrow 1 \rightarrow 2 \rightarrow \cdots \rightarrow n$$

for all non-negative integer n . And morphisms are functions of sets preserving the order of arrows. The category of *simplicial sets* means the category of presheaves on Δ with values over **Sets**. It is denoted by **sSets**.

In category of simplicial sets, we have following canonical objects

$$\Delta[n] := \Delta(-, [n])$$

They are call *standard simplicial sets*.

Proposition 1.1.1. $\Delta([m], [n])$ is generated by following morphisms

$$d^i: [k] \rightarrow [k+1]$$

$$d^i(0 \rightarrow 1 \rightarrow 2 \rightarrow \cdots \rightarrow k) = \begin{cases} 1 \rightarrow 2 \rightarrow \cdots \rightarrow k+1 & \text{if } i = 0 \\ 0 \rightarrow 1 \rightarrow \cdots \rightarrow i-1 \rightarrow i+1 \rightarrow \cdots \rightarrow k+1 & \text{if } 1 \leq i \leq k \\ 0 \rightarrow 1 \rightarrow \cdots \rightarrow k & \text{if } i = k+1 \end{cases}$$

and

$$s^i: [k] \rightarrow [k-1]$$

$$s^i(0 \rightarrow 1 \rightarrow 2 \rightarrow \cdots \rightarrow k) = \begin{cases} 0 \rightarrow 0 \rightarrow 1 \rightarrow \cdots \rightarrow k-1 & \text{if } i = 0 \\ 0 \rightarrow \cdots \rightarrow i \rightarrow i \rightarrow \cdots \rightarrow k-1 & \text{if } 1 \leq i \leq k-1 \end{cases}$$

where d^i is called *co-face map* and s^i is called *co-degeneracy map*.

Δ is small category. As diagram, Δ looks like

$$\begin{array}{ccc} & & \xrightarrow{d^1} \\ [0] & \xrightarrow[d^1]{d^0} & [1] \xrightarrow[d^2]{d^0} \cdots \\ & & \end{array}$$

$$\begin{array}{ccc} & & \xleftarrow{s^0} \\ [0] & \xleftarrow{s^1} & [1] \xleftarrow{s^1} \cdots \\ & & \end{array}$$

By Yoneda lemma, for simplicial set $X \in \mathbf{sSets}$, we have


$$X([n]) \simeq \mathbf{sSets}(\Delta[n], X)$$

For convenience, we always denote $X([n])$ by X_n for simplicial set X , $d_i = X(d^i)$ and $s_i = X(s^i)$.

Let Δ^n be standard simplex in topological space category. Then we can defined topological space respect to given simplicial set X as follows

$$|X| := \operatorname{colim}_{(\Delta_n \rightarrow X) \in \mathbf{sSets}/X} \Delta^n$$

Figure 1: Idea of simplicial sets (from ncatlab)



Furthermore, $|-|$ is covariant functorial in \mathbf{sSets} since co-representable $\mathbf{sSets}(\Delta_n, -)$ and colimit are functorial. $|-|$ is called *geometric realization functor*.

In classical simplicial homotopy theory, we can endow \mathbf{sSets} with model structure where fibrations are Kan fibrations and weak equivalence are morphisms which induce isomorphisms between fibrant objects on homotopy groups (can be defined as homotopy groups of corresponding geometric realizations).

1.1.1 Simplicial Objects

Let \mathcal{C} be arbitrary (concrete) category. We say X_* is a *simplicial object* on \mathcal{C} if X_* is a presheaf $X_*: \Delta^{op} \rightarrow \mathcal{C}$ and \mathbf{sC} denotes the category of simplicial objects on \mathcal{C} . Hence simplicial sets are simplicial objects on \mathbf{Sets} .

Let $\Delta_{\leq n}$ the full subcategory of Δ generated by objects $\{0, 1, \dots, n\}$ and $s_n\mathcal{C}$ the category of presheaves of objects in \mathcal{C} on $\Delta_{\leq n}$, these objects are called *n-truncated simplicial objects* in \mathcal{C} . Composite with inclusion $\Delta_{\leq n} \hookrightarrow \Delta$, we get functor

$$(-)_{\leq n}: \mathbf{sC} \rightarrow s_n\mathcal{C}$$

. It is called *n-truncation functor*. It has right adjoint functor

$$(-)_{\leq n}: \mathbf{sC} \rightleftarrows s_n\mathcal{C} : \text{cosk}_n$$

and left adjoint functor

$$\text{sk}_n: s_n\mathcal{C} \rightleftarrows \mathbf{sC} : (-)_{\leq n}$$

More concretely, $\text{sk}_n(X_{\leq n})$ is smallest subobject of X in $s\mathcal{C}$ agrees with X in dimension lower than n and $\text{cosk}_n(X_{\leq n})_k$ is $\mathcal{H}om(\text{sk}_n(\Delta([k])_{\leq n}), X)$ for all k .

If \mathcal{A} is an Abelian category, then we have following correspondence

Proposition 1.1.2 (Dold-Kan Correspondence).

$$N: s\mathcal{A} \rightarrow \mathbf{Ch}_{\geq 0}(\mathcal{A})$$

is equivalence of categories and is also Quillen equivalence (i.e. Quillen functor induce equivalence between homotopy categories) with respect to canonical model structure on $s\mathcal{A}$ and projective model on $\mathbf{Ch}_{\geq 0}(\mathcal{A})$. N is functor of normalized complex and $\mathbf{Ch}_{\geq 0}(\mathcal{A})$ is category of non-negative chain complexes. This equivalence is called Dold-Kan correspondence.

This proposition means that homological algebra over Abelian category in low bounded case is equivalent to homotopy theory for its simplicial objects. It is convenient to study homotopy theory for simplicial objects because we have geometric realization functor to make constructions more natural. So we can transplant properties from classical homotopy theory into homological algebra.

1.1.2 Simplicial Homotopy Theory for Presheaves and Sheafification

Suppose $\mathbf{PSh}^{\mathbf{Sets}}(\mathcal{C})$ be the category of presheaves of sets on \mathcal{C} . Then the associated category of simplicial objects $s\mathbf{PSh}^{\mathbf{Sets}}(\mathcal{C})$ is isomorphic to $\mathbf{PSh}^{s\mathbf{Sets}}(\mathcal{C})$ — the category of presheaves of simplicial sets on \mathcal{C} . Let $X_* \in s\mathbf{PSh}^{\mathbf{Sets}}(\mathcal{C})$

$$F_* \mapsto F$$

where

$$F(X)_n := F_n(X) \quad \forall X \in \mathcal{C}$$

In this case, $s\mathbf{PSh}^{\mathbf{Sets}}(\mathcal{C})$ or $\mathbf{PSh}^{s\mathbf{Sets}}(\mathcal{C})$ is called category of simplicial presheaves of sets on \mathcal{C} .

$\mathbf{PSh}^{s\mathbf{Sets}}(\mathcal{C})$ can be endowed with model structure objectwise by model structure of $s\mathbf{Sets}$.

- $\alpha: F \Rightarrow G$ is weak-equivalence if and only if for all $X \in \mathcal{C}$, $\alpha_X: F(X) \rightarrow G(X)$ is weak-equivalence in $s\mathbf{Sets}$.
- $\beta: F \Rightarrow G$ is fibration if and only if for all $X \in \mathcal{C}$, β_X is Kan fibration.

Then the model structure of $\mathbf{PSh}^{s\mathbf{Sets}}(\mathcal{C})$ is fibrantly generated. This model structure on $\mathbf{PSh}^{s\mathbf{Sets}}(\mathcal{C})$ is called *global model structure*.

If \mathcal{C} is a site with Grothendieck topology, we can define $\mathbf{Sh}_{\tau}^{\mathbf{Sets}}(\mathcal{C})$ be the category of τ -sheaves of sets on \mathcal{C} and $\mathbf{PSh}_{\tau}^{s\mathbf{Sets}}(\mathcal{C})$ the category of simplicial presheaves with model structure called *τ -local model structure* as follows.

Definition 1.1.1 (τ -local weak equivalence). A τ -local weak equivalence is a morphism between presheaves $f: F \Rightarrow G$ such that

- The morphism induces isomorphism $\tilde{\pi}_0(f): \tilde{\pi}_0 X \rightarrow \tilde{\pi}_0 Y$ in $\mathbf{Sh}^{\mathbf{Sets}}(\mathcal{C})$;

- For all $U \in \mathcal{C}$, $\tilde{\pi}_n(f) : \tilde{\pi}_n(X, x) \rightarrow \tilde{\pi}_n(Y, f(x))$ is isomorphism on \mathcal{C}/U for any choice of base point $x \in X(U)$.

The τ -local cofibrations are same as ordinary cofibrations of $\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C})$ and τ -fibrations are morphisms in $\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C})$ satisfy RLP for τ -acyclic cofibration (i.e. be both τ -cofibration and τ -weak equivalence). Jardine proved that these datum actually define a model structure.

In particular, if $\mathcal{C} = (Sm/S)_\tau$ is site of smooth S -schemes with Grothendieck topology τ , we denote $\mathbf{Spc}^{pre}(S)$ (resp. $\mathbf{Spc}_\tau^{pre}(S)$, resp. $\mathbf{Spc}_\tau(S)$) the category of simplicial presheaves (res. simplicial presheaves with τ -local model structure, resp. simplicial τ -sheaves) on $\mathcal{C} = (Sm/S)_\tau$. $\mathbf{Spc}_\tau(S)$ is full subcategory of $\mathbf{Spc}_\tau^{pre}(S)$ and left adjoint functor s of inclusion exists. This functor is called *sheafification functor* under topology τ . Hence $\mathbf{Spc}_\tau(S)$ is model category. Furthermore, we have

Proposition 1.1.3. *The pair (s, i) of adjunction is a pair of Quillen equivalence.*

Hence we have $\mathbf{Ho}(\mathbf{Spc}_\tau^{pre}(S)) \simeq \mathbf{Ho}(\mathbf{Spc}_\tau(S))$, denoted by $\mathcal{H}_\tau(S)$.

1.1.3 Hypercovers and Bousfield localization

More generally, we can realize τ -local model structure on $\mathbf{PSh}_\tau^{\mathbf{sSets}}(\mathcal{C})$ as Bousfield localization with respect to τ -hypercovers.

Let \mathcal{M} be a model category. A subcategory $\mathcal{S} \subseteq \mathcal{M}$ is called a *subcategory of weak equivalences* if all identity maps in \mathcal{M} are in \mathcal{S} , \mathcal{S} is closed under retracts and \mathcal{S} satisfies the two out of three property.

Definition 1.1.2. Suppose \mathcal{S} be a subcategory of weak equivalence. An object X of \mathcal{M} is called \mathcal{S} -local if

- X is fibrant object in \mathcal{M} ;
- For every morphism $Y \xrightarrow{f} Z \in \mathcal{S}$, the induced morphism

$$f^* : [Z, X] \rightarrow [Y, X]$$

is bijection, where $[-, -]$ is hom-set in homotopy category of \mathcal{M} .

In particular, if \mathcal{M} is a simplicial model category (we omit the definition and the only things we need to know are (1) $\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C})$ is simplicial model category; (2) Hom-sets in this category are in \mathbf{sSets}). Then $X \in \mathcal{M}$ is called \mathcal{S} -local if

- X is fibrant object in \mathcal{M} .
- For every morphism $Y \xrightarrow{f} Z \in \mathcal{S}$, the induced morphism

$$f^* : \mathcal{M}(Z, X) \rightarrow \mathcal{M}(Y, X)$$

is weak equivalence in \mathbf{sSets} .

And a morphism $X \xrightarrow{f} Y \in \mathcal{M}$ if and only if for all \mathcal{S} -local object Z , the induced morphism

$$f^*: \mathcal{M}(Y, Z) \rightarrow \mathcal{M}(X, Z)$$

is weak equivalence in **sSets**.

Definition 1.1.3 (Bousfield localization). Let $\mathbf{Fib}_{\mathcal{S}}$ be the class of morphisms in \mathcal{M} satisfy RLP for $\mathcal{S} \cap \mathbf{Cof}$. If tuple $(\mathcal{S}, \mathbf{Cof}, \mathbf{Fib}_{\mathcal{S}})$ forms a model structure on \mathcal{M} , then a \mathcal{S} -fibrant replacement functor $X \mapsto LX$ is called Bousfield localization with respect to \mathcal{S} and the new model category is denoted by $L_{\mathcal{S}}\mathcal{M}$.

The (left) Bousfield localization has following universal property: The (left) Bousfield localization $L_{\mathcal{S}}$ is a left Quillen functor whose total left derived functor sends all morphisms in \mathcal{S} to isomorphisms in $\mathbf{Ho}(L_{\mathcal{S}}\mathcal{M})$; Any left Quillen functor F whose total left derived functor sends all morphisms in \mathcal{S} into isomorphisms of $\mathbf{Ho}(L_{\mathcal{S}}\mathcal{M})$ can factors through the (left) Bousfield localization by a left Quillen functor.

Theorem 1.1.4. *For category of simplicial presheaves on essentially small category \mathcal{C} , the global model structure admits (left) Bousfield localization for all \mathcal{S} .*

Let \mathcal{C} be a site with Grothendieck topology τ .

Definition 1.1.4. Simplicial object U_* in \mathcal{C} is said to be τ -hypercov if

- $U_0 \rightarrow *$ is covering in τ .
- The unit of adjunction

$$id_{s\mathcal{C}} \Rightarrow \text{cosk}_n \circ (-)_{\leq n}$$

induces covering $\in \tau$.

$$X_{n+1} \rightarrow \text{cosk}_n(X_{\leq n})_{n+1}$$

If X is any object of \mathcal{C} , we will refer to a τ -hypercov in slice site \mathcal{C}/X as simply a τ -hypercov of X .

Then define \mathcal{W}_{τ} be the smallest subcategory of weak equivalences contains class of τ -hypercovers U_* of objects in \mathcal{C} . Denote $L_{\tau}\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C})$ the left Bousfield localization of $\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C})$. Then we have

Theorem 1.1.5. *The identity functor $L_{\tau}\mathbf{PSh}^{\mathbf{sSets}}(\mathcal{C}) \rightarrow \mathbf{PSh}_{\tau}^{\mathbf{sSets}}(\mathcal{C})$ is Quillen equivalence.*

Corollary 1.1.6. *The τ -homotopy category $\mathcal{H}_{\tau}(S)$ is equivalent to $\mathbf{Ho}(L_{\tau}\mathbf{PSh}^{\mathbf{sSets}}(Sm/S))$.*

1.2 Motivic Homotopy Category

Let k be a field.

Definition 1.2.1 (\mathbb{A}^1 -local model). The model structure of left Bousfield localization of $\mathbf{Spc}_{\tau}(k)$ with respect to class of maps $\mathbf{Spc}_{\tau}(k)(X \times_k \mathbb{A}^1, X)$, $X \in Sm_k$ is called \mathbb{A}^1 -local model.

- $X_* \in \mathbf{Spc}_\tau(k)$ is called \mathbb{A}^1 -local if X_* is hypercover of some smooth k -scheme X and the projection $X \times_k \mathbb{A}^1 \rightarrow X$ induce weak equivalence in $\mathbf{Spc}_\tau(k)$.
- Morphism $f : X_* \rightarrow Y_*$ is called \mathbb{A}^1 -weak equivalence if it induces isomorphism

$$f^* : \mathrm{Hom}_{\mathcal{H}_\tau(k)}(Y_*, Z) \rightarrow \mathrm{Hom}_{\mathcal{H}_\tau(k)}(X_*, Z)$$

for any \mathbb{A}^1 -local object Z .

Definition 1.2.2 (motivic homotopy category). The left Bousfield localization of $\mathbf{Spc}_{\mathbf{Nis}}(k)$ with respect to \mathbb{A}^1 -local model structure is denoted by $\mathbf{Spc}(k)$. Homotopy category of $\mathbf{Spc}(k)$ is called motivic homotopy category and is denoted by $\mathcal{H}(k)$.

The fibrant objects in $\mathbf{Spc}(k)$ are exactly \mathbb{A}^1 -local objects. \mathbb{A}^1 is \mathbb{A}^1 -local object and it is contractible in $\mathbf{Spc}(k)$, since

$$\mathbb{A}^1 \cong \mathrm{Spec} k \times_k \mathbb{A}^1 \rightarrow \mathrm{Spec} k$$

is \mathbb{A}^1 -weak equivalence.

Let τ be subcanonical topology with enough points. The τ -local model structure implies for every $X \in \mathbf{Sm}/k$, if $\{U, V\}$ is a τ -covering of X , then image of Yoneda embedding in $\mathbf{Spc}(k)$ of $U \coprod_{U \times_X V} V \rightarrow X$ is τ -local weak equivalence since it induces isomorphism at each τ -stalk. Hence we have *Mayer-Vietoris property*, i.e. X is homotopy push-out of

$$\begin{array}{ccc} U \times_X V & \longrightarrow & U \\ \downarrow & & \\ V & & \end{array}$$

Definition 1.2.3 (homotopy quotient). The quotient Y/X of morphisms of k -schemes $f : X \rightarrow Y$ is homotopy push-out of

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \\ \mathrm{Spec} k & & \end{array}$$

in $\mathbf{Spc}(k)$.

By M-V Property, we have the natural map $V/(U \times_X V) \rightarrow X/U$ is weak equivalence in $\mathbf{Spc}(k)$.

1.2.1 Realization to Motives

We want to construct a realization functor from $\mathbf{Spc}(k)$ to Voevodsky's category $DM_-^{\mathrm{eff}}(k)$. Since $\mathbf{Sh}_{\mathbf{Nis}}(\mathrm{Cor}(k))$ is Abelian category, the category of chain complex $\mathbf{Ch}_{\geq 0}(\mathbf{Sh}_{\mathbf{Nis}}(\mathrm{Cor}(k)))$

is category equivalent to $s\mathbf{Sh}_{\mathbf{Nis}}(\text{Cor}(k))$ and it is also Quillen equivalence. The embedding functor

$$i: s\mathbf{Sh}_{\mathbf{Nis}}(\text{Cor}(k)) \rightarrow s\mathbf{PSh}_{\mathbf{Nis}}(\text{Cor}(k))$$

is Quillen equivalence. Hence

$$DM_-^{\text{eff}}(k) \cong \mathbf{Ho}(L_{\mathbb{A}^1} C^-(\mathbf{Sh}_{\mathbf{Nis}}(\text{Cor}(k)))) \cong \mathbf{Ho}(L_{\mathbb{A}^1} s\mathbf{PSh}_{\mathbf{Nis}}(\text{Cor}(k))).$$

Since total left derived functor of

$$s\mathbf{PSh}_{\mathbf{Nis}}^{\mathbf{Sets}}(Sm/k) \xrightarrow{F} s\mathbf{PSh}_{\mathbf{Nis}}(Sm/k) \xrightarrow{i} L_{\mathbb{A}^1} s\mathbf{PSh}_{\mathbf{Nis}}(\text{Cor}(k))$$

, where F is induced functor from free generating functor $F: \mathbf{Sets} \rightarrow \mathbf{Abs}$ sends \mathbb{A}^1 -weak equivalences of $s\mathbf{PSh}_{\mathbf{Nis}}^{\mathbf{Sets}}(Sm/k)$ to isomorphisms in $DM_-^{\text{eff}}(k)$. This functor factors through Bousfield localization by universal property of Bousfield localization

$$s\mathbf{PSh}_{\mathbf{Nis}}^{\mathbf{Sets}}(Sm/k) \rightarrow \mathbf{Spc}(k)$$

So we get

$$\begin{array}{ccc} \mathbf{Spc}(k) & \longrightarrow & L_{\mathbb{A}^1} s\mathbf{PSh}_{\mathbf{Nis}}(\text{Cor}(k)) \\ & \searrow u & \downarrow \\ & & DM_-^{\text{eff}}(k) \end{array}$$

Hence u is the required realization functor.

1.2.2 Homotopy Purity

1.3 Properties of Motivic Homotopy Category

Definition 1.3.1. For $F_* \in \mathbf{Spc}_{\mathbf{Nis}}(S)$, one says F_* has Brown-Gersten property if for any distinguished square,

$$\begin{array}{ccc} F(X)_* & \longrightarrow & F(V)_* \\ \downarrow & & \downarrow \\ F(U)_* & \longrightarrow & F(U \times_X V)_* \end{array}$$

is homotopy push-out in $s\mathbf{Sets}$.

If $F_* \in \mathbf{Spc}_{\mathbf{Nis}}(\mathbf{S})$ is Nisnevich local object, then F_* has Brown-Gersten property. Yoneda lemma implies that this diagram is equal to

$$\begin{array}{ccc} \text{Hom}(X, F) & \longrightarrow & \text{Hom}(V, F) \\ \downarrow & & \downarrow \\ \text{Hom}(U, F) & \longrightarrow & \text{Hom}(U \times_X V, F) \end{array}$$

where X, U, V and $U \times_X V$ is viewed as the simplicial presheaf represented by them. If F_* is Nisnevich local object, then arrows in this diagram are all weak equivalences in **sSets**. Hence it is homotopy push-out.

Proposition 1.3.1. *If \mathcal{X} is motivic space in $\mathcal{H}(k)$, i.e. \mathcal{X} is \mathbb{A}^1 -local object in **Spc**(k), then we have*

$$\pi_0(\mathcal{X})(U) \simeq \mathcal{H}(k)(U, \mathcal{X}_{Nis})$$

Furthermore, if \mathcal{X} is pointed, then

$$\pi_n(\mathcal{X}, *) (U) \simeq \mathcal{H}_*(k)(\mathbb{S}^n \wedge U_+, \mathcal{X}_{Nis})$$

Proof. \mathcal{X} is motivic space implies \mathcal{X}_{Nis} is also fibrant object in **Spc**(k), hence by fundamental theorem of homotopical algebra, we have

$$\mathcal{H}(k)(U, \mathcal{X}_{Nis}) = [U, \mathcal{X}_{Nis}] \cong [U, \mathcal{X}] \cong \pi(U, \mathcal{X}) = \pi_0(\mathcal{X})(U)$$

□

2 Stable Motivic Homotopy Theory

In the context of unstable motivic homotopy category, we can develop the theory of \mathbb{T} -spectra and stable motivic homotopy category. Most of \mathbb{A}^1 -invariants in algebraic geometry can be proved to be representable in such stable homotopy category.

2.1 \mathbb{T} -Spectra

2.1.1 Motivic Spheres

Let **Spc** $_*(S)$ be the category of pointed \mathbb{A}^1 -presheaves, which means each object is \mathbb{A}^1 -presheaf together with base point

$$* \rightarrow F$$

We have wedge sum and smash product in **Spc** $_*(S)$,

- $\mathcal{X} \vee \mathcal{Y}$ is defined as push-out

$$\begin{array}{ccc} * & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \mathcal{Y} & \longrightarrow & \mathcal{X} \vee \mathcal{Y} \end{array}$$

- smash product $\mathcal{X} \wedge \mathcal{Y} := \frac{\mathcal{X} \times \mathcal{Y}}{\mathcal{X} \vee \mathcal{Y}}$.

We have following two kinds of circles in context of motivic homotopy theory.

- (simplicial circle) the constant sheaf $\mathbb{S}^1 = \Delta[1]/\partial\Delta[1]$. Its pointed sheaf $(\mathbb{S}^1, 0)$ is denoted by $S^{1,0}$ or S_s^1 ;

- (Tate circle) the presheaf represented by $\mathbb{G}_m = \mathbb{A}^1 - 0$, its pointed sheaf $(\mathbb{G}_m, 1)$ is denoted by $S^{1,1}$ or S_t^1 .

Hence we can define two kinds of suspension: (1) simplicial suspension $\Sigma_s X := S_s^1 \wedge X$; (2) Tate suspension $\Sigma_t X := S_t^1 \wedge X$. However, they not suspension we truly need to construct stable homotopy category. Let $\mathbb{T} = (\mathbb{P}^1, \infty)$, define $\Sigma X = \mathbb{T} \wedge X$. As we can show $\Sigma_s S_t^1$ is push-out of

$$\begin{array}{ccc} \mathbb{G}_m & \longrightarrow & * \\ \downarrow & & \\ * & & \end{array}$$

hence $\Sigma_s S_t^1$ is homotopy equivalent to $* \coprod_{\mathbb{A}^1 - 0} \mathbb{A}^1 \cong \mathbb{P}^1$. Hence $\Sigma \simeq \Sigma_s \Sigma_t$ in $\mathcal{H}_*(S)$.

Definition 2.1.1. In $\mathcal{H}_*(S)$, we define motivic spheres as following

$$S^{p,q} = \left(\bigwedge^{p-q} S_s^1 \right) \wedge \left(\bigwedge^q S_t^1 \right)$$

for all $p \geq q \geq 0$.

Since smash product \wedge can be viewed as internal tensor product in $\mathbf{Spc}_*(S)$, it can defined internal hom functor $\mathcal{H}om$ which is right adjoint to \wedge . Hence

$$\mathcal{H}_*(S)(\mathbb{T} \wedge \mathcal{X}, \mathcal{Y}) = \mathcal{H}_*(S)(\mathcal{X}, \mathcal{H}om(\mathbb{T}, \mathcal{Y}))$$

Denote $\Omega \mathcal{Y}$ the internal hom $\mathcal{H}om(\mathbb{T}, \mathcal{Y})$. In this way (Σ, Ω) is a pair of Quillen functors since smash product preserves cofibrations and acyclic cofibrations.

2.1.2 Stable Weak Equivalence

Definition 2.1.2. A \mathbb{T} -spectra is a sequence of pointed objects $\{E_0, E_1, \dots\}$ in $\mathbf{Spc}_*(S)$ together with bonding maps

$$l_n: \Sigma E_n \rightarrow E_{n+1}$$

. Morphisms between \mathbb{T} -spectrum are defined level-wise, i.e. $f_n: E_n \rightarrow K_n$ is morphism in $\mathbf{Spc}_*(S)$ and compatible with structure maps, diagram

$$\begin{array}{ccc} E_{n+1} & \xrightarrow{f_{n+1}} & K_{n+1} \\ \downarrow & & \downarrow \\ \Sigma E_n & \xrightarrow{\Sigma(f_n)} & \Sigma K_n \end{array}$$

commutes. This category is denoted by $\mathbf{Spt}^{\mathbb{T}}(S)$.

Category $\mathbf{Spt}^{\mathbb{T}}(S)$ is with canonical model structure called *projective model*: The weak equivalences and fibrations are defined level-wise and cofibrations are generated by LLP property. We have Quillen adjoint pair of suspension and loops functors $(\Sigma_{\mathbb{T}}, \Omega_{\mathbb{T}})$:

$$\begin{aligned}\Sigma_{\mathbb{T}}(E) &:= (E_1, E_2, \dots) \\ \Omega_{\mathbb{T}}(E) &:= (\Omega E_0, E_0, E_1, \dots)\end{aligned}$$

For $\mathcal{X} \in \mathcal{H}_*(S)$, we define

$$\Sigma^{\infty}(\mathcal{X}) = (\mathcal{X}, \Sigma\mathcal{X}, \Sigma^2\mathcal{X}, \dots)$$

It is functor from $\mathcal{H}_*(S)$ to $\mathbf{Spt}^{\mathbb{T}}(S)$, its right adjoint is

$$\Omega^{\infty}(E) = \operatorname{colim}_n (\Omega^n E_n)^{\text{fib}}$$

where $(-)^{\text{fib}}$ is fibrant replacement in $\mathbf{Spc}_*(S)$.

Definition 2.1.3. The morphism $f: E \rightarrow K$ is called a \mathbb{T} -stable weak equivalence if

$$\Omega^{\infty} \circ \Sigma_{\mathbb{T}}(f): \Omega^{\infty} \Sigma_{\mathbb{T}}(E) \rightarrow \Omega^{\infty} \Sigma_{\mathbb{T}}(K)$$

is isomorphism in $\mathcal{H}_*(S)$ for all $n \geq 0$. It is proper and combination which can be checked level-wisely.

The Bousfield localization of $\mathbf{Spt}^{\mathbb{T}}(S)$ with respect to \mathbb{T} -stable weak equivalences is denoted by $L_{\mathbb{T}}\mathbf{Spt}^{\mathbb{T}}(S)$. Its homotopy category is called *motivic stable homotopy category over S* , denoted by $\mathcal{SH}(S)$.

2.1.3 Bigraded Cohomology

For $X \in \mathbf{Sm}/S$ and $E \in \mathbf{Spt}^{\mathbb{T}}(S)$, we can define bigraded cohomology

$$E^{p,q}(X) := [\Sigma^{\infty} X_+, S^{p,q} \wedge E]_{\mathcal{SH}(S)}$$

2.1.4 Motivic Eilenberg-MacLane Spectrum

Definition 2.1.4 (free sheaf with transfers generated by X). Functor $L(X): \mathbf{Sm}/k \rightarrow \mathbf{Abs}$ is defined as

$$L(X)(Y) := \text{free abelian group generated by all finite correspondences } Y \rightarrow X$$

For commutative ring R , $L(X, R)(Y)$ is free R -module generated by $L(X)(Y)$.

Proposition 2.1.1. $L(X, R)$ is Nisnevich sheaf (viewed as sheaf of sets).

Let

$$K(R(2n), n) := L(\mathbb{A}^n, R)/L(\mathbb{A}^n - 0, R)$$

in $\mathbf{Spc}_*(k)$. Together with bonding maps

$$\mathbb{T} \wedge K(R(2n), n) \rightarrow K(R(2n+2), n+1)$$

, which is composition of

$$\mathbb{T} \wedge K(R(2n), n) \rightarrow K(R(2), 1) \wedge K(R(2n), n) \rightarrow K(R(2n+2), n+1)$$

induced by natural map $\mathbb{T} \simeq \mathbb{A}^1/\mathbb{A}^1 - 0 \rightarrow L(\mathbb{A}^1, R)/L(\mathbb{A}^1 - 0, R)$, we get \mathbb{T} -spectra. It is denoted by $M\mathbb{H}_R$ called *motivic Eilenberg-MacLane spectra* over R . It represents R -coefficient motivic cohomology

$$H_{\mathcal{M}}^{p,q}(X, R) = H^p(X, R(q)) \cong [\Sigma^\infty X_+, S^{p,q} \wedge M\mathbb{H}_R]_{\mathcal{SH}(k)}$$

for smooth schemes X over field k .