

DISSERTATION

BARIUM TAGGING IN SOLID XENON FOR THE NEXO NEUTRINOLESS DOUBLE BETA  
DECAY EXPERIMENT

Submitted by

Timothy Walton

Department of Physics

In partial fulfillment of the requirements

For the Degree of Doctor of Philosophy

Colorado State University

Fort Collins, Colorado

Fall 2015

Doctoral Committee:

Advisor: William M. Fairbank, Jr.

Robert Wilson  
someone  
Alan Van Orden

Copyright by Timothy Walton 2015

All Rights Reserved

## ABSTRACT

abztrakt

## ACKNOWLEDGEMENTS

Thanks of course goes to my advisor Bill Fairbank

This dissertation is typeset in L<sup>A</sup>T<sub>E</sub>X using a document class designed by Leif Anderson.

# TABLE OF CONTENTS

|  |     |
|--|-----|
| Abstract .....   | ii  |
| Acknowledgements .....   | iii |
| Chapter 1. Introduction .....  | 1   |
| 1.1. something like "Can We Do This?", but probably not that at all..... | 2   |
| 1.2. This is a subsection .....  | 2   |
| Chapter 2. Theory .....  | 3   |
| 2.1. Neutrinos .....   | 3   |
| 2.2. Barium Spectroscopy .....   | 8   |
| 2.3. Matrix Isolation Spectroscopy .....                                 | 8   |
| Chapter 3. Enriched Xenon Observatory .....                              | 9   |
| 3.1. EXO-200 .....   | 10  |
| 3.2. nEXO .....  | 10  |
| Chapter 4. Apparatus .....   | 11  |
| 4.1. Ion Beam .....  | 11  |
| 4.2. Ba Getter Source .....  | 13  |
| 4.3. Solid Xenon Matrix Deposition .....                                 | 13  |
| Chapter 5. Results .....   | 15  |
| Chapter 6. Conclusions .....   | 16  |
| Appendix A. Supplementary Material .....                                 | 17  |
| A.1. Some Sample Material .....  | 17  |

|                                      |    |
|--------------------------------------|----|
| Appendix B. Another Supplement ..... | 18 |
|--------------------------------------|----|

## CHAPTER 1

### INTRODUCTION

start with neutrino prediction by pauli in beta decay, then talk about neutrinos and maybe even mention your dumb "extremes in experimental technique thing", but that's dumb as the main point.

And while the first indication of neutrinos came from the "missing" energy they took away from single beta decays, we now look for the mode of double beta decay which looks like single beta decay was expected to look – where there are no neutrinos. No, this is pretty dumb.

or: we search for electrons coming out in double beta decay

maybe here you should talk about neutrinos' aid in glimpsing certain things, like inner star workings and super novae, and at the beginnings of the universe

#### **old stupid intro:**

The study of the neutrino has required extremes in experimental technique from the beginning. Neutrinos were described by W. Pauli, who first proposed their existence to remedy an apparent violation of energy conservation in beta decay, as being [impossible to detect] [ref.]. Rather, it requires a great deal of sensitivi, ingenuity, and hardship (just "ingenuity and hardship"? sensitivity may be redundant) to observe them, and it was [] years before they were first observed by [Reines and Cowan] in [], by [] [ref.]. (is "rather, ..." too demoting-sounding? it is absolutely not meant to be, of course)

Neutrino experiments of greater discovery power have been developed around the world, and command large collaborations of scientists. ummmmmmmmm this is supposed to kind of allude to barium tagging as an extreme technique

Neutrinoless Double Beta Decay experiments like EXO are a different kind of neutrino experiment, not detecting neutrinos directly, but searching for an effect (neutrinoless double beta decay itself) which would demonstrate the Majorana nature of neutrinos. A liquid xenon experiment like EXO provides a the challenging opportunity for another extreme experimental technique, barium tagging, where a single barium ion would be observed at a specific double beta decay site in the volume. This thesis is part of an exploration of one promising barium tagging technique. (these things may be saved for the EXO chapter... idk).

From the first formulation of beta decay theory by E. Fermi [ref.], neutrinos have provided an avenue into a world of new physics, and they continue to be such an avenue. Questions which may be answered by this up and coming generation of neutrino experiments are expected to help explain how the universe came to be this way.

[lead into barium tagging discussion]

#### 1.1. SOMETHING LIKE "CAN WE DO THIS?", BUT PROBABLY NOT THAT AT ALL.

A liquid xenon double beta decay detector allows unique access to the daughters of decays in the liquid volume. The feasibility of grabbing and detecting a single ion from the volume is what must be determined next.

#### 1.2. THIS IS A SUBSECTION

see original (Leif) for example of a figure and a table.



## CHAPTER 2

### THEORY

This chapter outlines the current theory for neutrinos and of double beta decay. Theory relevant to the spectroscopy of Ba in SXe is then discussed.

*See more Leif examples of some things.*

#### 2.1. NEUTRINOS

Neutrinos are chargeless leptons which only interact via the weak force (and gravity). There are three known “flavors” of neutrinos, each corresponding to one of the three known leptons:  $\nu_e$ ,  $\nu_\mu$ , and  $\nu_\tau$ . These are the eigenstates in the basis of the weak force, and so they are the states in which a neutrino will interact via the weak force.

*History:* Neutrinos were first proposed while studying beta decay. If the emitted electron and daughter nucleus in beta decay were the only products, the electron’s energy should be essentially the same for every observed beta decay for a given isotope, since there is a certain energy difference between the initial and final states of the nucleus, and since the final kinetic energy of the nucleus is negligible due to its mass. But instead of a sharp peak at this Q-value, a very broad electron energy spectrum is observed, all beneath and decreasing toward the Q-value.

In 1930, Wolfgang Pauli proposed that this variable “loss” in energy could be due to an additional particle being emitted along with the electron, but which is not observed.

[Fermi, massless]

[observation, and discovery of  $\nu_\mu$ ]

2.1.1. NEUTRINO OSCILLATION AND MASS. Neutrinos exhibit mixing between their energy eigenstates and their weak force eigenstates, and these are not the same basis.

**not really:** This means that a flavor eigenstate is not a stationary state [the fact is that quark mixing happens, and one theory was that neutrinos would do something similar, and then oscillation is discovered...etc.]— a neutrino which begins as a pure flavor state (as all neutrinos will, coming out of a quantum process involving one of the three leptons) will oscillate into the other two flavors as it evolves in time, i.e. the probability of measuring it to be one of the other two flavors is no longer zero.

*History:* The first indications of neutrino oscillation came around 1970 with the Ray Davis Experiment [ref.], which measured the flux (at Earth) of solar electron neutrinos. The flux measured was quite a bit lower than predicted by solar models, and this became known as the Solar Neutrino Problem. The discovery of neutrino oscillation in the late 1990s [ref.] solved this problem, as only a fraction of the sun's neutrinos, produced as pure electron neutrinos, would interact as such.

The very small mass of a neutrino, specifically relative to its momentum, lets one write its Hamiltonian in terms of mass squared differences  $\Delta m_{ij}^2 = m_i^2 - m_j^2$ , where  $i, j = 1, 2, 3$ , referring to what we then call mass states. The mass basis is really the energy basis with the small mass approximation, along with dropping some constant terms in the Hamiltonian (which do not affect time evolution). Writing the time evolution in terms of mass squared differences means that neutrino oscillation experiments can produce measurements of these differences. In fact, the discovery of neutrino oscillation was the first (and only, so far) demonstration that neutrinos have a non-zero mass. Without neutrino mass (particularly without differences between the masses of the mass states), neutrinos would not oscillate.

Neutrino oscillation experiments also provide measurements on the amount of mixing between the flavor basis and the mass basis. We define the mixing between them by a rotation in terms of three mixing angles,  $\theta_{12}$ ,  $\theta_{23}$ , and  $\theta_{13}$ . Transformation between the flavor and mass bases is done with the following unitary matrix, called the Pontecorvo–Maki-Nakagawa-Sakata (PMNS) matrix:

$$\begin{aligned}
 (1) \quad U &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha_1/2} & 0 \\ 0 & 0 & e^{i\alpha_2/2} \end{pmatrix} \\
 &= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha_1/2} & 0 \\ 0 & 0 & e^{i\alpha_2/2} \end{pmatrix}
 \end{aligned}$$

where  $c_{ij} = \cos \theta_{ij}$  and  $s_{ij} = \sin \theta_{ij}$ .  $\delta$  is a phase factor related to lepton CP violation, and  $\alpha_i$  are Majorana phases.

Studying oscillations of neutrinos from different kinds of sources, with different energies and path lengths, can isolate sensitivities to the different parameters (not really sure if this is the right thing to say). For example, the study of solar neutrinos (neutrinos emanating from nuclear fusion reactions in the core of the sun) provides sensitivity to  $\theta_{12}$  and  $\Delta m_{12}^2$  (*right?  $\theta_{12}$  may not be specifically solar...*). Beamline neutrino detectors can be designed for maximum sensitivity to parameters. The parameters so far measured are as follows in Table 2.1:

TABLE 2.1. up to date values with references, and denote “solar”, “atmos.”, etc.

| Parameter            | Measurement |
|----------------------|-------------|
| $\Delta m_{12}^2$    |             |
| $ \Delta m_{31}^2 $  |             |
| $\sin^2 \theta_{12}$ |             |
| $\sin^2 \theta_{23}$ |             |
| $\sin^2 \theta_{13}$ |             |

Note that only the absolute value of  $\Delta m_{31}^2$  is known. As a consequence, there are still two possibilities for the hierarchy of the three neutrino masses. These are called the Normal and Inverted Hierarchies, as shown in Fig. 2.1.




FIGURE 2.1. The two possible hierarchies of neutrino masses. The colors depict the mixing between the mass and flavor bases. [ref]

Neutrino oscillation demonstrates that neutrinos have non-zero mass, and though these experiments can measure the mass squared differences, we still do not have a measurement of the absolute masses of the three neutrinos. *History:* mention Pauli’s first statement of limit near electron mass? Then maybe say the 0 assumption, which I think came from Fermi’s beta decay theory, but if you mention that above, just refer to it here.

The current upper limits on the mass come from... Neutrinoless double beta decay experiments like EXO-200 can put upper limits on specifically the Majorana neutrino mass

(i.e., upper limits on the neutrino mass if neutrinos are indeed Majorana particles). As discussed in the next chapter, [EXO-200 and KamLAND (sp?) ZEN (sp?) together provide the strongest Majorana neutrino mass upper limit of [] (IS THAT TRUE?)].

Neutrino oscillation and non-zero neutrino mass are physics beyond the Standard Model (SM) of particle physics, and though much has been discovered through oscillation experiments, there is much yet to learn about neutrinos. Since they are chargeless, they may be Majorana particles, and their small mass could be explained by the See-saw Mechanism [ref]. Majorana particles are their own anti-particle, and this, along with the discovery that neutrinos have mass, allows for a unique test of the Majorana (vs. Dirac) nature of neutrinos: neutrinoless double beta decay.

2.1.2. NEUTRINOLESS DOUBLE BETA DECAY. Double beta decay is the simultaneous emission of two electrons from a nucleus. Two-neutrino double beta decay, shown in Fig. (figure), is allowed by the Standard Model and has been observed in several isotopes which are listed in Table (table). Similar to beta decay, a neutrino accompanies each electron in this decay, broadening the spectrum of the summed electron energy. This is a second-order process, making it a rare decay, and requiring low backgrounds to measure.

Neutrinoless double beta decay is another postulated mode of double beta decay, and is shown in Fig. (figure). In this case, the neutrino is exchanged as a virtual particle, and this requires that it is a Majorana particle. If discovered, neutrinos would not only be determined to be Majorana particles, but their mass could also be measured in the form of an effective electron neutrino mass, as the rate of neutrinoless double beta decay will depend on the absolute neutrino mass as shown in Eq. (equation).

## 2.2. BARIUM SPECTROSCOPY

Do we want this here? It flows more to have this theory after the proposition of the tagging technique, but maybe that's more appropriate for a talk.

## 2.3. MATRIX ISOLATION SPECTROSCOPY

(same thing)

## CHAPTER 3

# ENRICHED XENON OBSERVATORY

The Enriched Xenon Observatory (EXO) is a set of two experiments, each a LXe time projection chamber (TPC) designed to study the double beta decay of the isotope  $^{136}\text{Xe}$ , and ultimately to search for the zero-neutrino mode. There are several advantages to a LXe detector. Xe scintillates at [around?] [xxx] nm, which is [efficiently collected by [type that the APDs are]] [reference]; so the Xe acts as a detection medium in addition to being the source of the double beta decay [reference? I didn't make up that kind of sentence]. Xe can also be continuously purified to maintain large electron lifetimes in the LXe. Also, the ratio between observed scintillation light and remaining ionized electrons (drifted from the decay site by the TPC's electric field) exhibits a well-known microscopic anti-correlation [ref.], the understanding of which improves the energy resolution of the detector. Finally, a LXe TPC approach offers the opportunity, [fairly] unique in double beta decay, to "tag" the daughter, in this case Ba, at the site of the double beta decay event (specifically, of course, the neutrinoless ones).

EXO-200 is the first of the two experiments, and has been operational since April of 20[xx](?). It is a liquid xenon TPC designed to probe Majorana neutrino masses down to around 100 meV. [EXO instrum. paper part I] The following sections describe the EXO-200 experiment, as well as nEXO, the next-generation tonne-scale liquid xenon TPC which is now in the design stages. EXO-200 does not have Ba tagging implemented, but it is hoped that nEXO will.

3.1. EXO-200

3.2. nEXO



## CHAPTER 4

# APPARATUS

This chapter describes the apparatus at Colorado State University, which we have used for all described studies of Ba fluorescence in SXe after deposition in vacuum. Our main Ba source, the Ba<sup>+</sup> ion source/beam, is first described, as well as the measurements (using Faraday cups) used for determining the number of ions we deposit. A purely Ba neutral source is described. The co-deposit of Ba/Ba<sup>+</sup> with Xe gas onto a cold sapphire window, subsequent laser excitation, and finally the collection optics for the fluorescence, are described.

### 4.1. ION BEAM

4.1.1. BARIUM ION SOURCE/ACCELERATION. Barium ions are produced in a Colutron [type?] ion gun system [reference], as depicted in Fig. x. A solid barium charge is placed into the hollowed end of a stainless steel rod, which is then inserted into the discharge chamber, near the hot filament. The heated barium vaporizes, allowed to escape the hollowed rod around a loosely threaded set screw at the end of the rod. The discharge chamber then fills with barium vapor. A voltage is applied to the anode plate, which then creates a discharge, through the barium vapor, between the anode and the filament. The resulting plasma, containing barium ions, then escapes through the small hole in the anode plate, where it enters the acceleration potential.

The acceleration potential is 2 kV, between the ion source anode and an aperture, which constitutes the first element of the "acceleration lens" (Fig. 4.1). The acceleration lens is an Einzel lens, the voltages for which are chosen to approximately collimate the ion beam for passage through the  $E \times B$  velocity filter.



FIGURE 4.1. figyer

4.1.2. VELOCITY FILTER, LENSING. The  $E \times B$  velocity filter selects  $Ba^+$  by providing perpendicular electric and magnetic fields, which produce opposing forces on charged particles moving straight through the filter. Those fields are chosen such that those forces are equal for  $Ba^+$ , according to Eqn. 2:

$$(2) \quad \sigma = 1.$$

Other ions will be deflected, while  $Ba^+$  will continue along the beam path.

The full ion beam is shown in Fig. xxx. The Decelerator lens can be used to reduce the beam energy, but is not needed for 2 keV beams, which are used in this work. Einzel Lens 3 focuses the beam onto the main Faraday cup, which is used during experiments to measure

ion current. The final set of deflection plates, H2 and V2, are also used during experiments to steer the beam for deposits.

4.1.3. ION BEAM PULSING. To deposit small numbers of ions in a controlled manner, a set of pulsing plates can be used (Fig. xxxx). When running in this mode, the pulsing plates are first placed at 200 V and -200 V to deflect the beam, and are pulsed to 0 V for 1  $\mu$ s for each pulse.

The pulses can be detected by the Induction Plates. Since they use induction, they can be used to observe the pulses during an ion deposition (unlike using the Faraday cup to measure ion current during a DC deposit). An example of an oscilloscope readout of the pulsing plate signal, and subsequent induction plate signal, is shown in Fig. 5x.

## 4.2. BA GETTER SOURCE

Ba "getters" are typically used in vacuum systems to improve vacuum by emitting Ba atoms, which grab gas molecules and hold them to the chamber walls. We employ getters as a neutral Ba source in our system.

...

It is very helpful to have a completely different type of Ba source, to rule out any source-related quirks, e.g. source-produced impurities.

## 4.3. SOLID XENON MATRIX DEPOSITION

The final destination of the barium ions is in the solid xenon matrix, which is deposited onto a cold sapphire window. Sapphire has good thermal conductivity, good optical transparency in the visible, and does not fluoresce in the wavelength region where barium fluoresces.

Xenon freezes around 73 K (?) at our pressures ( $0.5 - 1 \times 10^{-7}$  Torr), so the window is cooled to temperatures below that. The window is held to a cold finger (Fig. 6x, picture of), cooled by a -brand- cryostat.

4.3.1. DEPOSITION PROCEDURE. Before barium ions are let through, xenon gas is allowed to flow, controlled by a leak valve, onto the cold sapphire window, where it freezes and begins growing the solid matrix. The Faraday cup is then retracted, to clear the path for barium ions. The cup serves as a shutter for DC deposition, or if pulsing is being used, they are performed at this time. Barium ions land in the solid xenon as the matrix continues to grow. The cup is then replaced, and the xenon leak stopped.

## CHAPTER 5

### RESULTS

## CHAPTER 6

# CONCLUSIONS

Ba<sup>+</sup>

## APPENDIX A

### SUPPLEMENTARY MATERIAL

make these also into separate files plz

#### A.1. SOME SAMPLE MATERIAL

Appendix is a strange name. Did the name for the written material come before the name of the organ? Here [?] is a citation in an appendix.

APPENDIX B

ANOTHER SUPPLEMENT