

Report: Mechanism of ATF3 Recruitment by USF1 in GM12878

1. Initial Observation

Context: An initial machine learning model for predicting ATF3 binding affinity identified the **USF1 motif** as the most significant feature, surpassing ATF3's own motif in importance.

Objective: To determine the biophysical mechanism underlying the statistical dependence of ATF3 binding on USF1.

2. Iterative Hypothesis Verification

Hypothesis 1.0: The "Pioneer-Assist" Model

Proposed Mechanism: USF1 acts as a pioneer factor to open chromatin, allowing ATF3 to bind nearby DNA.

- **Verification Method:**
 - **Spatial Analysis:** Measured distance between ATF3 and USF1 peak summits.
 - **Accessibility Analysis:** Compared DNase-seq signal at "ATF3-only" vs. "ATF3+USF1" sites.
- **Results:**
 - **Co-occurrence:** Validated. 61.2% of ATF3 peaks have a nearby USF1 peak. The median distance between them is **17.0 bp**, with an **82.5-fold enrichment** over random expectation ($p < 0.01$).
 - **Accessibility: Rejected.** There was negligible difference in chromatin accessibility between co-bound sites (DNase signal 1.14) and solo sites (1.10). This suggests USF1 is not "opening" closed chromatin for ATF3.

Hypothesis 2.0: The "Weak-Affinity Rescue" Model

Proposed Mechanism: USF1 stabilizes ATF3 binding at sites where the ATF3 motif is degenerate (weak affinity).

- **Verification Method:**
 - **Motif Scoring:** Compared PWM log-odds scores of ATF3 motifs in "Solo" vs. "Co-bound" peaks.
- **Results:**

- **Affinity Difference:** Validated. ATF3 motifs at Solo sites were significantly stronger (Mean Score: **12.0**) than at Co-bound sites (Mean Score: **10.3**; $p=4.35 \times 10^{-4}$).
- **The Anomaly:** While the hypothesis held for sites *with* motifs, a massive discrepancy was found: **90.3% of Co-bound peaks lacked a detectable ATF3 motif entirely**, despite having strong USF1 motifs.

Hypothesis 3.0: The "Indirect Tethering" Model

Proposed Mechanism: ATF3 does not bind DNA directly at these sites but is tethered to the chromatin via protein-protein interaction with USF1.

- **Verification Method:**
 - **Summit Displacement:** Measured the distance from the ATF3 ChIP-seq peak summit to the center of the USF1 motif in the "motif-less" peaks.
- **Results:**
 - **Geometry:** Validated. The mean displacement was **-1.3 bp** (Median: -0.5 bp). The ChIP signal for ATF3 is centered precisely on the USF1 binding site, confirming indirect recruitment.

Hypothesis 4.0: The "Enhancer-Specific Tethering" Model

Proposed Mechanism: This tethering mechanism is specific to distal enhancers, while promoters are refractory to such large complexes.

- **Verification Method:**
 - **Genomic Annotation:** Classified "Recruiters" (USF1+ATF3) and "Loners" (USF1-only) as Promoter ($\leq 2\text{kb TSS}$) or Distal.
- **Results:**
 - **Distribution: Inverted.** The hypothesis was incorrect. "Recruiters" are heavily enriched at **Promoters (61.1%)**, while "Loners" are primarily **Distal (60.5%)** ($p=1.26 \times 10^{-70}$).
 - **Activity:** Validated. When Recruiters *are* distal, they mark highly active regions (DNase signal 1.05) compared to Loners (0.63).

Hypothesis 5.0: The "Density-Dependent Licensing" Model

Proposed Mechanism: ATF3 recruitment occurs only at sites of high USF1 density (occupancy), which distinguishes active promoters from transient background binding.

- **Verification Method:**
 - **Signal Intensity:** Compared USF1 ChIP-seq signal strength between Recruiters and Loners.
 - **Gene Expression:** Compared TPM of genes driven by Recruiter vs. Loner promoters.
- **Results:**

- **Protein Density:** Validated. Recruiter sites have **4.11-fold higher** USF1 signal intensity than Loner sites ($p \approx 0.00$).
- **Function:** Validated. Genes associated with Recruiter promoters have **1.41-fold higher** median expression ($p = 2.6 \times 10^{-6}$), with 82.5% falling in the top expression quartile.

Hypothesis 6.0: The "E-box Cluster" Architecture

Proposed Mechanism: The high USF1 density required for recruitment is driven by clusters of E-box motifs (avidity) rather than single sites.

- **Verification Method:**
 - **Motif Counting:** Counted the number of USF1 motifs per peak in Recruiters vs. Loners.
 - **Results:**
 - **Multiplicity:** Validated. Recruiters contain significantly more motifs per peak (Mean: **1.53**) than Loners (Mean: **1.12**).
 - **Clustering:** **51.5%** of Recruiter peaks contain ≥ 2 motifs, compared to only 35.1% of Loners ($p = 3.55 \times 10^{-33}$).
 - **Recruitment Rate:** Clusters are **1.37x** more likely to recruit ATF3 than singleton motifs.
-

3. Final Scientific Conclusion

Mechanism: Promoter Licensing via High-Density Indirect Tethering. ATF3 binds to the GM12878 genome in two distinct modes:

1. **Direct Binding:** Independent binding to high-affinity consensus sites (minority).
2. **Indirect Tethering:** Recruitment to **USF1 complexes** at active promoters (majority).

Key Driver: Recruitment is strictly **density-dependent**. ATF3 is "licensed" to dock only when USF1 occupancy exceeds a critical threshold. This threshold is achieved genetically through **E-box clustering** (avidity) at promoter regions. Single/transient USF1 binding events at distal sites are insufficient to stabilize the ATF3 interaction.

Biological Function: This mechanism couples the stress-responsive factor ATF3 to the core transcriptional machinery of **highly expressed housekeeping genes**, allowing ATF3 to act as a co-factor without requiring its own evolved DNA binding sites at these essential loci.