

Non-Simply Connected Subsystems and the Reduced Density Matrices

Chen-Chih Wang

Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan
(Dated: September 24, 2026)

Abstract

We discuss the structure of reduced density matrices of non-simply connected subsystems. Subtle though the mathematical form may first appear, it turns out that the residual electric fields along the inner boundaries provide an additional source that organizes the reduced density as we did in the previous work. As a result, the organizing structure of the entanglement spectrum still applies, and the Yao-Qi formula for calculating entanglement entropy is still valid.

1 Patch Configurations

First, we consider a non-simply connected region A with an interior hole B to demonstrate a hidden structure that was not considered previously in simply connected subsystems. First, we have to take the non-contractible closed loop $\mathcal{L}$ inside the subsystem A encircling the interior hole B into account. The second subtle thing is the edge configuration $\mathfrak{H}$, which represents the *patch configuration* of the hole enclosed by A as illustrated in Fig. 1(a). Note that $\Sigma_{\mathfrak{D}\ominus\mathfrak{H}}$ is shorthand notation for the string operator of the combination of $\mathfrak{D}$ combined with the patch configuration, but with the local identity operators generated by the relation $\sigma^x\sigma^y\sigma^z \propto \mathbb{1}$ inside the hole B excluded. With these two additional parts, the reduced density matrix of this subsystem is as follows,

$$\rho_A = \frac{1}{2^{N^A - N_p^A}} \hat{\Pi}_A (\mathbb{1} + W_{\mathcal{L}}) \sum_{[\mathfrak{D}] \subset A} (\mathfrak{C}_{\mathfrak{D}} \Sigma_{\mathfrak{D}} + \mathfrak{C}_{\mathfrak{D}\ominus\mathfrak{H}} \Sigma_{\mathfrak{D}\ominus\mathfrak{H}}).$$




Figure 1: The patch configuration $\mathfrak{H}$ covering the interior hole of a non-simply connected subsystem. This configuration is devised in such a way that the corresponding string operator $\Sigma_{\mathfrak{H}}$ has all its local components inside the hole B equal to the identity. Its corresponding string operator $\Sigma_{\mathfrak{H}}$ commutes with other string operators lying within the shaded subsystem.

The non-contractible loop Wilson operator $W_{\mathcal{L}}$ with the identity forms a projector due to the 1-form Wilson symmetry. However, the expectation values of string operators $\Sigma_{\mathfrak{D} \oplus \mathfrak{H}}$ are not constant, and each operator $\Sigma_{\mathfrak{D} \oplus \mathfrak{H}}$ cannot be separated into $\Sigma_{\mathfrak{D}}$ and $\Sigma_{\mathfrak{H}}$ without phase factors either. As a result, the string operators containing the patch operator cannot be written as projectors, unlike the non-contractible loop $W_{\mathcal{L}}$. Therefore, we take the other way around to bypass this difficulty. We first decompose the string operator $\Sigma_{\mathfrak{D} \oplus \mathfrak{H}}$ into the form $\varphi(\mathfrak{H}, \mathfrak{D}) \Sigma_{\mathfrak{H}} \Sigma_{\mathfrak{D}}$, and then write the reduced density matrix as follows,

$$\rho_A = \frac{1}{2^{N^A - N_p^A - 1}} \hat{\Pi}_A \underbrace{\left(\frac{1 + W_{\mathcal{L}}}{2} \right)}_{\equiv \hat{\Pi}_{\mathcal{L}}} \sum_{[\mathfrak{D}] \subset A} \left(\mathbb{1} + \underbrace{\varphi(\mathfrak{H}, \mathfrak{D}) \frac{\mathfrak{C}_{\mathfrak{D} \oplus \mathfrak{H}}}{\mathfrak{C}_{\mathfrak{D}}}}_{\equiv 2p_{\mathfrak{D}}^{(+)} - 1} \Sigma_{\mathfrak{H}} \right) \mathfrak{C}_{\mathfrak{D}} \Sigma_{\mathfrak{D}}.$$

In the above equation, we defined a new quantity $p_{\mathfrak{D}}^{(+)}$ derived from the phase factor and the string coefficients. Applying the following relation

$$\frac{1 + aA}{2} = \left(\frac{1 + a}{2} \right) \left(\frac{1 + A}{2} \right) + \left(\frac{1 - a}{2} \right) \left(\frac{1 - A}{2} \right),$$

the matter part of the density matrix can be decomposed. As a result, we obtain

$$\rho_A = \frac{1}{2^{N^A - N_p^A - 2}} \hat{\Pi}_A \hat{\Pi}_{\mathcal{L}} \sum_{[\mathfrak{D}] \subset A} \left[p_{\mathfrak{D}}^{(+)} \left(\frac{1 + \Sigma_{\mathfrak{H}}}{2} \right) + p_{\mathfrak{D}}^{(-)} \left(\frac{1 - \Sigma_{\mathfrak{H}}}{2} \right) \right] \mathfrak{C}_{\mathfrak{D}} \Sigma_{\mathfrak{D}}$$

where $p_{\mathfrak{D}}^{(-)} \equiv 1 - p_{\mathfrak{D}}^{(+)}$ with $p_{\mathfrak{D}}^{(+)}$ defined at the beginning forms a binary probability distribution $\{p_{\mathfrak{D}}^{(+)}, p_{\mathfrak{D}}^{(-)}\}$ normalized as $p_{\mathfrak{D}}^{(+)} + p_{\mathfrak{D}}^{(-)} = 1$. The projectors $(\mathbb{1} \pm \Sigma_{\mathfrak{H}})/2$, which will be denoted by the shorthand notation $\hat{\Pi}_{\mathfrak{H}}^{(\pm)}$ later on, divide the local Hilbert space into two orthogonal sectors with the eigenvalues of $\Sigma_{\mathfrak{H}}$ being $+1$ and -1 , respectively. As a result, the matter part of the reduced density matrix becomes a direct sum of the two sectors.

$$\rho_A \equiv \frac{1}{2^{N^A - N_p^A - 2}} \hat{\Pi}_A \hat{\Pi}_{\mathcal{L}} \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \left[\left(p_{\mathfrak{D}}^{(+)} \hat{\Pi}_{\mathfrak{H}}^{(+)} \Sigma_{\mathfrak{D}} \right) \oplus \left(p_{\mathfrak{D}}^{(-)} \hat{\Pi}_{\mathfrak{H}}^{(-)} \Sigma_{\mathfrak{D}} \right) \right].$$

The reduced density matrix expressed in the above form implies that the coefficients $p_{\mathfrak{D}}^{(s)}$ represent the probability of finding Majorana fermions in each sector labeled by $\Sigma_{\mathfrak{H}}$. Note that this separation of Hilbert space is not only for the matter sector, but for the entire reduced density matrix because every string operator $\Sigma_{\mathfrak{D}}$ with its edge configuration $\mathfrak{D} \subset A$ commutes with the patch operator projector,

$$[\Sigma_{\mathfrak{D}}, \hat{\Pi}_{\mathfrak{H}}^{(s)}] = 0 \quad \forall \mathfrak{D} \subset A.$$

For subsystems with more than one hole, the density matrix takes a similar form.

$$\rho_A = \frac{1}{2^{N^A - N_p^A - 2g}} \hat{\Pi}_A \hat{\Pi}_g \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \bigoplus_{\{s\}} p_{\mathfrak{D}}^{(\{s\})} \hat{\Pi}_{\mathfrak{H}}^{(\{s\})} \Sigma_{\mathfrak{D}}. \quad (1)$$

In the above formula, g represents the number of holes inside the subsystem (genus number), and $\{s\} \equiv \{s_1, s_2, \dots, s_g\}$ is a set of eigenvalues of patch string operators $\Sigma_{\mathfrak{H}^{(n)}}$ covering different holes

$B^{(n)}$ that label different sectors. The projectors and the probability distribution are generalized as follows.

$$\begin{cases} \hat{\Pi}_g \equiv \prod_{n=1}^g \left(\frac{1 + W_{\mathcal{L}^{(n)}}}{2} \right) \\ \hat{\Pi}_{\mathfrak{H}}^{\{\{s\}\}} \equiv \prod_{n=1}^g \left(\frac{1 + s_n \Sigma_{\mathfrak{H}^{(n)}}}{2} \right) \\ p_{\mathfrak{D}}^{\{\{s\}\}} \equiv \prod_{n=1}^g p_{\mathfrak{D},n}^{(s_n)} = \frac{1}{2^g} \prod_{n=1}^g \left(1 + s_n \frac{\varphi(\mathfrak{H}^{(n)}, \mathfrak{D}) \mathfrak{C}_{\mathfrak{H}^{(n)} \ominus \mathfrak{D}}}{\mathfrak{C}_{\mathfrak{D}}} \right) \end{cases} \quad (2)$$

where $\mathcal{L}^{(n)}$ represents the non-contractible Wilson loop enclosing the n -th hole, and $\mathfrak{H}^{(n)}$ represents the patch configuration covering the n -th hole. The set $\{p_{\mathfrak{D}}^{\{\{s\}\}}\}$ of each configuration class $[\mathfrak{D}]$ forms a probability distribution, which normalizes as follows,

$$\sum_{\{s\}} p_{\mathfrak{D}}^{\{\{s\}\}} = \sum_{\{s\}} \left(\prod_{n=1}^g p_{\mathfrak{D},n}^{(s_n)} \right) = \prod_{n=1}^g \left(p_{\mathfrak{D},n}^{(+)} + p_{\mathfrak{D},n}^{(-)} \right) = 1. \quad (3)$$

The Hilbert space dimension of each sector can be computed straightforwardly

$$\text{Tr}_A \left(\hat{\Pi}_A \hat{\Pi}_g \hat{\Pi}_{\mathfrak{H}}^{\{\{s\}\}} \right) = 2^{N^A - N_p^A - 2g}.$$

We can further simplify the structure of the reduced density matrix to make it resemble the equivalent Majorana fermionic reduced density matrix. If this is doable, it may imply that the conventional matter-gauge separation picture in simply connected subsystems still applies for multiply-connected subsystems. First, we introduce the **projected string operators** as follows,

$$\hat{\mathcal{S}}_{\mathfrak{D}} \equiv \hat{\Pi}_A \hat{\Pi}_g \bigoplus_{\{s\}} \tilde{p}_{\mathfrak{D}}^{\{\{s\}\}} \hat{\Pi}_{\mathfrak{H}}^{\{\{s\}\}} \Sigma_{\mathfrak{D}}. \quad (4)$$

Since $[\Sigma_{\mathfrak{H}}, \Sigma_{\mathfrak{H}'}] = 0$ and $[\Sigma_{\mathfrak{H}}, \Sigma_{\mathfrak{D}}] = 0 \ \forall \mathfrak{D} \subset A$, the projected string operators also satisfy the following algebraic relation

$$\hat{\mathcal{S}}_{\mathfrak{D}} \hat{\mathcal{S}}_{\mathfrak{D}'} = \varphi(\mathfrak{D}, \mathfrak{D}') \hat{\mathcal{S}}_{\mathfrak{D} \ominus \mathfrak{D}'} \quad (5)$$

The fact that this relation still applies implies that the endpoints of the projected string operators defined in non-simply connected subsystems still obey fermionic statistics. Therefore, **the string operators are still equivalent to Majorana fermionic operators in non-simply connected subsystems.**

Together, the reduced density matrix can be organized into the following form that resembles the Majorana fermionic reduced density matrix of the equivalent Majorana fermionic system.

$$\rho_A = \frac{1}{2^{|\partial A|/2 - N_{\partial}^A}} \left[\frac{1}{2^{N^A/2}} \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \hat{\mathcal{S}}_{\mathfrak{D}} \right] \quad (6)$$

In the above equation, we have used the generalized geometric relation (13), and $N_{\partial}^A \equiv (1 + g)$ is the number of disconnected boundaries of the subsystem A .

On the other hand, the reduced density matrix of the equivalent Majorana fermionic system reads

$$\rho_A^F = \frac{1}{2^{N^A/2}} \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \hat{\mathcal{M}}_{\mathfrak{D}}$$

Accordingly, the Hilbert space dimension of the equivalent Majorana fermionic system is

$$\frac{\dim \mathcal{H}_S}{\dim \mathcal{H}_F} = \frac{\text{Tr}_A \hat{\mathcal{S}}_\emptyset}{\text{Tr}_{A,F} \hat{\mathcal{M}}_\emptyset} = 2^{|\partial A|/2 - N_\partial^A}$$

Therefore, the entanglement entropy computed using the spin reduced density matrix is

$$\begin{aligned} S_A &= -\text{Tr}_A \rho_A \ln \rho_A \\ &= -\frac{1}{2^{|\partial A|/2 - N_\partial^A}} \text{Tr}_A \left[\left(\frac{1}{2^{N^A/2}} \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \hat{\mathcal{S}}_{\mathfrak{D}} \right) \ln \left(\frac{1}{2^{N^A/2}} \sum_{[\mathfrak{D}] \subset A} \mathfrak{C}_{\mathfrak{D}} \hat{\mathcal{S}}_{\mathfrak{D}} \right) \right] + \left(\frac{1}{2} |\partial A| - N_\partial^A \right) \ln 2 \\ &= \underbrace{-\text{Tr}_{A,F} \rho_A^F \ln \rho_A^F}_{S_A^F} + \underbrace{\left(\frac{1}{2} |\partial A| - N_\partial^A \right) \ln 2}_{S_A^G} \end{aligned} \quad (7)$$

The above result reproduces the correct **topological entanglement entropy** $-N_\partial^A \ln 2$.

2 Patch String Operators Interpreted as Parity of Residual Electric Fields

The physical meaning of the patch string operator $\Sigma_{\mathfrak{H}}$ can be interpreted in the other way. The string operator $\Sigma_{\mathfrak{H}}$ of the patch configuration can be decomposed into the multiplication of all the ξ operators along the interior boundary and the Pauli spin operators that represent the residual electric field inside the hole along the boundary attached to the string operators covering the hole. This decomposition is pictorially illustrated as follows.

$$\begin{aligned} \Sigma \left(\text{Diagram 1} \right) &= \left(\text{Diagram 2} \right) \\ &= \left(\text{Diagram 3} \right) \underbrace{\left(\text{Diagram 4} \right)}_{\propto \mathbb{1}} \\ &= \left(\prod_{\hat{\xi} \in \partial B} \hat{\xi} \right) \underbrace{\left(\prod_{\langle ij \rangle \in B} \sigma_i^{\alpha_{ij}} \sigma_j^{\alpha_{ij}} \right)}_{\propto \mathbb{1}} \left(\prod_{i \in \partial B} \sigma_i^{\mu_i} \right) \propto \prod_{\hat{\xi} \in \partial B} \hat{\xi} \end{aligned} \quad (8)$$

From this result, we see that the string operator $\Sigma_{\mathfrak{H}}$ of the patch configuration is the product of all the ξ -operators along the interior boundary. Accordingly, the projector $\hat{\Pi}_{\mathfrak{H}}$ can be redefined as follows,

$$\hat{\Pi}_{\mathfrak{H}}^{\{\{s\}\}} = \prod_{n=1}^g \frac{1}{2} \left(\mathbb{1} + s_n \prod_{\hat{\xi} \in \partial B^{(n)}} \hat{\xi} \right). \quad (9)$$

This result clearly indicates that **the patch sector actually corresponds to the parity of electric fields surrounding each hole** $B^{(n)}$.

Note that **we don't have to consider this pattern in a simply connected subsystem** since we have already included the configurations satisfying $[\mathfrak{D} \ominus \mathfrak{D}_Z] \subset A$ when constructing the reduced density matrix. The string operator defined as $\prod_{\xi \in \partial A} \hat{\xi}$ in a simply connected region A is equivalent to the combination of the product of all the dimers lying outside A , the dimers crossing the boundary ∂A , and the gauge links forming ξ -operators along the boundary. This configuration is taken into account by $[\mathfrak{D} \ominus \mathfrak{D}_Z] \subset A$. For a similar reason, **we should not include the outer boundary either**, because otherwise the reduced density matrix ρ_A will contain the component $\prod_{\xi \in \partial A} \hat{\xi}$ consisting of all the ξ operators along all the boundaries of the region A as mentioned earlier. Such an operator is equivalent to multiplying all the Majorana fermions lying outside the subsystem A , and which has already been taken into account in ρ_A since such a string operator satisfies the condition $[\mathfrak{D} \ominus \mathfrak{D}_Z] \subset A$.

Following the same pattern, we can define the proper $\Sigma_{\mathfrak{H}}$ for subsystems whose **inner** boundary has **odd** length and **outer** boundary has **even** length.

$$\begin{aligned}
\Sigma_{\mathfrak{H}} &\equiv \left(\text{Diagram 1} \right) \underbrace{\hspace{1cm}}_{=\hat{\Xi}} \left(\text{Diagram 2} \right) \underbrace{\hspace{1cm}}_{=\prod_{\xi \in \partial B} \hat{\xi}} \left(\text{Diagram 3} \right) \underbrace{\hspace{1cm}}_{=(\prod_{\langle ij \rangle \in B} \sigma_i^{\alpha_{ij}} \sigma_j^{\alpha_{ij}})(\prod_{i \in \partial B} \sigma_i^{\mu_i}) \propto 1} \\
&= \hat{\Xi} \left(\prod_{\xi \in \partial B} \hat{\xi} \right) \underbrace{\left(\prod_{\langle ij \rangle \in B} \sigma_i^{\alpha_{ij}} \sigma_j^{\alpha_{ij}} \right) \left(\prod_{i \in \partial B} \sigma_i^{\mu_i} \right)}_{\propto 1} \propto \hat{\Xi} \prod_{\xi \in \partial B} \hat{\xi}
\end{aligned} \tag{10}$$




Figure 2: **An unsolved problem** is the pattern of patch sectors of subsystems with both their inner and outer boundaries having odd length. In this case, there is no well-defined Ξ operator, but the ξ operators surrounding the inner boundary cannot form a valid patch configuration either. A possible resolution is to connect the remaining sites at the inner and outer boundaries. One example is the Wilson operator enclosed by the dashed frame in the above illustration. However, this method also requires the ξ operators along the outer boundary, which should not be included in this framework.




Figure 3: The non-contractible Wilson operator W_2 with two endpoints attached to different boundaries anti-commutes with the patch operator $\Sigma_{\mathfrak{H}}$ covering the interior hole, which is proportional to the product of all the ξ operators surrounding the inner boundary, while the non-contractible Wilson operator W_1 with both endpoints attached to the patch commutes with $\Sigma_{\mathfrak{H}}$ as well as other string operators supported within the subsystem A , hence commutes with the reduced density matrix ρ_A .

3 Extensive Degeneracy in the Entanglement Spectrum

The non-contractible Wilson operators connecting two electric-field operators fall into two types. One type, denoted as $\mathfrak{W}_s$, has both endpoints attached to the same boundary, while the other, denoted as $\mathfrak{W}_d$, has endpoints attached to different boundaries. As illustrated in Fig. 1(b), the non-contractible Wilson operator W_1 is of type $\mathfrak{W}_s$, while the other Wilson operator W_2 is of type $\mathfrak{W}_d$.

$$W_2 \hat{\Pi}_{\mathfrak{H}}^{(\pm)} W_2^\dagger = \hat{\Pi}_{\mathfrak{H}}^{(\mp)}$$

As a consequence, the non-contractible Wilson operators in $\mathfrak{W}_d$ do not commute with the reduced density matrix ρ_A , making them not proper unitary operators for proving that the subblocks are identical. In contrast, the operators in $\mathfrak{W}_s$ let the patch projectors remain intact. Thus, they commute with the density matrix and become the only proper operators for our purpose. Our arguments lead to the conclusion: **the non-contracted Wilson operators that relate different subblocks are those with both endpoints attached to the same boundary**. As a result, the degree of degeneracy of the entanglement spectrum is equal to the number L of different $\{\xi; \Xi\}$ configurations divided by two to the power of the number of disconnected boundaries due to the conservation of residual electric field parity along each inner boundary, which is equal to

$$\text{Degree of degeneracy} = \frac{2^L}{2^{N_\partial^A}} = 2^{L-N_\partial^A}$$

If all the boundaries have even length, the number of residual electric field configurations L reduces to $L = |\partial A|/2$, which reproduces the conventional result

$$\text{Degree of degeneracy} = 2^{|\partial A|/2 - N_\partial^A} = \exp(S_G^A)$$




Figure 4: In the Kitaev model, assembling ξ -operators along a closed loop represents a 't Hooft loop of the $\mathbb{Z}_2$ gauge theory, which can be understood as the path of a $\mathbb{Z}_2$ gauge flux (vison) moving along the closed loop and eventually being annihilated by its partner created at the same spot at the very beginning. It is also equivalent to the subsystem fermion parity of the region enclosed by the 't Hooft loop.

4 Electric Field Parity, 't Hooft Loops, and Subsystem Fermion Parity

Aside from the Wilson loops, the 't Hooft loops should also be discussed since the low-energy effective theory of the Kitaev model is a TQFT. The 't Hooft loop in this note is operationally defined as the operator that creates a pair of $\mathbb{Z}_2$ gauge fluxes (visons) and then moves one of the visons around a loop until it gets back to the initial spot to annihilate its partner vison. An example of a 't Hooft loop is illustrated in Fig. 4 as represented by a dashed closed loop crossing a series of bonds at which the $\mathbb{Z}_2$ gauge fields are flipped by the electric fields. From the gauge theory perspective, the 't Hooft operator $\tilde{T}_{\mathcal{C}}$ along a closed loop $\mathcal{C}$ in the Kitaev model can be defined as follows

$$\tilde{T}_{\mathcal{C}} = \prod_{i \in \mathcal{C}} b_i^{\mu_i} = \exp \left(i\pi \oint_{\mathcal{C}} E_{\perp} d\ell \right),$$

where the electric fields are provided by the b -Majorana fermionic operators along the loop $\mathcal{C}$. The above interpretation, combined with our discussions on the relationship between the patch string operators and electric fields, leads to a picture that the 't Hooft loop, which is equivalent to the electric field parity along the loop, is also related to the subsystem fermion parity of the region enclosed by the loop. Such a relationship is formulated mathematically as follows,

$$\tilde{T}_{\partial\Gamma} = \exp \left(i\pi \oint_{\partial\Gamma} E_{\perp} d\ell \right) \rightarrow \prod_{\hat{\xi} \in \partial\Gamma} \hat{\xi} \rightarrow \prod_{i \in \Gamma} c_i \propto (-1)^{F_{\Gamma}}; \quad F_{\Gamma} = \sum_{i \in \Gamma} n_i, \quad (11)$$

where $E_{\perp}$ represents the electric field components perpendicular to the 't Hooft loop direction. According to the divergence theorem, the relationship between the electric fields along the loop

$\partial\Gamma$ and the subsystem fermion parity can be translated to the Gauss law of the gauge theory,

$$\oint_{\partial\Gamma} E_{\perp} d\ell = \sum_{i \in \Gamma} n_i \Rightarrow \nabla \cdot E = \rho.$$

In conclusion, the Gauss law of the gauge theory in the Kitaev model implies the equivalence between the 't Hooft loop and the subsystem fermion parity. We can accordingly rewrite the patch projectors in terms of 't Hooft loop operators enclosing each hole inside the multiply-connected subsystem.

$$\hat{\Pi}_{\mathfrak{H}}^{(\{s\})} = \prod_{n=1}^g \left(\frac{\mathbb{1} + s_n \tilde{T}_{\partial B^{(n)}}}{2} \right). \quad (12)$$

In Ref.[1], the topological entanglement entropy is interpreted as a consequence of **spontaneously broken 't Hooft 1-form symmetry**. Formally, the state $|\psi\rangle$ breaks the 't Hooft 1-form symmetry if $T|\psi\rangle$ is orthogonal to $|\psi\rangle$. In the Kitaev model, the 't Hooft loop is not a symmetry since it doesn't commute with the Hamiltonian. However, it does serve as a source to produce the topological entanglement entropy as discussed in previous sections.

A Generalized Geometric Relation

This formula can be generalized to multiply-connected regions. Suppose a connected region has a genus number g , and the n -th hole surrounded by the connected region A is denoted by $B^{(n)}$ such that $A \cap B^{(n)} = \emptyset$ and $A \cup (\bigcup_{n=1}^g B^{(n)})$ is simply-connected. Denoting by $\ell_{\text{out}} = |\partial(A \cup (\bigcup_{n=1}^g B^{(n)}))|$ the number of crossing edges of the outer boundary, and by $\ell_{\text{in}}^{(n)} = |\partial B^{(n)}|$ the length of the n -th inner boundary, we have

$$\begin{cases} \ell_{\text{out}} = N^A - 2N_p^A + \sum_{n=1}^g \left(N^{B^{(n)}} - 2N_p^{B^{(n)}} - 2N_p^{\partial B^{(n)}} \right) + 2 \\ \ell_{\text{in}}^{(n)} = N^{B^{(n)}} - 2N_p^{B^{(n)}} + 2 \end{cases}$$

where $N_p^{\partial B^{(n)}}$ is the number of plaquettes at the interface between A and B . It is obvious that $N_p^{\partial B^{(n)}} = \ell_{\text{in}}^{(n)}$. Thus, we have

$$\ell_{\text{out}} = N^A - 2N_p^A - \sum_{n=1}^g \ell_{\text{in}}^{(n)} - 2g + 2$$

At the same time, we know that the length of the boundary of the region A is the sum of the outer boundary and all inner boundaries

$$|\partial A| = \ell_{\text{out}} + \sum_{n=1}^g \ell_{\text{in}}^{(n)}$$

Combining this relation with the result we obtained gives a generalized geometric relation:

$$|\partial A| = N^A - 2N_p^A + 2(1 - g) \quad (13)$$

The above relation implies that the number of lattice sites N^A determines whether the total boundary length $|\partial A|$ is even or odd.

$$N^A = |\partial A| \pmod{2}$$

References

- [1] W.-T. Xu, T. Rakovszky, M. Knap, and F. Pollmann, *Entanglement Properties of Gauge Theories from Higher-Form Symmetries*, [Phys. Rev. X **15**, 011001 \(2025\)](#).