

Computer Simulations and Risk Assessment – Lecture 6

Fall 2019

Brandeis International Business School

Course Information - Schedule

Class Date	Text Chapters
Aug. 30, 2019 – L1	<ul style="list-style-type: none">• Course Introduction/Python Installation• Introduction to Quantitative Finance Career• Python basics
Sep. 6, 2019 – L2	<ul style="list-style-type: none">• Advanced Python Topics
Sep. 13, 2019 – L3	<ul style="list-style-type: none">• Advanced Python Topics
Sep. 20, 2019 – L4	<ul style="list-style-type: none">• Sourcing and handling Data• Stylized financial data analysis using Python
Sep. 27, 2019 – L5	<ul style="list-style-type: none">• Value at Risk
Oct. 4, 2019 – L6	<ul style="list-style-type: none">• Conditional Value at Risk (Expected Shortfall) + Mid-term Review
Oct. 11, 2019	<ul style="list-style-type: none">• Mid-term
Oct. 18, 2019 – L7	<ul style="list-style-type: none">• Modeling Volatility I
Oct. 25, 2019 – L8	<ul style="list-style-type: none">• Modeling Volatility II
Nov. 1, 2019 – L9	<ul style="list-style-type: none">• Practical application case Studies I
Nov. 8, 2019 – L10	<ul style="list-style-type: none">• Practical application case Studies II
Nov. 15, 2019 – L11	<ul style="list-style-type: none">• Back Testing + Conditional risk prediction
Nov. 22, 2019 – L12	<ul style="list-style-type: none">• Research project presentation
Dec. 6, 2019 – L13	<ul style="list-style-type: none">• Final Review


Expected Shortfall and the Impact of Holding Period

- **Expected Shortfall and Calculation**
- **VaR may violate risk subadditivity**
- **Holding periods and scaling**

Issues for VaR: It Doesn't Tell the Whole Story

- VaR is just a quantile marking a threshold point on the distribution
- VaR does not tell us anything about left tail beyond it
- Might ignore a lot of risk if the tail is fat or irregular
- What is the right probability level to use? 5%, 1%. It is not always the smaller the probability the better


VaR Ignores the Left Tail



- Doesn't tell us the magnitudes of loss and associated probabilities left of the cutoff point associated with VaR

Solution: Expected Shortfall

- CVaR (Expected shortfall): Mean loss given that VaR loss is exceeded – the mean of returns that are less than VaR, or the weighted average of areas under the cyan line



Expected Shortfall

- Definition of expected shortfall

$$p = \Pr(X \leq VaR(p)) = \int_{-\infty}^{-VaR(p)} f_q(x) dx$$

$$\text{Expected Shortfall} = -E(Q|Q \leq -VaR(p)) = -\frac{1}{p} \int_{-\infty}^{-VaR(p)} x f_q(x) dx$$

- Assuming x follows normal distribution, with mean of μ and standard deviation of σ , we have

$$f_q(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right]$$

Expected Shortfall under normal distribution

- Convert X from a normal distribution to a standard normal distribution, $z = \frac{x - \mu}{\sigma}$ (or $x = \mu + z \sigma$), which leads to $\phi(z) = \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{z^2}{2}\right]$, for z , the expected shortfall is

$$\begin{aligned} ES0 &= -\frac{1}{p} \int_{-\infty}^{-VaR0(p)} z \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{z^2}{2}\right] dz \\ &= \frac{1}{p} \left[-\frac{1}{\sqrt{2\pi}} \exp\left[-\frac{z^2}{2}\right] \right]_{-\infty}^{-VaR0(p)} \\ &= -\frac{\frac{1}{\sqrt{2\pi}} \exp\left[-\frac{VaR0(p)^2}{2}\right]}{p} - 0 = -\frac{\phi(-VaR0(p))}{p} \end{aligned}$$

- Converting back to x from z (ES is a x in this case, and $ES0$ is a z):

$$x = \mu + z \sigma \rightarrow ES = \mu + ES0 * \sigma = \mu - \frac{\phi(-VaR0(p))}{p} * \sigma$$

Where ϕ is the pdf, and $VaR0(p)$ is the VaR at $p\%$ for a standard normal distribution

Expected Shortfall

- For a more formal prove of the formula for expected shortfall, please see the following online prove:

<http://blog.smaga.ch/expected-shortfall-closed-form-for-normal-distribution/>

Expected Shortfall for Normal Distribution

- $R_{t+1} \sim N(\mu, \sigma^2)$, e.g., $R_{t+1} \sim N(\mu = 0.12, \sigma^2 = 0.2^2)$
- Assume $\Phi(x)$ and $\phi(x)$ are Normal(0,1) CDF and PDF
- VaR(5%) for a regular normal distribution:

$R^*(p) = \sigma * \Phi^{-1}(p) + \mu = \text{stats.norm.ppf}(p, \mu, \sigma)$, which means

$$\text{VaR}(p) = \Phi^{-1}(p) = \frac{R^*(p) - \mu}{\sigma}$$

- This means Expected shortfall

$$\tilde{R}(p) = E(R | R \leq R^*(p)) = \mu - \sigma \frac{\phi\left(\frac{R^*(p) - \mu}{\sigma}\right)}{p}$$


$$\text{ES}(p) = -((1 + \tilde{R}(p)) P_t - P_t) = -P_t \tilde{R}(p)$$

- Python Code in sample code L6_ES.py:
 - $R_star = \text{stats.norm.ppf}(p, \mu_R, \sigma_R)$
 - $R_Tilde = -\sigma_R * \text{stats.norm.pdf}((R_star - \mu_R) / \sigma_R) / p + \mu_R$


Expected Loss When Loss is worse than VaR

$$p = \Pr(X \leq \text{VaR}(p)) = \int_{-\infty}^{-\text{VaR}(p)} f_q(x) dx$$


$$\text{Expected Shortfall} = -E(Q|Q \leq -\text{VaR}(p)) = -\frac{1}{p} \int_{-\infty}^{-\text{VaR}(p)} x f_q(x) dx$$



(a) Density, $f(P/L)$, and VaR




(b) Tail of density, $f(P/L)$, and VaR



(c) Blow up the tail. The darker shading has area p , whilst the entire shaded area has area 1

Expected Loss When Loss is worse than VaR

- Expected shortfall: Expected loss given that VaR loss is exceeded – there is more information on the left tail
- Also known as Expected tail loss (ETL) or conditional Value-at-Risk (CVaR)



ES analytics for Normal Distribution

- A little more details on ES analytics

$$p = \Pr(X \leq VaR(p)) = \int_{-\infty}^{-VaR(p)} f_q(x) dx$$

$$\text{Rescale } p \text{ to } 100\%: 1 = \int_{-\infty}^{-VaR(p)} f_{VaR}(x) dx = \frac{1}{p} \int_{-\infty}^{-VaR(p)} f_q(x) dx$$

$$\text{Expected Shortfall} = -E(Q|Q \leq -VaR(p)) = -\frac{1}{p} \int_{-\infty}^{-VaR(p)} x f_q(x) dx$$

- **For a portfolio with a current value of 1 and standard deviation of 1:**

$$ES = -\frac{\phi(\Phi^{-1}(p))}{p}$$

VaR and ES Table for Normal/t Distribution

- For standard normal (mean = 0, std = 1) and student-t distribution (mean = 0, std = 1, dof = 4)

VaR table							
p	0.001	0.01	0.025	0.05	0.1	0.1584	0.5
normal	-3.090	-2.326	-1.960	-1.645	-1.282	-1.000	0.000
t (4 dof)	-7.173	-3.747	-2.776	-2.132	-1.533	-1.143	0.000
ES table							
p	0.001	0.01	0.025	0.05	0.1	0.1584	0.5
normal	-3.367	-2.665	-2.338	-2.063	-1.755	-1.526	-0.798
t (4 dof)	-8.952	-5.305	-4.018	-3.215	-2.507	-2.062	-1.010

Historical Expected Shortfall

- Two routes: Portfolio value based or return based
- Portfolio value (price) based approach
 - Produce historical distribution of the portfolio value (price)
 - Estimate p quantile of portfolio value distribution, or the VaR
 - Estimate the conditional mean for portfolio values below VaR
 - $ES(p) = E(P_{t+1} | P_{t+1} \leq P^*) - P_{t+1}$
- Returns based approach
 - Estimate p quantile for returns
 - Estimate the conditional mean for returns that is below return that is corresponding to VaR
 - $ES(p) = -P_t E(R_{t+1} | R_{t+1} \leq R^*)$

Expected Shortfall: Pros and Cons

- Pros
 - More/Extra information on extreme losses
 - Less easily manipulated
 - subadditive
- Cons
 - Needs tail data to do the calculation
 - Harder to explain
 - Less adoption by financial institutions
 - Harder to backtest

Impact of Holding Period

- We discussed the importance of knowing the pdf/cdf for loss/return for calculating risk
- One important determinant of the risk number we haven't discussed is the length of the holding period for the risk calculation
 - Normally, the longer the holding period for the investment, the higher the risk level, as there is more time for bad things to happen
 - But exactly how do we link risk with length of the investment holding period?

The Data Challenge for Long Holding Period

- Normally the VaR is calculated for daily holding period
 - For 1% VaR, need a few hundred daily return data to have at least a few data points on the left tail
- For 10-day (bi-weekly) VaR, the data requirement gets serious
 - For 1% VaR, need a few hundred 10-day periods, say 300, to have a decent tail coverage. This is 3000 daily data or about 12 years!
- Solution: Scaling
 - Based on assumptions that statistical theory governs how properties of distribution changes as data are added

The Square-Root-of-Time Rule of Scaling

- The statistical measurement a random variable, such as volatility and VaR, are obtained by multiplying a higher frequency measurement by the square root of the number of observations in the holding period
 - This rule applies to Volatility as long as the returns are IID (independent and identically distributed), regardless whether they are normally distributed or not
 - However, for VaR, this rule applies only if the returns are IID and normal
- E.g., $VaR(10 \text{ day}) = VaR(\text{daily}) * \sqrt{10}$

Expected Value for Random Variables

- Discrete variables:

$$E(X) = \sum_{i=1}^T p_i x_i$$

$A = \{i | x_i < z\}$ – think of z as VaR

$$E(X | X < z) = \frac{\sum_{i \in A} p_i x_i}{\sum_{i \in A} p_i}$$

- Continuous variables

$$E(X) = \int_{-\infty}^{\infty} x f_x(x) dx$$

$$p = \Pr(X \leq z) = \int_{-\infty}^z f_x(x) dx$$

$$E(X | X \leq z) = \frac{1}{p} \int_{-\infty}^z x f_x(x) dx$$

Non-standardized student-t

- Standard Student-t random variable $t(v)$: mean = 0, $Std = \sqrt{\frac{v}{v-2}}$
- Set non-standard $t^* = t * \frac{\sigma}{\sqrt{v/(v-2)}} + \mu$
- $Var(t^*) = v/(v-2) * (\frac{\sigma}{\sqrt{v/(v-2)}})^2 = \sigma^2$
- $E(t^*) = \mu$

Covariance & Correlation


- **Risk management Q&A**
- **Revisiting Covariance and Correlation**
- **Portfolio level risk measures**
- **Intuition and Applications**

Risk Management – the question

- Remember I challenged you with the interview question of “tell me how would you forecast the risk of a given portfolio?”
- Last lecture, I talked about the core of forecasting risk is to forecast the distribution of the profit & loss or the returns of the portfolio?
- The most common risk (measure) people talk about: **volatility** – how do we think about volatility in the risk management context? How does it relate to potential P&L or negative returns?
- We studied **VaR and CVaR**. How about them? How do we explain VaR and CVaR?
- Let’s look at the example of current portfolio=100, future mean=100, sigma=1,

Risk Management – The Answer

- Volatility: it measures the magnitude of loss in the sense that there is a **X%** probability that loss will be at or larger than $\mu - \sigma$ (or \$1)
 - X=15.9 % for normal distribution
 - X=19.6 % for student-t distribution with degree of freedom=3
- VaR (5%) Normal distribution: The probability of loss greater than $\mu - 1.65 * \sigma = -\$1.65$ is 5%
- ES(5%) Normal distribution: The mean of all loss that's higher than VaR(5) = - \$1.65 is -\$2.06



Standard Deviation, Correlation & Covariance

- Portfolio return mean and standard deviation

$$\mu_k = \frac{1}{N} \sum_{t=1}^N R_{t,k}$$

$$\sigma_k = \sqrt{\frac{1}{N-1} \sum_{t=1}^N (R_{t,k} - \mu_k)^2} \quad (\text{Also called volatility})$$

- Correlation and Covariance


$$\rho = \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^N (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^N (y_i - \bar{y})^2}}$$

$$\text{cov} = \sigma_{ij}^2 = \frac{1}{N-1} \sum_{i=1}^N (x_{ij} - \bar{x}_j)(x_{ik} - \bar{x}_k)$$

Correlation

- Correlation measures how returns move together

$$\rho = \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^N (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^N (y_i - \bar{y})^2}}$$



Covariance Matrix Σ_t

- The covariance between two assets i and j is :

$$\text{Cov}(R_k, R_j) = \sigma_{kj}^2$$

- In the case of a three assets, the conditional matrix takes the following form

$$\Sigma_t = \begin{pmatrix} \sigma_{11}^2 & \sigma_{12}^2 & \sigma_{13}^2 \\ \sigma_{12}^2 & \sigma_{22}^2 & \sigma_{23}^2 \\ \sigma_{13}^2 & \sigma_{23}^2 & \sigma_{33}^2 \end{pmatrix} \text{ where } \sigma_{ij}^2 = \rho_{ij} \sigma_i \sigma_j$$

- The diagonal terms measure the volatilities of each of the assets $\sigma_{ii}^2 = \rho_{ii} \sigma_i \sigma_i = \sigma_i^2$ (because $\rho_{ii}=1$)
- The off-diagonal terms is a product of the correlation of the assets and the volatilities of each of the asset $\sigma_{ij}^2 = \rho_{ij} \sigma_i \sigma_j$

Covariance Matrix Σ_t

- Observe the covariance: the diagonal terms are always positive and in general larger in magnitude relative to the off-diagonal terms

Index	Mkt-RF	SMB	HML	RF	MoM
Mkt-RF	0.0028482	0.000540799	0.000442601	-8.92293e-06	-0.00084904
SMB	0.000540799	0.00102323	0.000138314	-4.12695e-06	-0.000219559
HML	0.000442601	0.000138314	0.00121865	2.09176e-06	-0.000679714
RF	-8.92293e-06	-4.12695e-06	2.09176e-06	6.45226e-06	6.61262e-06
MoM	-0.00084904	-0.000219559	-0.000679714	6.61262e-06	0.00221179

Formulas for Portfolio Level Risk

- Asset level

$$E(R_t)=\mu_i, \text{ var}(R_i)=\sigma_i^2, \text{ corr}(R_1, R_2)=\rho$$

- Portfolio return, $R_p=w_1R_1 + w_2R_2$
- Portfolio variance: $\sigma_p^2=w_1^2\sigma_1^2+2w_1w_2\rho\sigma_1\sigma_2+w_2^2\sigma_2^2$
- In the general case, with time varying, for a multi-assets portfolio


$$\text{Return: } R_{p,t} = \sum_{i=1}^N w_{i,t} R_{i,t}$$

$$\text{Portfolio Variance: } \Sigma_{p,t} = \sigma_{p,t}^2 = \mathbf{W}'\Sigma\mathbf{W}$$


$$= \sum_{i=1}^N w_{i,t}^2 \sigma_{ii,t}^2 + \sum_{i=1}^N \sum_{j=1}^N w_{i,t} w_{j,t} \rho_{i,j,t} \sigma_{ij,t} \sigma_{ji,t}$$

Foundation of a Modern Risk System


1. For each asset (risk factor), develop assumptions about distribution of its returns



.....



2. Incorporate assumptions about correlations of these returns



3. Combining weights (exposure to risk factors)

$\mathbf{W}'\Sigma\mathbf{W}$ or through Monte Carlo Simulations

here \mathbf{W} is the weights vector and Σ is the covariance matrix

Portfolio Level Risk Example

$$W = [0.6, 0.4],$$
$$\Sigma = \begin{pmatrix} 0.001 & 0.0002 \\ 0.0002 & 0.002 \end{pmatrix}$$

$$\text{Portfolio Variance} = \sigma_{p,t}^2 = W' \Sigma W =$$
$$0.6^2 * 0.001 + 0.4^2 * 0.002 + 2 * 0.6 * 0.4 * 0.0002 = 0.000776$$

Tracking Error of a Portfolio

- **Realized** tracking error of a fund's returns vs. its benchmark:

$$\text{Tracking Error} = \text{std}(r_{\text{fund}} - r_{\text{benchmark}}) = \sqrt{\frac{1}{N-1} \sum_{t=1}^N (R_{t,\text{fund}} - r_{t,\text{benchmark}})^2}$$

- What about **forecasting TE for the future**? Tracking error of a portfolio, calculated using components of the portfolio

$$TE = \sqrt{W_a' \Sigma W_a}$$

where W_a is the active weights vector and Σ is the covariance matrix

- Example

$$W_a = [-0.1, 0.1],$$
$$\Sigma = \begin{pmatrix} 0.001 & 0.0002 \\ 0.0002 & 0.002 \end{pmatrix}$$

$$\text{Projected TE} = \sqrt{W_a' \Sigma W_a} = 0.0051$$

Units of the variables

- Return has the unit of %
- Variance has the unit of %²
- Correlation has no unit
- Volatility has the unit of %

For the rest of the this class

- We develop ways to forecast the covariance matrix, e.g., exponential moving average model
- Linkage to the foundation of the risk system: These models can also be extended to calculate VaR and CVaR based on normal or student-t distribution assumptions (e.g. utilizing the VaR and ES Table for Normal/t Distribution on page 12)
- We develop the conditional simulation approach to calculate conditional covariance and tail risk
- We apply these forecasts to real-time risk modeling applications
 - Forecasting of portfolio risk
 - Tracking error optimization for the construction of exchange traded funds (ETFs)
 - Risk parity investing strategies