

MATH 151A Homework 1

Zheng Wang (404855295)

January 22, 2019

Question 1

- (a) We will first show there is at least one solution for $f(x) = 0$.

Notice that $f(0.5) = -0.1 < 0$ and $f(1) = 0.3 > 0$. Also, since $f(x)$ is a polynomial function, so it is continuous for $x \in \mathbb{R}$, so $f(x)$ is continuous $\forall x \in [0.5, 1]$. By Intermediate Value Theorem, since $f(0.5) < 0 < f(1)$, it follows that there exist some $\xi \in (0.5, 1)$ such that $f(\xi) = 0$.

We then show that ξ is unique.

We find that $f'(x) = 2x - 0.7$ is a continuous function for all $x \in \mathbb{R}$. Thus for $x \in [0.5, 1]$, $f'(x) \in [0.3, 1.3]$. Suppose that there exist more than one solution for the equation $f(x) = 0$, then we have $f(\xi) = f(\xi') = 0$, where $\xi \neq \xi'$. Therefore, since $f(x)$ is continuous, differentiable on (ξ, ξ') (without loss of generality, assume that $\xi < \xi'$), and $f(\xi) = f(\xi') = 0$, by Rolle's Theorem, there exist some $x^* \in (\xi, \xi')$ such that $f'(x^*) = 0$, a contradiction. ■

- (b) Claim: $|p_n - p| \leq \frac{1-0.5}{2^n}$.

proof. Since $b_n - a_n = \frac{1}{2}(b_{n-1} - a_{n-1}) = \frac{1}{2^{n-1}}(b_1 - a_1)$. Also, since $p_n = \frac{1}{2}(a_n + b_n)$.

Thus,

$$|p_n - p| \leq \frac{1}{2} (b_n - a_n) = \frac{1}{2^n} (b_1 - a_1) = \frac{1 - 0.5}{2^n}$$

■

Thus, to have $p_k - p \leq 10^{-5}$, we must have $10^{-5} \geq \frac{1-0.5}{2^k}$. By solving the equation, we have $k \geq \frac{5}{\log 2} - 1 \approx 15.6$. Therefore, we must take $\boxed{k = 16}$ steps then the error will be less than 10^{-5} .

Question 2

Take $g(x) = f(x) - x$. Then if $g(x) = 0$, $f(x) = x$. There are two cases for $f(x)$, we will discuss them one by one.

Case 1

If $f(a) = a$ or $f(b) = b$. Then, we have at least one fixed point at $x = a$ or $x = b$, depending on which of the condition before is true.

Case 2

Otherwise, since $f(x) \in [a, b]$ for any $x \in [a, b]$, $f(a) > a$ and $f(b) < b$. Thus $g(a) = f(a) - a > 0$ and $g(b) = f(b) - b < 0$. Also, we define $f(x)$ such that it is continuous on $[a, b]$. Thus, by Intermediate Value Theorem, there exist some $\xi \in (a, b)$ such that $g(\xi) = 0$. In another word, there exist some $\xi \in (a, b)$ such that $f(\xi) = \xi$.

Thus, there is at least one fix point for $f(x)$. ■

Question 3

$$\begin{aligned} \text{(a)} \quad p_1 &= \frac{p_0^2 + 3}{2p_0} = \frac{3^2 + 3}{2 \times 3} = 2 \\ p_2 &= \frac{p_1^2 + 3}{2p_1} = \frac{2^2 + 3}{2 \times 2} = 1.75 \end{aligned}$$

(b) Suppose that the limit is L . Then we have the following:

$$L = \lim_{n \rightarrow \infty} p_{n+1} = \lim_{n \rightarrow \infty} \frac{p_n^2 + 3}{2p_n} = \frac{(\lim_{n \rightarrow \infty} p_n)^2 + 3}{2 \cdot (\lim_{n \rightarrow \infty} p_n)} = \frac{L^2 + 3}{2L}$$

By solving the equation above, we have

$$2L^2 = L^2 + 3 \quad \Rightarrow \quad L = \pm\sqrt{3}$$

Thus, all possible limits of the sequence $\{p_n\}_{n=0}^{\infty}$ are $\boxed{-\sqrt{3}}$ (when $p_0 < 0$) or $\boxed{\sqrt{3}}$ (when $p_0 > 0$).

(c) By the definition of the Newton's method, the sequence of solving the equation $x^2 - 3 = 0$ is

$$p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)} = p_n - \frac{p_n^2 - 3}{2p_n} = \frac{2p_n^2 - p_n^2 + 3}{2p_n} = \frac{p_n^2 + 3}{2p_n}, \quad n \geq 1$$

and p_0 is given. This is the same as the sequence mentioned in the question. ■

Question 4

- (a) The general formula for secant method is

$$p_{n+1} = p_n - \frac{f(p_n)(p_n - p_{n-1})}{f(p_n) - f(p_{n-1})}$$

Since we are using $f(x) = x^2 - 3$. Thus, we have

$$p_{n+1} = p_n - \frac{(p_n^2 - 3)(p_n - p_{n-1})}{p_n^2 - p_{n-1}^2}$$

If we are using $p_0 = \frac{1}{2}$ and $p_1 = 3$, we have $p_2 = 3 - \frac{(3^2-3)(3-\frac{1}{2})}{3^2-\frac{1}{2}^2} = \boxed{\frac{9}{7}}$.

Also, we have $p_3 = \frac{9}{7} - \frac{(\frac{9}{7}^2-3)(\frac{9}{7}-3)}{\frac{9}{7}^2-3^2} = \boxed{\frac{8}{5}}$.

- (b) Using the Method of False Position to solve $f(x) = x^2 - 3$. Since $f(p_0) \cdot f(p_1) = -\frac{33}{2} < 0$, we have

$$p_2 = 3 - \frac{(3^2 - 3)(3 - \frac{1}{2})}{3^2 - \frac{1}{2}^2} = \frac{9}{7}$$


Now, since $f(p_1) \cdot f(p_2) = -\frac{396}{49} < 0$, we have

$$p_3 = \frac{9}{7} - \frac{(\frac{9}{7}^2 - 3)(\frac{9}{7} - 3)}{\frac{9}{7}^2 - 3^2} = \frac{8}{5}$$

Notice that the result is the same for both method this far. This is due to the fact that $f(p_0) \cdot f(p_1) < 0$ and $f(p_1) \cdot f(p_2) < 0$. Otherwise, there should be differences.

Question 5 (Coding)


(a) The resulting graph I got is



I also got the following from the console. Thus the solution is 2.620×10^{-5} .

The solution using secant method is 2.619881e-05 at iteration #38

(b) The resulting graph I got is



I also got the following from the console. Thus the solution is 1.357×10^{-5} .

The solution of Newton's method is 1.356836e-05 at iteration #27

Code for Question 5

The following matlab codes are used.

```
% % MATH 151A HOMEWORK1
% % QUESTION 5
% % Wang, Zheng (404855295)

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Secant Method to solve  $f(x) = \sin(x) - x = 0$ 
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% initialize
i = 2;
p0 = pi/4;
p1 = 3*pi/8;
q0 = f(p0);
q1 = f(p1);
arr_p = [];
arr_it = [];

% Set the max iteration number
N = 10000;

% Set the tolerance
T0 = 10^(-5);

%iteration
while i <= N
    p = p1 - q1*(p1-p0)/(q1-q0);
    % stopping condition
    if abs(p - p1) < T0
        fprintf('The solution using secant method is %e at iteration %d\n', p, i)
        break;
    end
    i = i+1;
    p0 = p1;
    q0 = q1;
    p1 = p;
    q1 = f(p);
    arr_p = [arr_p p];
    arr_it = [arr_it i];
end

% output the failure message if needed
```

```

if i > N
fprintf('Secant Method failed after %d iteration, the value found is %e\n',N, p)
end

% plot the graph
figure;
plot(arr_it, arr_p, 'o-r', 'Linewidth', 1.1);
title('\{p_n\} Generated by Secant Method (for n >= 2)');
xlabel('iteration number')
ylabel('p_n')
grid on;

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Newton's Method to solve  $f(x) = \sin(x) - x = 0$ 
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% initialize
i = 1;
p0 = pi/4;
q0 = f(p0);
d0 = fprim(p0);
arr_p = [];
arr_it = [];

% Set the max iteration number
N = 10000;

% Set the tolerance
T0 = 10(-5);

%iteration
while i <= N
    p = p0 - q0/d0;
    if abs(p - p0) < T0
        fprintf('The solution of Newton's method is %e at iteration #%d\n', p, i)
        break;
    end
    i = i+1;
    p0 = p;
    arr_p = [arr_p p0];
    arr_it = [arr_it i];
    q0 = f(p0);
    d0 = fprim(p);
end

```

```

% output the failure message if needed
if i > N
fprintf('Newton''s Method failed after %d iteration, the value found is %e\n',N, p)
end

% plot the graph
figure;
plot(arr_it, arr_p, 'o-', 'Linewidth', 1.1);
title('\{p_n\} Generated by Newton''s Method (for n >= 1)');
xlabel('iteration number')
ylabel('p_n')
grid on;

% % function declaration
function y = f(x)
    y = sin(x) - x;
end

function y = fprim(x)
    y = cos(x) - 1;
end

```