

The botanist keeps tasting tea

*A gentle introduction to e-values  
and sequential statistical inference*



---

Ian Waudby-Smith

Department of Statistics, UC Berkeley

Miller retreat, 2024

# THE LADY TASTING TEA

HOW STATISTICS  
REVOLUTIONIZED SCIENCE  
IN THE  
TWENTIETH CENTURY



DAVID SALSBURG

---

"A fascinating description of the kinds of people who interacted,  
collaborated, disagreed, and were brilliant in the development of statistics."  
—Barbara A. Bailar, National Opinion research Center

## The botanist tasting tea (1920s)



Ronald Fisher



Muriel Bristol



## The botanist tasting tea (1920s)

Would you like some tea?



Ronald Fisher

Muriel Bristol



## The botanist tasting tea (1920s)

Would you like some tea?



Ronald Fisher



Muriel Bristol



## The botanist tasting tea (1920s)

Would you like some tea?



No,  $T$  in  $M \neq M$  in  $T$



Can you *really* tell them apart?



Ronald Fisher

Muriel Bristol



Ronald Fisher

## The botanist tasting tea (1920s)

Would you like some tea?



No,  $T$  in  $M \neq M$  in  $T$



Can you *really* tell them apart?



Indeed, yes!



Muriel Bristol



Ronald Fisher

## The botanist tasting tea (1920s)

Would you like some tea?



No, T in M ≠ M in T



Can you *really* tell them apart?



Indeed, yes!



Muriel Bristol





Ronald Fisher

## The botanist tasting tea (1920s)

Would you like some tea?



No, T in M  $\neq$  M in T



Can you *really* tell them apart?



Indeed, yes!



Muriel Bristol



T

M

T

T

M

M

M

T



Ronald Fisher

## The botanist tasting tea (1920s)

Would you like some tea?



No,  $T$  in  $M \neq M$  in  $T$



Can you *really* tell them apart?



Indeed, yes!



Muriel Bristol



T

M

T

T

M

M

M

T

What's the probability that a *chance* guess would be *perfect*?

$$1/70 \approx 0.014$$



Ronald Fisher

## The botanist tasting tea (1920s)

Would you like some tea?



No,  $T$  in  $M \neq M$  in  $T$



Can you *really* tell them apart?



Indeed, yes!



Muriel Bristol



T

M

T

T

M

M

M

T

What's the probability that a *chance* guess would be *perfect*?

$$1/70 \approx 0.014$$

This is a p-value for  $H_0$  : Muriel cannot distinguish btwn MT and TM.

# The botanist tasting tea (1920s)



## The botanist tasting tea (1920s)



However, the odds were stacked against Muriel from the start!

## The botanist tasting tea (1920s)



However, the odds were stacked against Muriel from the start!

The probability that a chance guess would yield at most one error is  $17/70 \approx 0.24$ , which is not so impressive.

## The botanist tasting tea (1920s)



However, the odds were stacked against Muriel from the start!

The probability that a chance guess would yield at most one error is  $17/70 \approx 0.24$ , which is not so impressive.

If she had made exactly one mistake, could we just let her keep tasting tea?  
No, this is blatant p-hacking.

An **p-value** is a function  $P_n$  of the data (MT or TM) so that

For any sample size  $n$ ,  $\mathbb{P}_{H_0}(P_n \leq \alpha) \leq \alpha$ .

An **p-value** is a function  $P_n$  of the data (MT or TM) so that

For any sample size  $n$ ,  $\mathbb{P}_{H_0}(P_n \leq \alpha) \leq \alpha$ .

But, it is *not* the case that

$$\mathbb{P}_{H_0}(P_{\tau} \leq \alpha) \leq \alpha$$

if  $\tau$  was chosen based on the data so far.

An **p-value** is a function  $P_n$  of the data (MT or TM) so that

For any sample size  $n$ ,  $\mathbb{P}_{H_0}(P_n \leq \alpha) \leq \alpha$ .

But, it is *not* the case that

$$\mathbb{P}_{H_0}(P_{\tau} \leq \alpha) \leq \alpha$$

if  $\tau$  was chosen based on the data so far.

In fact, it is typically the case that

$$\mathbb{P}_{H_0}(P_{\tau} \leq \alpha) \approx 1!$$

I have done some work on “*e*-values”.

I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:

I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:



I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:



I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:




I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:




I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:




I have done some work on “*e*-values”.

An *e*-value is a function  $E$  of the data (MT or TM) so that

$$E \geq 0 \quad \text{and} \quad \mathbb{E}_{H_0}(E) \leq 1.$$

Suppose Muriel keeps tasting tea:



Then,  $\mathbb{P}_{H_0}(P_{\tau}^{\star} \leq \alpha) \leq \alpha$ , regardless of how  $\tau$  is chosen!

where  $P_n^{\star} := (E_1 \cdot E_2 \cdots E_n)^{-1}$ .

Thank you!

[ianws.com](http://ianws.com)



---

Ian Waudby-Smith

Department of Statistics, UC Berkeley

Miller retreat, 2024