

Estimating means of bounded random variables by betting

Ian Waudby-Smith & Aaditya Ramdas
Carnegie Mellon University



Estimating means of bounded random variables by betting

Ian Waudby-Smith & Aaditya Ramdas
Carnegie Mellon University



JRSS-B 2023 (w/ discussion)

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Goal: produce a set $C_n \equiv C(X_1, \dots, X_n)$ s.t.

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Goal: produce a set $C_n \equiv C(X_1, \dots, X_n)$ s.t.

$$\mathbb{P}(\mu \notin C_n) \leq \alpha$$

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Goal: produce a set $C_n \equiv C(X_1, \dots, X_n)$ s.t.

$$\mathbb{P}(\mu \notin C_n) \leq \alpha \quad \equiv \quad \mathbb{P}(\mu \in C_n) \geq 1 - \alpha$$

Hoeffding's inequality (1963) provides one solution:

$$C_n^H := \frac{1}{n} \sum_{i=1}^n X_i \pm \sqrt{\frac{\log(2/\alpha)}{2n}}$$

Hoeffding's inequality (1963) provides one solution:

$$C_n^H := \frac{1}{n} \sum_{i=1}^n X_i \pm \sqrt{\frac{\log(2/\alpha)}{2n}}$$

No asymptotics, no parametric assumptions at all!

Hoeffding's inequality (1963) provides one solution:

$$C_n^H := \frac{1}{n} \sum_{i=1}^n X_i \pm \sqrt{\frac{\log(2/\alpha)}{2n}}$$

No asymptotics, no parametric assumptions at all!

The downside? Not very sharp,
especially for small variance $\sigma^2 := \text{Var}(X_i)$.

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Goal: produce a set $C_n \equiv C(X_1, \dots, X_n)$ s.t.

$$\mathbb{P}(\mu \notin C_n) \leq \alpha \quad \equiv \quad \mathbb{P}(\mu \in C_n) \geq 1 - \alpha,$$

Given bounded rvs: $X_1, X_2, \dots, X_n \in [0,1]$
with mean $\mathbb{E}(X_i) = \mu$,

Goal: produce a set $C_n \equiv C(X_1, \dots, X_n)$ s.t.

$$\mathbb{P}(\mu \notin C_n) \leq \alpha \quad \equiv \quad \mathbb{P}(\mu \in C_n) \geq 1 - \alpha,$$

and make C_n as sharp as possible.

Our bound:

$$C_n := \left\{ \textcolor{red}{m} : \prod_{i=1}^n (1 + \textcolor{blue}{\lambda}_i(X_i - \textcolor{red}{m})) < \frac{1}{\alpha} \right\}. \quad (\text{design } \textcolor{blue}{\lambda}_i \text{ later})$$

Our bound:

$$C_n := \left\{ \textcolor{red}{m} : \prod_{i=1}^n (1 + \textcolor{blue}{\lambda}_i(X_i - \textcolor{red}{m})) < \frac{1}{\alpha} \right\}. \quad (\text{design } \textcolor{blue}{\lambda}_i \text{ later})$$

Hoeffding:


Our bound:

$$C_n := \left\{ \textcolor{red}{m} : \prod_{i=1}^n (1 + \textcolor{blue}{\lambda}_i(X_i - \textcolor{red}{m})) < \frac{1}{\alpha} \right\}. \quad (\text{design } \textcolor{blue}{\lambda}_i \text{ later})$$


Hoeffding:

$$C_n^H := \left\{ \textcolor{red}{m} : \prod_{i=1}^n \exp \left\{ \lambda(X_i - \textcolor{red}{m}) - \lambda^2/8 \right\} < \frac{1}{\alpha} \right\}.$$

$$\sigma^2 = 1/4$$




$$\sigma^2 \approx 0.0046$$





$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



- H-CI [H63]
- - - Bentkus-CI [B04]
- - - EB-CI [MP09]
- - - VA-EB-CI [Rmk 1]
- - - Hedged-CI [Rmk 3]
- - - - - Anderson [A69]



$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



- | | | |
|------------------------|----------------------|-------------------------|
| — H-CI [H63] | - - - EB-CI [MP09] | - - - Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | --- VA-EB-CI [Rmk 1] | ... Anderson [A69] |



$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



- | | | |
|------------------------|------------------------|--------------------------|
| — H-CI [H63] | - - - EB-CI [MP09] | - · - Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | - - - VA-EB-CI [Rmk 1] | - · - · - Anderson [A69] |



$$\sigma^2 = 1/4$$

$$X_i \sim \text{Bernoulli}(1/2)$$




$$\sigma^2 \approx 0.0046$$

$$X_i \sim \text{Beta}(10, 30)$$




- | | | |
|------------------------|------------------------|--------------------------|
| — H-CI [H63] | - - - EB-CI [MP09] | - · - Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | - - - VA-EB-CI [Rmk 1] | - · - · - Anderson [A69] |

$\sigma^2 = 1/4$ $\sigma^2 \approx 0.0046$ $X_i \sim \text{Bernoulli}(1/2)$ 

- H-CI [H63]
- - - Bentkus-CI [B04]
- - - EB-CI [MP09]
- - - VA-EB-CI [Rmk 1]
- - - Hedged-CI [Rmk 3]
- - - Anderson [A69]
- Ours**
- Hoeffding**



$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



- | | | |
|------------------------|------------------------|--------------------------|
| — H-CI [H63] | - - - EB-CI [MP09] | - - - Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | - - - VA-EB-CI [Rmk 1] | - - - - - Anderson [A69] |




$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



— H-CI [H63]	- - - EB-CI [MP09]	- - - Hedged-CI [Rmk 3]
- - - Bentkus-CI [B04]	- - - VA-EB-CI [Rmk 1]	- - - Anderson [A69]




$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



— H-CI [H63]	- - - EB-CI [MP09]	- - - - Hedged-CI [Rmk 3]
- - - Bentkus-CI [B04]	- - - - VA-EB-CI [Rmk 1]	- - - - - Anderson [A69]




$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



— H-CI [H63]	- - - EB-CI [MP09]	- - - Hedged-CI [Rmk 3]
- - - Bentkus-CI [B04]	- - - VA-EB-CI [Rmk 1]	- - - Anderson [A69]




$$\sigma^2 = 1/4$$

$$X_i \sim \text{Bernoulli}(1/2)$$



$$\sigma^2 \approx 0.0046$$

$$X_i \sim \text{Beta}(10, 30)$$




- | | | |
|---|--|---|
| — H-CI [H63] | — EB-CI [MP09] | ··· Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | - - - VA-EB-CI [Rmk 1] | ... · · · Anderson [A69] |

$$\sigma^2 = 1/4$$


$$X_i \sim \text{Bernoulli}(1/2)$$

$$\sigma^2 \approx 0.0046$$






$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$

$X_i \sim \text{Beta}(10, 30)$



— H-CI [H63]	- - - EB-CI [MP09]	- - - Hedged-CI [Rmk 3]
- - - Bentkus-CI [B04]	- - - VA-EB-CI [Rmk 1]	- - - Anderson [A69]

Formal setup:

Formal setup:

1. Observe X_1, X_2, X_3, \dots

Formal setup:

1. Observe X_1, X_2, X_3, \dots
2. $X_t \in [0,1]$ almost surely.

Formal setup:

1. Observe X_1, X_2, X_3, \dots
2. $X_t \in [0,1]$ almost surely.
3. $\mathbb{E}(X_t | X_1^{t-1}) = \mu.$

$$X_1^{t-1} \equiv (X_1, \dots, X_{t-1})$$

Formal setup:

1. Observe X_1, X_2, X_3, \dots
2. $X_t \in [0,1]$ almost surely.
3. $\mathbb{E}(X_t | X_1^{t-1}) = \mu.$

$$X_1^{t-1} \equiv (X_1, \dots, X_{t-1})$$

Familiar special case: $X_1, X_2, \dots \stackrel{iid}{\sim} \mathbb{P}$, with $\mathbb{E}_{\mathbb{P}}(X_1) = \mu.$

Consider a “candidate mean” $m \in [0,1]$



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$



Consider a “candidate mean” $m \in [0,1]$

$K_0 \leftarrow \$1$

For $t = 1, 2, 3, \dots$:



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m), 1/m)$



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m), 1/m)$

Observe X_t



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m), 1/m)$

Observe X_t

$$K_t \leftarrow K_{t-1} + K_{t-1} \cdot \lambda_t \cdot (X_t - m)$$

EndFor



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m), 1/m)$

Observe X_t

$$K_t \leftarrow K_{t-1} + K_{t-1} \cdot \lambda_t \cdot (X_t - m)$$

EndFor



What if

$\mathbb{E}(X_t) \equiv \mu \gg m$?
or $\mu \ll m$?
or $\mu = m$?

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

$$\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) = K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1})$$

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

$$\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) = K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1})$$

◻

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

$$\begin{aligned}\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) &= K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1}) \\ &= \lambda_t (\mathbb{E}(X_t \mid X_1^{t-1}) - \mu)\end{aligned}$$

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

$$\begin{aligned}\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) &= K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1}) \\ &\quad \boxed{\phantom{\mathbb{E}(K_t(\mu) \mid X_1^{t-1})} = \lambda_t (\mathbb{E}(X_t \mid X_1^{t-1}) - \mu)} \\ &= \lambda_t (\mathbb{E}(X_t \mid X_1^{t-1}) - \mu) \\ &= 0\end{aligned}$$

The “capital process” $(K_t(\mu))_{t=0}^\infty$

is a nonnegative martingale starting at one.

$$\begin{aligned}\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) &= K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1}) \\ &\quad \boxed{\phantom{\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) =}} \\ &= \lambda_t (\mathbb{E}(X_t \mid X_1^{t-1}) - \mu) \\ &= 0\end{aligned}$$

$$\therefore \mathbb{E}(K_t(\mu) \mid X_1^{t-1}) = K_{t-1}(\mu)$$

The “capital process” $(K_t(\mu))_{t=0}^\infty$


is a nonnegative martingale starting at one.

$$\begin{aligned}\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) &= K_{t-1}(\mu) + K_{t-1}(\mu) \cdot \mathbb{E}(\lambda_t(X_t - \mu) \mid X_1^{t-1}) \\ &\quad \boxed{\phantom{\mathbb{E}(K_t(\mu) \mid X_1^{t-1}) =}} \\ &= \lambda_t (\mathbb{E}(X_t \mid X_1^{t-1}) - \mu) \\ &= 0\end{aligned}$$

$$\therefore \mathbb{E}(K_t(\mu) \mid X_1^{t-1}) = K_{t-1}(\mu)$$

Similar ideas from
Shafer & Vovk,
Hendriks,
Jun & Orabona, etc.

Wealth processes / nonnegative martingales






Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$




Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$






Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$






Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$






Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$






Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$





Jean Ville (1939): $\mathbb{P} \left(\exists t \geq 1 : K_t(\mu) \geq \frac{1}{\alpha} \right) \leq \alpha$



“Invert” Ville’s inequality:

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}.$$

Robbins et al. (1960s-1970s),
Shafer & Vovk (2001–2019)
Johari et al. (2015)
Jun & Orabona (2019–),
Howard et al. (2019–).
Grunwald (2019–)

“Invert” Ville’s inequality:

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}.$$

Robbins et al. (1960s-1970s),
Shafer & Vovk (2001–2019)
Johari et al. (2015)
Jun & Orabona (2019–),
Howard et al. (2019–).
Grunwald (2019–)

“The set of $\textcolor{red}{m}$ for which the gambler didn’t make much money”.

“Invert” Ville’s inequality:

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}.$$

Robbins et al. (1960s-1970s),
Shafer & Vovk (2001–2019)
Johari et al. (2015)
Jun & Orabona (2019–),
Howard et al. (2019–).
Grunwald (2019–)

“The set of $\textcolor{red}{m}$ for which the gambler didn’t make much money”.

$$\mathbb{P} (\exists t \geq 1 : \textcolor{violet}{\mu} \notin C_t) \leq \alpha.$$

“Invert” Ville’s inequality:

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}.$$

Robbins et al. (1960s-1970s),
Shafer & Vovk (2001–2019)
Johari et al. (2015)
Jun & Orabona (2019–),
Howard et al. (2019–).
Grunwald (2019–)

“The set of $\textcolor{red}{m}$ for which the gambler didn’t make much money”.

$$\mathbb{P} (\exists t \geq 1 : \textcolor{violet}{\mu} \notin C_t) \leq \alpha.$$

$(C_t)_{t=1}^\infty$ forms a **$(1 - \alpha)$ -confidence sequence**.

Detour: confidence sequences



Detour: confidence sequences



Herbert Robbins, 1960s/70s

+ Siegmund, Darling, & Lai

Confidence sequence

Confidence interval

Confidence sequence

$$\mathbb{P} \left(\exists t \geq 1 : \mu \notin C_t \right) \leq \alpha$$

Confidence interval

Confidence sequence

$$\mathbb{P} (\exists t \geq 1 : \mu \notin C_t) \leq \alpha$$

Confidence interval

$$\forall n, \mathbb{P} (\mu \notin C_n) \leq \alpha$$

Confidence sequence

$$\mathbb{P} (\exists t \geq 1 : \mu \notin C_t) \leq \alpha$$

$$\mathbb{P} (\forall t \geq 1, \mu \in C_t) \geq 1 - \alpha$$

Confidence interval

$$\forall n, \mathbb{P} (\mu \notin C_n) \leq \alpha$$

Confidence sequence


$$\mathbb{P} (\exists t \geq 1 : \mu \notin C_t) \leq \alpha$$



$$\mathbb{P} (\forall t \geq 1, \mu \in C_t) \geq 1 - \alpha$$


Confidence interval

$$\forall n, \mathbb{P} (\mu \notin C_n) \leq \alpha$$


$$\forall n, \mathbb{P} (\mu \in C_n) \geq 1 - \alpha$$







Confidence intervals are valid at a *single sample size*.



Confidence intervals are valid at a *single sample size*.
 Confidence sequences are valid at *all sample sizes simultaneously*.

Back to confidence sequences for
means of bounded random variables

Our confidence sequence:

$$C_t := \left\{ m \in [0,1] : \prod_{i=1}^t (1 + \lambda_i(m) \cdot (X_i - m)) < \frac{1}{\alpha} \right\}$$

is valid for any λ_i but what is a smart choice?

Maximize the Growth Rate Adapted to
the Particular Alternative (**GRAPA**).

$$\text{Choose } \lambda_t(\mathbf{\bar{m}}) = \arg \max_{\lambda} \frac{1}{t-1} \sum_{i=1}^{t-1} \log \left(1 + \lambda \cdot (\mathbf{X}_i - \mathbf{\bar{m}}) \right)$$

Maximize the Growth Rate Adapted to
the Particular Alternative (**GRAPA**).

$$\text{Choose } \lambda_t(\mathbf{m}) = \arg \max_{\lambda} \frac{1}{t-1} \sum_{i=1}^{t-1} \log \left(1 + \lambda \cdot (X_i - \mathbf{m}) \right)$$

$$\text{Or } \lambda_t(\mathbf{m}) \approx \frac{\hat{\mu}_{t-1} - \mathbf{m}}{\hat{\sigma}_{t-1}^2 + (\hat{\mu}_{t-1} - \mathbf{m})^2}$$

Maximize the Growth Rate Adapted to
the Particular Alternative (**GRAPA**).

$$\text{Choose } \lambda_t(\mathbf{m}) = \arg \max_{\lambda} \frac{1}{t-1} \sum_{i=1}^{t-1} \log \left(1 + \lambda \cdot (X_i - \mathbf{m}) \right)$$



$$\text{Or } \lambda_t(\mathbf{m}) \approx \frac{\hat{\mu}_{t-1} - \mathbf{m}}{\hat{\sigma}_{t-1}^2 + (\hat{\mu}_{t-1} - \mathbf{m})^2}$$

•
•
•

$$\sigma^2 = 1/4$$

$$X_i \sim \text{Bernoulli}(1/2)$$


$$\sigma^2 \approx 0.0046$$



- | | | | |
|-------------------|-----------------------|----------------------|----------------------|
| — PM-H [Prop 1] | — Hedged [Thm 3] | — BANCO [JO19] | — Bernoulli [HRMS20] |
| --- PM-EB [Thm 2] | hgKelly [Prop 4] | ... A-Bentkus [KZ21] | |



$$\sigma^2 = 1/4$$

$$X_i \sim \text{Bernoulli}(1/2)$$



$$\sigma^2 \approx 0.0046$$


$$X_i \sim \text{Beta}(10, 30)$$



- | | | | |
|---------------------|----------------------|----------------------|----------------------|
| — PM-H [Prop 1] | — Hedged [Thm 3] | — BANCO [JO19] | — Bernoulli [HRMS20] |
| - - - PM-EB [Thm 2] | ... hgKelly [Prop 4] | ... A-Bentkus [KZ21] | |


$$\sigma^2 = 1/4$$

$$X_i \sim \text{Bernoulli}(1/2)$$





$$\sigma^2 \approx 0.0046$$

$$X_i \sim \text{Beta}(10, 30)$$



- | | | | |
|-------------------|----------------------|----------------------|----------------------|
| — PM-H [Prop 1] | — Hedged [Thm 3] | — BANCO [JO19] | — Bernoulli [HRMS20] |
| --- PM-EB [Thm 2] | ... hgKelly [Prop 4] | ... A-Bentkus [KZ21] | |



Contributions

Contributions

1. Developed nonparametric confidence sets for means of bounded random variables,

Contributions

1. Developed nonparametric confidence sets for means of bounded random variables,
2. Valid at arbitrary stopping times, w/ no penalties for peeking at data early.

Contributions

1. Developed nonparametric confidence sets for means of bounded random variables,
2. Valid at arbitrary stopping times, w/ no penalties for peeking at data early.
3. Substantially outperform prior work on this problem.

Contributions

1. Developed nonparametric confidence sets for means of bounded random variables,
 2. Valid at arbitrary stopping times, w/ no penalties for peeking at data early.
 3. Substantially outperform prior work on this problem.
- + Closed-form empirical Bernstein confidence sets*

Contributions

1. Developed nonparametric confidence sets for means of bounded random variables,
 2. Valid at arbitrary stopping times, w/ no penalties for peeking at data early.
 3. Substantially outperform prior work on this problem.
- + Closed-form empirical Bernstein confidence sets*
- + Extensions to sampling without replacement*


Thank you.

ian.waudbysmith.com


Fixed-time confidence intervals

If $(C_t)_{t=1}^{\infty}$ is a confidence sequence, then $C_{\textcolor{green}{n}}$ is a confidence interval for any fixed $\textcolor{green}{n}$.


If $(C_t)_{t=1}^\infty$ is a confidence sequence, then $C_{\textcolor{violet}{n}}$ is a confidence interval for any fixed $\textcolor{violet}{n}$.




If $(C_t)_{t=1}^\infty$ is a confidence sequence, then $C_{\textcolor{green}{n}}$ is a confidence interval for any fixed $\textcolor{green}{n}$.



If $(C_t)_{t=1}^\infty$ is a confidence sequence, then $C_{\textcolor{green}{n}}$ is a confidence interval for any fixed $\textcolor{green}{n}$.



If $(C_t)_{t=1}^\infty$ is a confidence sequence, then $C_{\textcolor{green}{n}}$ is a confidence interval for any fixed $\textcolor{green}{n}$.



$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

After some heuristic calculations*, a good choice is

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

After some heuristic calculations*, a good choice is

$$\lambda_i^+(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{\textcolor{green}{n} \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{\textcolor{red}{m}}$$

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

After some heuristic calculations*, a good choice is

$$\lambda_i^+(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{\textcolor{green}{n} \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{\textcolor{red}{m}}$$

$$\lambda_i^-(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{\textcolor{green}{n} \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{1-\textcolor{red}{m}}$$

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

After some heuristic calculations*, a good choice is

$$\lambda_i^+(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{\textcolor{green}{n} \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{\textcolor{red}{m}}$$

$$\lambda_i^-(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{\textcolor{green}{n} \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{1-\textcolor{red}{m}}$$

$$K_n^\pm(\textcolor{red}{m}) := \max \left\{ \frac{1}{2} \prod_{i=1}^{\textcolor{green}{n}} (1 + \lambda_i^+ \cdot (X_i - \textcolor{red}{m})), \frac{1}{2} \prod_{i=1}^{\textcolor{green}{n}} (1 - \lambda_i^- \cdot (X_i - \textcolor{red}{m})) \right\}$$

$$C_t := \left\{ \textcolor{red}{m} \in [0,1] : K_t(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$


After some heuristic calculations*, a good choice is

$$\lambda_i^+(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{n \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{\textcolor{red}{m}}$$

$$\lambda_i^-(\textcolor{red}{m}) := \sqrt{\frac{2 \log(2/\alpha)}{n \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{1-\textcolor{red}{m}}$$

$$K_n^\pm(\textcolor{red}{m}) := \max \left\{ \frac{1}{2} \prod_{i=1}^n (1 + \lambda_i^+ \cdot (X_i - \textcolor{red}{m})), \frac{1}{2} \prod_{i=1}^n (1 - \lambda_i^- \cdot (X_i - \textcolor{red}{m})) \right\}$$

data-dependent tuning parameters without sample splitting!



So, given X_1, \dots, X_n bounded,

$$C_n^\pm := \left\{ \textcolor{red}{m} \in [0,1] : K_n^\pm(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

is a sharp confidence interval for μ .

So, given X_1, \dots, X_n bounded,



$$C_n^\pm := \left\{ \textcolor{red}{m} \in [0,1] : K_n^\pm(\textcolor{red}{m}) < \frac{1}{\alpha} \right\}$$

is a sharp confidence interval for μ .



There's one more modification we can make to get **strictly** tighter confidence intervals!

If $C_1, C_2, \dots, C_{\underline{n}}, \dots$ forms a $(1 - \alpha)$ -confidence sequence, then so does $\bigcap_{i \leq \underline{n}} C_i$



If $C_1, C_2, \dots, C_{\textcolor{red}{n}}, \dots$ forms a $(1 - \alpha)$ -confidence sequence, then so does $\bigcap_{i \leq \textcolor{red}{n}} C_i$





If $C_1, C_2, \dots, C_{\textcolor{red}{n}}, \dots$ forms a $(1 - \alpha)$ -confidence sequence, then so does $\bigcap_{i \leq \textcolor{red}{n}} C_i$



If $C_1, C_2, \dots, C_{\textcolor{red}{n}}, \dots$ forms a $(1 - \alpha)$ -confidence sequence, then so does $\bigcap_{i \leq \textcolor{red}{n}} C_i$




If $C_1, C_2, \dots, C_{\textcolor{red}{n}}, \dots$ forms a $(1 - \alpha)$ -confidence sequence, then so does $\bigcap_{i \leq \textcolor{red}{n}} C_i$




So, $\bigcap_{i \leq \textcolor{red}{n}} C_i^\pm$ is a strict improvement over $C_{\textcolor{red}{n}}^\pm$ for free!

$$\sigma^2 = 1/4$$




$$\sigma^2 \approx 0.0046$$





$$\sigma^2 = 1/4$$

$X_i \sim \text{Bernoulli}(1/2)$



$$\sigma^2 \approx 0.0046$$


$X_i \sim \text{Beta}(10, 30)$



- H-CI [H63]
- - - Bentkus-CI [B04]
- - - - EB-CI [MP09]
- - - - - VA-EB-CI [Rmk 1]
- - - - - - Hedged-CI [Rmk 3]
- - - - - - - Anderson [A69]



$X_i \sim \text{Bernoulli}(1/2)$

$$\sigma^2 = 1/4$$



$X_i \sim \text{Beta}(10, 30)$


$$\sigma^2 \approx 0.0046$$



- H-CI [H63]
- - - Bentkus-CI [B04]
- EB-CI [MP09]
- VA-EB-CI [Rmk 1]
- Hedged-CI [Rmk 3]
- ··· Anderson [A69]



$X_i \sim \text{Bernoulli}(1/2)$

$$\sigma^2 = 1/4$$



$X_i \sim \text{Beta}(10, 30)$


$$\sigma^2 \approx 0.0046$$



- | | | |
|---|--|--|
| — H-CI [H63] | — EB-CI [MP09] | ··· Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | --- VA-EB-CI [Rmk 1] | Anderson [A69] |




$X_i \sim \text{Bernoulli}(1/2)$

$$\sigma^2 = 1/4$$



$X_i \sim \text{Beta}(10, 30)$


$$\sigma^2 \approx 0.0046$$



- | | | |
|-------------------------------|--------------------------------|-----------------------------|
| H-CI [H63] | EB-CI [MP09] | Hedged-CI [Rmk 3] |
| $\text{--- Bentkus-CI [B04]}$ | $\text{---- VA-EB-CI [Rmk 1]}$ | $\text{... Anderson [A69]}$ |




$X_i \sim \text{Bernoulli}(1/2)$

$$\sigma^2 = 1/4$$



$X_i \sim \text{Beta}(10, 30)$


$$\sigma^2 \approx 0.0046$$



- | | | |
|------------------------|------------------------|------------------------|
| — H-CI [H63] | - - - EB-CI [MP09] | Hedged-CI [Rmk 3] |
| - - - Bentkus-CI [B04] | - - - VA-EB-CI [Rmk 1] | ... - - Anderson [A69] |




$X_i \sim \text{Bernoulli}(1/2)$

$$\sigma^2 = 1/4$$



$X_i \sim \text{Beta}(10, 30)$

$$\sigma^2 \approx 0.0046$$



— H-CI [H63]	- - - EB-CI [MP09]	··· Hedged-CI [Rmk 3]
- - - Bentkus-CI [B04]	- - - VA-EB-CI [Rmk 1]	··· ··· Anderson [A69]

Recap of fixed-time confidence intervals:

Recap of fixed-time confidence intervals:

Used *sequential* ideas for **fixed-time** inference.
Most notably,

Recap of fixed-time confidence intervals:

Used *sequential* ideas for **fixed-time** inference.
Most notably,

1. Data-dependent tuning parameters, and

Recap of fixed-time confidence intervals:


Used *sequential* ideas for **fixed-time** inference.
Most notably,

1. Data-dependent tuning parameters, and
2. A strictly-stronger-than-Markov inequality.


Confidence sets for sampling without replacement

Without-replacement (WoR) sampling:


Without-replacement (WoR) sampling:




Without-replacement (WoR) sampling:



Without-replacement (WoR) sampling:




Without-replacement (WoR) sampling:




$$(x_1, \dots, x_N) \in [0,1]^N, \quad \mu := \frac{1}{N} \sum_{i=1}^N x_i$$

Without-replacement (WoR) sampling:



$$(x_1, \dots, x_N) \in [0,1]^N, \quad \mu := \frac{1}{N} \sum_{i=1}^N x_i$$
$$X_1 \sim \text{Unif}((x_1, \dots, x_N))$$

Without-replacement (WoR) sampling:




$$(x_1, \dots, x_N) \in [0,1]^N, \quad \mu := \frac{1}{N} \sum_{i=1}^N x_i$$

$$X_1 \sim \text{Unif} \left((x_1, \dots, x_N) \right)$$

$$X_2 \sim \text{Unif} \left((x_1, \dots, x_N) \setminus X_1 \right)$$

Without-replacement (WoR) sampling:




$$(x_1, \dots, x_N) \in [0,1]^N, \quad \mu := \frac{1}{N} \sum_{i=1}^N x_i$$

$$X_1 \sim \text{Unif}\left((x_1, \dots, x_N)\right)$$

$$X_2 \sim \text{Unif}\left((x_1, \dots, x_N) \setminus X_1\right)$$

⋮

Without-replacement (WoR) sampling:



$$(x_1, \dots, x_N) \in [0,1]^N, \quad \mu := \frac{1}{N} \sum_{i=1}^N x_i$$
$$X_1 \sim \text{Unif}\left((x_1, \dots, x_N)\right)$$
$$X_2 \sim \text{Unif}\left((x_1, \dots, x_N) \setminus X_1\right)$$
$$\vdots$$
$$X_t \sim \text{Unif}\left((x_1, \dots, x_N) \setminus X_1^{t-1}\right)$$

Want to estimate $\mu := \frac{1}{N} \sum_{i=1}^N x_i$

Goal: construct a game so that $(K_t(\mu))_{t=0}^N$ is a martingale under WoR sampling.

$$X_t \sim \text{Unif}\left((x_1, \dots, x_N) \setminus X_1^{t-1}\right) \implies \mathbb{E}(X_t | X_1^{t-1}) = \underbrace{\frac{N\mu - \sum_{i=1}^{t-1} X_i}{N - t + 1}}_{=: \mu_t^{\text{WoR}}}$$

Consider a “candidate mean” $m \in [0,1]$



Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$



Consider a “candidate mean” $m \in [0,1]$

$K_0 \leftarrow \$1$

For $t = 1, 2, 3, \dots$:



Consider a “candidate mean” $\textcolor{red}{m} \in [0,1]$

$K_0 \leftarrow \$1$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m_t^{\text{WoR}}), 1/m_t^{\text{WoR}})$



$$m_t^{\text{WoR}} = \frac{N\textcolor{red}{m} - \sum_{i=1}^{t-1} X_i}{N - t + 1}$$

Consider a “candidate mean” $\textcolor{red}{m} \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-\textcolor{red}{m}_t^{\text{WoR}}), 1/\textcolor{red}{m}_t^{\text{WoR}})$

Observe X_t



$$\textcolor{red}{m}_t^{\text{WoR}} = \frac{N\textcolor{red}{m} - \sum_{i=1}^{t-1} X_i}{N - t + 1}$$

Consider a “candidate mean” $\textcolor{red}{m} \in [0,1]$

$K_0 \leftarrow \$1$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-\textcolor{red}{m}_t^{\text{WoR}}), 1/\textcolor{red}{m}_t^{\text{WoR}})$

Observe X_t

$K_t \leftarrow K_{t-1} + K_{t-1} \cdot \lambda_t \cdot (X_t - \textcolor{red}{m}_t^{\text{WoR}})$

EndFor



$$\textcolor{red}{m}_t^{\text{WoR}} = \frac{N\textcolor{red}{m} - \sum_{i=1}^{t-1} X_i}{N - t + 1}$$

Consider a “candidate mean” $m \in [0,1]$

$$K_0 \leftarrow \$1$$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m), 1/m)$

Observe X_t

$$K_t \leftarrow K_{t-1} + K_{t-1} \cdot \lambda_t \cdot (X_t - m)$$

EndFor



Consider a “candidate mean” $m \in [0,1]$

$K_0 \leftarrow \$1$

For $t = 1, 2, 3, \dots$:

Gambler chooses bet $\lambda_t \in (-1/(1-m_t^{\text{WoR}}), 1/m_t^{\text{WoR}})$

Observe X_t

$K_t \leftarrow K_{t-1} + K_{t-1} \cdot \lambda_t \cdot (X_t - m_t^{\text{WoR}})$

EndFor



$$m_t^{\text{WoR}} = \frac{N\mathbf{m} - \sum_{i=1}^{t-1} X_i}{N - t + 1}$$

Then,

$$K_t^{\text{WoR}}(\mu) := \prod_{i=1}^t (1 + \lambda_i \cdot (X_i - \mu_t^{\text{WoR}}))$$



forms a nonnegative martingale, and

$$C_t^{\text{WoR}} := \left\{ m \in [0,1] : K_t^{\text{WoR}}(m) < \frac{1}{\alpha} \right\}$$



forms a $(1 - \alpha)$ -confidence sequence.

Confidence sequences for sampling WoR

Discrete 0/1 high variance





Real-valued concentrated





Confidence sequences for sampling WoR

Discrete 0/1 high variance




Real-valued concentrated






Confidence sequences for sampling WoR

Discrete 0/1 high variance





Real-valued concentrated





Confidence sequences for sampling WoR

Discrete 0/1 high variance





Real-valued concentrated





Confidence sequences for sampling WoR





Discrete 0/1 high variance







Real-valued concentrated







Confidence intervals for sampling WoR







Confidence intervals for sampling WoR







Confidence intervals for sampling WoR



Confidence intervals for sampling WoR



Confidence intervals for sampling WoR



Closed-form empirical Bernstein confidence sequences & confidence intervals

$$C_t := \left\{ m \in [0,1] : \prod_{i=1}^t (1 + \lambda_i(m) \cdot (X_i - m)) \right\}$$

While C_t is easy to compute, it is not closed-form.

However,

$$C_t^{\text{PMEB}} := \left\{ m \in [0,1] : \prod_{i=1}^t \exp \left\{ \lambda_i(X_i - m) - 4(X_i - \hat{\mu}_{i-1})^2 \psi_E(\lambda_i) \right\} \right\} \quad \text{is!}$$

$$C_t^{\text{PMEB}} := \left(\frac{\sum_{i=1}^t \lambda_i X_i}{\sum_{i=1}^t \lambda_i} \pm \frac{\log(2/\alpha) + 4 \sum_{i=1}^t (X_i - \hat{\mu}_{i-1})^2 \psi_E(\lambda_i)}{\sum_{i=1}^t \lambda_i} \right)$$



where

$$\psi_E(\lambda) := -(\log(1-\lambda) - \lambda)/4,$$



$$\lambda_t := \sqrt{\frac{2 \log(2/\alpha)}{\hat{\sigma}_{t-1} t \log(1+t)}} \wedge \frac{1}{2},$$

$$\hat{\sigma}_t^2 := \frac{1/4 + \sum_{i=1}^t (X_i - \hat{\mu}_i^2)^2}{t+1}, \quad \text{and} \quad \hat{\mu}_t := \frac{1/2 + \sum_{i=1}^t X_i}{t+1}.$$

$$X_i \sim \text{Bernoulli}(1/2)$$



$$X_i \sim \text{Beta}(10, 30)$$







Similarly for fixed-time confidence intervals:

$$C_n^{\text{PMEB}} := \left(\frac{\sum_{i=1}^n \lambda_i X_i}{\sum_{i=1}^n \lambda_i} \pm \frac{\log(2/\alpha) + 4 \sum_{i=1}^n (X_i - \hat{\mu}_{i-1})^2 \psi_E(\lambda_i)}{\sum_{i=1}^n \lambda_i} \right)$$

but here, $\lambda_i := \sqrt{\frac{2 \log(2/\alpha)}{n \hat{\sigma}_{i-1}}} \wedge \frac{1}{2}$,

Final bound: $\bigcap_{i \leq n} C_i^{\text{PMEB}}$



Choice of $(\lambda_t^+)^n_{t=1}$ and $(\lambda_t^-)^n_{t=1}$ for
fixed-time confidence intervals

Why does perform so well?

$$\begin{aligned} K_n^+(\mu) &:= \prod_{i=1}^n (1 + \lambda \cdot (X_i - \mu)) \\ &\gtrsim \prod_{i=1}^n \exp \left\{ \lambda \cdot (X_i - \mu) - (X_i - \hat{\mu}_{i-1})^2 \lambda^2 / 2 \right\} \end{aligned}$$

$$\begin{aligned} \implies \text{Width}_n &:= \frac{\log(2/\alpha) + \sum_{i=1}^n (X_i - \hat{\mu}_{i-1})^2 / 2}{t\lambda} \\ &\approx \frac{\log(2/\alpha) + n\sigma^2 / 2}{t\lambda} \end{aligned}$$

Why does $\lambda_i^+(m) := \sqrt{\frac{2 \log(2/\alpha)}{n \hat{\sigma}_{i-1}^2}} \wedge \frac{1}{m}$ perform so well?

$$\begin{aligned}
 K_n^+(\mu) &:= \prod_{i=1}^n (1 + \lambda \cdot (X_i - \mu)) \\
 &\gtrsim \prod_{i=1}^n \exp \left\{ \lambda \cdot (X_i - \mu) - (X_i - \hat{\mu}_{i-1})^2 \lambda^2 / 2 \right\}
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow \text{Width}_n &:= \frac{\log(2/\alpha) + \sum_{i=1}^n (X_i - \hat{\mu}_{i-1})^2 / 2}{t\lambda} \\
 &\approx \frac{\log(2/\alpha) + n\sigma^2 / 2}{t\lambda}
 \end{aligned}$$

$$\operatorname*{argmin}_{\lambda} \text{Width}_n = \sqrt{\frac{2 \log(2/\alpha)}{n\sigma^2}}$$

$$\operatorname*{argmin}_{\lambda} \text{Width}_n = \sqrt{\frac{2 \log(2/\alpha)}{n\sigma^2}}$$

$$\lambda_t^+(m) := \sqrt{\frac{2 \log(2/\alpha)}{n\hat{\sigma}_{t-1}^2}} \wedge \frac{1}{m}$$

$$\operatorname*{argmin}_{\lambda} \text{Width}_n = \sqrt{\frac{2 \log(2/\alpha)}{n\sigma^2}}$$

$$\lambda_t^+(m) := \sqrt{\frac{2 \log(2/\alpha)}{n\hat{\sigma}_{t-1}^2}} \wedge \frac{1}{m} \quad \quad \lambda_t^-(m) := \sqrt{\frac{2 \log(2/\alpha)}{n\hat{\sigma}_{t-1}^2}} \wedge \frac{1}{1-m}$$

Brief selective history of betting ideas

