


\mathcal{P} -uniform anytime-valid inference

ICSIDS 2023 | Lisbon, Portugal

Ian Waudby-Smith

Carnegie Mellon University
Pittsburgh, PA, USA



Ian Waudby-Smith
CMU Statistics



Aaditya Ramdas
CMU Statistics

Some background: CLT-based confidence intervals

Some background: CLT-based confidence intervals

Given $X_1, \dots, X_n \sim P$ with mean $\mu_P \equiv \mathbb{E}_P(X)$ and finite variance,

Some background: CLT-based confidence intervals

Given $X_1, \dots, X_n \sim P$ with mean $\mu_P \equiv \mathbb{E}_P(X)$ and finite variance,

$$\lim_{n \rightarrow \infty} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha,$$

Some background: CLT-based confidence intervals

Given $X_1, \dots, X_n \sim P$ with mean $\mu_P \equiv \mathbb{E}_P(X)$ and finite variance,

$$\lim_{n \rightarrow \infty} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha,$$

where $\alpha \in (0, 1)$ and

$$\dot{C}_n := \frac{1}{n} \sum_{i=1}^n X_i \pm \frac{\hat{\sigma}_n \cdot \Phi^{-1}(1 - \alpha/2)}{\sqrt{n}}.$$

Some background: CLT-based confidence intervals

Given $X_1, \dots, X_n \sim P$ with mean $\mu_P \equiv \mathbb{E}_P(X)$ and finite variance,

$$\lim_{n \rightarrow \infty} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha,$$

where $\alpha \in (0, 1)$ and

$$\dot{C}_n := \frac{1}{n} \sum_{i=1}^n X_i \pm \frac{\hat{\sigma}_n \cdot \Phi^{-1}(1 - \alpha/2)}{\sqrt{n}}.$$

Only holds for a *fixed* and *prespecified* sample size n .

(Asymptotic) confidence sequences enable statistical inference at stopping times.

For example, continuously monitoring the CIs of a sequential experiment.

What does this mean *mathematically*?

What does this mean *mathematically*?

The guarantees of asymptotic confidence sequences $(\bar{C}_{\textcolor{red}{k}}^{(\textcolor{blue}{m})})_{\textcolor{red}{k}=\textcolor{blue}{m}}^{\infty}$ hold *uniformly* for all sufficiently large sample sizes $\textcolor{red}{k} \geq \textcolor{blue}{m}$.


What does this mean *mathematically*?

The guarantees of asymptotic confidence sequences $(\bar{C}_k^{(m)})_{k=m}^\infty$ hold *uniformly* for all sufficiently large sample sizes $k \geq m$.

Confidence interval	Confidence sequence
$\lim_{n \rightarrow \infty} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha$	$\lim_{m \rightarrow \infty} \mathbb{P}_P \left(\exists k \geq m : \mu_P \notin \bar{C}_k \right) = \alpha$

What does this mean *practically*?

What does this mean *practically*?



Note that asymptotic CIs have an even stronger \mathcal{P} -uniform guarantee.

Note that asymptotic CIs have an even stronger \mathcal{P} -uniform guarantee. If

$$\sup_{P \in \mathcal{P}} \mathbb{E}_P |X - \mu_P|^{2+\delta} < \infty,$$

then the guarantees of \dot{C}_n hold *uniformly* in \mathcal{P} :

$$\lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha.$$

This paper: \mathcal{P} -uniform anytime-valid inference

This paper: \mathcal{P} -uniform anytime-valid inference

*Can similar uniform statements be
made for confidence sequences?*

This paper: \mathcal{P} -uniform anytime-valid inference

Can similar uniform statements be made for confidence sequences?

	Confidence interval	Confidence sequence
P -pointwise	$\lim_{n \rightarrow \infty} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha$	$\lim_{m \rightarrow \infty} \mathbb{P}_P \left(\exists k \geq m : \mu_P \notin \bar{C}_k \right) = \alpha$
\mathcal{P} -uniform	$\lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} \mathbb{P}_P \left(\mu_P \notin \dot{C}_n \right) = \alpha$	$\lim_{m \rightarrow \infty} \sup_{P \in \mathcal{P}} \mathbb{P}_P \left(\exists k \geq m : \mu_P \notin \bar{C}_k \right) = \alpha$

The answer is "**Yes**", but this required first filling a gap in the probability literature (i.e. strong invariance principles).

The answer is **"Yes"**, but this required first filling a gap in the probability literature (i.e. strong invariance principles).

Informal: Need to show that for iid $(X_n)_{n=1}^\infty$, there exist standard Gaussians $(Z_n)_{n=1}^\infty$ so that

$$\sum_{i=1}^n \frac{X_i - \mu_P}{\sigma_P} - \sum_{i=1}^n Z_i = o\left(\sqrt{n \log \log n}\right),$$

P -almost surely uniformly in $P \in \mathcal{P}$.

Theorem: \mathcal{P} -uniform asymptotic confidence sequences

Suppose $(X_n)_{n=1}^\infty$ have \mathcal{P} -uniformly bounded $(2 + \delta)^{\text{th}}$ moments.

Letting $\bar{C}_{\mathbf{k}}^{(\mathbf{m})}$ be given by

$$\bar{C}_{\mathbf{k}}^{(\mathbf{m})} := \hat{\sigma}_{\mathbf{k}} \sqrt{\frac{\Psi^{-1}(1 - \alpha) + \log(\mathbf{k}/\mathbf{m})}{\mathbf{k}}}.$$

Theorem: \mathcal{P} -uniform asymptotic confidence sequences

Suppose $(X_n)_{n=1}^\infty$ have \mathcal{P} -uniformly bounded $(2 + \delta)^{\text{th}}$ moments.
Letting $\bar{C}_{\mathbf{k}}^{(\mathbf{m})}$ be given by

$$\bar{C}_{\mathbf{k}}^{(\mathbf{m})} := \hat{\sigma}_{\mathbf{k}} \sqrt{\frac{\Psi^{-1}(1 - \alpha) + \log(\mathbf{k}/\mathbf{m})}{\mathbf{k}}}.$$

Then the asymptotic guarantees of $(\bar{C}_{\mathbf{k}}^{(\mathbf{m})})_{\mathbf{k}=\mathbf{m}}^\infty$ hold *uniformly* in \mathcal{P} :

$$\lim_{\mathbf{m} \rightarrow \infty} \sup_{P \in \mathcal{P}} \mathbb{P}_P \left(\exists \mathbf{k} \geq \mathbf{m} : \mu_P \notin \bar{C}_{\mathbf{k}}^{(\mathbf{m})} \right) = \alpha.$$

Thank you!
ianws.com