

A Fluid Introduction to Condensed Mathematics

Wannes Malfait

MOTIVATION

Problem: Algebra and topology clash:

$$\text{id}: \mathbb{R}_{\text{discrete}} \rightarrow \mathbb{R}_{\text{Euclidean}}$$

is epi + mono, but not iso

Solution?

MOTIVATION

Problem: Algebra and topology clash:

$$\text{id}: \mathbb{R}_{\text{discrete}} \rightarrow \mathbb{R}_{\text{Euclidean}}$$

is epi + mono, but not iso

Solution?

\rightsquigarrow Condensed abelian groups

PRESHEAVES

Idea: Study better behaved objects.

Idea: Study better behaved objects.

Definition

A **presheaf** on a topological space X is a functor

$$\mathcal{F}: \mathbf{Open}(X)^{\mathrm{opp}} \rightarrow \mathbf{Set} / \mathbf{Rng} / \mathbf{Ab} / \dots$$

Notation:

- ▶ Elements of $\mathcal{F}(U)$ are called **sections**
- ▶ If $i: U \hookrightarrow V$ then $\mathcal{F}(i)$ is called the **restriction** map:

$$s \in \mathcal{F}(V) \mapsto s|_U \in \mathcal{F}(U).$$

Example

We always have the presheaf given by

- ▶ $\mathcal{F}(U) \mapsto Y$, and
- ▶ $\mathcal{F}(f) \mapsto \text{id}_Y$

Example

We always have the presheaf given by

- ▶ $\mathcal{F}(U) \mapsto Y$, and
- ▶ $\mathcal{F}(f) \mapsto \text{id}_Y$

Problem:

- ▶ Want more structure
- ▶ Want global things to be defined locally

\rightsquigarrow sheaf conditions:

SHEAF CONDITIONS

For an open cover $\{U_i\}_{i \in I}$ of U :

- ▶ **uniqueness/locality**: If $s, t \in \mathcal{F}(U)$ are sections such that $s|_{U_i} = t|_{U_i}$ for all $i \in I$, then $s = t$.

SHEAF CONDITIONS

For an open cover $\{U_i\}_{i \in I}$ of U :

- ▶ **uniqueness/locality**: If $\mathbf{s}, \mathbf{t} \in \mathcal{F}(U)$ are sections such that $\mathbf{s}|_{U_i} = \mathbf{t}|_{U_i}$ for all $i \in I$, then $\mathbf{s} = \mathbf{t}$.
- ▶ **gluing**: If $\{\mathbf{s}_i \in \mathcal{F}(U_i)\}_{i \in I}$ is a collection of sections such that

$$\mathbf{s}_i|_{U_i \cap U_j} = \mathbf{s}_j|_{U_i \cap U_j} \text{ for all } i, j,$$

then there is a section $\mathbf{s} \in \mathcal{F}(U)$ such that $\mathbf{s}_i = \mathbf{s}|_{U_i}$ for all $i \in I$.

EXAMPLES

Typical examples are rings of functions:

- ▶ $\text{Cont}(-, \mathbb{C})$
- ▶ $C^\infty(-, \mathbb{C})$

EXAMPLES

Typical examples are rings of functions:

- ▶ $\text{Cont}(-, \mathbb{C})$
- ▶ $\mathcal{C}^\infty(-, \mathbb{C})$

The constant functor $U \mapsto Y$ is **not** a sheaf!

Uniqueness axiom implies $\mathcal{F}(\emptyset) = \{*\}$

Proposition

Limits and colimits of presheaves can be computed section-wise, i.e. the functor $\mathcal{F} \mapsto \mathcal{F}(U)$ commutes with all limits and colimits.

Proposition

Limits and colimits of presheaves can be computed section-wise, i.e. the functor $\mathcal{F} \mapsto \mathcal{F}(U)$ commutes with all limits and colimits.

Example

$$(\mathcal{F} \times \mathcal{G})(U) = \mathcal{F}(U) \times \mathcal{G}(U)$$

SHEAFIFICATION

Can we do the same for sheaves?

SHEAFIFICATION

Can we do the same for sheaves?

- ▶ Limits: ok.
- ▶ Colimits: not always a sheaf.

There is a solution:

SHEAFIFICATION

Can we do the same for sheaves?

- ▶ Limits: ok.
- ▶ Colimits: not always a sheaf.

There is a solution:

Theorem

The fully faithful inclusion $\iota: \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$ admits an *exact left adjoint: sheafification*.

Example

Sheafification of $U \mapsto Y$, is $U \mapsto \text{Cont}(U, Y_{\text{discrete}})$.

Proposition

The category of abelian sheaves on a topological space X is abelian, and has enough injectives.

Example

If M is an injective object in \mathbf{Ab} , i.e. a divisible group, then for $x \in X$,

$$U \mapsto \begin{cases} M & x \in U \\ \{*\} & x \notin U \end{cases}$$

is an injective sheaf.

SHEAF COHOMOLOGY

If

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F}'' \longrightarrow 0$$

is exact, then globally we only get

$$0 \longrightarrow \mathcal{F}(X) \longrightarrow \mathcal{F}'(X) \longrightarrow \mathcal{F}''(X)$$

SHEAF COHOMOLOGY

If

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F}'' \longrightarrow 0$$

is exact, then globally we only get

$$0 \longrightarrow \mathcal{F}(X) \longrightarrow \mathcal{F}'(X) \longrightarrow \mathcal{F}''(X)$$

\rightsquigarrow Right derived functors $H^i(X, -)$ of $\mathcal{F} \mapsto \mathcal{F}(X)$

Definition

The sheaf cohomology of X with coefficients A is $H^i(X, \text{Cont}(-, A))$.

We have

$$0 \longrightarrow \text{Cont}(\mathbb{Z}, -) \longrightarrow \text{Cont}(\mathbb{R}, -) \longrightarrow \text{Cont}(\mathbb{R}/\mathbb{Z}, -) \longrightarrow 0$$

since every map $U \subseteq S^1 \rightarrow S^1$ can be lifted **locally** to a map $U \rightarrow \mathbb{R}$.

But, there is no global lift of $\text{id}: S^1 \rightarrow S^1$ to $S^1 \rightarrow \mathbb{R}$. So $H^1(S^1, \mathbb{Z}) \neq 0$.

We have

$$0 \rightarrow \text{Cont}(\mathbb{Z}, -) \rightarrow \text{Cont}(\mathbb{R}, -) \rightarrow \text{Cont}(\mathbb{R}/\mathbb{Z}, -) \rightarrow 0$$

since every map $U \subseteq S^1 \rightarrow S^1$ can be lifted **locally** to a map $U \rightarrow \mathbb{R}$.

But, there is no global lift of $\text{id}: S^1 \rightarrow S^1$ to $S^1 \rightarrow \mathbb{R}$. So $H^1(S^1, \mathbb{Z}) \neq 0$.

Proposition

If S is a CW-complex:

$$H_{\text{sheaf}}^i(S, \mathbb{Z}) = H_{\text{singular}}^i(S, \mathbb{Z}).$$

- ▶ Need a slight abstraction of sheaves on a space.
- ▶ Want sheaves on categories, not just spaces.
- ▶ Crucial ingredient: coverings.

- ▶ Need a slight abstraction of sheaves on a space.
- ▶ Want sheaves on categories, not just spaces.
- ▶ Crucial ingredient: coverings.

↪ Definition of a site \approx "Category + notion of coverings".

CONDENSED SETS

Definition

A **condensed set**^{*} is a sheaf on the site of profinite sets with coverings given by jointly surjective maps.

CONDENSED SETS

Definition

A **condensed set**^{*} is a sheaf on the site of profinite sets with coverings given by jointly surjective maps.

For a topological space X , we have an associated condensed set $\underline{X} = \text{Cont}(-, X)$.

Definition

A **profinite set** is a compact, Hausdorff, totally disconnected space.

Example

A finite discrete space

PROFINITE SETS

Definition

A **profinite set** is a compact, Hausdorff, totally disconnected space.

Example

A finite discrete space

Proposition

A limit of profinite sets is profinite. In particular, products of profinite sets are again profinite.

THE MOTIVATING PROBLEM

Claim: for condensed sets the map:

$$\underline{\text{id}}: \underline{\mathbb{R}}_{\text{discrete}} \rightarrow \underline{\mathbb{R}}_{\text{Euclidean}}$$

is no longer an epimorphism.

THE MOTIVATING PROBLEM

Claim: for condensed sets the map:

$$\text{id}: \underline{\mathbb{R}}_{\text{discrete}} \rightarrow \underline{\mathbb{R}}_{\text{Euclidean}}$$

is no longer an epimorphism.

Let $S \subset \mathbb{R}_{\text{Euclidean}}$ be the cantor set. $S \cong \prod_{n \in \mathbb{N}} \{0, 1\}$, so S is profinite. Hence:

$$\text{Cont}(S, \mathbb{R}_{\text{discrete}}) \subsetneq \text{Cont}(S, \mathbb{R}_{\text{Euclidean}})$$

CONDENSED COHOMOLOGY

For a condensed set T , let $\mathbb{Z}[T]$ be the sheafification of $S \mapsto \mathbb{Z}[T(S)]$.

Definition

The cohomology of S with coefficients A is

$$H^i(S, A) = \mathrm{Ext}^i(\mathbb{Z}[\underline{S}], \underline{A})$$

CONDENSED COHOMOLOGY

For a condensed set T , let $\mathbb{Z}[T]$ be the sheafification of $S \mapsto \mathbb{Z}[T(S)]$.

Definition

The cohomology of S with coefficients A is

$$H^i(S, A) = \operatorname{Ext}^i(\mathbb{Z}[\underline{S}], \underline{A})$$

Theorem ([Sch19], Theorem 3.2)

Let S be a compact Hausdorff space. Then

$$H_{cond}^i(S, A) \cong H_{sheaf}^i(S, A)$$

REFERENCES



Rodrigo Marlasca Aparicio.

Condensed mathematics: The internal hom of condensed sets and condensed abelian groups and a prismatic construction of the real numbers, 2021.



Dustin Clausen and Peter Scholze.

Masterclass in condensed mathematics.

<https://www.math.ku.dk/english/calendar/events/condensed-mathematics/>, November 2020.



Peter Scholze.

Lectures on condensed mathematics.

<https://www.math.uni-bonn.de/people/scholze/Condensed.pdf>, 2019.



The Stacks Project Authors.

The stacks project.

<https://stacks.math.columbia.edu/>, 2023.