

Applied Numerical Analysis (7th Edition)

Chapter 3, Problem 14E

Bookmark

Show all steps: ON

Problem

Use the divided-difference table from Exercise to interpolate for $f(0.4)$

a. Using the first three points.

b. Using the last three points.

c. Using the best set of three points. Which points should be used?

d. Using the best set of four points.

e. Using all of the points.

f. Explain why the results are not all the same.

Exercise

Construct the divided-difference table from these data:

x	-0.2	0.3	0.7	-0.3	0.1
$f(x)$	1.23	2.34	-1.05	6.51	-0.06

Step-by-step solution

Step 1 of 7

When using numerical approximation techniques, it is useful to have programs written to aid in numerical calculations to avoid calculations by hand. Using Matlab, create an

M-file named 'ddtinterp.m' which will contain Matlab code that implements the algorithm to create a divided difference table interpolation corresponding to a set of node data points and interpolation point.

Inside this file, write the code to implement the method. One example of a correctly executable routine based on the algorithm description in the text is as follows:

```
function fp = ddtinterp(x,y,p)

%given an input vector x of x data values
%and an input vector f of f data values,
%the divided difference table d is generated
%and outputed. Additionally, an interpolation is
%performed using x=p.

n = length(x);
a(1) = y(1);
for k = 1 : n - 1
    d(k, 1) = (y(k+1) - y(k))/(x(k+1) - x(k));
end
for j = 2 : n - 1
    for k = 1 : n - j
        d(k, j) = (d(k+1, j - 1) - d(k, j - 1))/(x(k+j) - x(k));
    end
end
for j = 2 : n
    a(j) = d(1, j-1);
end
D(1) = 1;
c(1) = a(1);
for j = 2 : n
    D(j) = (p - x(j-1)) .* D(j-1);
    c(j) = a(j) .* D(j);
end
d(n,:) = 0;
d = [x' y' d];
d
fp = sum(c);
```

[Comment](#)

Step 2 of 7

a) To interpolate $f(0.4)$ using the first three points, use the following Matlab commands

INPUT:

```
>> x=[-0.2,0.3,0.7];
>> f=[1.23,2.34,-1.05];
>> ddtinterp(x,f,0.4)
```

OUTPUT:

```
d =
-0.2000 1.2300 2.2200 -11.8833
0.3000 2.3400 -8.4750 0
0.7000 -1.0500 0 0
ans =
1.8490
```

[Comment](#)

Search



Post a question

Answers from our experts for your tough homework questions

Continue to post

20 questions remaining

My Textbook Solutions



Applied
Numerical...
7th Edition



Engineering
Mechanics
14th Edition



Numerical
Methods for...
7th Edition

[View all solutions](#)

Chegg tutors who can help right now



Grisha
PhD - Capella Univ... 250



Y
University of San Fr... 760



Matthew
University of Color... 707

FIND ME A TUTOR

b) To interpolate $f(0.4)$ using the last three points, use the following Matlab commands

```
INPUT:
>> x=[0,7,-0,3,0,1];
>> f=[-1,05,6,51,-0,06];
>> ddinterp(x,f,0,4)

OUTPUT:
d =
0,7000 -1,0500 -7,5600 14,7750
-0,3000 6,5100 -16,4250 0
0,1000 -0,0600 0 0

ans =
-1,8847
```

[Comment](#)

Step 4 of 7

c) To interpolate $f(0.4)$ using the best three points, use the following Matlab commands with the data values corresponding to $x = .1$, $x = .3$, and $x = .7$ since they are best centered around $x = 0.4$.

```
INPUT:
>> x=[0,3,0,7,0,1];
>> f=[2,34,-1,05,-0,06];
>> ddinterp(x,f,0,4)

OUTPUT:
d =
0,3000 2,3400 -8,4750 -34,1250
0,7000 -1,0500 -1,6500 0
0,1000 -0,0600 0 0

ans =
2,5162
```

[Comment](#)

Step 5 of 7

d) To interpolate $f(0.4)$ using the best four points, use the following Matlab commands with the data values corresponding to $x = .1$, $x = .3$, and $x = .7$ since they are best centered around $x = 0.4$, and $x = -0.2$ since it is the next closest eligible point

```
INPUT:
>> x=[-0,2,0,3,0,7,0,1];
>> f=[1,23,2,34,-1,05,-0,06];
>> ddinterp(x,f,0,4)

OUTPUT:
d =
-0,2000 1,2300 2,2200 -11,8833 -74,1389
0,3000 2,3400 -8,4750 -34,1250 0
0,7000 -1,0500 -1,6500 0 0
0,1000 -0,0600 0 0 0

ans =
3,1835
```

[Comment](#)

Step 6 of 7

e) To interpolate $f(0.4)$ using all the points, use the following Matlab commands

```
INPUT:
>> x=[-0,2,0,3,0,7,-0,3,0,1];
>> f=[1,23,2,34,-1,05,6,51,-0,06];
>> ddinterp(x,f,0,4)

OUTPUT:
d =
-0,2000 1,2300 2,2200 -11,8833 -103,5833 73,6111
0,3000 2,3400 -8,4750 -1,5250 -81,5000 0
0,7000 -1,0500 -7,5600 14,7750 0 0
-0,3000 6,5100 -16,4250 0 0 0
0,1000 -0,0600 0 0 0 0

ans =
2,7860
```

[Comment](#)

Step 7 of 7

f) The results will not all be the same because each divided difference table represents a different interpolating polynomial function. So when interpolating at the point $x = 0.4$ in the previous steps, a different polynomial function was used to evaluate $f(0.4)$.

[Comment](#)

Was this solution helpful?

0

0

Exercises 1 and 2 to get the quadratics in the form $ax^2 + bx + c$. How different are the values for a, b, and c? Exercise 1 Write out the Lagrangian polynomial from this table....

[View this solution](#)

of points: Point #
0123456789x10507590105150180190160130y101560
Draw the graph determined by the ten points.b.
Why is the graph smoothly connected at points 3 and 6?c....

[View this solution](#)

ABOUT CHEGG

Media Center
College Marketing
Privacy Policy
Your CA Privacy Rights
Terms of Use
General Policies
Intellectual Property Rights
Investor Relations
Corporate Development
Enrollment Services

RESOURCES

Site Map
Mobile
Publishers
Join Our Affiliate Program
Advertising Choices

TEXTBOOK LINKS

Return Your Books
Textbook Rental
eTextbooks
Used Textbooks
Cheap Textbooks
College Textbooks
Sell Textbooks

STUDENT SERVICES

Chegg Play
Study 101
Chegg Coupon
Scholarships
Career Search
Internships
College Search
College Majors
Scholarship Redemption

COMPANY

Jobs
Customer Service
Give Us Feedback
Chegg For Good
Become a Tutor

LEARNING SERVICES

Online Tutoring
Chegg Study Help
Solutions Manual
Tutors by City
GPA Calculator
Test Prep



© 2003-2018 Chegg Inc. All rights reserved.