VCE MATHEMATICAL METHODS · EXAM 2

Bound reference

Short edition: the CAS commands, formula must-knows and notes on every topic, with the worked examples that match recurring Exam 2 questions, and the hardest real Exam 2 multiple-choice questions (2023–2025, under 50% correct) after their topics. Section B questions are in the full edition. Question finder: page 2. Index: page 46.

START HERE

TI-Nspire CX II CAS commands **3**

Methods formula sheet & must-knows **8**

FUNCTIONS, RELATIONS & GRAPHS

Functions: domain, range, hybrids & modulus · exam question p. **11** **9**

Polynomials: factor theorem, cubics & quartics · exam question p. **13** **11**

Power functions, hyperbolas & root functions **13**

Transformations of graphs · exam question p. **16** **14**

Exponential & logarithmic functions **17**

Circular (trigonometric) functions · exam question p. **20** **18**

Composite, inverse & combined functions · exam question p. **23** **21**

Functional equations, odd & even functions **23**

ALGEBRA, NUMBER & STRUCTURE

Simultaneous equations & parameters **24**

Newton's method, bisection & pseudocode · exam question p. **26** **25**

CALCULUS

Limits, continuity & differentiability · exam question p. **28** **27**

Differentiation rules · exam question p. **29** **28**

Applications of differentiation · exam question p. **31** **29**

Optimisation (max/min problems) · exam question p. **32** **31**

Antidifferentiation **32**

Definite integrals, areas & average value · exam question p. **35** **33**

Numerical integration: trapezium & rectangle rules · exam question p. **36** **35**

DATA ANALYSIS, PROBABILITY & STATISTICS

Probability: conditional & independent events · exam question p. **38** **37**

Discrete random variables · exam question p. **39** **38**

The binomial distribution · exam question p. **40** **39**

Continuous random variables (pdfs) **40**

The normal distribution **42**

Sampling & sample proportions **43**

Confidence intervals for proportions **44**

Exam questions © Victorian Curriculum and Assessment Authority (VCAA) (the questions themselves are not reprinted in this copy: find them in the official papers). Answers, marking points and notes are paraphrased study notes built from the VCAA examiners' reports, not official marking schemes.

Question finder

Question type	Notes: page	CAS: page, row
FUNCTIONS, RELATIONS & GRAPHS		
Is it a function? One-to-one?	**9**	
Maximal (implied) domain	**9**	**4** C20
Range over a restricted domain	**10**	**5** C40
Hybrid function, continuity at a join	**10**	**4** C21
Sketch a cubic or quartic	**11–12**	**4** C12
Number of solutions as k varies	**12**	**4** C17
Distinct roots or intercepts with a parameter	**12**	
Cubic from its graph or turning points	**12**	**4** C18
Hyperbola, truncus, root: asymptotes, intercepts	**14**	**4** C13
Image of a graph under a transformation	**15**	**4** C25
Transformations that map points onto points	**15**	
Undo a transformation (inverse transformation)	**15**	
Family with a parameter: locus, common points, k conditions	**16**	**5** C28
Where f meets its inverse; tangency with a parameter	**16, 22**	**5** C27
Exp/log graph: asymptote, domain, intercepts	**17**	**4** C20
Exponential or log equation, hidden quadratic	**17**	**4** C7
Fit $y = Ae^{kx}$ through two points, change of base	**18**	
Trig graph: period, range, asymptotes, inflections	**18**	
Trig equation on an interval, general solution	**19**	**4** C8
Quadratic in sin x; show there are no solutions	**19–20**	
Quadratic or composite inequality	**19**	**4** C16
Periodic model: fit it, max, min, time above a level	**20**	**4** C8
Trend plus a periodic part (rise per period)	**20**	
Sum of the solutions of a trig equation	**20**	**4** C15
Composite: does it exist, its rule, its range	**21**	**4** C23
Graph of g(f(x)) when f is linear	**21**	
Inverse function, its domain, choosing the branch	**21**	**4** C24
Where f^{-1} meets g^{-1} (swap the points)	**22**	
Integral of an inverse function	**22**	
Functional equation, odd/even, periodic	**23**	**4** C19
Smooth join: find a and b	**27**	**5** C39
Derivative of a hybrid, domain of f′	**27**	
ALGEBRA, NUMBER & STRUCTURE		
Simultaneous equations: none or infinitely many	**24**	**4** C11
Line tangent to a curve (discriminant)	**24**	**4** C17
Newton's method iterations	**25**	**6** C51
Trace pseudocode, accumulator loops	**25–26**	

CALCULUS		
What follows from the sign of f′ (monotone, no inverse)	**21, 29–30**	
Chain, product and quotient rules	**28**	**5** C32
Derivatives from a table of values	**28**	
Average or instantaneous rate of change	**29**	**5** C32
Tangent and normal, tangent through a point	**29**	**5** C38
Stationary points, gradient table, strictly increasing	**29–30**	**5** C33
Inflection; greatest rate of change	**30**	**5** C36
Which graph could be f′ (or F)?	**30–31, 33**	**5** C29
Optimisation in context, endpoints	**31–32**	**5** C40
Greatest difference between two models	**32**	**5** C42
Properties of definite integrals	**33**	
f from f′ and a point	**33**	**5** C43
“Hence” antiderivative by recognition	**33**	
Area under a curve and between curves	**34**	**5** C44
One integral for an area with crossings	**34**	**5** C46
Average value; integral over a period	**34**	**5** C47
Total change from a rate, solve for a limit	**35**	**6** C48
Trapezium rule: estimate, over or under	**35–36**	**6** C53
DATA ANALYSIS, PROBABILITY & STATISTICS		
Conditional probability, trees, false positives	**37**	
Independent trials with different probabilities	**38**	**6** C54
DRV: k, mean, variance, E(aX + b)	**38**	**6** C55
DRV: conditional probability from a table	**38**	
Binomial: show that, at least one, smallest n	**39**	**6** C59
Binomial: conditional, n and p, rounded values	**39–40**	**6** C60
pdf: k, mean, median, mode, variance, percentile	**40–41**	**6** C63
pdf, then binomial; integer k from a rounded probability	**40**	**6** C65
Normal: probability and percentile	**42**	**6** C68
Normal: unknown σ, or μ and σ	**42**	**7** C73
Normal, then binomial (the Exam 2 chain)	**42**	**7** C70
Sample proportion probabilities, simulation	**43**	**7** C74
Confidence interval, sample size, level	**44**	**7** C77
How many intervals contain p?	**44**	**6** C57

START HERE

TI-Nspire CX II CAS commands

Typed in a Calculator page on the TI-Nspire CX II CAS. Each example has a checked output (rounded: the screen shows 6 significant figures by default), so you can test your syntax against an answer you already know; the page number in each row is the notes page for that question type, and the row numbers (C1, C2, …) are what the question finder points to. On paper, always write the equation you solved and the answer in the form asked, not just “by CAS”.

Set up

To do this	Type or press
C1 Radians, real answers	doc ▸ Settings: Angle **Radian**, Real or Complex **Real**, Calculation Mode **Auto** (Home ▸ Settings ▸ Document Settings ▸ Make Default for new documents)
C2 Start each question clean	insert a new Problem (doc ▸ Insert ▸ Problem), or `ClearAZ` (menu ▸ Actions ▸ Clear a–z; single-letter names only)
C3 Decimal answer	**ctrl + enter**, or type a decimal point (`2.`): decimals in give decimals out. `exact(0.25)` gives 1/4; `approx(π/4)` gives 0.785398
C4 Last answer; “such that”	`ans` (ctrl + (−)); `|` from the ctrl + = palette, e.g. `f(x)|x=2`
C5 Define; store a value	`Define f(x)=x^3-3·x` (menu ▸ Actions ▸ Define); store an unrounded result with **ctrl + var** (→), e.g. `normCdf(180,∞,170,8)→pr`, then use `pr` (pick a name you won’t solve for later)

To do this	Type or press
C6 Special symbols	`e^(x)` from the e^x key (the letter e is a variable); ∞ from the catalog's symbols, or `1E99` (the E from the **EE** key, not the letter); `<=` and `>=` become ≤ and ≥

Algebra and parameters

To do this	Type	Gives
C7 Solve	`solve(x^2-5·x+6=0,x)`	x = 2 or x = 3
C8 Solve on an interval → p. 19	`solve(2·sin(2·x)=1,x)|0≤x≤π`	x = π/12 or x = 5π/12
C9 Solve numerically → p. 25	`nSolve(x^3+x-1=0,x)|0≤x≤1` or with a guess `nSolve(x^3+x-1=0,x=0.5)`	0.682328 (one solution only)
C10 Simultaneous equations → p. 24	`solve(2·x+y=5 and x-y=1,{x,y})`	x = 2 and y = 1
C11 None or infinitely many solutions → p. 24	`solve(det([a,3;1,a-2])=0,a)`, then solve the system with each value of a	a = −1 or a = 3; `false` means no solution, an answer in terms of c1 means infinitely many
C12 Factorise, expand; roots → p. 11	`factor(2·x^3-3·x^2-11·x+6)`; `expand((x+1)^3)`; `zeros(x^3-2·x^2-5·x+6,x)`	(x − 3)(2x − 1)(x + 2); $x^3 + 3x^2 + 3x + 1$; {−2, 1, 3}
C13 Remainder; hyperbola form; turning-point form → pp. 11, 14	`polyRemainder(x^3+2·x+1,x-1,x)`; `propFrac((2·x+3)/(x+1),x)`; `completeSquare(x^2-6·x+5,x)`	4; 1/(x + 1) + 2; $(x - 3)^2 - 4$
C14 Inequality → p. 39	`solve(1-0.9^n>0.95,n)`	n > 28.4332, so n = 29
C15 Sum of the solutions → p. 20	`zeros(2·sin(2·x)-1,x)|0≤x≤π`, then `sum(ans)`	{π/12, 5π/12}, then π/2
C16 Quadratic inequality (then use the range) → p. 19	`solve(2·a^2+a-1>0,a)` with a = sin x	a < −1 or a > 1/2, so sin x > 1/2
C17 k for a number of solutions → pp. 12, 24	`solve(k^2-16>0,k)` (Δ of $x^2 + kx + 4 > 0$)	k < −4 or k > 4
C18 Cubic from turning points (coefficients from conditions) → p. 12	`Define p(x)=a·x^3+b·x^2+c·x+d`, `Define dp(x)=derivative(p(x),x)`, `solve(p(1)=5 and p(3)=1 and dp(1)=0 and dp(3)=0,{a,b,c,d})`	a = 1, b = −6, c = 9, d = 1
C19 Check a functional equation → p. 23	`Define f(x)=e^(3·x)`, then `f(x+y)=f(x)·f(y)`	true (anything else isn't a proof that it fails: test values)

Functions and graphs

To do this	Type	Gives
C20 Maximal domain → p. 9	`domain(ln(x-2)+√(5-x),x)`	2 < x ≤ 5
C21 Hybrid function → p. 10	`Define g(x)=piecewise(x+3,x<-1,x^2+1,-1≤x≤1,4-2·x,x>1)` (or the template)	use g like any function
C22 Restricted domain → p. 21	`Define r(x)=(x-2)^2-3|x≤2`, then `solve(r(x)=x,x)`	x = (5 − √21)/2 ≈ 0.209 (the other root is outside the domain)
C23 Composite → p. 21	`f(g(x))`	the rule of f∘g
C24 Inverse, choosing the branch → p. 21	`solve(x=(y-2)^2-3,y)|y≤2`	y = 2 − √(x + 3)
C25 Image under a transformation → p. 15	y = √x under (x, y) → (1 − 2x, 3y − 2): `solve((y+2)/3=√((1-x)/2),y)`	y = 3√((1 − x)/2) − 2
C26 Where f meets its inverse → p. 22	`Define f(x)=1-x^3`, then `solve(f(f(x))=x,x)`	all five x-values (`f(x)=x` finds only the one on y = x)

To do this	Type	Gives
C27 Tangency with a parameter → p. 16	`solve(e^(x)=k·x and e^(x)=k,{x,k})` (equal values and equal gradients)	x = 1 and k = e
C28 A family of graphs → p. 16	on a Graphs page: `f1(x)=x^2-2·{1,2,3}·x+3`	draws all three members
C29 Graph of f′ → p. 30	on a Graphs page: `f2(x)=derivative(f1(x),x)`	f′ drawn under f, for "which graph could be f′" questions
C30 Graphs page	menu ▸ Analyze Graph ▸ Zero / Minimum / Maximum / Intersection / Inflection; ctrl + T for a table; `f1(x)=x^2|0≤x≤3` restricts the domain	readings are approximate: confirm in Calculator
C31 Other functions	`abs(x)`, `ln(x)`, `log(x)` (base 10), `log(x,2)` (base 2), `root(x,3)`	

Calculus

The examples below use `Define f(x)=x^3-3·x^2-9·x+5` and `Define df(x)=derivative(f(x),x)` unless they say otherwise.

To do this	Type	Gives
C32 Derivative; at a point; second → p. 28	`derivative(f(x),x)`; `df(2)`; `derivative(f(x),x,2)`	$3x^2 - 6x - 9$; −9; 6x − 6
C33 Stationary points → p. 29	`zeros(df(x),x)`, then `f(ans)`	{−1, 3}, then {10, −22}: (−1, 10) and (3, −22)
C34 Where decreasing → p. 29	`solve(df(x)<0,x)`	−1 < x < 3 (write [−1, 3] for "strictly decreasing")
C35 Inflection point → p. 30	`solve(derivative(f(x),x,2)=0,x)`, then `df(1)`	x = 1; gradient −12 there
C36 Greatest rate of change on an interval → p. 30	steepest descent `fMin(df(x),x)|0≤x≤5`; steepest ascent `fMax(df(x),x)|0≤x≤5`, then `df(x)|ans` for each (periodic? add multiples of the period)	x = 1, −12; x = 5 (an endpoint), 36
C37 Tangent, normal → p. 29	`tangentLine(x^3-2·x,x,1)`, `normalLine(x^3-2·x,x,1)`	x − 2 and −x: write y = x − 2, y = −x
C38 Tangent through an outside point → p. 29	for $y = x^2$ through (1, −3): `Define w(x)=x^2`, `Define dw(x)=derivative(w(x),x)`, then `solve(-3-w(a)=dw(a)·(1-a),a)`	a = −1 or a = 3
C39 Smooth join of a hybrid (join at x = 1) → p. 27	`Define p(x)=a·x^2+b`, `Define q(x)=4·x-1`, `Define dp(x)=derivative(p(x),x)`, `Define dq(x)=derivative(q(x),x)`, then `solve(p(1)=q(1) and dp(1)=dq(1),{a,b})`	a = 2 and b = 1
C40 Max or min on an interval → p. 32	`fMax(f(x),x)|0≤x≤5` (or `fMin`), then `f(x)|ans`; periodic? add multiples of the period for every time	x = 5 (an endpoint), then 10
C41 Minimum distance to a curve → p. 31	from (3, 0) to y = √x: `fMin((x-3)^2+x,x)|x≥0`, then `√((x-3)^2+x)|ans` (minimise the squared distance)	x = 5/2, then √11/2 ≈ 1.6583
C42 Greatest difference of two models → p. 32	with u(x) = 2sin(x) + 3 and v(x) = 0.5x + 2 defined: `fMax(abs(u(x)-v(x)),x)|0≤x≤5`, then `abs(u(x)-v(x))|ans`	x = 4.965, then 3.419
C43 Antiderivative; definite integral → p. 32	`integral(f(x),x)` (add + c yourself); `integral(x^2-1,x,0,2)`	2/3 (signed: split at the x-intercepts for area)
C44 Area between curves → p. 34	`integral(x-x^2,x,0,1)` (upper − lower, split where they cross)	1/6
C45 Intersections as exact limits → p. 34	`zeros(x^2-(x+1),x)→xs`, then `integral(x+1-x^2,x,xs[1],xs[2])` (check xs[1] < xs[2])	{(1 − √5)/2, (1 + √5)/2}, then 5√5/6 ≈ 1.8634
C46 Area when curves cross (one integral) → p. 34	for f(x) = x³ − x and g(x) = 3x: `integral(abs((x^3-x)-3·x),x,-2,2)` (ctrl + enter if it stays unevaluated)	8
C47 Average value → p. 34	`1/2·integral(e^(x),x,0,2)`	$(e^2 - 1)/2$

To do this	Type	Gives
C48 Solve for an integral's limit → p. 35	`solve(integral(10·t,t,0,k)=45,k)|k>0`	k = 3
C49 Integrate a hybrid → p. 10	with the hybrid g defined by `piecewise` (above): `integral(g(x),x,-2,2)`; or split the integral at each join	31/6
C50 Limit → p. 27	`limit((x^2-9)/(x-3),x,3)`; from the left/right add `,-1` or `,1`	6
C51 Newton's method → p. 25	`Define f(x)=x^3-2·x-5`; type `2.` and press enter; then `x-f(x)/derivative(f(x),x)|x=ans` and re-run it (▲, enter, enter) for each step	2.1, 2.09457, 2.09455
C52 Left and right rectangles → p. 35	on [0, 2], width 0.5, with `Define f(x)=x^2`: `0.5·sum(seq(f(0.5·i),i,0,3))`; `0.5·sum(seq(f(0.5·i),i,1,4))`	1.75 (left); 3.75 (right)
C53 Trapezium rule → p. 35	`Define f(x)=x^2`, then `0.5/2·(f(0)+2·sum(seq(f(0.5·i),i,1,3))+f(2))`	2.75

Probability and statistics

Distributions: menu ▸ Probability ▸ Distributions. Confidence intervals: menu ▸ Statistics ▸ Confidence Intervals.

To do this	Type	Gives
C54 Independent trials, different probabilities → p. 38	`expand((0.5·x+0.5)·(0.4·x+0.6)·(0.2·x+0.8))` (one bracket per trial: p·x + (1 − p))	$0.04x^3 + 0.26x^2 + 0.46x + 0.24$: the coefficient of x^k is P(k successes)
C55 DRV mean and variance → p. 38	`sum({0,1,2,3}·{0.1,0.3,0.4,0.2})`; `sum({0,1,2,3}^2·{0.1,0.3,0.4,0.2})-1.7^2`	1.7; 0.81
C56 Binomial P(X = x); nCr; the whole distribution → p. 39	`binomPdf(10,0.3,3)`; `nCr(4,3)`; `binomPdf(10,0.3)`	0.2668; 4; {0.0282, 0.1211, 0.2335, 0.2668, 0.2001, ...}: the mode is 3
C57 Binomial P(a ≤ X ≤ b), P(X ≤ b) → p. 39	`binomCdf(10,0.3,3,10)`; `binomCdf(10,0.3,2)`	0.6172 (P(X ≥ 3)); 0.3828
C58 Conditional (binomial) → p. 39	`binomCdf(10,0.3,3,10)/binomCdf(10,0.3,1,10)`	0.6352 = P(X ≥ 3 | X ≥ 1)
C59 Smallest n (table of values) → p. 39	`seq(1-0.9^n,n,27,30)`	0.9419, 0.9477, 0.9529, 0.9576: n = 29
C60 Integer from a rounded probability → p. 40	`seq(1-0.9^n,n,13,19)`, keep the values in [0.75, 0.85) for "0.8 to 1 d.p."	0.7458, 0.7712, ..., 0.8499, 0.8649: n = 14 to 18
C61 n and p from the mean and variance → p. 40	`solve(n·p=6 and n·p·(1-p)=4.2,{n,p})`	n = 20 and p = 0.3
C62 Unknown p → p. 40	`solve((1-p)^10=0.2,p)|0<p<1` (write the formula out)	p = 0.1487
C63 CRV: k, mean, variance → p. 40	`solve(integral(k·x^2,x,0,3)=1,k)`; `integral(x·x^2/9,x,0,3)`; `integral(x^2·x^2/9,x,0,3)-(9/4)^2`	k = 1/9; 9/4; 27/80 (= 0.3375)
C64 CRV: median, percentile → p. 40	`solve(integral(t^2/9,t,0,m)=0.5,m)|0≤m≤3` (0.25 for the 25th percentile)	m = 2.381 (1.890)
C65 A parameter inside an integral, tabulated → p. 40	`seq(integral(t^2/9,t,k,3),k,0,3,0.5)` (low, high, step)	1, 0.9954, 0.9630, 0.875, 0.7037, 0.4213, 0: P(X > k) for each k
C66 pdf, then binomial (chain) → p. 40	P(at least one of 2 values exceeds k), k = 1, 1.5, ..., 3: `seq(1-(1-integral(t^2/9,t,k,3))^2,k,1,3,0.5)`	0.9986, 0.9844, 0.9122, 0.6651, 0
C67 Piecewise pdf, conditional → p. 41	`Define f(x)=piecewise(x,0≤x≤1,2-x,1<x≤2,0)` (the last 0 is "otherwise"), then `integral(f(x),x,1.5,2)/integral(f(x),x,1,2)`	0.25 = P(X > 1.5 | X > 1)
C68 Normal probability → p. 42	`normCdf(180,∞,170,8)` (lower, upper, μ, σ: not σ^2)	0.1056

To do this	Type	Gives
C69 Conditional (normal) → p. 42	`normCdf(180,∞,170,8)/normCdf(170,∞,170,8)`	0.2113
C70 Normal, then binomial (keep it unrounded) → p. 42	`normCdf(180,∞,170,8)→pr`, then `binomCdf(10,pr,2,10)`	0.2858 = P(at least 2 of 10 are over 180)
C71 Inverse normal; z-value → p. 42	`invNorm(0.9,170,8)` (area to the **left**); `invNorm(0.1)`	180.25; −1.2816
C72 Unknown σ → p. 42	`solve((40-50)/s=invNorm(0.1),s)`	s = 7.80
C73 Unknown μ and σ → p. 42	`solve((30-m)/s=invNorm(0.2) and (60-m)/s=invNorm(0.9),{m,s})`	m = 41.89, s = 14.13
C74 Sample proportion, exact → p. 43	`binomCdf(100,0.3,36,100)` for P(p̂ > 0.35), n = 100	0.1161
C75 Sample proportion, normal approximation → p. 43	`normCdf(0.35,∞,0.3,√(0.3·0.7/100))`	0.1376
C76 Simulate sample proportions → p. 43	`randBin(100,0.3,50)/100`	50 simulated values of p̂
C77 Confidence interval for p → p. 44	`zInterval_1Prop 60,200,0.95`	stat.CLower = 0.2365, stat.CUpper = 0.3635
C78 z for a C% interval → p. 44	`invNorm((1+0.9)/2)`; `invNorm(0.975)`; `invNorm(0.995)`	1.6449 (90%); 1.9600 (95%); 2.5758 (99%)
C79 Sample size for a margin of error → p. 44	`solve(invNorm(0.975)·√(0.5·0.5/n)=0.03,n)`	n = 1067.07, so n = 1068 (always round up)
C80 Confidence level from an interval → p. 44	z = margin ÷ standard error, then `normCdf(-1.645,1.645)`	0.90, a 90% interval

CAS traps that cost marks

- **Type the multiplication.** `kx` is one variable called "kx": type `k·x`. And `x(x+1)` is read as a function: type `x·(x+1)`.
- **Old definitions leak.** A new Problem for each question is safest; `ClearAZ` clears only single-letter names. Define stores the expression as it was, so redefine anything built from f after changing f. `DelVar p` (menu ▸ Actions ▸ Delete Variable) clears one name you stored.
- **nSolve, fMax and fMin return one value.** Restrict the domain to find each one, and for periodic models add the period to list every time in the domain.
- **Check every solution against the domain.** solve can return endpoint or extra roots (e.g. t = 0 where two models both start): keep only the ones the question allows.
- **Trig without a domain** gives general solutions with `n1` (any integer): add the question's domain, e.g. `|0≤x≤2·π`.
- **normCdf and invNorm take σ**, not the variance, and invNorm wants the area to the left.
- **Rounding.** If Pr = 0.2 "correct to 1 decimal place", the true value is in [0.15, 0.25). Don't solve = 0.2 exactly for an integer like n or k: tabulate the candidates with `seq` and keep every one in the interval. Store unrounded values (→) and round only at the end.
- **No rounding stated means exact.** In Section B, if no decimal places are given, the answer must be exact: enter exact inputs (fractions, π, √, not decimals) so CAS stays exact; `exact()` only turns a decimal into a simple fraction and can't recover √ or π.
- **CAS answers come in their own form.** Convert to the exact form asked, and write the setting-out a marker needs.

START HERE

Methods formula sheet & must-knows

VCAA provides a Methods formula sheet in both exams (mensuration, the derivative and antiderivative rules, and the probability and statistics formulas). Many things you use constantly aren't on it: exact values, log laws, transformations and the discriminant. Treat this page as a revision summary. Always check the current official sheet from the VCAA website before your exam.

Formulas already on the official VCAA sheet (derivative and antiderivative rules, the trapezium rule, Newton's method, probability rules, binomial, random-variable and sample-proportion formulas) are left out here: use the sheet. What's below is what the sheet doesn't give you.

Formulas

Algebra and functions

Quadratics

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \qquad \Delta = b^2 - 4ac$$

$\Delta > 0$: two roots; $= 0$: one; < 0: none. Turning point at $x = -\frac{b}{2a}$. Discriminant problems → p. 24

Polynomials

$$P(a) = 0 \iff (x - a) \text{ is a factor}$$

$$a^3 \pm b^3 = (a \pm b)(a^2 \mp ab + b^2)$$

Polynomials → p. 11

Index and log laws

$$a^m a^n = a^{m+n} \quad (a^m)^n = a^{mn}$$

$$\log(xy) = \log x + \log y \quad \log x^n = n \log x \quad \log_a x = \frac{\log_e x}{\log_e a}$$

Exponentials & logs → p. 17

Transformations

$$y = af(n(x - h)) + k$$

$$(x, y) \to \left(\tfrac{x}{n} + h,\ ay + k\right)$$

Dilation from the x-axis = vertical. Transformations → p. 14

Circular functions

$$y = a\sin(n(x - h)) + k: \text{ period } \tfrac{2\pi}{|n|},\ \text{amplitude } |a|$$

$$\tan: \text{ period } \tfrac{\pi}{|n|},\ \text{asymptotes where } n(x - h) = \tfrac{\pi}{2} + m\pi \qquad \sin^2 x + \cos^2 x = 1$$

$$\sin x = \sin\alpha: \ x = \alpha + 2m\pi \text{ or } \pi - \alpha + 2m\pi$$

$$\cos x = \cos\alpha: \ x = \pm\alpha + 2m\pi \qquad \tan x = \tan\alpha: \ x = \alpha + m\pi$$

Circular functions → p. 18

Composite and inverse

$$f \circ g \text{ exists} \iff \text{ran } g \subseteq \text{dom } f$$

$$\text{dom } f^{-1} = \text{ran } f$$

Composite & inverse → p. 21

Calculus

Tangents and normals

$$y - f(a) = f'(a)(x - a) \qquad m_{\text{normal}} = -\tfrac{1}{f'(a)}\ (f'(a) \neq 0)$$

Applications → p. 29

Integrals

$$\int_a^b f(x)\,dx = F(b) - F(a) \qquad f_{\text{avg}} = \tfrac{1}{b-a}\int_a^b f(x)\,dx$$

Area between curves: $\int(\text{upper} - \text{lower})$. Integration → p. 33

Probability and statistics

Normal

$$Z = \frac{X - \mu}{\sigma} \qquad 68\%/95\%/99.7\% \text{ within } 1/2/3\sigma$$

$$z = 1.645\ (90\%), \quad 1.960\ (95\%), \quad 2.576\ (99\%)$$

Normal → p. 42

Exact values (not on the sheet)

θ	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	undefined

FUNCTIONS, RELATIONS & GRAPHS

Functions: domain, range, hybrids & modulus

Relations and functions

A **relation** is any set of ordered pairs (x, y). A **function** is a relation where every x-value has *exactly one* y-value.

- **Vertical line test:** a graph is a function if no vertical line cuts it more than once.
- **One-to-one:** every y-value comes from exactly one x-value (passes the horizontal line test too). Examples: $y = x^3$, $y = e^x$, $y = \log_e x$.
- **Many-to-one:** some y-values come from more than one x-value. Examples: $y = x^2$, $y = \cos x$, $y = |x|$.

Function notation

$$f : D \to \mathbb{R}, \quad f(x) = \ldots$$

D is the **domain** (allowed inputs). The **range** is the set of outputs actually produced. $\mathbb{R}$ here is the codomain, not the range.

Interval notation: $[a, b]$ includes both ends, (a, b) excludes both, $[a, b)$ includes a only. $\mathbb{R} \setminus \{2\}$ means "all reals except 2"; $\mathbb{R}^+ = (0, \infty)$.

Maximal (implied) domain

If no domain is given, use the largest set of real numbers for which the rule makes sense:

Rule contains	Condition
$\sqrt{g(x)}$	$g(x) \geq 0$
$\frac{1}{g(x)}$	$g(x) \neq 0$
$\log_e(g(x))$	$g(x) > 0$ (strictly)
$\tan(x)$	$x \neq \frac{\pi}{2} + k\pi,\ k \in \mathbb{Z}$

When a rule is a sum or product of pieces, the maximal domain is the **intersection** of the pieces' domains.

Worked example 1 **Maximal domain of a combination** 2 marks

Find the maximal domain of $f(x) = \sqrt{4 - x} + \frac{1}{x + 1}$.

1 **Square root:** $4 - x \geq 0 \Rightarrow x \leq 4$. 2 **Fraction:** $x + 1 \neq 0 \Rightarrow x \neq -1$.

✓ $(-\infty, -1) \cup (-1, 4]$, which can also be written $(-\infty, 4] \setminus \{-1\}$.

Finding the range on a restricted domain

Sketch the graph over the domain only, then read off the lowest and highest y-values. Check three things: **turning points** inside the domain, the values at the **endpoints**, and whether each endpoint is included.

Worked example 2 **Range with an open endpoint (trick)** — 3 marks


Find the range of $f : [-1, 3) \to \mathbb{R},\ f(x) = x^2 - 2x$.

1 **Turning point:** $f(x) = (x-1)^2 - 1$, so the minimum is -1 at $x = 1$, which is in the domain. 2 **Endpoints:** $f(-1) = 1 + 2 = 3$ (included, since -1 is in the domain); $f(3) = 9 - 6 = 3$ ($x = 3$ is excluded). 3 **Is 3 in the range?** Yes: even though $x = 3$ is excluded, the value 3 is still produced at $x = -1$.

Range $= [-1, 3]$**.** Many students write $[-1, 3)$ because of the open endpoint. Always check whether the "missing" value is produced somewhere else.

Hybrid (piecewise) functions

A hybrid function uses different rules on different parts of the domain. Sketch each piece only over its own interval, with a closed dot where an endpoint is included and an open dot where it isn't.



$f(x) = \begin{cases} x + 3, & x < -1 \\ x^2 + 1, & -1 \le x \le 1 \\ 4 - 2x, & x > 1 \end{cases}$: the pieces meet at both joins ($f = 2$ at $x = -1$ and at $x = 1$), so f is continuous, but the corners there mean it isn't differentiable at $x = \pm 1$.

Key idea

A hybrid function is **continuous** at a join if the pieces give the same value there. Continuity problems usually become "set the two rules equal at the join and solve for the unknown constant". For differentiability (a smooth join), see Limits, continuity & differentiability → p. 27.

Worked example 3 **Making a hybrid continuous** — 2 marks

Find a so that $g(x) = \begin{cases} x^2 + a, & x \le 2 \\ 3x - 1, & x > 2 \end{cases}$ is continuous at $x = 2$.

1 Both rules must give the same value at $x = 2$: $2^2 + a = 3(2) - 1$.


$4 + a = 5 \Rightarrow a = 1$.

The modulus function

Modulus (absolute value)

$$|x| = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}$$

$|x|$ is the distance from 0. The graph of $y = |f(x)|$ is the graph of $y = f(x)$ with every part below the x-axis reflected up.



$y = x^2 - 4$ (dashed) and $y = |x^2 - 4|$ (solid). The part between $x = -2$ and $x = 2$ is flipped up, creating sharp corners at $x = \pm 2$.

Worked example 4 **Solving a modulus equation** 2 marks

Solve $|x^2 - 4| = 3$.

1 Split into two equations: $x^2 - 4 = 3$ or $x^2 - 4 = -3$. 2 $x^2 = 7 \Rightarrow x = \pm\sqrt{7}$; $x^2 = 1 \Rightarrow x = \pm 1$.

$x = \pm 1, \pm\sqrt{7}$ (four solutions: check on the graph, where the line $y = 3$ cuts the W shape four times).

Trick alert: domain and range traps

- **Logs need strictly positive arguments.** $\log_e(x^2)$ has domain $\mathbb{R} \setminus \{0\}$, but $2\log_e x$ has domain $(0, \infty)$. They are only equal for $x > 0$.
- **Open endpoints aren't automatically open in the range** (see the worked example above).
- **The basic graphs never reach their asymptotes.** The range of $y = 3 - \dfrac{2}{(x-1)^2}$ is $(-\infty, 3)$, not $(-\infty, 3]$. Combined functions can cross a horizontal asymptote, though: $y = xe^{-x}$ has asymptote $y = 0$ and equals 0 at $x = 0$, so check.

Hard exam questions

2023 Exam 2 · Section A · Question 3 © VCAA — 47% correct

The question (not reprinted in this copy): Domain of the sum of two functions with domains [−2, 3) and (−1, 5]
Full question: 2023 Exam 2, Section A, Question 3 (page 3 of the paper).

Answer: E · A 22% · B 12% · C 11% · D 7% · E 47%

The sum's domain is the intersection: $(-1, 3)$.

FUNCTIONS, RELATIONS & GRAPHS

Polynomials: factor theorem, cubics & quartics

Division, remainders and factors

Dividing a polynomial $P(x)$ by $(x - a)$ gives a quotient $Q(x)$ and a remainder R (a number):

Division statement

$$P(x) = (x - a)\,Q(x) + R$$

Substituting $x = a$ kills the first term, which gives the two theorems you use every week.

Remainder theorem

$$\text{The remainder when } P(x) \text{ is divided by } (x - a) \text{ is } P(a).$$

For a divisor $(bx - a)$ the remainder is $P\left(\frac{a}{b}\right)$.

Factor theorem

$$(x - a) \text{ is a factor of } P(x) \iff P(a) = 0.$$

Key idea

To factorise a cubic by hand: try the factors of the constant term ($\pm 1, \pm 2, \pm 3, \ldots$) until $P(a) = 0$, then divide by $(x - a)$ and factorise the quadratic that is left. With a non-monic leading coefficient, also try fractions $\pm\dfrac{\text{factor of constant}}{\text{factor of leading coeff}}$.

Sum and difference of cubes

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2) \qquad a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$


The quadratic factor never has real roots, so $x^3 - 27 = 0$ has only one real solution, $x = 3$.

Shapes of polynomial graphs

A factor tells you what the graph does at that x-intercept:

Factor	Behaviour at $x = a$
$(x - a)$	Cuts straight through the axis
$(x - a)^2$	Touches the axis and turns (a turning point on the axis)
$(x - a)^3$	Flattens out and passes through: a **stationary point of inflection**

End behaviour depends on the degree and the sign of the leading coefficient. Odd degree, positive: from bottom-left to top-right. Even degree, positive: both ends up. A negative leading coefficient flips everything.



$y = (x+2)(x-1)^2$ cuts at $x = -2$ and touches at $x = 1$. $y = (x-1)^3 - 2$ is the basic cubic translated: its stationary point of inflection is at $(1, -2)$.

Cubic in "turning point" (inflection) form

$$y = a(x-h)^3 + k$$

Stationary point of inflection at (h, k). No other turning points.

Worked example 3 **Rule from a graph's features** — 2 marks

A cubic touches the x-axis at $x = 2$, crosses it at $x = -1$ and passes through $(0, 8)$. Find its rule.

1 The touch gives a squared factor: $y = a(x+1)(x-2)^2$. 2 Substitute $(0, 8)$: $8 = a(1)(4)$, so $a = 2$.

✓ $y = 2(x+1)(x-2)^2$ (expanded: $y = 2x^3 - 6x^2 + 8$).

Worked example 4 **A cubic from its turning points (Exam 2 staple)** — 4 marks

A cubic $p(x) = ax^3 + bx^2 + cx + d$ has a local maximum at $(1, 5)$ and a local minimum at $(3, 1)$. Find a, b, c and d.

1 $p'(x) = 3ax^2 + 2bx + c$ is zero at $x = 1$ and $x = 3$, so $p'(x) = 3a(x-1)(x-3) = 3ax^2 - 12ax + 9a$. Matching coefficients: $b = -6a$, $c = 9a$. 2 $p(3) = 27a - 54a + 27a + d = d$, so $d = 1$. Then $p(1) = a - 6a + 9a + 1 = 4a + 1 = 5$, so $a = 1$.

✓ $a = 1,\ b = -6,\ c = 9,\ d = 1$, so $p(x) = x^3 - 6x^2 + 9x + 1$. On CAS: `Define p(x)=a·x^3+b·x^2+c·x+d`, `Define dp(x)=derivative(p(x),x)`, then `solve(p(1)=5 and p(3)=1 and dp(1)=0 and dp(3)=0,{a,b,c,d})`.

Worked example 5 **Biquadratic (quartic in disguise)** — 2 marks

Solve $x^4 - 5x^2 + 4 = 0$.

1 Let $u = x^2$: $u^2 - 5u + 4 = 0 \Rightarrow (u-1)(u-4) = 0$, so $u = 1$ or $u = 4$.

✓ $x = \pm 1, \pm 2$. Don't stop at u: the question asked for x.

Worked example 6 **Sketching a quartic** — 3 marks

Sketch $y = x^4 - 4x^3$, labelling the axis intercepts and stationary points.

1 $y = x^3(x-4)$: intercepts $(0, 0)$ (a triple root, so a stationary point of inflection) and $(4, 0)$ (crosses). 2 $y' = 4x^3 - 12x^2 = 4x^2(x-3) = 0$: $x = 0$ and $x = 3$, $y(3) = -27$. The gradient table is in Applications of differentiation → **p. 30**.

✓ **Positive x^4: up at both ends. Flattens through $(0, 0)$ going down, falls to a local minimum at $(3, -27)$, then rises through $(4, 0)$.**

Distinct roots when the roots depend on a parameter

When the roots are in terms of a, set each pair of roots equal to find the values of a where two of them merge into a repeated root. E.g. $f(x) = (x-1)(x-a)(x-2a)$ has roots $1, a, 2a$: pairs merge when $a = 1$, $2a = 1$ or $a = 2a$, so there are three distinct roots unless $a \in \{0, \frac{1}{2}, 1\}$, and two distinct roots (one repeated) at each of those. Check each special value: one value can merge more than one pair.

How many solutions? Polynomials with a parameter

Exam 2 loves "find the values of k for which the equation has exactly one / two / three solutions". Rewrite so the parameter sits alone, then think of it as a horizontal line $y = k$ moving up and down against a fixed graph.

Worked example 7 **Number of solutions of $x^3 - 3x + k = 0$** — 4 marks

Find the values of k for which $x^3 - 3x + k = 0$ has (a) exactly three, (b) exactly two, (c) exactly one real solution.

1 Rearrange: $-k = x^3 - 3x$. Count intersections of $y = x^3 - 3x$ with the horizontal line $y = -k$. 2 $y = x^3 - 3x$ has turning points where $3x^2 - 3 = 0$, i.e. $x = \pm 1$: a local max at $(-1, 2)$ and a local min at $(1, -2)$. 3 The line cuts three times when it is strictly between the turning points: $-2 < -k < 2$, i.e. $-2 < k < 2$. It touches a turning point (two solutions) when $-k = \pm 2$.

✓ **(a)** $-2 < k < 2$**; (b)** $k = \pm 2$**; (c)** $k < -2$ **or** $k > 2$**.** Check (b): $k = 2$ gives $x^3 - 3x + 2 = (x-1)^2(x+2)$, two distinct solutions.

Worked example 8 **How many solutions? A difference of squares (hard)** 4 marks

Let $f(x) = (x^2 - 1)^2$ and $k > 0$. Find the number of solutions of $f(x) = k^2$ for each value of k.

1 $(x^2 - 1)^2 - k^2 = 0 \Rightarrow (x^2 - 1 - k)(x^2 - 1 + k) = 0$, so $x^2 = 1 + k$ or $x^2 = 1 - k$. 2 $x^2 = 1 + k$: since $k > 0$, $1 + k > 0$, giving always two solutions $x = \pm\sqrt{1 + k}$. 3 $x^2 = 1 - k$: two solutions if $0 < k < 1$; one solution ($x = 0$) if $k = 1$; none if $k > 1$. The two equations never share a solution, because $1 + k \neq 1 - k$ for $k > 0$.

4 solutions for $0 < k < 1$; 3 solutions for $k = 1$; 2 solutions for $k > 1$. Graphically: $y = (x^2 - 1)^2$ has minimums at $(\pm 1, 0)$ and a local maximum at $(0, 1)$. The line $y = k^2$ cuts four times below the maximum, touches it (3 solutions) when $k^2 = 1$, and cuts twice above it.

Trick alert: polynomial traps

- **"Solutions" means distinct solutions.** $(x - 1)^2(x + 2) = 0$ has two solutions, not three.
- **A factor isn't always a root you can see.** If $(x - 1)$ is a factor of $x^3 + x - 2$, the other factor $x^2 + x + 2$ has a negative discriminant, so the graph has only one x-intercept.
- **Remainder with $(x + a)$:** substitute $x = -a$, not $x = a$.
- **The turning point at a double root isn't always a minimum.** For $y = (x + 1)^2(x - 3)$ (positive leading coefficient) the double root is on the left, so it's a local *maximum*.

Hard exam questions

2023 Exam 2 · Section A · Question 19 © VCAA 32% correct

The question (not reprinted in this copy): Values of k giving one positive and one negative root of a quadratic
Full question: 2023 Exam 2, Section A, Question 19 (page 9 of the paper).

Answer: D · A 27% · B 11% · C 13% · D 32% · E 16%


Opposite signs need the product of the roots to be negative: $4k^2 - \frac{9}{4} < 0$, so $-\frac{3}{4} < k < \frac{3}{4}$. A negative product already forces two real roots ($\Delta > 0$), so no separate discriminant check is needed.

FUNCTIONS, RELATIONS & GRAPHS

Power functions, hyperbolas & root functions

The families at a glance

Rule	Name	Domain	Range	Key features
x^n, n even positive	even power (x^2, x^4)	$\mathbb{R}$	$[0, \infty)$	U-shaped, symmetric in the y-axis; flatter near 0 and steeper outside $[-1, 1]$ as n grows
x^n, n odd positive	odd power (x^3, x^5)	$\mathbb{R}$	$\mathbb{R}$	Stationary point of inflection at the origin, point-symmetric
$x^{-1} = \frac{1}{x}$	hyperbola	$\mathbb{R} \setminus \{0\}$	$\mathbb{R} \setminus \{0\}$	Asymptotes $x = 0,\ y = 0$; branches in quadrants 1 and 3
$x^{-2} = \frac{1}{x^2}$	truncus	$\mathbb{R} \setminus \{0\}$	$(0, \infty)$	Both branches above the x-axis
$x^{1/2} = \sqrt{x}$	square root	$[0, \infty)$	$[0, \infty)$	Starts at the origin (an endpoint), increasing
$x^{1/3} = \sqrt[3]{x}$	cube root	$\mathbb{R}$	$\mathbb{R}$	Vertical tangent at the origin
$x^{2/3} = (\sqrt[3]{x})^2$	"seagull"	$\mathbb{R}$	$[0, \infty)$	Sharp cusp at the origin



On $(0, 1)$ the order flips: $x^{1/3} > x^{1/2} > x > x^2$. On $(1, \infty)$ it's the other way round. Every power function passes through $(1, 1)$.

Functions

Transformed forms

Hyperbola, truncus and root in transformed form

$$y = \frac{a}{x-h} + k \qquad y = \frac{a}{(x-h)^2} + k \qquad y = a\sqrt{\pm(x-h)} + k$$

For the hyperbola and truncus: asymptotes $x = h$ and $y = k$. For the root: the endpoint is (h, k).

Key idea

For every sketch, find: (1) asymptotes or endpoint, (2) the y-intercept (put $x = 0$), (3) the x-intercept(s) (put $y = 0$), (4) the domain and range. The sign of a tells you which way the branches go.

Worked example 1 **Sketching a hyperbola** 3 marks

Sketch $y = \frac{2}{x-1} + 3$, labelling asymptotes and intercepts.

1 **Asymptotes:** $x = 1$ and $y = 3$. 2 **y-intercept:** $y = \frac{2}{-1} + 3 = 1$, so $(0, 1)$. 3 **x-intercept:** $0 = \frac{2}{x-1} + 3 \Rightarrow x - 1 = -\frac{2}{3} \Rightarrow x = \frac{1}{3}$.

Since $a = 2 > 0$, the branches sit top-right and bottom-left of the asymptote intersection $(1, 3)$. **Asymptotes $x = 1$, $y = 3$; intercepts $(0, 1)$ and $(\frac{1}{3}, 0)$.** Domain $\mathbb{R} \setminus \{1\}$, range $\mathbb{R} \setminus \{3\}$.

Worked example 3 **A reflected square root** 3 marks

Sketch $y = 2\sqrt{3-x} - 4$, stating the domain, range and intercepts.

1 Rewrite as $y = 2\sqrt{-(x-3)} - 4$: reflected in the y-axis, so the graph goes *left* from the endpoint $(3, -4)$. Domain: $3 - x \geq 0 \Rightarrow x \leq 3$. 2 x-intercept: $\sqrt{3-x} = 2 \Rightarrow 3 - x = 4 \Rightarrow x = -1$. y-intercept: $2\sqrt{3} - 4 \approx -0.54$.

Domain $(-\infty, 3]$, range $[-4, \infty)$; intercepts $(-1, 0)$ and $(0, 2\sqrt{3} - 4)$.

Trick alert: power function traps

- **Cube roots of negatives exist.** $x^{1/3}$ and $x^{2/3}$ have domain $\mathbb{R}$. CAS sometimes shows only the $x \geq 0$ half. Don't trust it blindly.
- $x^{2/3} = 4$ **has two solutions:** $x^{1/3} = \pm 2$, so $x = \pm 8$.
- **Asymptotes are not in the range.** $y = \frac{a}{x-h} + k$ never equals k.
- $\sqrt{3-x}$ **goes left**, not right. Factor out the negative to see the reflection.

FUNCTIONS, RELATIONS & GRAPHS

Transformations of graphs

The three moves

Transformation	Mapping	Effect on the rule $y = f(x)$
Dilation by factor a **from the x-axis** (parallel to the y-axis)	$(x, y) \to (x, ay)$	$y = af(x)$
Dilation by factor b **from the y-axis** (parallel to the x-axis)	$(x, y) \to (bx, y)$	$y = f\left(\frac{x}{b}\right)$
Reflection in the x-axis	$(x, y) \to (x, -y)$	$y = -f(x)$
Reflection in the y-axis	$(x, y) \to (-x, y)$	$y = f(-x)$
Translation h right and k up	$(x, y) \to (x + h, y + k)$	$y = f(x - h) + k$

VCAA wording: "a translation of h units in the positive direction of the x-axis" means h right; "in the negative direction of the y-axis" means down.

"From the x-axis" means vertical

A dilation *from* the x-axis pushes points away from the x-axis, so it stretches vertically. VCAA also writes "parallel to the y-axis" for the same thing.

The general form, and the order that works

Standard form

$$y = a\,f\big(n(x-h)\big) + k$$

Read it as: dilate by $|a|$ from the x-axis (reflect in the x-axis if $a < 0$); dilate by $\frac{1}{|n|}$ from the y-axis (reflect in the y-axis if $n < 0$); *then* translate h right and k up. The mapping is $(x, y) \to \left(\frac{x}{n} + h,\ ay + k\right)$.

Key idea

Always factor out the coefficient of x first. $f(2x+6) = f(2(x+3))$ is a translation of 3 left, not 6, when the dilation is done first. With dilations first and translations last you never go wrong.

Finding the image equation: the dash method

This is the method VCAA's solutions use, and it works for any transformation (even ones you can't picture):

1. Write the mapping as $x' = \ldots$ and $y' = \ldots$.
2. Rearrange to get x and y in terms of x' and y'.
3. Substitute into the original rule, then drop the dashes.

Worked example 1 **Image of a square-root graph** 2 marks

Find the image of $y = \sqrt{x}$ under the transformation $(x, y) \to (-2x+1,\ 3y-2)$.

1 $x' = -2x+1 \Rightarrow x = \dfrac{1-x'}{2}$, and $y' = 3y-2 \Rightarrow y = \dfrac{y'+2}{3}$. 2 Substitute into $y = \sqrt{x}$: $\dfrac{y'+2}{3} = \sqrt{\dfrac{1-x'}{2}}$.

$y = 3\sqrt{\dfrac{1-x}{2}} - 2$, with domain $x \leq 1$.

Worked example 2 **Describing a sequence of transformations** 3 marks

Describe a sequence of transformations that maps $y = e^x$ onto $y = 3e^{2x-4} + 1$, and give the mapping.

1 Factor the exponent: $y = 3e^{2(x-2)} + 1$. So $a = 3,\ n = 2,\ h = 2,\ k = 1$. 2 Dilation by factor 3 from the x-axis; dilation by factor $\dfrac{1}{2}$ from the y-axis; translation 2 units right and 1 unit up.

Mapping $(x, y) \to \left(\dfrac{x}{2} + 2,\ 3y+1\right)$. Check with the dash method: $x = 2x' - 4$, $y = \dfrac{y'-1}{3}$, giving $\dfrac{y'-1}{3} = e^{2x'-4}$. ✓

Worked example 3 **Transforming a point, domain and range** 3 marks

The function f has domain $[-2, 4]$ and range $[0, 6]$, and $f(2) = 5$. For $g(x) = -f(2x) + 3$, find (a) the domain and range of g, and (b) the image of the point $(2, 5)$.

1 **Domain:** need $2x \in [-2, 4]$, so $x \in [-1, 2]$. 2 **Range:** $f \in [0, 6] \Rightarrow -f \in [-6, 0] \Rightarrow -f + 3 \in [-3, 3]$. 3 **Point:** the mapping is $(x, y) \to \left(\dfrac{x}{2},\ -y+3\right)$, so $(2, 5) \to (1, -2)$.

Domain $[-1, 2]$**, range** $[-3, 3]$**; image point** $(1, -2)$. Check: $g(1) = -f(2) + 3 = -5 + 3 = -2$. ✓

Worked example 4 **Choosing transformations to match points** 3 marks

The graph of $y = f(x)$ has turning points at $(0, 0)$ and $(2, -4)$. Find a dilation from the y-axis and a translation that map them to $(1, 3)$ and $(7, -1)$ respectively.

1 Horizontal gap: 2 before, 6 after, so dilate by factor 3 from the y-axis. Vertical gap: 4 before and after, so no vertical dilation. 2 After the dilation $(0, 0) \to (0, 0)$ and $(2, -4) \to (6, -4)$. Translating 1 right and 3 up gives $(1, 3)$ ✓ and $(7, -1)$ ✓.

Dilation by factor 3 from the y-axis, then a translation 1 unit right and 3 units up: $(x, y) \to (3x+1,\ y+3)$, so the image is $y = f\left(\dfrac{x-1}{3}\right) + 3$.

When the order of the points flips, reflect

If the image points are in the opposite left–right order, the map needs a **reflection in the y-axis**; opposite up–down order needs a **reflection in the x-axis**. E.g. $A(1, 2)$, $B(3, -2)$ to $A'(4, 2)$, $B'(0, -2)$: A was left of B and is now right of it, and the gap doubles, so reflect in the y-axis, dilate by 2 from the y-axis and translate 6 right: $(x, y) \to (6 - 2x,\ y)$, image $y = f\left(\dfrac{6-x}{2}\right)$.

Worked example 5 **The inverse transformation: mapping g back to f** 2 marks

The graph of $y = g(x)$, where $g(x) = 2f(3x-6) + 1$, is the image of $y = f(x)$. Find the transformation that maps g back to f.

1 $f \to g$: $y' = 2y + 1$ and $3x' - 6 = x$, so $(x, y) \to \left(\dfrac{x+6}{3},\ 2y+1\right)$. 2 Reverse: $x = 3x' - 6$, $y = \dfrac{y'-1}{2}$, so $g \to f$ is $(x, y) \to \left(3x - 6,\ \dfrac{y-1}{2}\right)$.

Dilate by factor 3 from the y-axis, then translate 6 units left; translate 1 unit down, then dilate by factor $\dfrac{1}{2}$ from the x-axis. Check: $f(x) = \dfrac{g\left(\frac{x+6}{3}\right) - 1}{2}$.

Functions

Families of functions with a parameter

A rule with a parameter, such as $y = x^2 - 2kx + 3$, describes a whole family of graphs. Typical questions ask where a feature (a turning point, an intercept) lies as k varies, or which values of k give a property (one-to-one, no x-intercepts, a given number of solutions).

Worked example 6 **Where the turning points go** 3 marks

For $y = x^2 - 2kx + 3$, find the turning point in terms of k and the curve it lies on as k varies. For which k are there no x-intercepts?

1 Turning point $(k,\ 3 - k^2)$. 2 $x = k$ and $y = 3 - k^2$, so $y = 3 - x^2$.

Turning point $(k, 3 - k^2)$, on the parabola $y = 3 - x^2$. No x-intercepts when the turning point is above the axis: $3 - k^2 > 0$, so $-\sqrt{3} < k < \sqrt{3}$ (the same as $\Delta = 4k^2 - 12 < 0$).

Worked example 7 **Parameter conditions: increasing, common points, stationary points** 4 marks

(a) For which k is $f(x) = x^3 + kx^2 + 3x$ strictly increasing for all x? (b) Find the points that every member of $y = kx^2 - (k + 2)x + 2$ passes through. (c) How many stationary points does $g(x) = x^3 - 3kx$ have, as k varies?

1 (a) $f'(x) = 3x^2 + 2kx + 3$; $\Delta = 4k^2 - 36 \leq 0 \Rightarrow -3 \leq k \leq 3$. 2 (b) $y = k(x^2 - x) + 2 - 2x$. Independent of k when $x^2 - x = 0$: $x = 0, y = 2$ and $x = 1, y = 0$. 3 (c) $g'(x) = 3x^2 - 3k = 0 \Rightarrow x^2 = k$.

(a) $-3 \leq k \leq 3$ (the ends are included: f' touches 0 once). (b) $(0, 2)$ and $(1, 0)$. (c) Two if $k > 0$, one (a stationary point of inflection at the origin) if $k = 0$, none if $k < 0$.

Worked example 8 **When does f meet its inverse? (tangency)** 3 marks

For which values of k does the graph of $f(x) = e^x + k$ intersect the graph of its inverse?

1 $e^x = 1 \Rightarrow x = 0$; then $e^0 + k = 0 \Rightarrow k = -1$. CAS: `solve(e^(x)+k=x and e^(x)=1,{x,k})`. 2 Lowering the curve (smaller k) makes it cross $y = x$ twice; raising it lifts it clear of the line.

$k \leq -1$ (one meeting point at $k = -1$, two for $k < -1$). The same "touch $y = x$" idea finds the smallest or largest parameter in Section B questions on inverses.

Trick alert: transformation traps

- **Horizontal changes act "backwards".** $f(x - 3)$ moves right 3; $f(2x)$ squashes by factor $\frac{1}{2}$ (dilation factor $\frac{1}{2}$ from the y-axis, not 2).
- **Order matters when dilations and translations mix.** "Translate 1 right, then dilate by 2 from the y-axis" gives $f\left(\frac{x}{2} - 1\right)$, not $f\left(\frac{x - 1}{2}\right)$.
- **Sequences aren't unique.** $y = (2x)^2 = 4x^2$: a dilation by $\frac{1}{2}$ from the y-axis gives the same graph as a dilation by 4 from the x-axis. Similarly $e^{x-2} = e^{-2}e^x$.
- **Transform the asymptotes too.** The image of $y = \log_e x$ under a translation 3 left has asymptote $x = -3$.

Hard exam questions

2025 Exam 2 · Section A · Question 20 © VCAA **36% correct**

The question (not reprinted in this copy): Which transformation sequence does not map a^x to a^{2x+2}
Full question: 2025 Exam 2, Section A, Question 20 (page 12 of the paper).

Answer: C · A 24% · B 18% · C 36% · D 21%

Apply each sequence in order. The sequence "dilate by a from the x-axis, dilate by ½ from the y-axis, then translate 1 right" gives a^{2x-1}.

FUNCTIONS, RELATIONS & GRAPHS

Exponential & logarithmic functions

Laws you must know cold

Index laws

$$a^m a^n = a^{m+n} \qquad \frac{a^m}{a^n} = a^{m-n} \qquad (a^m)^n = a^{mn}$$

$$a^0 = 1 \qquad a^{-n} = \frac{1}{a^n}$$

Log laws ($a, x, y > 0,\ a \neq 1$)

$$\log_a(xy) = \log_a x + \log_a y \qquad \log_a \frac{x}{y} = \log_a x - \log_a y$$


$$\log_a x^n = n \log_a x$$

Definition and inverse pair

$$y = a^x \iff x = \log_a y \qquad e^{\log_e x} = x\ (x > 0) \qquad \log_e(e^x) = x$$

$\log_e$ is written $\ln$ on most calculators. Change of base: $\log_a x = \dfrac{\log_e x}{\log_e a}$.

Graphs



$y = e^x$ and $y = \log_e x$ are inverses: each is the reflection of the other in $y = x$. e^x has horizontal asymptote $y = 0$ and passes through $(0, 1)$; $\log_e x$ has vertical asymptote $x = 0$ and passes through $(1, 0)$. A log graph has **no horizontal asymptote**: it keeps rising (slowly) without bound.

Worked example 1 **Sketching a transformed log graph** — 3 marks

For $y = 2\log_e(3 - x)$, state the maximal domain and the asymptote, and find the axis intercepts.

1 Need $3 - x > 0$: domain $(-\infty, 3)$, asymptote $x = 3$. It's reflected in the y-axis, so the graph rises to the *left*. 2 x-intercept: $\log_e(3 - x) = 0 \Rightarrow 3 - x = 1 \Rightarrow x = 2$.

✓ **Domain $(-\infty, 3)$, asymptote $x = 3$, intercepts $(2, 0)$ and $(0, 2\log_e 3)$.**

Solving equations

Worked example 2 **Hidden quadratic in eˣ (tech-free classic)** — 3 marks

Solve $e^{2x} - 5e^x + 6 = 0$.

1 Let $u = e^x$, so $e^{2x} = u^2$: $u^2 - 5u + 6 = 0 \Rightarrow (u - 2)(u - 3) = 0$. 2 $e^x = 2$ or $e^x = 3$.

✓ $x = \log_e 2$ **or** $x = \log_e 3$. (If a factor gave $e^x = -1$, you'd reject it: $e^x > 0$.)

Worked example 3 **Log equation with a rejected solution** — 3 marks

Solve $\log_2 x + \log_2(x - 2) = 3$.

1 Combine: $\log_2\big(x(x - 2)\big) = 3 \Rightarrow x^2 - 2x = 2^3 = 8$. 2 $x^2 - 2x - 8 = 0 \Rightarrow (x - 4)(x + 2) = 0$, so $x = 4$ or $x = -2$.

✓ $x = 4$ **only.** $x = -2$ makes $\log_2 x$ undefined in the original equation. Checking the original equation is worth a mark.

Growth and decay models

Exponential model

$$A = A_0 e^{kt}$$

$k > 0$: growth; $k < 0$: decay. Half-life (decay) $= \dfrac{\log_e 2}{|k|}$.

Functions

Worked example 6 **Newton's law of cooling (Exam 2 style)** 4 marks

A cup of coffee cools according to $T = 20 + 70e^{-kt}$ (°C, t in minutes). After 10 minutes it is 55 °C. (a) Find k exactly. (b) How long until it reaches 30 °C?

1 (a) $55 = 20 + 70e^{-10k} \Rightarrow e^{-10k} = \frac{1}{2} \Rightarrow k = \frac{\log_e 2}{10}$. 2 (b) $30 = 20 + 70e^{-kt} \Rightarrow e^{-kt} = \frac{1}{7} \Rightarrow t = \frac{\log_e 7}{k} = \frac{10\log_e 7}{\log_e 2}$.

(a) $k = \frac{\log_e 2}{10} \approx 0.0693$**; (b)** $t \approx 28.1$ **minutes.** The long-run temperature is 20 °C (room temperature), the horizontal asymptote.

Worked example 7 **Fitting y = Ae^kx through two points** 3 marks

The curve $y = Ae^{kx}$ passes through $(2, 12)$ and $(5, 96)$. Find A and k exactly.

1 $12 = Ae^{2k}$ and $96 = Ae^{5k}$. Divide: $e^{3k} = 8$, so $k = \frac{1}{3}\log_e 8 = \log_e 2$. 2 $A = \frac{12}{e^{2\log_e 2}} = \frac{12}{4} = 3$.

$A = 3$, $k = \log_e 2$, so $y = 3e^{x\log_e 2} = 3 \times 2^x$.

Changing the base

$a^x = e^{x\log_e a}$, and $e^{kx} = (e^k)^x$. To write e^{kx} as 2^{cx}, match exponents: $kx = cx\log_e 2$, so $c = \frac{k}{\log_e 2}$.

Trick alert: exp and log traps

- $e^x = 0$ **has no solution.** In $e^{2x} = 3e^x$, divide by e^x (it's never 0): $x = \log_e 3$ only.
- $\log_e(x^2) = 2$ **has two solutions** ($x = \pm e$), but $2\log_e x = 2$ has one ($x = e$). Log laws quietly change domains.
- $\log_e(a + b) \neq \log_e a + \log_e b$**.** The product law is for products only.
- $\log_e(2x) = \log_e 2 + \log_e x$: a horizontal dilation of a log graph is the same as a vertical translation.

FUNCTIONS, RELATIONS & GRAPHS

Circular (trigonometric) functions

Radians, exact values and symmetry

π radians $= 180°$. On the unit circle, the point at angle θ is $(\cos\theta, \sin\theta)$, and $\tan\theta = \frac{\sin\theta}{\cos\theta}$.

θ	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$\sin\theta$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
$\cos\theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\tan\theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undefined

Symmetry (quadrants 2, 3, 4)

$$\sin(\pi - \theta) = \sin\theta \quad \sin(\pi + \theta) = -\sin\theta \quad \sin(2\pi - \theta) = -\sin\theta$$
$$\cos(\pi - \theta) = -\cos\theta \quad \cos(\pi + \theta) = -\cos\theta \quad \cos(2\pi - \theta) = \cos\theta$$
$$\sin\left(\tfrac{\pi}{2} - \theta\right) = \cos\theta \qquad \sin^2\theta + \cos^2\theta = 1$$

Memory aid for the positive ratio in each quadrant, going anticlockwise from quadrant 1: **A**ll, **S**in, **T**an, **C**os.

Graphs

$m \in \mathbb{Z}$	$y = \sin x$	$y = \cos x$	$y = \tan x$
Period	2π	2π	π
Range	$[-1, 1]$	$[-1, 1]$	$\mathbb{R}$
Zeros	$m\pi$	$\frac{\pi}{2} + m\pi$	$m\pi$
Maximum 1 at	$\frac{\pi}{2} + 2m\pi$	$2m\pi$	none
Minimum −1 at	$\frac{3\pi}{2} + 2m\pi$	$\pi + 2m\pi$	none
Vertical asymptotes	none	none	$x = \frac{\pi}{2} + m\pi$
Points of inflection	at the zeros	at the zeros	at the zeros (gradient 1)

Transformed tan

$$y = a\tan\left(n(x - h)\right) + k: \quad \text{asymptotes } x = h + \frac{\pi}{2n} + \frac{m\pi}{n}, \quad \text{inflections } \left(h + \frac{m\pi}{n},\ k\right), \quad m \in \mathbb{Z}$$

Period $\frac{\pi}{|n|}$. E.g. $y = 3\tan(2x) + 1$: asymptotes $x = \frac{\pi}{4} + \frac{m\pi}{2}$, inflections at $\left(\frac{m\pi}{2}, 1\right)$, period $\frac{\pi}{2}$.

Transformed sin, cos and tan

$$y = a\sin\big(n(x-h)\big) + k$$


Amplitude $|a|$, period $\dfrac{2\pi}{|n|}$, horizontal shift h, centre line $y = k$, range $[k-|a|,\ k+|a|]$. For $y = a\tan(n(x-h)) + k$: period $\dfrac{\pi}{|n|}$, with vertical asymptotes wherever $n(x-h) = \dfrac{\pi}{2} + m\pi$ and points of inflection at $x = h + \dfrac{m\pi}{n}$ on $y = k$, halfway between the asymptotes. The period of a sum of trig terms is the lowest common multiple of their periods: $\sin\left(\dfrac{\pi t}{6}\right) + \cos\left(\dfrac{\pi t}{3}\right)$ has periods 12 and 6, so its period is 12.

Worked example 1 **Sketch over a restricted domain** 4 marks

Sketch $y = 2\sin\left(2x - \dfrac{\pi}{3}\right) + 1$ for $x \in [0, \pi]$, labelling endpoints and x-intercepts with exact coordinates.

1 Factor: $y = 2\sin\left(2\left(x - \dfrac{\pi}{6}\right)\right) + 1$. Amplitude 2, period π, shifted $\dfrac{\pi}{6}$ right, centre $y = 1$, range $[-1, 3]$. 2 Endpoints: $y(0) = 2\sin\left(-\dfrac{\pi}{3}\right) + 1 = 1 - \sqrt{3}$; $y(\pi) = 2\sin\left(\dfrac{5\pi}{3}\right) + 1 = 1 - \sqrt{3}$. 3 x-intercepts: $\sin\left(2x - \dfrac{\pi}{3}\right) = -\dfrac{1}{2}$. With $2x - \dfrac{\pi}{3} \in \left[-\dfrac{\pi}{3}, \dfrac{5\pi}{3}\right]$: $2x - \dfrac{\pi}{3} = -\dfrac{\pi}{6}$ or $\dfrac{7\pi}{6}$.

Endpoints $(0, 1-\sqrt{3})$ **and** $(\pi, 1-\sqrt{3})$**; x-intercepts** $\left(\dfrac{\pi}{12}, 0\right)$ **and** $\left(\dfrac{3\pi}{4}, 0\right)$; maximum $\left(\dfrac{5\pi}{12}, 3\right)$, minimum $\left(\dfrac{11\pi}{12}, -1\right)$.



$y = 2\sin\left(2x - \dfrac{\pi}{3}\right) + 1$ on $[0, \pi]$. Always find where the argument $2x - \dfrac{\pi}{3}$ starts and ends. That's the step that finds all the solutions.

Solving trig equations

1. Isolate the trig function: $\sin(\ldots) = c$.
2. **Adjust the domain** to match the argument (e.g. $x \in [0, 2\pi] \Rightarrow 2x \in [0, 4\pi]$).
3. Find the reference angle from $|c|$, then use the quadrants for the sign.
4. List every solution in the adjusted domain, then solve for x.

Worked example 3 **sin = cos, and general solutions** 3 marks

(a) Solve $\sin(2x) = \cos(2x)$ for $x \in [0, \pi]$. (b) Find the general solution of $\sqrt{3}\sin(2x) = \cos(2x)$.

1 (a) Divide by $\cos(2x)$: $\tan(2x) = 1$. $2x \in [0, 2\pi]$: $2x = \dfrac{\pi}{4}, \dfrac{5\pi}{4}$, so $x = \dfrac{\pi}{8}, \dfrac{5\pi}{8}$. 2 (b) $\tan(2x) = \dfrac{1}{\sqrt{3}}$, so $2x = \dfrac{\pi}{6} + n\pi$, $n \in \mathbb{Z}$.

(a) $x = \dfrac{\pi}{8}, \dfrac{5\pi}{8}$**; (b)** $x = \dfrac{\pi}{12} + \dfrac{n\pi}{2}$, $n \in \mathbb{Z}$.

General solutions ($n \in \mathbb{Z}$)

$$\sin x = a:\ x = 2n\pi + \sin^{-1} a \ \text{ or } \ x = (2n+1)\pi - \sin^{-1} a$$

$$\cos x = a:\ x = 2n\pi \pm \cos^{-1} a \qquad \tan x = a:\ x = n\pi + \tan^{-1} a$$

Worked example 4 **A hidden quadratic in sin x** 3 marks

Solve $2\sin^2 x - \sin x - 1 = 0$ for $x \in [0, 2\pi]$.

1 $2a^2 - a - 1 = (2a+1)(a-1) = 0 \Rightarrow \sin x = -\dfrac{1}{2}$ or $\sin x = 1$. 2 $\sin x = 1$: $x = \dfrac{\pi}{2}$. $\sin x = -\dfrac{1}{2}$: third and fourth quadrants, $x = \dfrac{7\pi}{6}, \dfrac{11\pi}{6}$.

$x = \dfrac{\pi}{2}, \dfrac{7\pi}{6}, \dfrac{11\pi}{6}$. The same move works for $e^{2x} - 3e^x + 2 = 0$ ($a = e^x > 0$) and for $\cos^2 x$ or $\tan x$ quadratics.

Inequalities in a hidden quadratic

Solve the quadratic inequality in a first, then intersect with the range of a. E.g. $2\sin^2 x + \sin x - 1 > 0$: $(2a-1)(a+1) > 0$ gives $a < -1$ or $a > \dfrac{1}{2}$. Since $-1 \le \sin x \le 1$, only $\sin x > \dfrac{1}{2}$ is possible, so on $[0, 2\pi]$: $\dfrac{\pi}{6} < x < \dfrac{5\pi}{6}$. CAS: `solve(2·a^2+a-1>0,a)` gives a < −1 or a > 1/2.

Functions

Showing there are no solutions: use the range

To show an equation has no solutions, show the two sides can never be equal. E.g. $\cos x = x^2 + 1.5$ has none, because $\cos x \le 1 < 1.5 \le x^2 + 1.5$. Similarly, $\sin(\sin 2x) = 1$ has none: $|\sin 2x| \le 1 < \frac{\pi}{2}$, and $\sin\theta = 1$ needs $\theta = \frac{\pi}{2} + 2n\pi$.

Periodic models

Worked example 5 **Tide model (Exam 2 style)** — 4 marks

The depth of water in a harbour is $D(t) = 3 + 2\cos\left(\frac{\pi t}{6}\right)$ metres, t hours after midnight. (a) State the maximum and minimum depths and the period. (b) For how long in the first 12 hours is the depth at least 4 m?

1 (a) Max $3 + 2 = 5$ m, min $3 - 2 = 1$ m, period $\frac{2\pi}{\pi/6} = 12$ hours. 2 (b) $\cos\left(\frac{\pi t}{6}\right) \ge \frac{1}{2}$ with $\frac{\pi t}{6} \in [0, 2\pi]$: $\frac{\pi t}{6} \in \left[0, \frac{\pi}{3}\right] \cup \left[\frac{5\pi}{3}, 2\pi\right]$. 3 So $t \in [0, 2] \cup [10, 12]$.

(a) 5 m, 1 m, 12 h. (b) 4 hours in total (midnight to 2 am and 10 am to noon).

Trend plus a periodic part

If $f(t) = mt + g(t)$ where g has period P, then $f(t + P) = f(t) + mP$: each period the graph repeats, shifted up by mP. E.g. $f(t) = 5t + 20\cos\left(\frac{\pi t}{4}\right)$ has period-8 wiggles and rises 40 every 8 units. To sketch, draw one period and repeat it, stepping up by mP each time; label the endpoints.

Worked example 6 **Fitting a sinusoidal model to its features** — 3 marks

The temperature in a town varies sinusoidally over a day, from a minimum of 8 °C at 3 am ($t = 3$) to a maximum of 18 °C at 3 pm ($t = 15$). Find a rule $T(t) = k - a\cos\big(n(t - h)\big)$, $0 \le t \le 24$.

1 $k = \frac{18 + 8}{2} = 13$, $a = \frac{18 - 8}{2} = 5$. 2 Min to max takes 12 h, so the period is 24 h and $n = \frac{2\pi}{24} = \frac{\pi}{12}$. The minimum is at $t = 3$, so $h = 3$.

$T(t) = 13 - 5\cos\left(\frac{\pi}{12}(t - 3)\right)$. Check: $T(15) = 13 - 5\cos\pi = 18$ ✓. The same curve is $13 + 5\sin\left(\frac{\pi}{12}(t - 9)\right)$: VCAA accepts any correct form.

Worked example 7 **Sum of the solutions, by symmetry** — 2 marks

Find the sum of the solutions of $\sin x = 0.3$ for $x \in [0, 4\pi]$.

1 Two cycles, so four solutions: one pair sums to $2 \times \frac{\pi}{2} = \pi$, the other to $2 \times \frac{5\pi}{2} = 5\pi$.

6π. CAS check: `sum(zeros(sin(x)-0.3,x)|0≤x≤4·π)` gives 18.8496 ≈ 6π.

Trick alert: trig traps

- **Adjust the domain for the argument.** Forgetting this is the number-one way students lose solutions.
- $\sin^2 x = \frac{1}{4}$ **means** $\sin x = \pm\frac{1}{2}$: four solutions in $[0, 2\pi]$, not two.
- **Endpoints count.** $\sin x = 0$ on $[0, 2\pi]$ has three solutions: $0, \pi, 2\pi$.
- **Calculator in radians!** Degree mode is an instant zero in Methods.
- **Tan's period is** $\frac{\pi}{|n|}$, **not** $\frac{2\pi}{|n|}$.

Hard exam questions

2023 Exam 2 · Section A · Question 18 © VCAA — 29% correct

The question (not reprinted in this copy): Number of local minima of $\sin(ax)$ on $[-a\pi, a\pi]$
Full question: 2023 Exam 2, Section A, Question 18 (page 9 of the paper).

Answer: E · A 12% · B 9% · C 22% · D 26% · E 29%

Interval length $2a\pi$ ÷ period $\frac{2\pi}{a}$ = a^2 periods, each with one minimum.

Functions

FUNCTIONS, RELATIONS & GRAPHS

Composite, inverse & combined functions

Combining functions

For $f+g$, $f-g$ and $f\times g$, the domain is the **intersection** of the two domains: $\text{dom}(f+g)=\text{dom}f\cap\text{dom}g$. To sketch $y=f(x)+g(x)$ by **addition of ordinates**, add the heights of the two graphs at key x-values (intercepts, asymptotes, where one graph is zero).

Composite functions

Composite

$$(f\circ g)(x)=f\big(g(x)\big)$$

$f\circ g$ exists if and only if $\text{ran}\,g\subseteq\text{dom}\,f$. Then $\text{dom}(f\circ g)=\text{dom}\,g$.

Key idea

Think of g as the first machine and f as the second. Every output of g must be an acceptable input for f. If it isn't, restrict the domain of g until it is.

Worked example 1 **Does the composite exist?** 4 marks

Let $f(x)=\sqrt{x}$ (maximal domain) and $g:\mathbb{R}\to\mathbb{R},\ g(x)=x^2-4$. (a) Explain why $f\circ g$ does not exist. (b) Find the maximal domain of g for which $f\circ g$ exists. (c) Does $g\circ f$ exist? If so, state its rule and domain.

1 (a) $\text{ran}\,g=[-4,\infty)$ but $\text{dom}\,f=[0,\infty)$. Since $[-4,\infty)\not\subseteq[0,\infty)$, $f\circ g$ does not exist. 2 (b) Need $g(x)\geq 0$: $x^2\geq 4\Rightarrow x\leq -2$ or $x\geq 2$. 3 (c) $\text{ran}\,f=[0,\infty)\subseteq\mathbb{R}=\text{dom}\,g$, so yes. $g(f(x))=(\sqrt{x})^2-4=x-4$, with domain $=\text{dom}\,f=[0,\infty)$.

(b) $(-\infty,-2]\cup[2,\infty)$**. (c)** $g\circ f(x)=x-4$**, domain** $[0,\infty)$**.** It's not the whole line $y=x-4$: it's only the part with $x\geq 0$.

Worked example 2 **Range of a composite** 2 marks

Let $f:[-1,2]\to\mathbb{R},\ f(x)=2-x^2$ and $g(x)=e^x$. Find the range of $g\circ f$.

1 f has its maximum 2 at $x=0$ (inside the domain) and its minimum at an endpoint: $f(-1)=1$, $f(2)=-2$. So $\text{ran}\,f=[-2,2]$. 2 $g(x)=e^x$ is increasing, so it maps $[-2,2]$ onto $[e^{-2},e^2]$.

$\text{ran}(g\circ f)=[e^{-2},e^2]$. Using only the endpoints of the domain gives $g(f(-1))=e$ and $g(f(2))=e^{-2}$, which misses the maximum e^2 from $x=0$. If the outer function is **decreasing**, the ends swap: e^{-x} maps $[-4,\infty)$ onto $(0,e^4]$, and an end that's never reached stays open.

A linear inside is a transformation

If $f(x)=mx+c$, then $g(f(x))=g(mx+c)$ is just a transformation of g: a dilation by factor $\dfrac{1}{|m|}$ from the y-axis (a reflection in the y-axis too if $m<0$) and a horizontal translation. For example, $y=g(c-x)=g\big(-(x-c)\big)$ is g reflected in the y-axis, then moved c units right.

Inverse functions

Only **one-to-one** functions have inverse functions. To find f^{-1}: swap x and y, make y the subject, and pick the correct branch using the range of f^{-1}. If f is continuous and f' takes both signs on an interval in the domain, f turns around, so it isn't one-to-one and has no inverse function (until the domain is restricted). The inverse of a hyperbola $y=\dfrac{a}{x-h}+k$ is another hyperbola with the asymptotes swapped: $y=\dfrac{1}{x-3}-2$ (asymptotes $x=3$, $y=-2$) has inverse $y=\dfrac{1}{x+2}+3$ (asymptotes $x=-2$, $y=3$).

Inverse facts

$$\text{dom}\,f^{-1}=\text{ran}\,f\qquad \text{ran}\,f^{-1}=\text{dom}\,f\qquad f\big(f^{-1}(x)\big)=x$$

The graph of f^{-1} is the reflection of f in the line $y=x$.

Worked example 3 **Restrict, then invert** 4 marks

Let $f:(-\infty,a]\to\mathbb{R},\ f(x)=(x-2)^2-3$. (a) Find the largest value of a for which f^{-1} exists. (b) With this a, find f^{-1} and its domain.

1 (a) The turning point is at $x=2$. To the left of it, f is one-to-one (decreasing). So $a=2$. 2 (b) Swap: $x=(y-2)^2-3\Rightarrow(y-2)^2=x+3\Rightarrow y-2=\pm\sqrt{x+3}$. 3 The range of f^{-1} is the domain of f: $y\leq 2$. So take the **negative** root.

$a=2$**;** $f^{-1}(x)=2-\sqrt{x+3}$**, domain** $[-3,\infty)$ (the range of f).

Where does a function meet its inverse?


If f is **increasing**, every intersection of $y = f(x)$ and $y = f^{-1}(x)$ lies on $y = x$, so you can solve $f(x) = x$ instead. That's much easier.

Worked example 4 **The decreasing-function trap** 3 marks

Find all points of intersection of $f(x) = 1 - x^3$ and $y = f^{-1}(x)$.

1 Solving $f(x) = x$: $1 - x^3 = x$, which has one real solution $x \approx 0.682$. That's one point, $(0.682, 0.682)$. 2 But f is *decreasing*, so there can be intersections off the line $y = x$. Look for pairs with $f(a) = b$ and $f(b) = a$: $f(0) = 1$ and $f(1) = 0$ is one pair. Solving $f(f(x)) = x$ on CAS finds another: $f(-0.536) \approx 1.154$ and $f(1.154) \approx -0.536$.

Five points: $(0.682, 0.682)$, $(0, 1)$, $(1, 0)$, $(-0.536, 1.154)$ **and** $(1.154, -0.536)$ (to 3 d.p.). Here $f^{-1}(x) = \sqrt[3]{1 - x}$, and a CAS solve of $f(x) = f^{-1}(x)$ confirms all five. Spotting one easy pair isn't enough: solve $f(f(x)) = x$ to be sure you have them all.



$y = 1 - x^3$ and its inverse $y = \sqrt[3]{1 - x}$ meet at five points. Only one is on $y = x$; the other four are two pairs swapped across it.

Worked example 5 **Where two inverses meet** 2 marks

$f(x) = x^3$ and $g(x) = 4x$ meet at $(-2, -8)$, $(0, 0)$ and $(2, 8)$. Where do f^{-1} and g^{-1} meet?

1 Swap each point: $(-8, -2)$, $(0, 0)$, $(8, 2)$. Check: $f^{-1}(x) = \sqrt[3]{x}$ and $g^{-1}(x) = \dfrac{x}{4}$ give $\sqrt[3]{8} = 2 = \dfrac{8}{4}$ ✓.

$(-8, -2)$, $(0, 0)$ **and** $(8, 2)$.

Worked example 6 **Integral of an inverse: the rectangle trick** 2 marks

Use $\displaystyle\int_0^1 e^x \, dx$ to find $\displaystyle\int_1^e \log_e y \, dy$.

1 $f(x) = e^x$, $a = 0$, $b = 1$, $c = 1$, $d = e$: $\displaystyle\int_1^e \log_e y \, dy = 1 \times e - 0 \times 1 - (e - 1)$.

1. Check: $\left[y \log_e y - y\right]_1^e = 0 - (-1) = 1$ ✓. Sketch it: the trick is just areas of rectangles.

Area between f and its inverse

If f and f^{-1} meet only on $y = x$, at $x = a$ and $x = b$, the area between them is twice the area between f and $y = x$: $2\displaystyle\int_a^b |x - f(x)| \, dx$.

Trick alert: composite and inverse traps

- $f^{-1}(x) \neq \dfrac{1}{f(x)}$. The inverse undoes f; the reciprocal is something else.
- **The domain of a composite comes from the inner function**, even if the simplified rule looks as if it's defined everywhere.
- **Check that the range fits inside the domain, value by value.** For $f(x) = \dfrac{1}{x - 1}$ and $g(x) = x^2 + 1$, $\operatorname{ran} g = [1, \infty)$ includes 1, which isn't in $\operatorname{dom} f$. So $f \circ g$ doesn't exist on $\mathbb{R}$.
- **Choose the right branch** when you take a square root, using the range of the inverse.

Hard exam questions

2023 Exam 2 · Section A · Question 20 © VCAA — 30% correct

The question (not reprinted in this copy): Largest interval where both $f \circ g$ and $g \circ f$ exist (log and sine)
Full question: 2023 Exam 2, Section A, Question 20 (page 9 of the paper).

Answer: A · A 30% · B 19% · C 26% · D 14% · E 9%

$g \circ f$ needs $x > -\frac{1}{\sqrt{2}}$. $f \circ g$ needs $\sin x > -\frac{1}{\sqrt{2}}$, i.e. $x < \frac{5\pi}{4}$ here. Combined: $\left(-\frac{1}{\sqrt{2}}, \frac{5\pi}{4}\right)$.

FUNCTIONS, RELATIONS & GRAPHS

Functional equations, odd & even functions

What's a functional equation?

A functional equation describes a *property* a function has for all inputs, such as $f(x+y) = f(x)f(y)$. VCAA usually asks "which function satisfies…" in multiple choice, or "show that f satisfies…" in short answer.

Property (for all x, y in the domain)	Satisfied by	Why
$f(x+y) = f(x) + f(y)$	$f(x) = kx$	$k(x+y) = kx + ky$
$f(x+y) = f(x)\,f(y)$	$f(x) = e^{kx}$ (or a^x)	index law $a^{x+y} = a^x a^y$
$f(xy) = f(x) + f(y)$	$f(x) = k\log_e x$	log law $\log(xy) = \log x + \log y$
$f(xy) = f(x)\,f(y)$	$f(x) = x^n$ (also $\lvert x\rvert^n$)	$(xy)^n = x^n y^n$
$f(2x) = 2f(x)$	$f(x) = kx$, and also $k\lvert x\rvert$ or anything linear on each side of 0	$k(2x) = 2(kx)$; $\lvert 2x\rvert = 2\lvert x\rvert$
$f(-x) = f(x)$	**even**: $x^2, \lvert x\rvert, \cos x, e^x + e^{-x}$	symmetric in the y-axis
$f(-x) = -f(x)$	**odd**: $x^3, \sin x, \tan x, \frac{1}{x}$	point symmetry about the origin
$f(x+p) = f(x)$	**periodic** with period p: $\sin x, \cos x$ ($p = 2\pi$), $\tan x$ ($p = \pi$)	the graph repeats every p units
$f(f(x)) = x$	self-inverse: $\frac{1}{x}$, $c - x$, some hyperbolas	graph symmetric in $y = x$

The number-test strategy

In multiple choice, don't do algebra first. Pick easy values ($x = 1$, $y = 2$, or $x = y = 0$) and test each option. Usually three options fail on the first test. Then prove the survivor in your head.

Worked example 1 **Which function satisfies f(x + y) = f(x)f(y)?** — 1 mark

Which one of $f(x) = 3x$, $f(x) = x^3$, $f(x) = e^{3x}$, $f(x) = \log_e(3x)$ satisfies $f(x+y) = f(x)f(y)$ for all real x, y?

1 Test $x = y = 0$: we need $f(0) = f(0)^2$, so $f(0) = 0$ or 1. $\log_e(0)$ is undefined, so D is out. 2 Test $x = y = 1$: $3x$ gives 6 vs 9 ×; x^3 gives 8 vs 1 ×; e^{3x} gives e^6 vs $e^3 \cdot e^3 = e^6$ ✓.

✓ $f(x) = e^{3x}$: $e^{3(x+y)} = e^{3x}e^{3y}$ for all x, y.

Worked example 3 **Showing a symmetry property** — 2 marks

Let $f(x) = x^2 - 2x$. Show that $f(1+h) = f(1-h)$ for all h, and interpret this.

1 $f(1+h) = (1+h)^2 - 2(1+h) = 1 + 2h + h^2 - 2 - 2h = h^2 - 1$. 2 $f(1-h) = (1-h)^2 - 2(1-h) = 1 - 2h + h^2 - 2 + 2h = h^2 - 1$. Equal. ✓

✓ Points the same distance either side of $x = 1$ have the same height: **the graph is symmetric about the line** $x = 1$ (its axis of symmetry).

Trick alert

- **"For all x" means one counter-example kills an option**, but one success doesn't prove it.
- **Watch the domain.** $f(x) = \log_e x$ satisfies $f(xy) = f(x) + f(y)$ only for $x, y > 0$.
- **Neither odd nor even is the usual case.** e^x, $x^2 + x$ and $\log_e x$ are neither.
- **A function can be odd and still have $f(0)$ undefined** (e.g. $\frac{1}{x}$). If an odd function is defined at 0, then $f(0) = 0$.

ALGEBRA, NUMBER & STRUCTURE

Simultaneous equations & parameters

Two linear equations, two unknowns

Two lines in the plane either cross once (**unique solution**), are parallel (**no solution**) or are the same line (**infinitely many solutions**).

The parameter test

$$\begin{aligned} ax + by &= e \\ cx + dy &= f \end{aligned} \qquad \text{unique solution} \iff ad - bc \neq 0$$

If $ad - bc = 0$ the lines have the same gradient. Then check the constants: if one equation is a multiple of the other, there are infinitely many solutions; otherwise there are none.

Key idea

$ad - bc$ is the determinant of the coefficient matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$. You don't need matrix theory. Just remember "unique ⇔ determinant ≠ 0", then test the special values by substituting them back in.

Worked example 1 **Classifying by a parameter** 4 marks

Consider $\begin{aligned} ax + 3y &= 3 \\ x + (a-2)y &= 1 \end{aligned}$. Find the values of a for which the system has (a) a unique solution, (b) no solution, (c) infinitely many solutions.

1 Determinant: $a(a-2) - 3 \cdot 1 = a^2 - 2a - 3 = (a-3)(a+1)$. Zero when $a = 3$ or $a = -1$. 2 $a = 3$: $3x + 3y = 3$ and $x + y = 1$. The first is 3 × the second, so they're the same line: infinitely many solutions. 3 $a = -1$: $-x + 3y = 3$ and $x - 3y = 1$, i.e. $x - 3y = -3$ and $x - 3y = 1$. Parallel, different lines: no solution.

(a) $a \in \mathbb{R} \setminus \{-1, 3\}$**; (b)** $a = -1$**; (c)** $a = 3$**.**

Three equations, three unknowns

Use elimination to reduce to two equations in two unknowns, or use CAS in Exam 2 (solve the three equations together). Geometrically each equation is a plane; they may meet at a point, along a line (infinitely many solutions) or not at all.

Worked example 3 **Infinitely many solutions in 3D (Exam 2 style)** 4 marks

For the system $x + y + z = 1$, $x + 2y + 3z = 2$, $x + 4y + kz = c$: (a) find k for which there is no unique solution; (b) for that k, find c so there are infinitely many solutions and give them in terms of a parameter.

1 (a) Subtract: (2) − (1) gives $y + 2z = 1$; (3) − (1) gives $3y + (k-1)z = c - 1$. 2 Triple the first: $3y + 6z = 3$. The two equations have the same left side when $k - 1 = 6$, i.e. $k = 7$. 3 (b) With $k = 7$: infinitely many solutions need $c - 1 = 3$, so $c = 4$ (otherwise no solution). Let $z = t$: $y = 1 - 2t$, $x = 1 - y - z = t$.

(a) $k = 7$**. (b)** $c = 4$**;** $(x, y, z) = (t,\ 1 - 2t,\ t),\ t \in \mathbb{R}$**.**

Lines meeting curves: the discriminant

Substitute the line into the curve to get a quadratic. Its discriminant $\Delta = b^2 - 4ac$ counts the intersections: $\Delta > 0$ two, $\Delta = 0$ one (the line is a **tangent**), $\Delta < 0$ none.

Worked example 4 **Tangent line condition** 3 marks

Find the values of m for which the line $y = mx + 1$ is a tangent to $y = x^2 + 2x + 2$.

1 Equate: $x^2 + 2x + 2 = mx + 1 \Rightarrow x^2 + (2 - m)x + 1 = 0$. 2 Tangent ⇔ $\Delta = 0$: $(2 - m)^2 - 4 = 0 \Rightarrow 2 - m = \pm 2$.

$m = 0$ **or** $m = 4$**.** (Check $m = 0$: the line $y = 1$ touches the parabola's vertex at $(-1, 1)$. ✓)

Literal equations

A literal equation has letters as well as the unknown: solve $ax + b = cx + d$ for x to get $x = \dfrac{d - b}{a - c}$, *provided* $a \neq c$. Always state the restriction: that's where the marks for "consider all cases" are.

Trick alert: parameter traps

- **Determinant = 0 doesn't tell you which case you're in.** Substitute each value back and test for "same line" vs "parallel".
- **Dividing by an expression containing a parameter** silently assumes it isn't 0. Handle that value separately.
- **"At least one solution"** means unique *or* infinitely many, so it includes the coincident case.
- **For x^2-coefficient parameters** (e.g. $kx^2 + 4x + 1 = 0$), check $k = 0$ separately: the equation becomes linear and has exactly one solution.

ALGEBRA, NUMBER & STRUCTURE

Newton's method, bisection & pseudocode

Newton's method

Start with a guess x_0 near a root. Draw the tangent at $(x_0, f(x_0))$; where it hits the x-axis is the next, usually better, estimate. Repeat.

Newton's method

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Derivation: the tangent $y - f(x_n) = f'(x_n)(x - x_n)$ crosses $y = 0$ at $x = x_n - \frac{f(x_n)}{f'(x_n)}$.

Worked example 1 **Two iterations by hand** 3 marks

Use Newton's method with $x_0 = 2$ to find x_2 for $f(x) = x^3 - 2x - 5$. Give x_2 correct to four decimal places.

1 $f'(x) = 3x^2 - 2$. $f(2) = 8 - 4 - 5 = -1$, $f'(2) = 10$. So $x_1 = 2 - \frac{-1}{10} = 2.1$. 2 $f(2.1) = 9.261 - 4.2 - 5 = 0.061$, $f'(2.1) = 13.23 - 2 = 11.23$. So $x_2 = 2.1 - \frac{0.061}{11.23} = 2.09457\ldots$

$x_2 \approx 2.0946$. (The true root is 2.094551…: two steps already give four correct decimal places.)

When Newton's method fails

Problem	What happens	Example
$f'(x_n) = 0$	Horizontal tangent never meets the axis: division by zero	$x^2 - 2$ with $x_0 = 0$
Near a stationary point	Tangent is nearly flat, so the next estimate shoots far away	$x^3 - 3x + 1$ with $x_0 = 1.01$
Cycling	Estimates bounce between values forever	$x^3 - 2x + 2$ with $x_0 = 0$: 0, 1, 0, 1, …
Wrong root	Converges, but to a different root from the one you wanted	Starting on the far side of a turning point

Worked example 3 **Show that it cycles** 2 marks

Show that Newton's method applied to $f(x) = x^3 - 2x + 2$ with $x_0 = 0$ does not converge.

1 $f'(x) = 3x^2 - 2$. $x_1 = 0 - \frac{2}{-2} = 1$. 2 $x_2 = 1 - \frac{1 - 2 + 2}{3 - 2} = 1 - 1 = 0$.

The estimates alternate 0, 1, 0, 1, … and never approach the actual root (which is near $x \approx -1.77$).

Pseudocode, VCAA style

VCAA exams show algorithms in a simple pseudocode and ask you to trace them (fill in a table of values), find an output, or complete a missing line. The conventions: `←` means "assign", `define` creates a function, and loops are `for … end for` and `while … end while`.

```
define f(x)
    return x^3 - 2x - 5
define df(x)
    return 3x^2 - 2

x ← 2
for i from 1 to 3
    x ← x - f(x) / df(x)
end for
print x
```

This prints $x_3 \approx 2.094551$ (three iterations of Newton's method from $x_0 = 2$).

Algebra

Worked example 4 **Trace a while loop** 3 marks

Trace this algorithm and state the output.

```
a ← 1
b ← 2
while b - a > 0.2
    m ← (a + b) / 2
    if m^2 - 2 < 0 then
        a ← m
    else
        b ← m
    end if
end while
print a, b
```

1 It's bisection on $x^2 - 2$. Start $a = 1, b = 2$ (width 1). 2 $m = 1.5$: $2.25 - 2 > 0$, so $b = 1.5$. $m = 1.25$: $1.5625 - 2 < 0$, so $a = 1.25$. Width now $0.25 > 0.2$: continue. 3 $m = 1.375$: $1.890625 - 2 < 0$, so $a = 1.375$. Width $0.125 \leq 0.2$: stop.

Output: 1.375, 1.5 (an interval containing $\sqrt{2} \approx 1.414$).

Worked example 5 **An accumulator loop (trapezium sum)** 3 marks

State the output, and the purpose of line 7.

```
define f(x)
    return x^2
a ← 0, b ← 2, n ← 4
h ← (b − a) / n
sum ← f(a) + f(b)
for i from 1 to n − 1
    sum ← sum + 2 × f(a + i × h)
end for
print h / 2 × sum
```

1 $h = 0.5$; start sum $= f(0) + f(2) = 4$. $i = 1$: $+2(0.25) = 4.5$; $i = 2$: $+2(1) = 6.5$; $i = 3$: $+2(2.25) = 11$.

Output $0.25 \times 11 = 2.75$: the trapezium estimate of $\int_0^2 x^2\,dx$ with 4 strips. Line 7 adds twice each interior height. If an exam blanks out a line like this, ask what the formula needs that the loop isn't yet doing.

Trick alert: algorithm traps

- **Count loop iterations carefully.** `for i from 1 to 3` runs 3 times; a `while` loop checks its condition *before* each pass.
- **Use the updated value.** Once `x ← …` runs, every later line uses the new x.
- **Newton needs f', bisection needs a sign change.** Don't mix up the input requirements.
- **Keep full precision** in the calculator between iterations; round only the final answer.

Hard exam questions

2024 Exam 2 · Section A · Question 17 © VCAA 27% correct

The question (not reprinted in this copy): Output of an algorithm that searches for integer roots of a cubic
Full question: 2024 Exam 2, Section A, Question 17 (page 9 of the paper).

Answer: D · A 14% · B 26% · C 32% · D 27%

The roots are 3, −3 and 2. The loop counts c down from 18, checking c then −c, so it prints 3, −3, 2.

CALCULUS

Limits, continuity & differentiability

Limits

$\lim_{x \to a} f(x) = L$ means $f(x)$ gets as close as we like to L as x gets close to a (from both sides). It doesn't matter what happens *at* $x = a$ itself.

Limits at infinity: divide top and bottom by the highest power. $\lim_{x \to \infty} \dfrac{2x+1}{x-3} = \lim_{x \to \infty} \dfrac{2+\frac{1}{x}}{1-\frac{3}{x}} = 2$: the horizontal asymptote.

Continuity

f is continuous at $x = a$ if $f(a)$ is defined and $\lim_{x \to a} f(x) = f(a)$. In practice: you can draw through $x = a$ without lifting your pen. For a hybrid function, the left and right rules must give the same value at the join.

Differentiability

f is differentiable at $x = a$ if the left and right gradients exist and are equal. A function that is **not continuous** at a point is **not differentiable** there. But continuity alone isn't enough:

Feature	Continuous?	Differentiable?	Example
Jump or hole	No	No	Hybrid with mismatched pieces
Corner (sharp point)	Yes	No	$\lvert x \rvert$ at $x = 0$; $\lvert x^2 - 4 \rvert$ at $x = \pm 2$
Cusp	Yes	No	$x^{2/3}$ at $x = 0$
Vertical tangent	Yes	No	$x^{1/3}$ at $x = 0$
Endpoint of a closed domain	Yes	No (only one-sided)	$\sqrt{x}$ at $x = 0$

Exam tip

VCAA convention: a function is differentiable on an **open** interval. When asked for the domain of f', exclude endpoints and every corner, cusp, jump and vertical tangent.

Worked example 3 **A smooth join (continuous and differentiable)** 3 marks

Find a and b such that $f(x) = \begin{cases} ax^2 + b, & x \leq 1 \\ 4x - 1, & x > 1 \end{cases}$ is differentiable at $x = 1$.

1 **Continuous:** $a(1)^2 + b = 4(1) - 1 \Rightarrow a + b = 3$. 2 **Gradients match:** left derivative $2ax$ at $x = 1$ is $2a$; right derivative is 4. So $2a = 4$.

$a = 2,\ b = 1$**.** Both conditions are needed. Matching gradients without continuity doesn't make the function differentiable.

Worked example 4 **Domain of the derivative** 2 marks

State the domain of f' for $f : [-3, 3] \to \mathbb{R},\ f(x) = \lvert x^2 - 4 \rvert$.

1 Corners where the graph is reflected: $x^2 - 4 = 0 \Rightarrow x = \pm 2$. Endpoints ± 3 are also excluded.

$(-3, -2) \cup (-2, 2) \cup (2, 3)$**.**

Worked example 5 **Writing the derivative of a hybrid** 2 marks

Let $f(x) = \begin{cases} x^2, & 0 \leq x \leq 2 \\ 8 - 2x, & 2 < x \leq 4 \end{cases}$. Write $f'(x)$ as a hybrid function, with its domain.

1 Left piece: $2x$, giving 4 at $x = 2$. Right piece: -2. The pieces meet (both 4) but the gradients differ, so there's a corner at $x = 2$.

$f'(x) = \begin{cases} 2x, & 0 < x < 2 \\ -2, & 2 < x < 4 \end{cases}$, with domain $(0, 2) \cup (2, 4)$. If the join were smooth, you'd include $x = 2$ in one piece.

Trick alert: limit and differentiability traps

- **Continuous doesn't mean differentiable.** Every corner is continuous.
- **A limit can exist even where $f(a)$ doesn't** (a hole), and $f(a)$ can exist without the limit (a jump).
- **For hybrid gradients, differentiate each piece separately** and compare the values at the join. Don't differentiate across the join.
- **Endpoints:** derivatives are only defined on open intervals in VCAA's convention.

Hard exam questions

2023 Exam 2 · Section A · Question 9 © VCAA — 42% correct

The question (not reprinted in this copy): Value of a that makes a tan/sin hybrid continuous and smooth at x = 2π
Full question: 2023 Exam 2, Section A, Question 9 (page 5 of the paper).

Answer: C · A 7% · B 9% · C 42% · D 31% · E 10%

Continuity needs $\sin(2\pi a) = 0$, so $a = \frac{n}{2}$ for an integer n. Smoothness needs $a\cos(2\pi a) = \frac{1}{2}$ (the gradient of $\tan\frac{x}{2}$ at 2π is $\frac{1}{2}\sec^2\pi = \frac{1}{2}$), i.e. $\frac{n}{2}(-1)^n = \frac{1}{2}$, which holds only for $n = -1$. So $a = -\frac{1}{2}$ ($a = \frac{1}{2}$ gives $-\frac{1}{2}$).

CALCULUS

Differentiation rules

The standard derivatives

$f(x)$	$f'(x)$	$f(x)$	$f'(x)$
x^n	nx^{n-1}	$\sin(kx)$	$k\cos(kx)$
$(ax+b)^n$	$an(ax+b)^{n-1}$	$\cos(kx)$	$-k\sin(kx)$
e^{kx}	ke^{kx}	$\tan(kx)$	$\frac{k}{\cos^2(kx)} = k\sec^2(kx)$
$\log_e(kx)$	$\frac{1}{x}$	$\log_e(g(x))$	$\frac{g'(x)}{g(x)}$

Rewrite roots and fractions as powers before differentiating: $\sqrt{x} = x^{1/2}$ and $\frac{3}{x^2} = 3x^{-2}$.

The three rules

Chain, product and quotient rules

$$\frac{dy}{dx} = \frac{dy}{du}\cdot\frac{du}{dx} \qquad (uv)' = u'v + uv' \qquad \left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

Function notation for the chain rule: $\big(f(g(x))\big)' = f'(g(x))\cdot g'(x)$.

Key idea

Decide which rule to use from the **outermost** structure. x^2e^{3x} is a product. $e^{\sin x}$ is a composite (chain). $\frac{\sin x}{x}$ is a quotient. Many questions need two rules in a row.

Worked example 2 **Chain rule with a log** — 2 marks

Find $\frac{dy}{dx}$ for (a) $y = \log_e(x^2+1)$, (b) $y = \log_e(\cos x)$.

1 Use $\frac{d}{dx}\log_e(g(x)) = \frac{g'(x)}{g(x)}$.

(a) $\frac{2x}{x^2+1}$**; (b)** $\frac{-\sin x}{\cos x} = -\tan x$.

Worked example 5 **Derivatives from a table (Exam 1 style)** — 3 marks

The table gives values of f, g and their derivatives.

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	3	−2	2	5
2	4	7	1	−3

Find the derivatives of (a) $f(g(x))$ at $x = 1$, (b) $f(x)g(x)$ at $x = 2$, (c) $[f(x)]^2$ at $x = 1$.

1 (a) $f'(g(1))\cdot g'(1) = f'(2)\cdot 5 = 7\times 5 = 35$. 2 (b) $f'(2)g(2) + f(2)g'(2) = 7(1) + 4(-3) = -5$. 3 (c) $2f(1)f'(1) = 2(3)(-2) = -12$.

(a) 35, (b) −5, (c) −12. In (a), f' is evaluated at $g(1) = 2$, not at 1. That's the trap.

Trick alert: differentiation traps

- $\frac{d}{dx}\log_e(2x) = \frac{1}{x}$, not $\frac{2}{x}$: the 2 cancels ($\frac{2}{2x}$).
- $\sin^2 x$ **vs** $\sin(x^2)$: $2\sin x\cos x$ vs $2x\cos(x^2)$.
- e^{x^2} **differentiates to** $2xe^{x^2}$, not $x^2e^{x^2-1}$. The power rule doesn't apply to exponentials.
- **The derivative of a constant is 0:** $\frac{d}{dx}e^2 = 0$ and $\frac{d}{dx}\pi^2 = 0$.
- **Quotient rule sign:** it's $u'v - uv'$ on top, so the order matters.

Calculus

Hard exam questions

2025 Exam 2 · Section A · Question 16 © VCAA

18% correct

The question (not reprinted in this copy): Conditions on a, b if $h(x) = a\log_e(bx)$ has $h'(x) > 0$ everywhere
Full question: 2025 Exam 2, Section A, Question 16 (page 9 of the paper).

Answer: D · A 20% · B 16% · C 46% · D 18%

$h'(x) = \frac{a}{x}$, with domain $bx > 0$. If b > 0 then x > 0 and you need a > 0. If b < 0 then x < 0 and you need a < 0. Either way ab > 0.

CALCULUS

Applications of differentiation

Tangents and normals

Tangent and normal at $x = a$

$$\text{tangent: } y - f(a) = f'(a)(x - a) \qquad \text{normal: } y - f(a) = -\frac{1}{f'(a)}(x - a)$$

The normal is perpendicular to the tangent: $m_1 m_2 = -1$.

Worked example 2 **Tangents through a point not on the curve (hard)** 4 marks

Find the equations of the tangents to $y = x^2$ that pass through the point $(1, -3)$.

1 Let the point of contact be (a, a^2). The gradient there is $2a$, so the tangent is $y - a^2 = 2a(x - a)$, i.e. $y = 2ax - a^2$. 2 It must pass through $(1, -3)$: $-3 = 2a - a^2 \Rightarrow a^2 - 2a - 3 = 0 \Rightarrow (a - 3)(a + 1) = 0$.

$a = 3$: $y = 6x - 9$; $a = -1$: $y = -2x - 1$. The external point isn't on the curve, so you can't use it as the point of contact. Use a general point instead.

Worked example 3 **A common tangent to two curves (hard)** 4 marks

Find the equation of the straight line that is a tangent to both $y = x^2$ and $y = x^2 - 4x + 8$.

1 With $y = x^2$: $x^2 - mx - c = 0$, so $\Delta = m^2 + 4c = 0$, giving $c = -\frac{m^2}{4}$. 2 With $y = x^2 - 4x + 8$: $x^2 - (4 + m)x + (8 - c) = 0$, so $(4 + m)^2 - 4(8 - c) = 0$. 3 Substitute $4c = -m^2$: $(4 + m)^2 - 32 - m^2 = 0 \Rightarrow 16 + 8m - 32 = 0 \Rightarrow m = 2$, so $c = -1$.

$y = 2x - 1$. Check: it touches $y = x^2$ at $(1, 1)$ and $y = x^2 - 4x + 8$ at $(3, 5)$. If both curves have a turning point at the same height, that horizontal line is a common tangent too (gradient 0), so look for it first.

The angle a tangent makes with the x-axis

A line with gradient m makes an angle θ with the positive direction of the x-axis, where $\tan\theta = m$. A negative gradient gives an obtuse angle ($\theta = 180^\circ + \tan^{-1} m$, or $\pi + \tan^{-1} m$ in radians).

Rates of change

Average rate of change over $[a, b]$ = $\frac{f(b) - f(a)}{b - a}$ (gradient of a secant). The **instantaneous** rate at $x = a$ is $f'(a)$ (gradient of the tangent). Read the question carefully: VCAA asks for both, often in the same question.

Worked example 4 **Average vs instantaneous rate (Exam 2 style)** 4 marks

A drink cools so that its temperature after t minutes is $T(t) = 80e^{-0.2t} + 20$ °C. Find (a) the average rate of change over the first 5 minutes, (b) the instantaneous rate at $t = 5$, and (c) when the instantaneous rate equals the average rate in (a).

1 (a) $\frac{T(5) - T(0)}{5} = \frac{80e^{-1} - 80}{5} = -16\left(1 - e^{-1}\right) \approx -10.11$ °C/min. 2 (b) $T'(t) = -16e^{-0.2t}$, so $T'(5) = -16e^{-1} \approx -5.89$ °C/min. 3 (c) $-16e^{-0.2t} = -16\left(1 - e^{-1}\right) \Rightarrow t = -5\log_e\left(1 - e^{-1}\right) \approx 2.29$.

(a) –10.11 °C/min; (b) –5.89 °C/min; (c) $t \approx 2.29$ **min.** Units on rates: °C per minute.

Stationary points and their nature

Stationary points occur where $f'(x) = 0$. Classify with a **sign diagram of** f' or with the **second derivative**:

Sign of f' (left → right)	$f''(a)$	Nature
$+ \to -$	< 0	local maximum
$- \to +$	> 0	local minimum
$+ \to +$ or $- \to -$	usually 0, but $f''(a) = 0$ on its own decides nothing: use the sign test	stationary point of inflection

f''(a) = 0 doesn't prove an inflection

$f(x) = x^4$ has $f'(0) = 0$ and $f''(0) = 0$, but $(0, 0)$ is a local *minimum*. When the second derivative is 0, go back to the sign test.

Calculus


Worked example 5 **A gradient table, VCAA style** 3 marks

For $f(x) = x^4 - 4x^3$, find the stationary points and use a gradient table to state their nature.

1 $f'(x) = 4x^3 - 12x^2 = 4x^2(x-3) = 0 \Rightarrow x = 0$ or $x = 3$. $f(0) = 0$, $f(3) = 81 - 108 = -27$. 2 Gradient table:

x	-1	0	1	3	4
$f'(x)$	-16	0	-8	0	64
slope	↘	→	↘	→	↗

$(0, 0)$ is a stationary point of inflection (negative on both sides); $(3, -27)$ is a local minimum. Here $f''(0) = 0$ as well, which on its own proves nothing.



$f(x) = x^3 - 3x^2 - 9x + 5$ (solid) and its gradient function $f'(x) = 3x^2 - 6x - 9$ (dashed). Where f turns, f' crosses the axis; where f inflects, f' has its turning point.

"Hence verify there's a stationary point in the interval"

If f' is continuous on $[a, b]$ and $f'(a)$ and $f'(b)$ have **opposite signs**, then $f'(x) = 0$ somewhere in (a, b), so f has a stationary point there. It's the same sign-change idea as bisection, applied to f'. "Verify" means evaluate and conclude; don't solve $f'(x) = 0$ exactly.

Concavity and points of inflection

$f''(x) > 0$: concave up (gradient increasing). $f''(x) < 0$: concave down. A **point of inflection** is where the concavity changes; it may or may not be stationary. At a non-stationary inflection point the gradient is at a local max or min, so it's where a quantity grows fastest or slowest (a favourite Exam 2 context).

Worked example 8 **Greatest rate of change: inflection or endpoint** 3 marks

The depth of water in a channel is $h(t) = -t^3 + 6t^2 + 15$ m, $0 \le t \le 5$ hours. When is the depth increasing most rapidly, and at what rate? When is it decreasing most rapidly?

1 $h'(t) = -3t^2 + 12t$, $h''(t) = -6t + 12 = 0 \Rightarrow t = 2$, and $h'(2) = 12$. 2 Endpoints: $h'(0) = 0$, $h'(5) = -75 + 60 = -15$. CAS: `fMax(derivative(h(t),t),t)|0≤t≤5` and `fMin(…)`.

Increasing most rapidly at $t = 2$ h, at 12 m/h; decreasing most rapidly at $t = 5$ h (the endpoint), at 15 m/h.

Sketching the gradient function

Read across a row to match a graph with its gradient graph (or, reading right to left, f' back to f; the same table links an antiderivative F to f).

On the graph of f	On the graph of f'
increasing / decreasing	above / below the x-axis
local max / local min	crosses the axis $+ \to -$ / $- \to +$
stationary point of inflection	touches the axis without crossing
non-stationary point of inflection	a turning point not on the axis
concave up / concave down	increasing / decreasing
straight-line section, gradient m	horizontal section at height m
corner	a jump (open circles), f' undefined there
cusp	a vertical asymptote, with opposite signs on each side (e.g. $y = x^{2/3}$)
vertical tangent	vertical asymptote
vertical asymptote	vertical asymptote at the same x
horizontal asymptote	$f' \to 0$ (approaches the x-axis)

"Which graph could be f'?"

Mark the x-values of every stationary point of f first and rule out any option that isn't zero exactly there. Then check the sign between them. That usually leaves one option; use concavity (where f' turns) only to break a tie.

Trick alert: curve-sketching traps

- **Stationary point ≠ turning point.** Turning points are only the maxima and minima; stationary inflections also have $f' = 0$.
- **Local ≠ absolute.** The absolute maximum on a closed interval might be at an endpoint.
- **Normal gradient:** negative reciprocal. If the tangent is horizontal, the normal is vertical ($x = a$).
- **A factor squared in f'** (e.g. $f'(x) = (x-2)^2(x+1)$) gives no sign change at $x = 2$: stationary inflection.

Hard exam questions

2023 Exam 2 · Section A · Question 14 © VCAA — 29% correct

The question (not reprinted in this copy): Number of tangents to a cubic that pass through its positive x-intercept
Full question: 2023 Exam 2, Section A, Question 14 (page 7 of the paper).

Answer: D · A 6% · B 33% · C 22% · D 29% · E 10%

A tangent at $x = a$ passes through $\left(\frac{1}{3}, 0\right)$ when $-f(a) = f'(a)\left(\frac{1}{3} - a\right)$. This simplifies to $(3a-1)^2(3a^2+8a+3) = 0$, so $a = \frac{1}{3}$ (the tangent at the intercept itself counts) or $a = \frac{-4 \pm \sqrt{7}}{3}$: three tangents.

CALCULUS

Optimisation (max/min problems)

The method

1. **Draw a diagram** and label the variable(s).
2. **Write the quantity to optimise** as a function of *one* variable (use the constraint to eliminate the others).
3. **State the domain** (lengths positive, can't cut more than you have…).
4. **Differentiate and solve** $= 0$; check the nature.
5. **Check the endpoints** and answer the question actually asked (the maximum volume? the dimensions? the time?).

Exam tip

VCAA marking schemes award a mark for a correct expression in one variable and often for the domain, before any calculus. Show them.

Worked example 1 **The open box** — 4 marks

Squares of side x cm are cut from each corner of a 24 cm × 24 cm sheet, and the sides are folded up to make an open box. Find the maximum volume.

1 $V = x(24-2x)^2$, domain $0 < x < 12$. 2 $V'(x) = (24-2x)^2 + x \cdot 2(24-2x)(-2) = (24-2x)(24-6x)$. 3 $V' = 0 \Rightarrow x = 12$ (outside the domain; zero volume) or $x = 4$. Sign: $V' > 0$ before 4, < 0 after, so it's a maximum.

$V_{\max} = 4(16)^2 = 1024 \text{ cm}^3$ when $x = 4$.

Worked example 4 **Closest point on a curve** — 4 marks

Find the minimum distance from the point $(0, 3)$ to the parabola $y = x^2$.

1 A point on the curve is (x, x^2). Minimise the *square* of the distance (easier, same minimising x): $D = x^2 + (x^2-3)^2$. 2 $D'(x) = 2x + 4x(x^2-3) = 2x(2x^2-5) = 0 \Rightarrow x = 0$ or $x^2 = \frac{5}{2}$. 3 $D(0) = 9$ (local max of D); D at $x^2 = \frac{5}{2}$: $\frac{5}{2} + \frac{1}{4} = \frac{11}{4}$.

Minimum distance $= \sqrt{\frac{11}{4}} = \frac{\sqrt{11}}{2}$, at $\left(\pm\sqrt{\frac{5}{2}}, \frac{5}{2}\right)$.

Worked example 5 **Swim then run (Exam 2 style)** — 4 marks

A swimmer is 3 km offshore, directly opposite point P on a straight beach. She wants to reach a café on the beach 8 km from P. She swims at 2 km/h and runs at 5 km/h. Where should she land to minimise the total time?

1 Land x km from P towards the café. $T(x) = \frac{\sqrt{9+x^2}}{2} + \frac{8-x}{5}$, $0 \le x \le 8$. 2 $T'(x) = \frac{x}{2\sqrt{9+x^2}} - \frac{1}{5} = 0 \Rightarrow 5x = 2\sqrt{9+x^2} \Rightarrow 21x^2 = 36$. 3 $x = \frac{6}{\sqrt{21}} \approx 1.31$ km, giving $T \approx 2.9748$ h. Endpoints: $T(0) = 3.1$ h, $T(8) \approx 4.27$ h.

Land about 1.31 km from P; the minimum time is about 2.97 h (2 h 58.5 min).

Endpoints: absolute max and min

On a closed interval, the absolute maximum and minimum are among the stationary points **and the endpoints**. Evaluate all of them and compare.

Worked example 6 **Absolute extrema on [−2, 3]** — 3 marks

Find the absolute maximum and minimum of $f(x) = x^3 - 3x$ on $[-2, 3]$.

1 $f'(x) = 3x^2 - 3 = 0 \Rightarrow x = \pm 1$. $f(-1) = 2$, $f(1) = -2$. 2 Endpoints: $f(-2) = -8 + 6 = -2$, $f(3) = 27 - 9 = 18$.

Absolute max 18 (at $x = 3$); absolute min -2 (at $x = -2$ and $x = 1$). The local max of 2 isn't the absolute max.

Worked example 7 **Greatest difference between two models** — 3 marks

Two models are $u(x) = 2\sin x + 3$ and $v(x) = 0.5x + 2$ for $0 \le x \le 5$. Find the greatest difference between them.

1 $d(x) = 2\sin x - 0.5x + 1$, $d'(x) = 2\cos x - 0.5 = 0 \Rightarrow \cos x = 0.25$: $x \approx 1.318$, $d \approx 2.277$; $x \approx 4.965$, $d \approx -3.419$. 2 Endpoints: $d(0) = 1$, $d(5) \approx -3.418$.

The greatest difference is about 3.419, at $x \approx 4.965$ (v above u there). The endpoint gives 3.418, so close that skipping either check could cost the mark. CAS: `fMax(abs(u(x)-v(x)),x)|0≤x≤5`.

Trick alert: optimisation traps

- **Forgetting the endpoints.** Especially when a stationary point is outside the domain.
- **Answering the wrong question.** "Find the maximum area" wants the area, not x.
- **Two variables left in the function.** Use the constraint to eliminate one before differentiating.
- **Rounding too early.** Keep the exact x (or full calculator value) until the final answer.

Hard exam questions

2025 Exam 2 · Section A · Question 19 © VCAA — 14% correct

The question (not reprinted in this copy): Value of c when the shortest distance from $y = x + c$ to $y = \log_e(x - 1)$ is $\sqrt{2}$
Full question: 2025 Exam 2, Section A, Question 19 (page 12 of the paper).

Answer: D · A 35% · B 25% · C 24% · D 14%

The closest point on the curve is where its gradient equals the line's: $\frac{1}{x-1} = 1$ at (2, 0). The normal there is y = 2 − x, and the points $\sqrt{2}$ units along it are (1, 1) and (3, −1). Putting each on y = x + c gives c = 0 or c = −4, but y = x − 4 crosses the curve (distance 0), so c = 0.

CALCULUS

Antidifferentiation

Standard antiderivatives

Antidifferentiation reverses differentiation: if $F'(x) = f(x)$ then $\int f(x)\,dx = F(x) + c$.

$f(x)$	$\int f(x)\,dx$	$f(x)$	$\int f(x)\,dx$
$x^n,\ n \neq -1$	$\frac{x^{n+1}}{n+1} + c$	e^{kx}	$\frac{1}{k}e^{kx} + c$
$(ax+b)^n,\ n \neq -1$	$\frac{(ax+b)^{n+1}}{a(n+1)} + c$	$\sin(kx)$	$-\frac{1}{k}\cos(kx) + c$
$\frac{1}{x}$	$\log_e\lvert x\rvert + c$	$\cos(kx)$	$\frac{1}{k}\sin(kx) + c$
$\frac{1}{ax+b}$	$\frac{1}{a}\log_e\lvert ax+b\rvert + c$	$\frac{1}{\cos^2(kx)}$	$\frac{1}{k}\tan(kx) + c$

Key idea

Linear inside, divide by the coefficient of x. That's the reverse chain rule, and it only works when the inside is **linear**. $\int e^{x^2}\,dx$ is *not* $\frac{e^{x^2}}{2x}$.

Worked example 1 **A mixed antiderivative** — 2 marks

Find $\int \left(6x^2 - \frac{4}{x^3} + \frac{2}{2x+1} + e^{-3x}\right) dx$.

1 Term by term: $2x^3$; $-4 \cdot \frac{x^{-2}}{-2} = \frac{2}{x^2}$; $\frac{2}{2}\log_e\lvert 2x+1\rvert = \log_e\lvert 2x+1\rvert$; $-\frac{1}{3}e^{-3x}$.

$2x^3 + \frac{2}{x^2} + \log_e\lvert 2x+1\rvert - \frac{1}{3}e^{-3x} + c$.

Calculus

Worked example 2 **Function from derivative and a point** — 2 marks

$f'(x) = 3x^2 - 4x + 1$ and $f(1) = 2$. Find $f(x)$.

1. $f(x) = x^3 - 2x^2 + x + c$.

✓ $f(1) = 1 - 2 + 1 + c = 2 \Rightarrow c = 2$. $f(x) = x^3 - 2x^2 + x + 2$.

Antidifferentiation by recognition

This is an Exam 1 staple: "Differentiate $g(x)$. Hence find an antiderivative of $h(x)$." Differentiate, rearrange so the thing you want is on its own, then antidifferentiate both sides.

Worked example 4 **Hence find ∫ log x dx** — 3 marks

Find $\frac{d}{dx}(x \log_e x)$. Hence find $\int \log_e x \, dx$.

1. Product rule: $\frac{d}{dx}(x \log_e x) = \log_e x + x \cdot \frac{1}{x} = \log_e x + 1$. 2. Antidifferentiate both sides: $x \log_e x = \int \log_e x \, dx + \int 1 \, dx$.

✓ $\int \log_e x \, dx = x \log_e x - x + c$.

One constant, one condition

An antiderivative has one constant c, so it can usually be made to pass through only one given point. For two points, find c from each: if the values differ, no antiderivative of f passes through both, and the question will change something else (a dilation, or a parameter in f) to make it work.

Graphs of antiderivatives

If you're given the graph of f and asked to sketch F (where $F' = f$): where f is positive, F increases; where f crosses the axis, F has a stationary point; turning points of f are inflection points of F. There's a whole family of possible F graphs, each a vertical translation of the others.

Trick alert: antiderivative traps

- $\int \frac{1}{x} \, dx \neq \frac{x^0}{0}$. The power rule excludes $n = -1$; use $\log_e |x|$.
- **Sign flip for sine:** $\int \sin x \, dx = -\cos x + c$.
- $\int \frac{1}{3x} \, dx = \frac{1}{3} \log_e |x| + c$, which equals $\frac{1}{3} \log_e |3x| + c'$: both are correct (the constants differ).
- **Don't forget** $+c$ when a question asks for "the general antiderivative" or "$\int \ldots dx$". When it asks for "an antiderivative", $c = 0$ is fine.

CALCULUS

Definite integrals, areas & average value

The fundamental theorem

Fundamental theorem of calculus

$$\int_a^b f(x) \, dx = F(b) - F(a) \quad \text{where } F' = f \qquad \frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

The definite integral gives the **signed area** between the graph and the x-axis: area above counts positive, area below counts negative.

Properties

$$\int_a^a f = 0 \qquad \int_b^a f = -\int_a^b f \qquad \int_a^b f + \int_b^c f = \int_a^c f \qquad \int_a^b (kf + g) = k\int_a^b f + \int_a^b g$$

Constants integrate to (constant) × (interval width): $\int_a^b k \, dx = k(b - a)$.

Worked example 1 **Using properties (Exam 1 classic)** — 2 marks

Given $\int_1^4 f(x) \, dx = 5$, find $\int_1^4 (2f(x) + 1) \, dx$ and $\int_4^1 f(x) \, dx$.

1. $2\int_1^4 f + \int_1^4 1 \, dx = 2(5) + (4 - 1) = 13$.

✓ **13 and** -5. The common error is $2(5) + 1 = 11$: the 1 must be integrated over an interval of width 3.

Calculus


Signed area vs actual area

Worked example 2 **When part of the region is below the axis** 3 marks

For $f(x) = x^2 - 1$ on $[0, 2]$, find (a) $\int_0^2 f(x)\,dx$ and (b) the total area between the graph and the x-axis.

1 (a) $\left[\frac{x^3}{3} - x\right]_0^2 = \frac{8}{3} - 2 = \frac{2}{3}$. 2 (b) The graph crosses at $x = 1$. Area $= -\int_0^1 (x^2 - 1)\,dx + \int_1^2 (x^2 - 1)\,dx = \frac{2}{3} + \frac{4}{3}$.

(a) $\frac{2}{3}$ (signed); (b) 2 square units. Always sketch first and split at the x-intercepts.



$y = x^2 - 1$ on $[0, 2]$: the region below the axis has area $\frac{2}{3}$ and the region above has area $\frac{4}{3}$. The integral subtracts them ($\frac{4}{3} - \frac{2}{3} = \frac{2}{3}$); the area adds them (2).

Area between curves

Area between curves

$$A = \int_a^b (\text{upper} - \text{lower})\,dx$$

Find the intersection points first. If the curves swap over, split the integral where they cross. Upper − lower works whether the region is above or below the axis. When a question asks for **one** integral (Exam 2), write $A = \int_a^b |f(x) - g(x)|\,dx$: the modulus handles every crossing at once.

Worked example 5 **A region where the curves swap over** 3 marks

Find the total area enclosed between $y = x^3 - x$ and $y = 3x$.

1 Intersections: $x^3 - x = 3x \Rightarrow x(x^2 - 4) = 0 \Rightarrow x = -2, 0, 2$. 2 On $(0, 2)$, $3x$ is on top; on $(-2, 0)$, the cubic is on top. By symmetry (both are odd functions), the two regions are equal.

$A = 2\int_0^2 (3x - (x^3 - x))\,dx = 2\left[2x^2 - \frac{x^4}{4}\right]_0^2 = 2(8 - 4) = \mathbf{8}$. $\int_{-2}^2 (3x - x^3 + x)\,dx = 0$: don't integrate straight across a crossing. On CAS, one integral does it: `integral(abs(x^3-x-3·x),x,-2,2)` gives 8.

Average value

Average value of f on $[a, b]$

$$f_{\text{avg}} = \frac{1}{b - a}\int_a^b f(x)\,dx$$

The height of the rectangle on $[a, b]$ with the same (signed) area as the region under f.

Worked example 6 **Average value of e^x** 2 marks

Find the average value of $f(x) = e^x$ on $[0, 2]$.

1 $\frac{1}{2}\int_0^2 e^x\,dx = \frac{1}{2}\left[e^x\right]_0^2$.

$\frac{e^2 - 1}{2} \approx 3.19$. This is not the same as the average of $f(0)$ and $f(2)$ (≈ 4.19).

Periodic functions: any full period gives the same integral

If $f(x + p) = f(x)$ for all x (period p), then $\int_a^{a+p} f(x)\,dx = \int_0^p f(x)\,dx$ for any a, and over n full periods the integral is n times that. So the average value over any whole number of periods is the same: e.g. $\sin^2 x$ (period π) averages $\frac{1}{2}$ over $[0, \pi]$, $[3, 3 + \pi]$ or $[0, 5\pi]$.

Average value vs average rate of change

Average value of f on $[a,b]$ is $\dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx$: a height, and it needs an integral. **Average rate of change** of f on $[a,b]$ is $\dfrac{f(b)-f(a)}{b-a}$: a gradient, with no integral. If f is itself a rate (litres per minute, say), its average value is the average rate.

From rates to totals

If $R(t)$ is a rate (litres/min, cars/hour, m/s), then $\displaystyle\int_a^b R(t)\,dt$ is the **net change** in the quantity from $t=a$ to $t=b$.

Worked example 7 **Total change, then solve for the time (Exam 2 style)** 3 marks

A tank holds 50 L. Water flows in at $R(t)=30e^{-0.1t}$ L/min. (a) How much water is in the tank after 10 minutes? (b) When does it hold 200 L?

1 (a) $V(10)=50+\displaystyle\int_0^{10}30e^{-0.1t}\,dt=50+300\left(1-e^{-1}\right)\approx 239.6$ L. 2 (b) $50+300\left(1-e^{-0.1T}\right)=200\Rightarrow e^{-0.1T}=\dfrac{1}{2}\Rightarrow T=10\log_e 2$. CAS: `solve(50+integral(30·e^(-0.1·t),t,0,T)=200,T)`.

(a) about 239.6 L; (b) $T=10\log_e 2\approx 6.93$ **min.** Don't forget the starting 50 L.

Trick alert: integration traps

- **"Area" is never negative.** Split at the x-intercepts and add magnitudes.
- **Constant terms integrate to (constant) × width**, not the constant itself.
- **Upper – lower**, not "curve – axis", for regions between curves.
- **Average value divides by $b-a$**, not by 2.
- **Exact vs approximate:** if no rounding is asked for, an area like $\dfrac{e^2-1}{2}$ must stay exact.

Hard exam questions

2025 Exam 2 · Section A · Question 17 © VCAA 38% correct

The question (not reprinted in this copy): Which graph fits $\displaystyle\int_1^2 f\,dx > \int_1^3 f\,dx$

Full question: 2025 Exam 2, Section A, Question 17 (page 10 of the paper).

Answer: A · A 38% · B 20% · C 17% · D 24%

The condition means $\displaystyle\int_2^3 f(x)\,dx<0$: the net signed area on [2, 3] is negative, so there is more area below the axis than above it there.

CALCULUS

Numerical integration: trapezium & rectangle rules

Rectangles and trapeziums

Split $[a,b]$ into n strips of width $h=\dfrac{b-a}{n}$, with $x_0=a, x_1,\ldots,x_n=b$.

The three estimates

$$L=h\left[f(x_0)+f(x_1)+\cdots+f(x_{n-1})\right]\qquad R=h\left[f(x_1)+\cdots+f(x_n)\right]$$

$$T=\frac{h}{2}\left[f(x_0)+2f(x_1)+\cdots+2f(x_{n-1})+f(x_n)\right]=\frac{L+R}{2}$$

The trapezium rule is on the formula sheet. L and R aren't, but they're easy: left uses every height except the last, right every height except the first.

Worked example 1 **All three estimates for x^2 on [0, 2]** 3 marks

Estimate $\displaystyle\int_0^2 x^2\,dx$ with four strips using (a) left rectangles, (b) right rectangles, (c) the trapezium rule. Compare with the exact value.

1 $h=0.5$. Heights at $x=0,0.5,1,1.5,2$: $0,0.25,1,2.25,4$. 2 (a) $L=0.5(0+0.25+1+2.25)=1.75$. (b) $R=0.5(0.25+1+2.25+4)=3.75$. 3 (c) $T=\dfrac{0.5}{2}\left(0+2(0.25+1+2.25)+4\right)=0.25\times 11=2.75$.

$L=1.75,\ R=3.75,\ T=2.75$; **exact** $\dfrac{8}{3}\approx 2.667$. x^2 is increasing, so L under- and R over-estimates. It's concave up, so the trapeziums sit above the curve and T is an overestimate.

Calculus

Over or under?

Function on the interval	Left rectangles	Right rectangles	Trapezium rule
Increasing	under	over	depends on concavity
Decreasing	over	under	depends on concavity
Concave up ($f'' > 0$)	–	–	over
Concave down ($f'' < 0$)	–	–	under

Key idea

Picture the tops: a trapezium's slanted top is a chord. Chords lie *above* a concave-up curve and *below* a concave-down one.

Function	Concavity	Trapezium rule
e^{kx}, x^2, x^n $(n \geq 2, x > 0)$, $\dfrac{1}{x}$ $(x > 0)$	concave up	overestimates
$\log_e x$, $\sqrt{x}$, $\sqrt[3]{x}$ $(x > 0)$	concave down	underestimates
$\sin x$	down on $(0, \pi)$, up on $(\pi, 2\pi)$	depends on the interval
$\cos x$	down on $\left(-\dfrac{\pi}{2}, \dfrac{\pi}{2}\right)$, up on $\left(\dfrac{\pi}{2}, \dfrac{3\pi}{2}\right)$	depends on the interval

Worked example 3 **From a table of data** 2 marks

A car's speed (m/s) is recorded every 2 seconds: $t = 0, 2, 4, 6, 8$; $v = 0, 6, 10, 12, 13$. Estimate the distance travelled in 8 s using the trapezium rule.

1 $h = 2$: $T = \dfrac{2}{2}\big(0 + 2(6 + 10 + 12) + 13\big) = 56 + 13.$

About 69 m. Distance is the area under a speed–time graph.

Trick alert: numerical-integration traps

- **Count strips, not points.** Five x-values make four strips.
- **End heights appear once** in the trapezium rule; the middle ones are doubled.
- **Left rectangles don't use the last height**, and right rectangles don't use the first.
- **Over/under for a function that changes concavity** can't be decided by the table above. Compute both, or compare with the exact value.

Hard exam questions

2025 Exam 2 · Section A · Question 6 © VCAA — 50% correct

The question (not reprinted in this copy): For which function the trapezium rule overestimates the area on [0, 1]
Full question: 2025 Exam 2, Section A, Question 6 (page 4 of the paper).

Answer: B · A 18% · B 50% · C 19% · D 11%

The trapezium rule overestimates when the curve is concave up on the interval. $x^3 + 1$ is concave up on [0, 1].

DATA ANALYSIS, PROBABILITY & STATISTICS

Probability: conditional & independent events

The core rules

Addition, conditional and multiplication rules

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) \qquad P(A') = 1 - P(A)$$

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} \qquad P(A \cap B) = P(A \mid B)\,P(B)$$

Mutually exclusive

A and B can't both happen: $P(A \cap B) = 0$, so $P(A \cup B) = P(A) + P(B)$.

Independent

One happening doesn't change the other's probability: $P(A \mid B) = P(A)$, or equivalently $P(A \cap B) = P(A)P(B)$.

Mutually exclusive ≠ independent

If A and B are mutually exclusive and both have non-zero probability, they're *dependent*: knowing A happened tells you B definitely didn't. $P(A \cap B) = 0 \neq P(A)P(B)$.

Karnaugh maps (probability tables)

A 2 × 2 table with totals is the fastest tool when there are two events. Fill in what you know; every row and column must add up.

Worked example 1 **Karnaugh map and independence** 3 marks

$P(A) = 0.6$, $P(B) = 0.3$ and $P(A \cap B') = 0.45$. (a) Find $P(A \cap B)$. (b) Are A and B independent? (c) Find $P(B \mid A)$.

1

	B	B'	total
A	**0.15**	0.45	0.6
A'	0.15	0.25	0.4
total	0.3	0.7	1

2 (b) $P(A)P(B) = 0.18 \neq 0.15 = P(A \cap B)$: not independent.

(a) 0.15; (b) not independent; (c) $P(B \mid A) = \dfrac{0.15}{0.6} = 0.25$.

Tree diagrams and the law of total probability

Law of total probability

$$P(B) = P(B \mid A)P(A) + P(B \mid A')P(A')$$

Multiply along branches; add the branches that give the outcome you want.

Trees with letters

With $P(A) = m$, $P(B \mid A) = p$ and $P(B \mid A') = q$: $P(B) = mp + (1 - m)q$ and $P(A \mid B) = \dfrac{mp}{mp + (1 - m)q}$. Multiple-choice options are often these expressions, so build the tree with letters rather than guessing.

Worked example 3 **The false-positive problem (hard)** 4 marks

2% of a population has a disease. A test gives a positive result for 95% of people with the disease and for 10% of people without it. A randomly chosen person tests positive. Find the probability that they have the disease, correct to four decimal places.

1 Tree: $P(D \cap +) = 0.02 \times 0.95 = 0.019$; $P(D' \cap +) = 0.98 \times 0.10 = 0.098$. 2 $P(+) = 0.019 + 0.098 = 0.117$. 3 $P(D \mid +) = \dfrac{P(D \cap +)}{P(+)} = \dfrac{0.019}{0.117}$.

≈ 0.1624. Only about 16% of positives actually have the disease, because the disease is rare and the false positives swamp the true ones. This is the classic "reverse conditional".

Worked example 4 **Without replacement** 2 marks

A bag holds 5 red and 3 blue marbles. Two are drawn without replacement. Find the probability of getting one of each colour.

1 $P(RB) + P(BR) = \dfrac{5}{8} \times \dfrac{3}{7} + \dfrac{3}{8} \times \dfrac{5}{7}$.

$= \dfrac{30}{56} = \dfrac{15}{28}$. Don't forget both orders.

Worked example 5 **Independent trials with different probabilities** 3 marks

Three independent traffic lights are green with probabilities 0.5, 0.4 and 0.2. Y is the number of green lights. Find the probability distribution of Y and $E(Y)$.

1 $P(Y=0) = 0.5 \times 0.6 \times 0.8 = 0.24$; $P(Y=3) = 0.5 \times 0.4 \times 0.2 = 0.04$. 2 $P(Y=1) = 0.5 \times 0.6 \times 0.8 + 0.5 \times 0.4 \times 0.8 + 0.5 \times 0.6 \times 0.2 = 0.24 + 0.16 + 0.06 = 0.46$; then $P(Y=2) = 1 - 0.24 - 0.46 - 0.04 = 0.26$.

✓ $P(Y=0,1,2,3) = 0.24,\ 0.46,\ 0.26,\ 0.04$; $E(Y) = 0.5 + 0.4 + 0.2 = 1.1$ (expected values add even though Y isn't binomial). CAS check: `expand((0.5+0.5·t)·(0.6+0.4·t)·(0.8+0.2·t))` gives $0.24 + 0.46t + 0.26t^2 + 0.04t^3$: the coefficients are the distribution.

Trick alert: probability traps

- $P(A \mid B) \neq P(B \mid A)$ in general, as the disease example shows.
- **"Given" shrinks the sample space.** Divide by the probability of the condition.
- **Independence must be checked with numbers**, not intuition.
- **Without replacement:** the second-draw probabilities change.

Hard exam questions

2024 Exam 2 · Section A · Question 11 © VCAA **40% correct**

The question (not reprinted in this copy): Choosing 3 of 12 students: probability exactly two are from the front row of seven
Full question: 2024 Exam 2, Section A, Question 11 (page 5 of the paper).

Answer: B · A 12% · B 40% · C 10% · D 37%

$$\frac{\binom{7}{2}\binom{5}{1}}{\binom{12}{3}} = \frac{105}{220} = \frac{21}{44}$$

DATA ANALYSIS, PROBABILITY & STATISTICS

Discrete random variables

Distributions

A **random variable** assigns a real number to each outcome in the sample space, e.g. the number of heads when three coins are tossed. A discrete random variable X takes separate values $x_1, x_2, \ldots$ with probabilities $p_1, p_2, \ldots$. Every probability satisfies $0 \leq p_i \leq 1$, and together they sum to 1.

Mean, variance, standard deviation

$$E(X) = \mu = \sum x\,p(x) \qquad \mathrm{Var}(X) = E\left[(X-\mu)^2\right] = E(X^2) - \mu^2 \qquad \mathrm{sd}(X) = \sqrt{\mathrm{Var}(X)}$$

$$E(aX+b) = aE(X) + b \qquad \mathrm{Var}(aX+b) = a^2\,\mathrm{Var}(X)$$

Worked example 1 **Full workout on a distribution table** 4 marks

x	0	1	2	3
$P(X=x)$	0.1	0.3	0.4	0.2

Find $E(X)$, $\mathrm{Var}(X)$, $\mathrm{sd}(X)$, and the mean and standard deviation of $Y = 3X - 2$.

1 $E(X) = 0 + 0.3 + 0.8 + 0.6 = 1.7$. 2 $E(X^2) = 0 + 0.3 + 1.6 + 1.8 = 3.7$, so $\mathrm{Var}(X) = 3.7 - 1.7^2 = 0.81$ and $\mathrm{sd}(X) = 0.9$. 3 $E(Y) = 3(1.7) - 2 = 3.1$; $\mathrm{Var}(Y) = 9 \times 0.81 = 7.29$, so $\mathrm{sd}(Y) = 2.7$.

✓ $E(X) = 1.7,\ \mathrm{Var}(X) = 0.81,\ \mathrm{sd}(X) = 0.9$; $E(Y) = 3.1,\ \mathrm{sd}(Y) = 2.7$. The -2 shifts the mean but doesn't change the spread.

Conditional probability from a distribution table

Add the probabilities of the values in both events, then divide by the probability of the given event. With the table above: $P(X \geq 2 \mid X \geq 1) = \dfrac{0.4 + 0.2}{0.9} = \dfrac{2}{3}$, and $P(X < 3 \mid X > 0) = \dfrac{P(1 \leq X \leq 2)}{P(X \geq 1)} = \dfrac{0.7}{0.9} = \dfrac{7}{9}$. The numerator is the overlap, not the first event on its own.

Worked example 2 **Unknown probabilities** 3 marks

$P(X = x) = kx$ for $x = 1, 2, 3, 4$. Find k, $E(X)$ and $P(X > E(X))$.

1 $k(1+2+3+4) = 1 \Rightarrow k = 0.1$. Probabilities: 0.1, 0.2, 0.3, 0.4. 2 $E(X) = 0.1 + 0.4 + 0.9 + 1.6 = 3$.

✓ $k = 0.1$, $E(X) = 3$, $P(X > 3) = P(X = 4) = 0.4$. "Greater than" is strict: don't include $X = 3$.

Probability

Trick alert: random variable traps

- $\operatorname{Var}(aX+b) = a^2\operatorname{Var}(X)$. A negative a never makes the variance negative: $\operatorname{Var}(-2X) = 4\operatorname{Var}(X)$, not $-2\operatorname{Var}(X)$.
- $E(X^2) \neq [E(X)]^2$. Their difference *is* the variance.
- **sd scales by** $|a|$, variance by a^2.
- **The expected value needn't be a possible value** (1.7 children...).
- **Symmetric distribution?** The mean is the centre of symmetry: no calculation needed.

Hard exam questions

2023 Exam 2 · Section A · Question 12 © VCAA — 29% correct

The question (not reprinted in this copy): Maximum possible mean of a pmf with parameter k
Full question: 2023 Exam 2, Section A, Question 12 (page 6 of the paper).

Answer: E · A 11% · B 14% · C 26% · D 18% · E 29%

All probabilities must lie in [0, 1], so $0 \leq k \leq \sqrt{5} - 2$. The mean $-3k^2 - 7k + 2$ is largest at k = 0, giving 2.

DATA ANALYSIS, PROBABILITY & STATISTICS

The binomial distribution

Bernoulli trials

A **Bernoulli trial** has exactly two outcomes: success, scored $X = 1$, with probability p, and failure, scored $X = 0$, with probability $1 - p$.

Bernoulli random variable

$$P(X=1) = p \qquad P(X=0) = 1-p \qquad E(X) = p \qquad \operatorname{Var}(X) = p(1-p)$$

A binomial variable counts the successes in n independent Bernoulli trials with the same p, which is why its mean and variance are n times these: np and $np(1-p)$.

When is it binomial?

$X \sim \operatorname{Bi}(n, p)$ counts the number of successes when:

1. there's a fixed number n of trials;
2. each trial has two outcomes (success/failure);
3. the probability of success p is the same every trial;
4. the trials are independent.

Binomial distribution

$$P(X=x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \ldots, n \qquad E(X) = np \qquad \operatorname{Var}(X) = np(1-p)$$

$\binom{n}{x} = \dfrac{n!}{x!(n-x)!}$ counts the arrangements of x successes among n trials. For example $\binom{5}{2} = 10$.

How n and p change the graph

- $p = 0.5$: symmetric about the mean np.
- $p < 0.5$: positively skewed (tail to the right). $p > 0.5$: negatively skewed (tail to the left).
- Larger n: the peak moves right (mean np), the spread grows (sd $\sqrt{np(1-p)}$) and the shape becomes more symmetric.
- The most likely value (the mode) is close to the mean np.

Worked example 1 **Exactly, at least, at most** — 4 marks

30% of customers buy a coffee. Ten customers are chosen at random. Find the probability that (a) exactly 3 buy coffee, (b) at least 3 buy coffee, (c) given at least one buys coffee, at least 3 do.

1 (a) $\binom{10}{3}(0.3)^3(0.7)^7 = 120 \times 0.027 \times 0.0823543 \approx 0.2668$. 2 (b) $P(X \geq 3) = 1 - P(X \leq 2) \approx 1 - 0.3828 = 0.6172$ (CAS: binomCdf). 3 (c) $P(X \geq 3 \mid X \geq 1) = \dfrac{P(X \geq 3)}{P(X \geq 1)} = \dfrac{0.61722}{1 - 0.7^{10}} = \dfrac{0.61722}{0.97175} \approx 0.6352$ (keep unrounded values until the end).

(a) 0.2668, (b) 0.6172, (c) 0.6352. In (c), "at least 3" is already inside "at least 1", so the intersection is just $P(X \geq 3)$.

Worked example 2 **Find the smallest n (Exam 2 staple)** — 3 marks

A machine produces faulty items with probability 0.1. How many items must be tested so that the probability of finding at least one faulty item exceeds 0.95?

1 $P(X \geq 1) = 1 - 0.9^n > 0.95 \Rightarrow 0.9^n < 0.05$. 2 $n > \dfrac{\log_e 0.05}{\log_e 0.9} \approx 28.43$. (The inequality flips because $\log_e 0.9 < 0$.)

$n = 29$. Always round *up* for "smallest n". Check: $1 - 0.9^{28} \approx 0.948$ (too small), $1 - 0.9^{29} \approx 0.953$ ✓.

Probability

Worked example 3 **An integer from a rounded probability** 2 marks

$X \sim \text{Bi}(n, 0.1)$. For which values of n is $P(X \geq 1) = 0.8$, correct to one decimal place?

1 $P(X \geq 1) = 1 - 0.9^n$. CAS: `seq(1-0.9^n,n,13,19)` gives 0.7458, 0.7712, 0.7941, 0.8147, 0.8332, 0.8499, 0.8649.

$n = 14, 15, 16, 17, 18$. $n = 18$ counts: 0.8499 rounds to 0.8. Give every value unless the question asks for the smallest or largest.

Worked example 4 **Find n and p from the mean and variance** 2 marks

$X \sim \text{Bi}(n, p)$ with $E(X) = 6$ and $\text{Var}(X) = 4.2$. Find n and p.

1 Divide: $\dfrac{np(1-p)}{np} = \dfrac{4.2}{6} \Rightarrow 1 - p = 0.7$.

$p = 0.3$, $n = \dfrac{6}{0.3} = 20$.

Finding p from a probability

Write the probability out with p and solve. E.g. $X \sim \text{Bi}(10, p)$ with $P(X = 0) = 0.2$: $(1-p)^{10} = 0.2$, so $p = 1 - 0.2^{0.1} \approx 0.1487$. CAS: `solve((1-p)^10=0.2,p)|0<p<1`. If CAS gives more than one value, keep those in $(0, 1)$.

Trick alert: binomial traps

- **"At least 1" =** $1 - P(X = 0)$. Never add up 1, 2, 3, … by hand.
- **Discrete inequalities:** $P(X < 4) = P(X \leq 3)$, and $P(X > 4) = 1 - P(X \leq 4)$. Off-by-one errors are the most common binomial mistake.
- **Without replacement from a small population isn't binomial** (the probability changes each trial), and neither are independent trials with different probabilities: multiply along each arrangement instead (see Probability → **p. 38**).
- **Conditional binomial:** the numerator is the intersection. $P(X = 2 \mid X \geq 1) = \dfrac{P(X = 2)}{P(X \geq 1)}$ because $X = 2$ is inside $X \geq 1$.

Hard exam questions

2023 Exam 2 · Section A · Question 8 © VCAA **49% correct**

The question (not reprinted in this copy): Probability of at least one green in 8 draws with replacement
Full question: 2023 Exam 2, Section A, Question 8 (page 4 of the paper).

Answer: C · A 12% · B 14% · C 49% · D 16% · E 9%

$1 - \Pr(\text{no green}) = 1 - \left(\dfrac{m}{n+m}\right)^8$.

DATA ANALYSIS, PROBABILITY & STATISTICS

Continuous random variables (pdfs)

Probability density functions

A continuous random variable takes any value in an interval. Its **probability density function** f satisfies:

Conditions for a pdf

$$f(x) \geq 0 \text{ for all } x \qquad \int_{-\infty}^{\infty} f(x)\,dx = 1 \qquad P(a < X < b) = \int_a^b f(x)\,dx$$

$P(X = a) = 0$ for any single value, so $<$ and $\leq$ give the same probability. (For discrete variables they don't!)

Summary statistics

$$E(X) = \int x f(x)\,dx \qquad \text{Var}(X) = \int x^2 f(x)\,dx - \mu^2 \qquad \int_{-\infty}^{m} f(x)\,dx = \frac{1}{2}\ (\text{median } m)$$

The pth percentile q satisfies $\int_{-\infty}^{q} f(x)\,dx = \dfrac{p}{100}$. $E(aX + b) = aE(X) + b$ and $\text{Var}(aX + b) = a^2\,\text{Var}(X)$, exactly as for discrete variables. If no rounding is asked for, give exact values: for $f(x) = \dfrac{x^2}{9}$ on $[0, 3]$, $\text{Var}(X) = \dfrac{27}{80}$ and $\text{sd}(X) = \dfrac{3\sqrt{15}}{20}$, not 0.581.


Probability

Worked example 1 **Everything about f(x) = kx² on [0, 3]** 6 marks

$f(x) = kx^2$ for $0 \le x \le 3$, and 0 elsewhere. Find (a) k, (b) $P(X < 1)$, (c) $E(X)$, (d) the median, (e) $\text{Var}(X)$.

1 (a) $\int_0^3 kx^2\,dx = 9k = 1 \Rightarrow k = \frac{1}{9}$. 2 (b) $\int_0^1 \frac{x^2}{9}dx = \frac{1}{27}$. 3 (c) $E(X) = \int_0^3 \frac{x^3}{9}dx = \frac{81}{36} = \frac{9}{4}$. 4 (d) $\int_0^m \frac{x^2}{9}dx = \frac{m^3}{27} = \frac{1}{2} \Rightarrow m = \frac{3}{\sqrt[3]{2}} \approx 2.381$. 5 (e) $E(X^2) = \int_0^3 \frac{x^4}{9}dx = \frac{243}{45} = 5.4$; $\text{Var} = 5.4 - \left(\frac{9}{4}\right)^2 = 0.3375$.

✓ $k = \frac{1}{9}$; $P(X < 1) = \frac{1}{27}$; $E(X) = 2.25$; **median** ≈ 2.381; $\text{Var} = \frac{27}{80} = 0.3375$. The median is bigger than the mean here because the distribution is skewed left (bunched at high values).



$f(x) = \frac{x^2}{9}$ on $[0, 3]$. The shaded area is $P(X < 2) = \frac{8}{27}$ (the worked example finds $P(X < 1) = \frac{1}{27}$); the dashed lines mark the mean (2.25) and the median (≈ 2.38).

Worked example 2 **A piecewise (triangular) pdf** 4 marks

$f(x) = \begin{cases} x, & 0 \le x \le 1 \\ 2 - x, & 1 < x \le 2 \\ 0, & \text{elsewhere} \end{cases}$. Find $P(X < 0.5)$, $E(X)$ and $P(X > 1.5 \mid X > 1)$.

1 $P(X < 0.5) = \int_0^{0.5} x\,dx = 0.125$. 2 The graph is symmetric about $x = 1$, so $E(X) = 1$ (and the median is 1). 3 $P(X > 1.5) = \int_{1.5}^2 (2 - x)\,dx = 0.125$ and $P(X > 1) = 0.5$.

✓ **0.125; 1;** $\frac{0.125}{0.5} = 0.25$**.** For a piecewise pdf, integrate each piece over its own interval.

Worked example 3 **Exponential-type pdf (CAS)** 3 marks

The time T (minutes) between customer arrivals has pdf $f(t) = 0.5e^{-0.5t}$, $t \ge 0$. Find $P(T > 2)$ and the median time.

1 $P(T > 2) = \int_2^\infty 0.5e^{-0.5t}dt = \left[-e^{-0.5t}\right]_2^\infty = e^{-1}$. 2 Median: $\int_0^m 0.5e^{-0.5t}dt = 1 - e^{-0.5m} = \frac{1}{2} \Rightarrow m = 2\log_e 2$.

✓ $P(T > 2) = e^{-1} \approx 0.368$**; median** $= 2\log_e 2 \approx 1.386$ **min** (less than the mean of 2 min: the long right tail pulls the mean up).

Worked example 4 **Median of a piecewise pdf, and the mode** 3 marks

X has pdf $f(x) = \frac{2x}{3}$ for $0 \le x \le 1$, $f(x) = \frac{3 - x}{3}$ for $1 < x \le 3$, and 0 elsewhere. Find the median and the mode.

1 $\int_0^1 \frac{2x}{3}\,dx = \frac{1}{3} < \frac{1}{2}$, so the median is in the second piece. 2 $\frac{1}{3} + \int_1^m \frac{3 - x}{3}\,dx = \frac{1}{2} \Rightarrow 3m - \frac{m^2}{2} - \frac{5}{2} = \frac{1}{2} \Rightarrow m^2 - 6m + 6 = 0 \Rightarrow m = 3 \pm \sqrt{3}$.

✓ **Median** $3 - \sqrt{3} \approx 1.27$ (the other root is outside $[1, 3]$); **mode 1**, where f peaks at $\frac{2}{3}$.

Transforming a pdf

If f is a pdf, then $g(x) = k\,f\left(\frac{x}{n}\right)$ (with $n > 0$) is a pdf only if $kn = 1$, because the dilation by n from the y-axis multiplies the area by n. Then g is the pdf of nX: its mean is n times the mean of f, and its standard deviation is n times the standard deviation of f.

Trick alert: pdf traps

- $f(x)$ **isn't a probability.** A pdf value can be bigger than 1; only areas are probabilities.
- **The median is where the area is** $\frac{1}{2}$, not where f is $\frac{1}{2}$, and not halfway along the domain.
- **For a pdf, "find** k**" needs both conditions:** total area 1 *and* $f \ge 0$ (sometimes this rules out a solution).
- **Conditional:** $P(X > a \mid X > b) = \frac{P(X > a)}{P(X > b)}$ only when $a \ge b$ (so the first event is inside the second).

DATA ANALYSIS, PROBABILITY & STATISTICS

The normal distribution

The normal curve

$X \sim \mathrm{N}(\mu, \sigma^2)$ has a symmetric bell-shaped pdf centred at μ (mean = median = mode), with spread set by the standard deviation σ. Note that the second parameter is the **variance**: $\mathrm{N}(170, 64)$ means $\sigma = 8$.

Standardising

$$Z = \frac{X - \mu}{\sigma} \sim \mathrm{N}(0, 1)$$

A z-score counts how many standard deviations a value is from the mean.

Interval	$\mu \pm \sigma$	$\mu \pm 2\sigma$	$\mu \pm 3\sigma$
Approx. probability	68%	95%	99.7%

How μ and σ change the graph

Changing μ translates the curve left or right without changing its shape. A larger σ gives a lower, wider curve and a smaller σ a taller, narrower one: the total area is always 1. The peak height is $\frac{1}{\sigma\sqrt{2\pi}}$, and the points of inflection are at $x = \mu \pm \sigma$. The standard normal $Z \sim \mathrm{N}(0, 1)$ has $\mu = 0$ and $\sigma = 1$.

Worked example 1 **Probability and percentile** — 3 marks

Heights of adult men are $X \sim \mathrm{N}(170, 8^2)$ cm. Find (a) $P(X > 180)$, (b) the height exceeded by only 10% of men.

1 (a) $z = \frac{180 - 170}{8} = 1.25$; $P(Z > 1.25) \approx 0.1056$. 2 (b) Need $P(X > h) = 0.1$, i.e. $P(X < h) = 0.9$. Inverse normal: $z = 1.2816$, so $h = 170 + 1.2816 \times 8$.

(a) 0.1056, (b) $h \approx 180.25$ **cm.** CAS inverse normal uses the area to the *left* by default: convert "top 10%" into 0.9.

Worked example 3 **Finding σ** — 2 marks

The mass of a packet is normally distributed with mean 50 g, and 10% of packets weigh less than 40 g. Find the standard deviation.

1 $P(X < 40) = 0.1 \Rightarrow \frac{40 - 50}{\sigma} = -1.2816$.

$\sigma = \frac{10}{1.2816} \approx 7.80$ **g.**

Worked example 4 **Finding μ and σ together (hard)** — 4 marks

For a normal variable, $P(X < 30) = 0.2$ and $P(X > 60) = 0.1$. Find μ and σ.

1 $\frac{30 - \mu}{\sigma} = z_{0.2} = -0.8416$ and $\frac{60 - \mu}{\sigma} = z_{0.9} = 1.2816$. 2 So $30 = \mu - 0.8416\sigma$ and $60 = \mu + 1.2816\sigma$. Subtract: $30 = 2.1232\sigma$.

$\sigma \approx 14.13$, $\mu \approx 30 + 0.8416 \times 14.13 \approx 41.89$.

Worked example 5 **Conditional normal, then binomial (Exam 2 chain)** — 4 marks

Using $X \sim \mathrm{N}(170, 8^2)$: (a) find $P(X > 180 \mid X > 170)$; (b) ten men are chosen at random. Find the probability that at least two are taller than 180 cm.

1 (a) $\frac{P(X > 180)}{P(X > 170)} = \frac{0.1056}{0.5} \approx 0.2113$. 2 (b) $Y \sim \mathrm{Bi}(10, 0.1056)$: $P(Y \geq 2) = 1 - P(Y = 0) - P(Y = 1)$.

(a) 0.2113; (b) ≈ 0.2858**.** Use the unrounded probability from the normal part in the binomial part.

Trick alert: normal traps

- $\mathrm{N}(\mu, \sigma^2)$**: the second number is the variance.** $\mathrm{N}(10, 9)$ has $\sigma = 3$.
- **Inverse normal = area to the left.** "Top 5%" means use 0.95.
- **Symmetry shortcuts:** $P(Z < -a) = P(Z > a)$, and $P(X > \mu) = 0.5$.
- **The 68–95–99.7 rule is approximate**, so use it only when told to (or tech-free).

Probability

DATA ANALYSIS, PROBABILITY & STATISTICS

Sampling & sample proportions

Populations, samples and bias

A **parameter** describes a population (e.g. the true proportion p of voters who support a policy). A **statistic** is calculated from a sample (e.g. the sample proportion $\hat{p}$). Different samples give different $\hat{p}$ values, so $\hat{p}$ is a random variable.

- **Simple random sample:** every group of n members is equally likely to be chosen.
- **Bias** is a systematic tendency to over- or under-estimate. Examples: surveying only people at a gym about exercise habits (selection bias); voluntary online polls (self-selection bias); leading questions.
- A bigger sample reduces random variability, but it *doesn't* fix bias.

The distribution of $\hat{p}$

Sample proportion

$$\hat{P} = \frac{X}{n},\ X \sim \text{Bi}(n, p) \qquad E(\hat{P}) = p \qquad \text{sd}(\hat{P}) = \sqrt{\frac{p(1-p)}{n}}$$

For large n (a common guide is $np \geq 10$ and $n(1-p) \geq 10$), $\hat{P}$ is approximately $\text{N}\left(p, \frac{p(1-p)}{n}\right)$.

Key idea

Quadrupling the sample size **halves** the standard deviation of $\hat{P}$ (because of the $\sqrt{n}$). Diminishing returns: going from $n = 100$ to $n = 400$ helps as much as going from 400 to 1600.

Simulating sample proportions

A simulation draws many random samples of size n from a population with proportion p and records $\hat{p}$ for each. A dot plot or histogram of the results shows the distribution of $\hat{P}$: centred near p, with spread close to $\sqrt{\frac{p(1-p)}{n}}$, and more bell-shaped as n grows. A bigger n shrinks the spread; with p near 0 or 1, small samples give a skewed distribution.

Simulated confidence intervals show the other key idea: every sample gives a different interval, and about 95% of 95% intervals contain the true p. On the TI-Nspire CX II CAS, `randBin(100,0.3,50)/100` gives 50 simulated sample proportions for $n = 100$ and $p = 0.3$.

Reading a simulation

From a dot plot of simulated $\hat{p}$ values: **estimate a probability** by counting, e.g. if 7 of 50 simulated values are at least 0.4, $P(\hat{P} \geq 0.4) \approx \frac{7}{50} = 0.14$; **estimate** p by the centre of the plot; **compare sample sizes** by spread (quadrupling n halves the standard deviation). A simulation only estimates: the exact value comes from the binomial distribution of $X = n\hat{P}$.

Worked example 1 **Mean and sd of p̂** 2 marks

30% of a town's households own a dog. Random samples of 100 households are taken. Find the mean and standard deviation of $\hat{P}$.

1 $E(\hat{P}) = p = 0.3$.

$\text{sd}(\hat{P}) = \sqrt{\frac{0.3 \times 0.7}{100}} \approx 0.0458$.

Worked example 2 **Exact vs approximate (a favourite trap)** 4 marks

With $p = 0.3$ and $n = 100$, find $P(\hat{P} > 0.35)$ (a) exactly, using the binomial distribution, and (b) using the normal approximation.

1 (a) $\hat{P} > 0.35 \iff X > 35 \iff X \geq 36$. With $X \sim \text{Bi}(100, 0.3)$: $P(X \geq 36) \approx 0.1161$. 2 (b) $\hat{P} \approx \text{N}(0.3, 0.0458^2)$: $P(\hat{P} > 0.35) = P(Z > 1.091) \approx 0.1376$.

(a) 0.1161 (exact); (b) 0.1376 (approximate). Read which one the question asks for. For the exact version, the strict ">" matters: $X \geq 36$, not $X \geq 35$.

Worked example 3 **Exact distribution of p̂ for a small sample (tech-free)** 3 marks

A fair coin is tossed 4 times and $\hat{P}$ is the proportion of heads. Find $P(\hat{P} \geq 0.75)$.

1 $\hat{P} \geq 0.75 \iff X \geq 3$ with $X \sim \text{Bi}(4, \frac{1}{2})$. 2 $P(X = 3) + P(X = 4) = \frac{4}{16} + \frac{1}{16}$.

$\frac{5}{16}$. Small samples: convert to a count and use the binomial.

Probability

Trick alert: sampling traps

- p **vs** $\hat{p}$**:** the formula for $\mathrm{sd}(\hat{P})$ uses the population p when it's known.
- **Convert proportions to counts carefully:** $\hat{P} \leq 0.2$ with $n = 50$ is $X \leq 10$.
- **Bigger samples don't remove bias.** A huge biased sample is still biased.
- **sd of** $\hat{P}$ **depends on** $\sqrt{n}$**, not** n**.**

DATA ANALYSIS, PROBABILITY & STATISTICS

Confidence intervals for proportions

The interval

Approximate confidence interval for p

$$\left(\hat{p} - z\sqrt{\frac{\hat{p}(1-\hat{p})}{n}},\ \hat{p} + z\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}\right)$$

$z = 1.96$ for 95% (≈ 2 tech-free), $z \approx 2.576$ for 99%, $z \approx 1.645$ for 90%. The **margin of error** is $M = z\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$, half the width.

What "95% confident" means

If we took many samples and built a 95% interval from each, about 95% of those intervals would contain the true p. The true p is fixed: any single interval either contains it or doesn't.

Worked example 1 **Building a 95% interval** 3 marks

In a random sample of 200 students, 60 walk to school. Find an approximate 95% confidence interval for the proportion of all students who walk, to 4 decimal places.

1 $\hat{p} = \frac{60}{200} = 0.3$. Standard error $= \sqrt{\frac{0.3 \times 0.7}{200}} \approx 0.03240$. 2 $M = 1.96 \times 0.03240 \approx 0.0635$.

$(0.2365,\ 0.3635)$.

Worked example 3 **Working backwards from an interval (hard)** 3 marks

An approximate 95% confidence interval for p is $(0.52, 0.64)$. Find $\hat{p}$, the margin of error and the sample size.

1 $\hat{p}$ is the midpoint: $\frac{0.52 + 0.64}{2} = 0.58$. M is half the width: 0.06. 2 $0.06 = 1.96\sqrt{\frac{0.58 \times 0.42}{n}} \Rightarrow n = \frac{1.96^2 \times 0.2436}{0.06^2} \approx 259.9$.

$\hat{p} = 0.58$, $M = 0.06$, $n = 260$.

Worked example 4 **Finding the confidence level** 2 marks

A sample of 200 gives $\hat{p} = 0.3$ and the interval $(0.2467,\ 0.3533)$. Find the level of confidence.

1 $M = 0.0533$; SE $= \sqrt{\frac{0.3 \times 0.7}{200}} \approx 0.03240$; $z = \frac{0.0533}{0.03240} \approx 1.645$.

$P(-1.645 < Z < 1.645) \approx 0.90$: **a 90% confidence interval**. On CAS: `normCdf(-1.645,1.645)`.

Worked example 5 **Choosing a sample size** 2 marks

A pollster wants a 95% margin of error of at most 3 percentage points and has no idea of p. How many people should be surveyed?

1 Use $\hat{p} = 0.5$ (which gives the largest possible margin): $1.96\sqrt{\frac{0.25}{n}} \leq 0.03 \Rightarrow n \geq \frac{1.96^2 \times 0.25}{0.03^2} \approx 1067.1$.

$n = 1068$ (round up). This is why national polls often survey about 1000 people.

Using an interval to judge a claim

If a claimed value of p lies **inside** the interval, the data are consistent with the claim. If it's **outside**, the claim is doubtful (at that confidence level). An interval never *proves* the claim.

Worked example 6 **How many intervals contain p?** 2 marks

Twenty independent samples each give an approximate 95% confidence interval for p. Find the probability that all 20 contain p, and that at least 18 do.

1 $P(Y = 20) = 0.95^{20}$; $P(Y \geq 18)$ = `binomCdf(20,0.95,18,20)`.

$0.95^{20} \approx 0.3585$; $P(Y \geq 18) \approx 0.9245$. On average 19 of the 20 contain p.

Probability

Trick alert: confidence interval traps

- **Don't say "there's a 95% probability that p is in (0.24, 0.36)".** VCAA wants the "95% of such intervals" interpretation.
- **Higher confidence → wider interval; larger n → narrower.** To halve the width, quadruple n.
- **Round sample sizes up when designing for a margin of at most M**, even from 1067.1. When recovering n from a given (rounded) interval, round to the nearest whole number instead.
- **The interval is about the population**, not about future samples.

Index

A

Algorithms
tracing pseudocode 25
accumulator loop 26
Antiderivatives (antidifferentiation)
standard antiderivatives 32
by recognition 33
f from f′ and a point 33
graphs of 33
Area
between curves 34
one integral, curves cross 34
signed vs actual 34
Asymptotes
exponential and log 17
hyperbola, truncus 14
tan 18
Average rate of change 29
Average value 34
over a period 34

B

Bayes' rule (false positives) 37
Bernoulli trials 39
Bias (sampling) 43
Binomial distribution
probabilities 39
smallest n 39
integer from a rounded probability 40
n and p from mean, variance 40
shape 39
CAS 6

C

CAS command names
solve, nSolve 4
zeros 4
factor, expand 4
propFrac, completeSquare 4
domain 4
piecewise 4
derivative 5
tangentLine, normalLine 5
fMax, fMin 5
integral 5
limit 5
seq, sum 6
det 4
exact, approx 3
binomPdf, binomCdf, nCr 6
normCdf, invNorm 6
randBin 7
zInterval_1Prop 7
ClearAZ 3
CAS commands
set up 3
algebra and parameters 4
functions and graphs 4
calculus 5
probability and statistics 6
traps 7
Chain, product, quotient rules 28
Change of base 18
Circular (trigonometric) functions
graphs 18
equations 19
models 20
exact values 18
Closest point, minimum distance 31
CAS 5
Common tangent 29
Composite functions
existence, rule 21
range 21
linear inside 21
Concavity 30
Conditional probability
rules 37
tree diagrams 37
from a DRV table 38
binomial 39
normal 42
pdf 41
pdf (CAS) 6
Confidence intervals
formula, margin of error 44
finding the level 44
sample size 44
judging a claim 44
z-values (CAS) 7
Confidence intervals containing p 44
Continuity 27
hybrid functions 10
Cubic
from turning points 12
factorising 11
shapes 11

D

Definite integrals
fundamental theorem, properties 33
signed vs actual area 34
Derivative
standard derivatives 28
from a table 28
of a hybrid 27
domain of f′ 27
graph of f′ 30
Differentiability 27
Dilations 14
Discrete random variables 38
Discriminant
tangent lines, number of solutions 24
parameter conditions 16
Domain
maximal (implied) 9
of a composite 21
of an inverse 21
of f′ 27

E

Endpoints (absolute max, min) 32
Even and odd functions 23
Exact answers (no rounding stated) 7
pdf 40
Exact values 18
table 9
Exact vs normal approximation ($\hat{p}$) 43
Expected value (mean)
DRV 38
binomial 40
pdf 40
sample proportion 43
Exponential functions
graphs 17
equations 17
growth and decay 17
fit $y = Ae^{kx}$ 18
Exponential pdf 41

F

Factor and remainder theorems 11
Families of functions
locus of a turning point 16
parameter conditions 16
Functional equations 23
Functions and relations, one-to-one 9

G

General solutions (trig) 19
Gradient table 30
Greatest difference between two models 32

H

Hidden quadratics
exponential 17
trig 19
biquadratic 12
Hybrid functions
continuity 10
smooth join 27
derivative 27
CAS 4

I

Independent events and trials 37
different probabilities 38
Inequalities
quadratic, hidden quadratic 19
smallest n 39
Inflection points 30
tan graphs 19
stationary 29
greatest rate of change 30
Integral of an inverse 22
Integration
fundamental theorem 33
areas 34
totals from rates 35
Inverse functions
rule, domain, branch 21
meeting the inverse 22
where two inverses meet 22
tangency with a parameter 16
Inverse transformation 15

K

Karnaugh maps 37

L

Law of total probability 37
Limits 27
Literal equations 24

Logarithms
laws 17
graphs 17
equations 17

M

Maximum and minimum
stationary points 29
on an interval 32
optimisation 31
Mean
DRV 38
binomial 40
pdf 40
normal 42
sample proportion 43
Median
pdf 41
CAS 6
Mode
binomial (CAS) 6
pdf 41
Modulus function 10

N

Net change from a rate 35
Newton's method 25
when it fails 25
Normal distribution
probability, percentile 42
unknown σ 42
unknown μ and σ 42
then binomial 42
shape 42
Number of solutions
polynomial with a parameter 12
simultaneous equations 24
show there are none 20

O

One-to-one
definition 9
needed for an inverse 21
Optimisation 31
Ordered pairs 9

P

Percentiles
normal 42
pdf 40
pdf (CAS) 6
Period and amplitude 18
periodic models 20
Periodic functions
f(x + p) = f(x) 23
integrals 34
Polynomials
shapes 11
sketching a quartic 12
Power functions 13
Probability density functions 40
transforming a pdf 41
Pseudocode (algorithms) 25
while loop 26

Q

Quadratic formula 8

R

Range
restricted domain 10
of a composite 21
sinusoid 18
Rates of change
average, instantaneous 29
greatest 30
totals from rates 35
Rectangle rule (left, right) 35
CAS 6
Reflections 14
when the order of points flips 15
Repeated roots
shapes 11
with a parameter 12
Rounding
integer from a rounded probability 40
CAS traps 7

S

Sample proportions
distribution of $\hat{p}$ 43
simulation 43
Sample size 44
Show that
no solutions 20
a symmetry property 23
Simulation
reading a simulation 43
CAS 7
Simultaneous equations
two unknowns 24
three unknowns 24
Sinusoidal models
reading a model 20
fitting a model 20
Sketching
cubic, quartic 11
log graph 17
trig graph 19
hyperbola 14
gradient function 30
Smooth join 27
Standard deviation
DRV 38
pdf (exact) 40
normal 42
sample proportion 43
Stationary points 29
number of, as k varies 16
Strictly increasing, decreasing 29
for all x (parameter) 16
CAS 5
Sum of solutions (trig) 20

T

Tangents and normals 29
through a point not on the curve 29
angle with the x-axis 29
Transformations
mapping, order 14
dash method 15
matching points 15
inverse transformation 15
of a pdf 41
Translations 14
Trapezium rule 35
over or under 36
Tree diagrams 37
Trend plus periodic models 20
Trig equations 19
Turning points (local max, min) 29
in terms of a parameter 16

V

Variance
DRV, Var(aX + b) 38
binomial 40
pdf 40

W

While loop 26
Without replacement 37

Z

z-values
normal 42
confidence levels 44
CAS 7