

## Heterotrophic Mass Cultures of *Chlorella Vulgaris* with Glucose Feeding in Fermenters<sup>\*</sup>

LIANG Shi-zhong<sup>1</sup>, ZHU Ming-jun<sup>1</sup>, MENG Hai-hua<sup>2</sup>, YIN Jian-yun<sup>2</sup>, CHEN Feng<sup>3</sup>

( 1. College of Food & Biological Eng., South China Univ. of Tech., Guangzhou 510640, China ;

2. Guangdong Jiangmen Center for Biotechnology, Jiangmen 529081, China ;

3. Dept. of Botany, The University of Hong Kong, Pokfulam Road, Hong Kong, China )

**Abstract :** *Chlorella vulgaris* was heterotrophically cultured in 5, 50, 200, 800, and 4 000 L stirred tank fermenters with glucose feeding, which gave significant improvement in cell density and productivity compared to batch culture. The highest cell dry weight mass concentration reached 43.31 g/L. The specific growth rate was 0.069 h<sup>-1</sup>. The cell productivity was 0.62 g/(L·h). The results suggest *Chlorella vulgaris* can be heterotrophically produced in stirred tank fermenters for commercial use.

**Key words :** *Chlorella vulgaris* ; stirred tank fermenter ; heterotrophic mass culture

**CLC number :** TQ 920.6      **Document code :** A      **Article ID :** 1000 - 565X(2000)12 - 0066 - 05

### 0 Introduction

Microalgae have been produced as health food, and have found applications in agri-culture, wastewater treatment and aquaculture and for the production of bioactives<sup>[1,2]</sup>. So there are considerable demands in market.

The production of microalgae has mainly employed photoautotrophic culture system. There are some drawbacks in such system, such as difficulty of maintaining monoculture, light limitation and lack of control of important environmental factors. Technical developments and improvements in photobioreactor design may overcome these problems, but this invariably requires significant extra initial capital investment<sup>[1,3]</sup>.

Heterotrophic culture system, which can overcome the above problems encountered in the photoautotrophic system and improve growth and productivity, may be used as a substitute for photoautotrophic culture system. Up to now, several kinds of microalgae strains with heterotrophic function and higher yield have been screened, among which *Chlorella vulgaris* is a typical one<sup>[4]</sup>. This paper reports the application of heterotrophic fed-batch culture of *Chlorella vulgaris* as a means for increasing

**Received date :** 2000 - 04 - 28

**\* Foundation item :** Supported by the Guangdong Scientific Research Funds

**Biography :** LIANG Shi-zhong ( born in 1943 ), male, professor, doctoral tutor; current research interests: microbial fermentation and biochemical engineering.

cell density and productivity.

## 1 Materials and Methods

### 1.1 Microorganism and Medium

The strains used were *Chlorella vulgaris* HKU-1 and 4 and the mutants generated in our laboratory. Cells were cultured in basal medium with 10 g/L glucose<sup>[5]</sup>.

### 1.2 Fed-batch Cultures with Glucose Feeding

The preculture broth was inoculated into fermenters ( 5 L Biostat B , 50 , 200 , 800 and 4 000 L were equipment of Guangdong Jiangmen Centre for Biotechnology ). When the residual glucose mass concentration decreased to 3 ~ 5 g/L , glucose was provided. The nutrients except for glucose were supplied at the same time. Dissolved oxygen concentration was controlled by increasing agitation speed and airflow. Aeration rate and the agitation speed were variable and initially set at 0.8 L/( L · min )( 0.8vvm ) and 300 r/min , respectively. Temperature was controlled ( 30 ± 1 ) °C .

### 1.3 Analytical Methods

Biomass concentration was determined by centrifuging a fixed volume of the culture , washing the pellet twice by resuspending and recentrifuging , and drying the final cell pellet to constant weight at 105 °C .

Glucose was analyzed with the method of 3,5-disalicylic acid<sup>[6]</sup>.

## 2 Results and Discussion

### 2.1 Fed-batch Culture in 5 L Fermenter

According to the result of flask culture , scale-up culture of *Chlorella vulgaris* was performed in 5 L fermenter. The preculture broth ( 300 mL ) was inoculated into 5 L fermenter containing 2.7 L of the medium. When the residual glucose mass concentration decreased to 3 ~ 5 g/L after 18 h , 600 g/L glucose was batch-fed.

As shown in Fig. 1 , with the fed-batch culture by glucose feeding , the cell growth reached maximum value 35.84 g/L in 64 h culture , which decreased to 32.87 g/L in the subsequent 6 h culture. The maximum specific growth rate was 0.057 h<sup>-1</sup>. The productivity was 0.56 g/( L · h ).

The growth yield coefficient  $Y_{X/S}$  based on glucose was 0.41 , which was comparable to other microbial systems.

### 2.2 Fed-batch Cultures in 50 L and 200 L Fermenter




Fig. 1 Growth and glucose consumption curve of the *Chlorella vulgaris* in 5 L stirred tank reactor culture

■—Glucose ; ◆—Biomass.

The good results stimulated us to perform cultures in 50 L and 200 L fermenter. The results of the cultures were shown in Fig.2 and Fig.3.




Fig.2 Growth and glucose consumption curve of the *Chlorella vulgaris* in 50 L stirred tank reactor culture

■—Glucose ; ◆—Biomass.




Fig.3 Growth and glucose consumption curve of the *Chlorella vulgaris* in 200 L stirred tank reactor culture

■—Glucose ; ◆—Biomass.

As shown in Fig.2, the growth reached 37.6 g/L in 68 h culture in 50 L bioreactor, which decreased to 34.75 g/L in the subsequent 4 h culture. The maximum specific growth rate was  $0.049 \text{ h}^{-1}$ . The productivity was  $0.55 \text{ g/(L} \cdot \text{h)}$ . The results cultured in 50 L bioreactor were comparable to that of 5 L bioreactor. The growth yield coefficient  $Y_{x/s}$  based on glucose was 0.47.

As shown in Fig.3, though the maximum specific growth rate was  $0.069 \text{ h}^{-1}$  which was lower than that of 5 L and 50 L bioreactor, the cell dry weight reached 43.31 g/L in 70 h culture which was higher than that of 5 L and 50 L bioreactor. The cell dry weight decreased to 39.31 g/L in the 4 h culture subsequently. The productivity was  $0.62 \text{ g/(L} \cdot \text{h)}$ , which was higher than that of 5 L and 50 L bioreactor. The growth yield coefficient  $Y_{x/s}$  based on glucose reached 0.48.

The results indicated that heterotrophic culture of *Chlorella vulgaris* was promising.

### 2.3 Mass Cultures in 800 L and 4 000 L Fermenter

As shown in Fig.4, cell mass concentration reached 32.42 g/L in 67 h culture, which was slightly lower than that of 5, 50 and 200 L. The maximum specific growth rate was  $0.047 \text{ h}^{-1}$ . The productivity was  $0.48 \text{ g/(L} \cdot \text{h)}$ . The growth yield coefficient  $Y_{x/s}$  based on glucose was 0.40.

As shown in Fig.5, cell mass concentration reached only 30.62 g/L in 63 h culture. The maximum specific growth rate was  $0.045 \text{ h}^{-1}$ . The productivity was  $0.37 \text{ g/(L} \cdot \text{h)}$  and the growth yield coefficient  $Y_{x/s}$  based on glucose was only 0.33, which was slightly lower than that of 5, 50, 200 and 800 L. It might result from inconvenient controls of the important environmental factors. Lower as the cell concentration was, it was still practical in large-scale industrial production.




Fig. 4 Growth and glucose consumption curve of the *Chlorella vulgaris* in 800 L stirred tank reactor culture

■—Glucose ; ◆—Biomass.




Fig.5 Growth and glucose consumption curve of the *Chlorella vulgaris* in 4 000 L stirred tank reactor culture

■—Glucose ; ◆—Biomass.

### 3 Conclusion

The heterotrophic culture of *Chlorella vulgaris* in 5 , 50 , 200 , 800 , and 4 000 L stirred tank reactors was successful and promising. The results demonstrate that the heterotrophic fed-batch culture of *Chlorella vulgaris* can increase the alga cell mass concentration and productivity on a large-scale. A cost-effective fermentation process has been established through high cell density strategies for the growth of selected *Chlorella vulgaris*. The results suggest that the development of industrial heterotrophic process for the production of *Chlorella vulgaris* is practical.

### References :

- [ 1 ] CHEN Feng , JOHNS M R. Heterotrophic growth of chlamydomonas reinhardtii on acetate in chemostat culture[ J ]. Process Biochemistry , 1996 , 31( 6 ) : 601 – 604.
- [ 2 ] CHEN Feng. High cell density culture of microalgae in heterotrophic growth[ J ]. Tibtech , 1996 , 14 , 421 – 426.
- [ 3 ] JOHN G D , ALEXANDER J T. An investigation of the heterotrophic culture of the green alga Tetraselmis[ J ]. Journal of Applied Phycology , 1996 , 8 : 73 – 77.
- [ 4 ] LIU Shi-ming , CHEN Feng , LIANG Shi-zhong. Researches on the heterotrophic culture of *Chlorella Vulgaris* [ J ]. Journal of South China University of Technology ( Natural Science Edition ) , 1999 , 27( 4 ) , 111 – 115.
- [ 5 ] OH-HAMA T , MIYACHI S. Microalgal biotechnology : Chlorella [ M ]. London : Cambridge University Press , 1988.
- [ 6 ] HUANG W K. Food test and analysis( in Chinese ) : Assay of reductional saccharide with the method of disalicylic acid[ M ]. Beijing : Light Industry Press , 1989.

# 发酵罐葡萄糖流加大规模异养培养小球藻<sup>\*</sup>

梁世中<sup>1</sup>, 朱明军<sup>1</sup>, 孟海华<sup>2</sup>, 尹建云<sup>2</sup>, 陈 峰<sup>3</sup>

(1. 华南理工大学 食品与生物工程学院, 广东 广州 510640; 2. 广东省江门生物技术开发中心, 广东 江门 529081; 3. 香港大学植物学系, 香港薄扶林道)

**摘要:**应用流加工艺在 5 50 200 800, 4 000 L 机械搅拌发酵罐中大规模异养培养小球藻。与间歇异养或光照自养相比, 流加培养大大地提高了细胞密度和生产率。最高细胞质量浓度达到 43.31 g/L, 比生长速率达到  $0.069\text{ h}^{-1}$ , 细胞生产率达到了  $0.62\text{ g/(L}\cdot\text{h)}$ 。结果表明, 利用传统的机械搅拌发酵罐大规模生产小球藻有良好的商业化前景。

**关键词:** 小球藻; 搅拌发酵罐; 大规模异养培养

中图分类号: TQ 920.6 文献标识码: A 文章编号: 1000-565X(2000)12-0066-05

收稿日期: 2000-04-28

<sup>\*</sup> 基金项目: 广东省科研基金资助项目

作者简介: 梁世中(1943-), 男, 教授, 博士生导师; 主要研究方向: 微生物发酵与生化工程。

# 发酵罐葡萄糖流加大规模异养培养小球藻

作者: [梁世中](#), [朱明军](#), [孟海华](#), [尹建云](#), [陈峰](#), [LIANG Shi-zhong](#), [ZHU Ming-jun](#), [MENG Hai-hua](#), [YIN Jian-yun](#), [CHEN Feng](#)

作者单位: [梁世中, 朱明军, LIANG Shi-zhong, ZHU Ming-jun \(华南理工大学, 食品与生物工程学院, 广东, 广州, 510640\)](#), [孟海华, 尹建云, MENG Hai-hua, YIN Jian-yun \(广东省江门生物技术开发中心, 广东, 江门, 529081\)](#), [陈峰, CHEN Feng \(香港大学植物学系, 香港薄扶林道\)](#)

刊名: [华南理工大学学报 \(自然科学版\)](#) 

英文刊名: [JOURNAL OF SOUTH CHINA UNIVERSITY OF TECHNOLOGY \(NATURAL SCIENCE EDITION\)](#)

年, 卷(期): 2000, 28 (12)

被引用次数: 3次

## 参考文献(6条)

1. [CHEN Feng;JOHNS M R](#) [Heterotrophic growth of chlamydomonas reinhardtii on acetate in chemostat culture](#)[外文期刊] 1996(06)
2. [CHEN Feng](#) [High cell density culture of microalgae in heterotrophic growth](#)[外文期刊] 1996(11)
3. [JOHN G D;ALEXANDER J T](#) [An investigation of the heterotrophic culture of the green alga Tetraselmis](#) 1996
4. [LIU Shi-ming;CHEN Feng;LIANG Shi-zhong](#) [Researches on the heterotrophic culture of Chlorella Vulgaris](#) 1999(04)
5. [OH-HAMA T;MIYACHI S](#) [Microalgal biotechnology:Chlorella](#) 1988
6. [HUANG W K](#) [Food test and analysis \(in Chinese\):Assay of reductional saccharide with the method of disalicylic acid](#) 1989

## 本文读者也读过(2条)

1. [景建克](#). [许倩倩](#). [刘硕](#). [陈灏](#). [闫海](#). [JING Jian-ke](#). [XU Qian-qian](#). [LIU Shuo](#). [CHEN Hao](#). [YAN Hai](#) [大规模异养发酵培养小球藻USTB-01研究](#)[期刊论文]-[现代化工](#)2008, 28(12)
2. [刘世名](#). [孟海华](#). [梁世中](#). [尹建云](#). [麦平珍](#) [生物反应器高密度异养培养小球藻](#)[期刊论文]-[华南理工大学学报](#)2000, 28(2)

## 引证文献(3条)

1. [韩兴梅](#). [李元广](#). [魏晓东](#). [李兴武](#). [孙勇如](#). [王义琴](#) [转免防御素基因小球藻培养的营养模式研究](#)[期刊论文]-[海洋科学](#) 2006(7)
2. [闫海](#). [张可毅](#). [王子敬](#). [高郁晨](#). [贾璇](#). [何欢聚](#) [小球藻培养与应用的研究进展](#)[期刊论文]-[中国科技成果](#) 2010(1)
3. [王素琴](#) [小球藻USTB-01的异养培养和叶黄素的提取](#)[学位论文]硕士 2007

本文链接: [http://d.wanfangdata.com.cn/Periodical\\_hnlgdxxb200012012.aspx](http://d.wanfangdata.com.cn/Periodical_hnlgdxxb200012012.aspx)