



Organoid | All MEKi | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
20
40
60
0
25
50
75
100
0
25
50
75
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Granularity
Neighbors
Other


Organoid | All MEKi | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Organoid
Organoid
Higher in MEKi
Lower in MEKi
0
5
10
15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Organoid | All MEKi | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
MPNST
Other
0
10
20
30
40
60
80
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Granularity
Neighbors
Other

Organoid | All MEKi | Across tumor types (shared)

No features with q < 0.05


Organoid | All MEKi | Across tumor types (differ)

No features with q < 0.05


Organoid | Binimetinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
2
4
6
8
0.0
2.5
5.0
7.5
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Granularity
Neighbors
Other


Organoid | Binimetinib 1 uM | Across patients


No features with q < 0.05



Organoid | Binimetinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
cNF
pNF
Higher in MEKi
Lower in MEKi
0
5
10
15
0.00
0.25
0.50
0.75
1.00
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Organoid | Binimetinib 1 uM | Across tumor types (shared)


No features with q < 0.05



Organoid | Binimetinib 1 uM | Across tumor types (differ)
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Differs by tumor type
Organoid
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Granularity



Organoid | Binimetinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
3
6
9
0
2
4
6
0.0
2.5
5.0
7.5
Features with q < 0.05
Feature category
Granularity
Intensity
Neighbors
Texture
Other


Organoid | Binimetinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Organoid
Organoid
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Other


Organoid | Binimetinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
MPNST
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Correlation
Granularity
Neighbors
Texture



Organoid | Binimetinib 10 uM | Across tumor types (shared)
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Organoid
Higher in MEKi
0.00
0.25
0.50
0.75
1.00
Features with q < 0.05
Feature category
Intensity


Organoid | Binimetinib 10 uM | Across tumor types (differ)
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Differs by tumor type
Organoid
0.00
0.25
0.50
0.75
1.00
Features with q < 0.05
Feature category
Granularity


Organoid | Mirdametinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
1
2
3
4
5
6
Features with q < 0.05
Feature category
Granularity
Intensity
Neighbors
Texture
Other

Organoid | Mirdametinib 1 uM | Across patients

No features with q < 0.05



Organoid | Mirdametinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
MPNST
Other
0
2
4
6
0
1
2
3
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Other


Organoid | Mirdametinib 1 uM | Across tumor types (shared)


No features with q < 0.05

Organoid | Mirdametinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Organoid | Mirdametinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
0
1
2
3
4
5
6
Features with q < 0.05
Feature category
Granularity
Intensity
Neighbors
Texture
Other


Organoid | Mirdametinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Organoid
Organoid
Higher in MEKi
Lower in MEKi
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Intensity
Texture
Other



Organoid | Mirdametinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
Other
0.00
0.25
0.50
0.75
1.00
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Granularity
Intensity
Texture
Other

Organoid | Mirdametinib 10 uM | Across tumor types (shared)

No features with q < 0.05


Organoid | Mirdametinib 10 uM | Across tumor types (differ)

No features with q < 0.05


Organoid | Trametinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
1
2
3
4
5
6
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Granularity
Neighbors
Other

Organoid | Trametinib 1 uM | Across patients

No features with q < 0.05



Organoid | Trametinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
MPNST
0.00
0.25
0.50
0.75
1.00
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Texture
Other

Organoid | Trametinib 1 uM | Across tumor types (shared)

No features with q < 0.05


Organoid | Trametinib 1 uM | Across tumor types (differ)




Organoid | Trametinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
1
2
3
4
5
6
Features with q < 0.05
Feature category
Granularity
Intensity
Neighbors
Texture
Other


Organoid | Trametinib 10 uM | Across patients


No features with q < 0.05



Organoid | Trametinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
cNF
Higher in MEKi
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Texture
Other


Organoid | Trametinib 10 uM | Across tumor types (shared)

No features with q < 0.05


Organoid | Trametinib 10 uM | Across tumor types (differ)
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Differs by tumor type
Organoid
0.00
0.25
0.50
0.75
1.00
Features with q < 0.05
Feature category
Granularity


Organoid | Selumetinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
0
1
2
3
4
Features with q < 0.05
Feature category
Granularity
Intensity
Neighbors
Texture
Other


Organoid | Selumetinib 1 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Organoid
Organoid
Higher in MEKi
Lower in MEKi
0
1
2
3
Features with q < 0.05
Feature category
Granularity
Intensity
Other



Organoid | Selumetinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
cNF
Higher in MEKi
cNF
Lower in MEKi
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Other


Organoid | Selumetinib 1 uM | Across tumor types (shared)


No features with q < 0.05

Organoid | Selumetinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Organoid | Selumetinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
5
10
15
0
1
2
3
4
5
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Granularity
Neighbors
Other


Organoid | Selumetinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Organoid
Organoid
Higher in MEKi
Lower in MEKi
0
1
2
3
4
5
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Organoid | Selumetinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
Other
0.00
0.25
0.50
0.75
1.00
0
1
2
3
4
5
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Organoid | Selumetinib 10 uM | Across tumor types (shared)


No features with q < 0.05

Organoid | Selumetinib 10 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | All MEKi | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
25
50
75
100
0
50
100
150
200
0
50
100
150
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | All MEKi | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
5
10
15
20
25
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Single cell | All MEKi | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
MPNST
Other
0
20
40
60
80
0
50
100
150
0
50
100
150
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | All MEKi | Across tumor types (shared)


No features with q < 0.05

Single cell | All MEKi | Across tumor types (differ)

No features with q < 0.05


Single cell | Binimetinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
2
4
6
0
5
10
0
5
10
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Binimetinib 1 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0.0
2.5
5.0
7.5
10.0
12.5
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Single cell | Binimetinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
Other
0
20
40
60
1
2
3
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other

Single cell | Binimetinib 1 uM | Across tumor types (shared)

No features with q < 0.05


Single cell | Binimetinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Binimetinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0.0
2.5
5.0
7.5
10.0
12.5
0
5
10
15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other

Single cell | Binimetinib 10 uM | Across patients

No features with q < 0.05



Single cell | Binimetinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
MPNST
Other
0.0
0.5
1.0
1.5
2.0
0
10
20
30
40
0
5
10
15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Binimetinib 10 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Binimetinib 10 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Mirdametinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0 1 2 3 4 5
0 3 6 9
0 5 10 15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Mirdametinib 1 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
5
10
15
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Single cell | Mirdametinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
Other
0.0
0.5
1.0
1.5
2.0
0
20
40
60
0
10
20
30
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Mirdametinib 1 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Mirdametinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Mirdametinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
1
2
3
4
5
0
2
4
6
8
0.0
2.5
5.0
7.5
Features with q < 0.05
Feature category
Granularity
Intensity
Texture
Other


Single cell | Mirdametinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
2
4
6
8
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Other



Single cell | Mirdametinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
MPNST
Other
0
10
20
30
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Mirdametinib 10 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Mirdametinib 10 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Trametinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
2
4
6
0.0
2.5
5.0
7.5
10.0
0
3
6
9
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Trametinib 1 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Intensity
Texture
Other



Single cell | Trametinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
Other
0
10
20
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Trametinib 1 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Trametinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Trametinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0.0
2.5
5.0
7.5
0
2
4
6
8
0.0
2.5
5.0
7.5
10.0
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Trametinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
5
10
15
0.0
0.5
1.0
1.5
2.0
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Single cell | Trametinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
Other
0
1
2
3
4
5
Features with q < 0.05
Feature category
Texture
Other


Single cell | Trametinib 10 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Trametinib 10 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Selumetinib 1 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0018_T6 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
2
4
6
0.0
2.5
5.0
7.5
10.0
12.5
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Selumetinib 1 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
5
10
15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other



Single cell | Selumetinib 1 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
pNF
0
20
40
60
0
5
10
15
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Selumetinib 1 uM | Across tumor types (shared)


No features with q < 0.05

Single cell | Selumetinib 1 uM | Across tumor types (differ)

No features with q < 0.05


Single cell | Selumetinib 10 uM | Within patient
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
NF0014_T1 (cNF)
NF0021_T1 (cNF)
NF0035_T1 (cNF)
NF0037_T1 (cNF)
NF0014_T2 (pNF)
NF0016_T1 (pNF)
NF0055_T1 (pNF)
SARCO219_T2 (MPNST)
SARCO361_T1 (MPNST)
NF0030_T1 (Other)
NF0040_T1 (Other)
0
2
4
6
8
0.0
2.5
5.0
7.5
0
3
6
9
Features with q < 0.05
Feature category
Granularity
Intensity
Texture
Other


Single cell | Selumetinib 10 uM | Across patients
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
Cell
Cytoplasm
Nuclei
0
5
10
15
20
0
1
2
3
4
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other


Single cell | Selumetinib 10 uM | Within tumor type
MEKi vs DMSO
MEKi vs Staurosporine
MEKi vs Digoxin
Higher in MEKi
Lower in MEKi
cNF
MPNST
Other
0.00
0.25
0.50
0.75
1.00
0
20
40
60
0
2
4
6
Features with q < 0.05
Feature category
Correlation
Granularity
Intensity
Texture
Other

Single cell | Selumetinib 10 uM | Across tumor types (shared)

No features with q < 0.05

Single cell | Selumetinib 10 uM | Across tumor types (differ)

No features with q < 0.05