
Organoid
one point per patient x treatment
Morphology mean |normalized difference| vs DMSO
0.6
0.4
0.2
−4
−2
0
Viability log2FC vs DMSO
Tumor type
cNF
pNF
MPNST
Treatment
Other
MEK inhibitor


Single cell
one point per patient x treatment
Morphology mean |normalized difference| vs DMSO
0.9
0.6
0.3
−4
−2
0
Viability log2FC vs DMSO
Tumor type
cNF
pNF
MPNST
Treatment
Other
MEK inhibitor


Organoid
one point per treatment x tumor type, mean over patients
Mean morphology mean |normalized difference| vs DMSO
0.5
0.4
0.3
0.2
0.1
Trabectedin 1 uM
Fimepinostat 1 uM
Staurosporine 10 nM
Onalespib 1 uM
Digoxin 1 uM
Copanlisib 1 uM
Sapanisertib 1 uM
−5
−4
−3
−2
−1
0
1
Mean viability log2FC vs DMSO
Tumor type
cNF
pNF
MPNST
Treatment
Other
MEK inhibitor


Single cell
one point per treatment x tumor type, mean over patients
Mean morphology mean |normalized difference| vs DMSO
1.25
1.00
0.75
0.50
0.25
Trabectedin 1 uM
Staurosporine 10 nM
Onalespib 1 uM
Digoxin 1 uM
Fimepinostat 1 uM
Copanlisib 1 uM
Sapanisertib 1 uM
−5
−4
−3
−2
−1
0
1
Mean viability log2FC vs DMSO
Tumor type
cNF
pNF
MPNST
Treatment
Other
MEK inhibitor