**Lemma B.2.** Let  $\pi: \mathcal{Y} \to \mathcal{W}$  be a morphism of ringed topoi. Let  $\mathcal{B}$  be a sheaf of  $\mathcal{O}_{\mathcal{W}}$ -algebras. Let  $\mathcal{W}'$  and  $\mathcal{Y}'$  be the ringed topoi  $(\mathcal{W}, \mathcal{B})$  and  $(\mathcal{Y}, \pi^*\mathcal{B})$ , respectively. There is an induced 2-commutative diagram of ringed topoi:

$$\begin{array}{ccc}
\mathcal{Y}' & \xrightarrow{\pi'} \mathcal{W}' \\
\downarrow^{j'} & & \downarrow^{j} \\
\mathcal{Y} & \xrightarrow{\pi} \mathcal{W}.
\end{array}$$

If  $\pi$  and j are tor-independent and  $\mathbb{N} \in D(\mathcal{W}')$ , then there is a natural isomorphism:

$$L\pi^*Rj_*\mathcal{N} \simeq Rj'_*L\pi'^*\mathcal{N}.$$

In particular, if  $Q \in \langle Rj_*D_{pc}^-(W')\rangle$ , then  $L\pi^*Q \in \langle Rj_*'D_{pc}^-(Y')\rangle$ .

*Proof.* Now  $j_*, j'_*$  are exact and  $\pi^{-1}j_* = j'_*\pi'^{-1}$ . By tor-independence of  $\pi$  and j:

$$\begin{split} \mathsf{L}\pi^*\mathsf{R} j_* & \mathcal{N} = \mathcal{O}_{\mathcal{Y}} \otimes_{\pi^{-1}\mathcal{O}_{\mathcal{W}}}^{\mathsf{L}} \pi^{-1} j_* \mathcal{N} \simeq (\mathcal{O}_{\mathcal{W}} \otimes_{\pi^{-1}\mathcal{O}_{\mathcal{Y}}}^{\mathsf{L}} \pi^{-1} j_* \mathcal{O}_{\mathcal{Y}'}) \otimes_{\pi^{-1} j_* \mathcal{O}_{\mathcal{Y}'}}^{\mathsf{L}} \pi^{-1} j_* \mathcal{N} \\ & \simeq j_*' \mathcal{O}_{\mathcal{Y}'} \otimes_{\pi^{-1} j_* \mathcal{O}_{\mathcal{Y}'}}^{\mathsf{L}} \pi^{-1} j_* \mathcal{N} \\ & \simeq j_*' \mathcal{O}_{\mathcal{Y}'} \otimes_{j_*' \pi'^{-1} \mathcal{O}_{\mathcal{W}'}}^{\mathsf{L}} j_*' \pi'^{-1} \mathcal{N} \\ & \simeq j_*' (\mathcal{O}_{\mathcal{Y}'} \otimes_{\pi'^{-1} \mathcal{O}_{\mathcal{W}'}}^{\mathsf{L}} \pi'^{-1} \mathcal{N}) = \mathsf{R} j_*' \mathsf{L} \pi'^* \mathcal{N}. \end{split}$$

The latter claim follows from a similar argument to that in Lemma B.1.  $\Box$ 

## References

- [AT18] D. Abramovich and M. Temkin, Functorial factorization of birational maps for qe schemes in characteristic 0, Algebra Number Theory (2018), accepted for publication.
- [BB03] A. Bondal and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, Mosc. Math. J. 3 (2003), no. 1, 1–36, 258
- [BJ14] B. Bhatt and A. J. de Jong, Lefschetz for local Picard groups, Ann. Sci. Éc. Norm. Supér. (4) 47 (2014), no. 4, 833–849.
- [BZNP17] D. Ben-Zvi, D. Nadler, and A. Preygel, Integral transforms for coherent sheaves, J. Eur. Math. Soc. (JEMS) 19 (2017), no. 12, 3763–3812.
- [CLO12] B. Conrad, M. Lieblich, and M. Olsson, Nagata compactification for algebraic spaces, J. Inst. Math. Jussieu 11 (2012), no. 4, 747–814.
- [Con06] B. Conrad, Relative ampleness in rigid geometry, Ann. Inst. Fourier (Grenoble) 56 (2006), no. 4, 1049–1126.
- [Con10] B. Conrad, Universal property of non-archimedean analytification, available on homepage, August 2010.
- [CT09] B. Conrad and M. Temkin, Non-Archimedean analytification of algebraic spaces, J. Algebraic Geom. 18 (2009), no. 4, 731–788.
- [EGA] A. Grothendieck, Éléments de géométrie algébrique, I.H.E.S. Publ. Math. 4, 8, 11, 17, 20, 24, 28, 32 (1960, 1961, 1961, 1963, 1964, 1965, 1966, 1967).
- [FF18] L. Fargues and J.-M. Fontaine, Courbes et fibrés vectoriels en théorie de Hodge padique, Astérisque (2018), no. 406, xiii+382, With a preface by Pierre Colmez.
- [FHM03] H. Fausk, P. Hu, and J. P. May, Isomorphisms between left and right adjoints, Theory Appl. Categ. 11 (2003), No. 4, 107–131.
- [FK13] K. Fujiwara and F. Kato, Foundations of Rigid Geometry I, August 2013, arXiv:1308.4734.
- [FL85] W. Fulton and S. Lang, Riemann-Roch algebra, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 277, Springer-Verlag, New York, 1985.
- [Gab62] P. Gabriel, Des catégories abéliennes, Bull. Soc. Math. France 90 (1962), 323-448.
- [GAGA] J. P. Serre, Géométrie algébrique et géométrie analytique, Ann. Inst. Fourier, Grenoble 6 (1955–1956), 1–42.