

- [14]. Emara, W. M., Moez, K. K., & Elkhoully, A. H. (2014). Topical versus intravenous Tranexamic Acid as a blood conservation intervention for reduction of post-operative bleeding in Hemiarthroplasty. *Anesthesia, Essays and Researches*, 8(1), 48–53.
- [15]. Petajl J, Myllynen I, Myllyla G, Vahtera E. Fibrinolysis after application of a pneumatic tourniquet. *Acta Chir Stand* 1987; 153:647-51.
- [16]. Ahn J, Man L-X, Park S, Sodl JF, Esterhai JL. Systematic review of cemented and uncemented Hemiarthroplasty outcomes for femoral neck fractures. *Clin Orthop Relat Res* 2008;466 (10):2513–8.
- [17]. Bierbaum, B. E. et al: An analysis of blood management in patients having a total hip or knee arthroplasty. *JBJS [Am]*1999, 81-A(1): 2-10.
- [18]. Sehat KR, Evans RL, Newman JH. Hidden blood loss following hip and knee arthroplasty. Correct management of blood loss should take hidden loss into account. *J Bone Joint Surg Br.* 2004;86:561–5.
- [19]. Na HS, Shin HJ, Lee YJ, Kim JH, Koo KH, Do SH. The effect of Tranexamic Acid on blood coagulation in Total Hip Arthroplasty arthroplasty: Rotational thromboelastographic analysis. *Anaesthesia.* 2016;71(1):67–75.
- [20]. Georgiadis AG, Muh SJ, Silverton CD et al: A prospective double-blind placebo Controlled trial of topical Tranexamic Acid in Total Knee Arthroplasty. *J Arthroplasty*, 2013; 28(8): S78–82.
- [21]. Iwai T, Tsuji S, Tomita T, Sugamoto K, Hideki Y, Hamada M. Repeat-dose intravenous Tranexamic Acid further decreases blood loss in Total Knee Arthroplasty. *Int Orthop.* 2013;37(3):441-5.
- [22]. Cap AP, Baer DG, Orman JA, Aden J, Ryan K, Blackburne LH. Tranexamic Acid for trauma patients: a critical review of the literature. *J Trauma.* 2011;71:S9-14.
- [23]. Hynes MC, Calder P, Rosenfeld P, Scott G. The use of Tranexamic Acid to reduce blood loss during Total Hip Arthroplasty: An observational study. *Ann R Coll Surg Engl.* 2005;87(2):99–101.
- [24]. Sarzaem MM, Razi M, Kazemian G, Moghaddam ME, Rasi AM, Karimi M. Comparing efficacy of three methods of Tranexamic Acid administration in reducing hemoglobin drop following Total Knee Arthroplasty. *J Arthroplasty.* 2014;29(8):1521–4.