

AN ALEKSANDROV ESTIMATE FOR KINETIC EQUATIONS WITH ROUGH COEFFICIENTS

AMÉLIE LOHER, CONNOR MOONEY, CLÉMENT MOUHOT, LUKAS NIEBEL, LUIS SILVESTRE, AND YUZHE ZHU

ABSTRACT. We give an exposition of a proof, discovered by Astra, of an Aleksandrov estimate for kinetic equations with rough coefficients.

1. INTRODUCTION

Consider the parabolic equation

$$(1) \quad \partial_t u - \operatorname{tr}(A(t, x) D_x^2 u) = 0.$$

Here $(t, x) \in (-1, 0) \times B_1 \subset \mathbb{R}^{n+1}$ and $A \in \operatorname{Sym}_{n \times n}$ satisfies

$$(2) \quad \lambda I \leq A \leq \Lambda I \quad \text{for some } 0 < \lambda \leq \Lambda < \infty.$$

Krylov and Safonov proved that a Harnack inequality holds for solutions to (1), despite the fact that no regularity is assumed on the coefficients [12].

An intriguing question is whether this result can be extended to hypoelliptic equations of the type studied by Hörmander [10], but with rough coefficients. A positive answer to this question was recently given by Astra, for the important special case of the Kolmogorov kinetic equation with rough coefficients. Namely,

$$(3) \quad Ku := \partial_t u + v \cdot \nabla_x u - \operatorname{tr}(A(t, x, v) D_v^2 u) = 0.$$

Here $(t, x, v) \in (-1, 0) \times B_1 \times B_1 \subset \mathbb{R}^{2n+1}$ and the same assumptions as above hold on A . The purpose of this note is to give an exposition of the proof.

It is well-known to experts that most of the Krylov–Safonov proof goes through for (3). The key missing ingredient was the following Aleksandrov estimate:

Theorem 1.1. *If $Ku = f \geq 0$ in $Q_1 = (-1, 0) \times B_1 \times B_1$ and $u|_{\partial_{in} Q_1} = 0$, then*

$$(4) \quad \sup_{Q_1} u \leq C \|f\|_{L^p(Q_1)}$$

for some $C(n, \lambda, \Lambda) > 0$ and $p = p(n, \lambda, \Lambda) \in (1, \infty)$.

Here ∂_{in} denotes the incoming boundary. In particular, $\partial_{in} Q_1 = \{t = -1\} \cup \{|x| = 1, v \cdot x < 0\} \cup \{|v| = 1\}$.

In the parabolic case, the corresponding estimate can be established with $p = n + 1$. This is accomplished using convex analysis methods and the study of certain contact sets. However, these methods do not seem to adapt easily to the hypoelliptic setting. The approach to Theorem 1.1 is instead by duality. The estimate (4) is equivalent to a higher-integrability estimate for solutions to the adjoint equation. Namely, if

$$\begin{cases} \partial_t \rho + v \cdot \nabla_x \rho - \operatorname{div}_v \operatorname{div}_v (A(t, x, v) \rho) = f \geq 0 \text{ in } Q_1, \\ \rho|_{\partial_{in} Q_1} = 0, \end{cases}$$

then

$$(5) \quad \|\rho\|_{L^q(Q_1)} \leq C \|f\|_{L^1(Q_1)} \text{ for some } q(n, \lambda, \Lambda) > 1.$$

The exponent p in (4) is the Hölder conjugate of q . The adjoint viewpoint was also used in [7].

Remark 1.2. Technically, we wrote the backward (reflected in time and velocity) adjoint equation above. However, we will abuse terminology and simply refer to it as the adjoint equation in the sequel.

For the purposes of proving the Harnack inequality, it actually suffices to prove the weaker estimate

$$(6) \quad \int_{Q_1} \eta(\rho) \leq C(n, \lambda, \Lambda),$$

under the additional assumption that f has mass 1, for some universal function η satisfying

$$(7) \quad \lim_{s \rightarrow \infty} \frac{\eta(s)}{s} = \infty.$$

We briefly explain why in the next section. Most of our exposition will focus on proving (6) for a choice $\eta(s)$ that is essentially the same as $s \log(s)$, which for reasons explained in Section 3 is particularly convenient.

The proof uses divergence-form methods. The starting point is to study energy estimates for the adjoint equation. These contain a bad term involving a derivative of the coefficients, which at first seems insurmountable. However, it turns out that one can evolve in (x, v) by an auxiliary flow in such a way that this bad term is the dissipation rate of a bounded quantity. After integrating in the auxiliary time, the term becomes harmless.

We will first illustrate how the proof works for elliptic and parabolic equations, which are simpler because the auxiliary flow (standard heat flow) commutes with the adjoint equation. We will then discuss the proof of (6) for kinetic equations, which is more complicated because one must account for new terms in the “flowed” adjoint equation coming from commutators with the drift. Finally, at the end we will discuss how, with a small adaptation of the method, one can improve to the L^q estimate (5). We conclude with a discussion of the exponent.

Let us also mention related literature on the regularity theory of equations in nondivergence form. Classical works on Krylov–Safonov regularity theory are [19, 15, 3, 2, 12, 17]. The kinetic analogue of the Krylov–Safonov theorem was recorded as an open problem in [18]. In [1] a Harnack inequality under a Cordes–Landis condition or continuity assumption was proven. The one-dimensional stationary problem was treated in [20]. Under restrictions on the variables on which the coefficients depend, [13] proves a kinetic Aleksandrov estimate. A proof of the kinetic Krylov–Safonov theorem was recorded in [8].

Declaration of AI Use. The proof of the kinetic Krylov–Safonov theorem was found by OpenAI’s GPT-6 Astra Ultra on September 6, 2026. A link to the chat and the initial output is available here [14]. We do not claim credit for this result. Anyone could have obtained it by prompting the AI about it.

Remark 1.3. A similar proof of the Aleksandrov estimate was made public in [8] on September 14, 2026. The precise involvement of AI in their proof was not made public.

Acknowledgments. C. Mooney is supported by NSF Grant DMS-2143668. L. Silvestre is supported by NSF grant DMS-2350263. L. Niebel is supported by SNSF Starting Grant TMSGI2_226018.

2. FROM $L \log L$ ESTIMATE TO HARNACK INEQUALITY

Here we briefly explain why (6)–(7) suffice for proving the Harnack inequality. None of the key ideas are lost by restricting our discussion to the simpler case of elliptic equations. Below, we will always assume that A satisfies (2).

The crucial estimate in the proof of the Harnack inequality that uses Aleksandrov, sometimes called the “density estimate,” is the following:

$$\text{If } \operatorname{tr}(A(x)D^2u) \leq 0 \text{ in } B_1 \subset \mathbb{R}^n, \quad u \geq 0, \quad \text{and} \quad u(0) \leq 1,$$

then

$$(8) \quad |\{u < C\}| \geq C^{-1} \text{ for some } C(n, \lambda, \Lambda) > 0.$$

To see how this follows from (the elliptic analogue of) (6)–(7), note that by the maximum principle, $v := u + 2(|x|^2 - 1)$ is bounded from below by the solution w to

$$\begin{cases} \operatorname{tr}(A(x)D^2w) = 4n\Lambda\chi_{\{v < 0\}}, \\ w|_{\partial B_1} = 0. \end{cases}$$

Moreover, $|w(0)| \geq 1$. Thus, if ρ is the Dirichlet Green's function with pole at 0, we have

$$1 \leq C(n, \Lambda) \int_{\{v < 0\}} \rho.$$

It is a simple exercise in real analysis to conclude from (6) and (7) that

$$1 \leq C(n, \lambda, \Lambda) |\{v < 0\}| \leq C(n, \lambda, \Lambda) |\{u < 2\}|.$$

Remark 2.1. Technically we are applying (6) with measure data, but this can be justified by approximation.

3. OVERVIEW

We now give an informal overview of the proof. We do so in the simplest setting of elliptic equations, and we ignore technical issues like solution class, domains, etc.

Assume that $A \in [\lambda I, \Lambda I]$ and that $\rho > 0$ solves

$$(9) \quad -\operatorname{div} \operatorname{div}(A(x)\rho) = f.$$

Let η and ω be functions on $[0, \infty)$ such that η is convex and

$$(10) \quad \eta''(s)s^2 \leq \omega(s), \quad s > 0.$$

If we test (9) with $\eta'(\rho)$, integrate by parts, and use Cauchy–Schwarz, we get

$$(11) \quad \int \eta''(\rho) |\nabla \rho|^2 \leq C(\lambda) \int \omega(\rho) |\operatorname{div} A|^2 + C(\lambda) \int f \eta'(\rho).$$

At first sight this estimate seems useless, since we have no control on the derivatives of A . The idea is to identify the left side and the first term on the right side as dissipation rates of entropy-like quantities along an auxiliary heat flow.

Let the subscript h denote evolution by heat flow for time h . Flowing both sides of (9) gives

$$(12) \quad -\operatorname{div} \operatorname{div}(B^h \rho_h) = f_h, \quad B^h := \frac{(A\rho)_h}{\rho_h} \in [\lambda I, \Lambda I].$$

The idea of introducing this regularized diffusion coefficient goes back to [11, Proof of Lemma 3.3]. Thus, by the same calculations leading to (11), we have

$$(13) \quad \int \eta''(\rho_h) |\nabla \rho_h|^2 \leq C(n, \lambda) \int \omega(\rho_h) |\nabla B^h|^2 + C(\lambda) \int f_h \eta'(\rho_h).$$

The left side is $-\frac{d}{dh} \int \eta(\rho_h)$, since

$$(14) \quad (\partial_h - \Delta)(\eta(\rho_h)) = -\eta''(\rho_h) |\nabla \rho_h|^2.$$

Recovering entropy by integrating its dissipation along an auxiliary flow/semigroup is a classical tool in the context of proving logarithmic Sobolev inequalities. We refer to [4, Proposition 5 and its proof, Corollary 1].

The main issue is to understand how the first term on the right arises as a dissipation. Note that the entries of $\rho_h B^h$ are caloric. The term $|\nabla B^h|^2$ will thus appear naturally in dissipations of quantities involving $|B^h|^2$. This leads us to the following question:

Assume that α and β are functions such that α and $\alpha\beta$ are both caloric. For which ω can we guarantee that

$$(15) \quad (\partial_h - \Delta)(\omega(\alpha)\beta^2) \leq -\kappa\omega(\alpha)|\nabla\beta|^2$$

for some $\kappa > 0$?

A relatively short calculation reveals that the condition

$$(16) \quad (2 - \kappa)\omega\omega'' \geq 4(\omega' - \omega/s)^2, \quad s > 0$$

suffices. In this case, by applying (15) with $\alpha = \rho_h$ and β any entry of B^h , we control the bad term by $-C(n, \lambda, \kappa) \frac{d}{dh} \int \omega(\rho_h) |B^h|^2$. Thus, after integrating (13) in h , we can hope to bound $\int \eta(\rho)$ provided

$$(17) \quad \omega(s) \text{ is small compared to } \eta(s), \text{ at least for large } s.$$

There is some flexibility in choices of η and ω satisfying (10), (16), (17) for which the argument will close. The choice

$$\eta(s) = s \log(s), \quad \omega(s) = s$$

is convenient because the bad term is then the dissipation rate of a bounded quantity. We will use (a variant of) this choice to obtain the $L \log L$ estimate (6). Similarly, the choice

$$\eta(s) = s^q, \omega(s) = s^2 \eta''(s)$$

works, provided $q > 1$ is sufficiently close to 1. In a later section we will sketch how this choice gives the L^q estimate (5).

4. ENTROPIES

We now record a couple of entropy formulae that will be useful in the sequel, motivated by the discussion in the previous section. Assume that $\rho > 0$, and that both ρ and $\beta\rho$ satisfy the parabolic PDE

$$(18) \quad P_M u := \partial_h u - \operatorname{div}_y(M(h, y) \nabla_y u) = 0.$$

Here $(h, y) \in \mathbb{R}_+ \times \mathbb{R}^d$, $M \in \operatorname{Sym}_{d \times d}$, and $M \geq 0$. Note that for any choice of η we have

$$(19) \quad P_M(\eta(\rho)) = -\eta''(\rho) |\nabla_y \rho|_M^2.$$

Here and below, $|\xi|_M^2 := M\xi \cdot \xi$. From here on out we fix

$$(20) \quad \eta(s) := \begin{cases} s \log(s) - s + 1, & s > 1 \\ 0, & \text{otherwise.} \end{cases}$$

Notice that $\eta = 0$ for $s \leq 1$. The corresponding choice of ω from the discussion in the previous section is $\omega(s) = s$. This clearly satisfies (10), (16) with $\kappa = 2$, and (17), and moreover (15) becomes an equality:

$$(21) \quad P_M(\rho\beta^2) = -2\rho |\nabla_y \beta|_M^2.$$

5. ELLIPTIC CASE

In this section and below, we will assume that all ingredients are smooth in order to emphasize ideas. The important thing is to obtain estimates that are independent of the regularity of the coefficients. The general case then follows by approximation.

Assume that $A(x) \in \operatorname{Sym}_{n \times n}$ has eigenvalues in $[\lambda, \Lambda]$ and that ρ solves

$$(22) \quad \begin{cases} -\operatorname{div} \operatorname{div}(A(x)\rho) = f \text{ in } B_1, & f \geq 0, \quad \int_{B_1} f = 1, \\ \rho|_{\partial B_1} = 0. \end{cases}$$

Then $\rho \geq 0$, by duality and the maximum principle. Extend ρ, f to be zero outside B_1 , and extend A outside B_1 preserving (2). Using that $A \nabla \rho \cdot \nu \leq 0$ on ∂B_1 (here ν is the outer unit normal), we get

$$(23) \quad -\operatorname{div} \operatorname{div}(A(x)\rho) \leq f \text{ in } \mathbb{R}^n.$$

Let the subscript h denote evolution by heat flow for time h . Then

$$B^h := \frac{(A\rho)_h}{\rho_h} \in [\lambda I, \Lambda I],$$

and we have for all $h > 0$ that

$$(24) \quad -\operatorname{div} \operatorname{div}(B^h \rho_h) \leq f_h.$$

It is easy to see that $\int_{B_1} \rho \leq C(n, \lambda)$ by testing (22) with $1 - |x|^2$, so we have the $L^1 - L^\infty$ estimate

$$(25) \quad \|\rho_h\|_{L^\infty} \leq C(n, \lambda) h^{-n/2}.$$

Testing (24) with $\eta'(\rho_h)$, integrating by parts, applying Cauchy-Schwarz, and using the fact that f has mass 1 gives

$$(26) \quad \int \eta''(\rho_h) |\nabla \rho_h|^2 \leq C(n, \lambda) \int \rho_h |\nabla B^h|^2 + C(\lambda) \|\eta'(\rho_h)\|_{L^\infty}.$$

Integrating (26) in h until $\|\rho_h\|_{L^\infty} < 1$ and applying (19) and (21) with $d = n$, $M = I$, along with (25), gives the desired estimate

$$\int_{B_1} \eta(\rho) < C(n, \lambda, \Lambda).$$

6. PARABOLIC CASE

The parabolic case is nearly identical to the elliptic one. We will evolve by an auxiliary heat flow, but on each fixed physical time slice $\{t = \text{const.}\}$.

Assume that $A(t, x) \in \text{Sym}_{n \times n}$ has eigenvalues in $[\lambda, \Lambda]$ and that ρ solves

$$(27) \quad \begin{cases} \partial_t \rho - \text{div}_x \text{div}_x(A(t, x)\rho) = f \text{ in } Q_1, & f \geq 0, \quad \int_{Q_1} f = 1, \\ \rho|_{\partial_p Q_1} = 0. \end{cases}$$

Here $Q_1 = (-1, 0) \times B_1$ is a parabolic cylinder, and $\partial_p Q_1$ is the parabolic boundary.

Again, $\rho \geq 0$. Extending ρ, f to be zero outside Q_1 and extending A outside Q_1 while preserving (2), we get in the same way as above that

$$(28) \quad \partial_t \rho - \text{div}_x \text{div}_x(A(t, x)\rho) \leq f \text{ in } \mathcal{S} := (-1, 0) \times \mathbb{R}^n.$$

Let the subscript h denote evolution by heat flow in x for time h . For a physical time slice on which ρ is not identically zero, let

$$B^h := \frac{(A\rho)_h}{\rho_h} \in [\lambda I, \Lambda I].$$

Otherwise, let $B^h = \lambda I$. Then

$$(29) \quad \partial_t \rho_h - \text{div}_x \text{div}_x(B^h \rho_h) \leq f_h.$$

It is easy to see that $\int_{B_1} \rho(t, x) dx \leq 1$ for all $t \in (-1, 0)$ by integrating (28), hence

$$\|\rho_h\|_{L^\infty} \leq C(n)h^{-n/2}.$$

Testing (29) with $\eta'(\rho_h)$, integrating by parts, and using Cauchy–Schwarz yields

$$(30) \quad \int \eta''(\rho_h) |\nabla \rho_h|^2 \leq C(n, \lambda) \int \rho_h |\nabla B^h|^2 + C(\lambda) \eta'(\|\rho_h\|_{L^\infty}).$$

Here the integrals are over $\mathcal{S}$, and we have discarded a final-time integral with a favorable sign. The desired $L \log L$ estimate follows in the same way as in the elliptic case after integrating (30) in h .

7. KINETIC CASE

Now assume that $A(t, x, v) \in \text{Sym}_{n \times n}$ has eigenvalues in $[\lambda, \Lambda]$ and that ρ solves

$$(31) \quad \begin{cases} \partial_t \rho + v \cdot \nabla_x \rho - \text{div}_v \text{div}_v(A\rho) = f \text{ in } Q_1, & f \geq 0, \quad \int_{Q_1} f = 1, \\ \rho|_{\partial_{in} Q_1} = 0. \end{cases}$$

Here $Q_1 = (-1, 0) \times B_1 \times B_1$ is a unit kinetic cylinder, and

$$\partial_{in} Q_1 = \{t = -1\} \cup \{|x| = 1, v \cdot x < 0\} \cup \{|v| = 1\}.$$

Again, $\rho \geq 0$. Extend ρ, f to be zero outside Q_1 and extend A outside Q_1 while preserving (2). Arguing as above and using in addition that ρ vanishes on the incoming spatial boundary (so the boundary contribution from the drift term has a favorable sign), we get

$$(32) \quad \partial_t \rho + v \cdot \nabla_x \rho - \text{div}_v \text{div}_v(A(t, x, v)\rho) \leq f \text{ in } \mathcal{S} := (-1, 0) \times \mathbb{R}^n \times \mathbb{R}^n.$$

We will evolve for time h by an auxiliary flow in x and v of the type (18), where $y = (x, v)$ and $d = 2n$, for some carefully chosen $M(h, x, v)$. Velocity heat alone leaves position untouched, whereas adding position heat independently produces gradient terms that the equation does not control. The transport term provides a way to couple the two. A change in velocity changes the position reached by transport, so we allow a mixed position–velocity derivative in the smoothing. It is natural to choose a flow that respects the kinetic scaling $(h, x, v) \mapsto (r^2 h, r^3 x, rv)$. For this reason, we consider diffusion matrices of the form

$$(33) \quad M(h, x, v) = \begin{pmatrix} ah^2 I & bhI \\ bhI & cI \end{pmatrix} := M^h, \quad a, b, c \in \mathbb{R}.$$

Similar matrices appear in [9] in the context of hypocoercivity and in [5] in the context of a kinetic Bakry–Émery calculus. We let the subscript h denote evolution for time h by the flow

$$P_M = \partial_h - \text{tr}(M^h D_{x,v}^2).$$

As above, we will let $B^h := (A\rho)_h/\rho_h$ on physical time slices for which ρ is not identically zero, and define it to be λI otherwise.

Unlike in the elliptic and parabolic cases, new terms will appear in the “flowed” version of (32) due to nontrivial commutators with the drift term:

$$(34) \quad \partial_t \rho_h + v \cdot \nabla_x \rho_h - \operatorname{div}_{x,v}(G^h \nabla_{x,v} \rho_h) - \operatorname{div}_v(\rho_h \operatorname{div}_v B^h) \leq f_h,$$

where

$$(35) \quad G^h = \begin{pmatrix} -bh^2 I & -chI \\ -chI & B^h \end{pmatrix}.$$

We delay this calculation to the end of the section, and instead we immediately note that everything works as in the elliptic and parabolic cases provided the following key inequalities hold:

$$(36) \quad ac > b^2, \quad a, c > 0$$

and

$$(37) \quad -\lambda b > c^2.$$

The matrix M^h determines the smoothing and the gradient terms in the entropy identities. The matrix G^h determines what the smoothed equation can control. We therefore need M^h to control velocity derivatives and G^h to control M^h . Both requirements are satisfied by enforcing (36) and (37).

Indeed, by integrating (32) we see that $\int_{B_1 \times B_1} \rho(t, x, v) dx dv \leq 1$ for all $t \in (-1, 0)$. The condition (36) thus implies the $L^1 - L^\infty$ estimate

$$(38) \quad \|\rho_h\|_{L^\infty} \leq C(a, b, c, n) h^{-2n},$$

as well as decay in x and v of ρ_h . It also implies the ellipticity preservation $B^h \in [\lambda I, \Lambda I]$. These facts can be seen by using the fundamental solution of P_M , given by the Gaussian

$$(39) \quad C(a, b, c, n) h^{-2n} \exp \left[-\frac{3c}{4ac - 3b^2} \frac{|x - \frac{b}{2c} hv|^2}{h^3} - \frac{|v|^2}{4ch} \right].$$

Moreover, it is not hard to see that (36) implies the existence of some $\kappa > 0$ such that

$$(40) \quad M^h \geq \begin{pmatrix} 0 & 0 \\ 0 & \kappa I \end{pmatrix},$$

and (37) implies in addition that

$$(41) \quad G^h \geq \kappa M^h,$$

up to taking κ smaller if needed. Here we use that $B^h \geq \lambda I$.

Test (34) with $\eta'(\rho_h)$. What comes from the last term on the left side (which involves only v derivatives) can be bounded above using (40), and the $|\nabla_{x,v} \rho_h|_{G^h}^2$ coming from the third term on the left side can be bounded below using (41), giving

$$(42) \quad \int \eta''(\rho_h) |\nabla \rho_h|_{M^h}^2 \leq C \int \rho_h |\nabla B^h|_{M^h}^2 + C \eta'(\|\rho_h\|_{L^\infty}).$$

The integrals are taken over $\mathcal{S}$, the constant $C = C(n, \lambda, a, b, c)$, and we have again discarded a final-time integral with a favorable sign. Integrating (42) in h and applying (19), (21), and (38) finishes the argument as before.

Remark 7.1. There are many choices of a, b, c satisfying (36) and (37), e.g. $a = \lambda, b = -\lambda/2, c = \lambda/2$ (which is similar to what Astra originally chose). More generally, take any $b < 0, c > 0$ satisfying (37), and then take $a > 0$ large enough that (36) is satisfied.

Remark 7.2. By the formula (39), we still have the $L^1 - L^\infty$ bound (38) if M^h is rank 1 (when viewed as a 2×2 matrix), i.e. $ac = b^2$, provided $a, c > 0$ as well. This corresponds to auxiliary flow by a kinetic equation of Kolmogorov type, which is tempting as an initial guess. However, in this case, the inequality (40), which is vital for controlling derivatives of B^h in the energy estimate, is not satisfied.

To conclude the section we give a brief derivation of the flowed adjoint equation (34). Let $\mathcal{T} := \partial_t + v \cdot \nabla_x$. It suffices to prove, for any g , that

$$(43) \quad (\mathcal{T}g)_h = \mathcal{T}g_h + bh^2\Delta_x g_h + 2ch\nabla_x \cdot \nabla_v g_h.$$

The sides of (43) coincide at $h = 0$, so it suffices to show that the right side solves $P_M(\cdot) = 0$, i.e. that

$$[\partial_h - (ah^2\Delta_x + 2bh\nabla_x \cdot \nabla_v + c\Delta_v), \mathcal{T} + bh^2\Delta_x + 2ch\nabla_x \cdot \nabla_v] = 0.$$

This follows quickly from the commutator relations

$$[\Delta_v, \mathcal{T}] = 2\nabla_x \cdot \nabla_v, \quad [[\Delta_v, \mathcal{T}], \mathcal{T}] = [2\nabla_x \cdot \nabla_v, \mathcal{T}] = 2\Delta_x, \quad [\Delta_x, \mathcal{T}] = 0.$$

Remark 7.3. When studying kinetic equations in non-divergence form we work with $\partial_t + v \cdot \nabla_x$ and Δ_v and spatial diffusion is recovered through two commutators. By contrast, for kinetic equations in divergence form we have access to ∇_v on its own and

$$[\nabla_v, \partial_t + v \cdot \nabla_x] = \nabla_x$$

already recovers the missing spatial directions.

Remark 7.4. The key coercivity term $-bh^2\Delta_x$ in the flowed equation (34) comes from the commutator of $\nabla_x \cdot \nabla_v$ with the drift. The bad mixed term $-2ch\nabla_x \cdot \nabla_v$ in (34) comes from the commutator of Δ_v with the drift. We are forced to take c small to guarantee (37). However, the fact that Δ_x commutes with the drift allows us to choose a freely to compensate for this and guarantee (36).

8. ALEKSANDROV ESTIMATE

In this section, we indicate how to get from (6) to the stronger L^q estimate (5). The overall strategy is exactly the same as above, with slightly modified entropies and energy estimates. We sketch how it works in the elliptic setting; the parabolic and kinetic cases are entirely analogous.

Entropies. First, enforce the same assumptions on ρ and β as in Section 4. We will update our choice of η as follows:

$$\eta(s) = \begin{cases} s^q - qs + q - 1, & s > 1, \\ 0 & \text{otherwise.} \end{cases}$$

Here $q > 1$. Motivated by the discussion in Section 3, we let

$$\omega(s) = q(q-1)s^q.$$

This choice clearly satisfies (10). It also satisfies (16) with constant

$$\kappa = \kappa_q = \frac{2(2-q)}{q},$$

for any $q \in (1, 2)$. We record the associated entropy inequality, analogous to (15):

$$(44) \quad P_M(\omega(\rho)\beta^2) \leq -\kappa_q\omega(\rho)|\nabla_y\beta|_M^2.$$

Finally, ω clearly satisfies (17) provided q is close to 1. Below we will assume in particular that $q \in (1, 3/2)$, which prevents κ_q from degenerating (precisely, $\kappa_q > 2/3$).

Elliptic Case. Assume the same on A, ρ, f as in Section 5. Test (24) with $\eta'(\rho_h)$. Integrating by parts and applying Cauchy–Schwarz gives

$$(45) \quad \int \eta''(\rho_h)|\nabla\rho_h|^2 \leq C(n, \lambda) \int \omega(\rho_h)|\nabla B^h|^2 + C(\lambda)\|\eta'(\rho_h)\|_{L^\infty}.$$

Recalling the $L^1 - L^\infty$ estimate (25) and that $\eta(s) = 0$ for $s < 1$, we see that the integral from 0 to ∞ in h of the last term is bounded provided $q < 1 + 2/n$, which we now enforce. Performing this integration and recalling (19) and (44), we get

$$(46) \quad \int_{B_1} \eta(\rho) \leq C(n, \lambda, \Lambda) \int_{B_1} \omega(\rho) + C(n, \lambda, q).$$

Finally, we recall that for $s \geq 0$,

$$s^q \leq \eta(s) + qs, \quad \omega(s) = q(q-1)s^q.$$

Combining with (46) we get

$$\int_{B_1} \rho^q \leq C(n, \lambda, \Lambda) q(q-1) \int_{B_1} \rho^q + q \int_{B_1} \rho + C(n, \lambda, q).$$

Taking $q(n, \lambda, \Lambda) < 1 + 2/n$ sufficiently close to 1 we can absorb the first term on the right, and recalling the L^1 bound $\int_{B_1} \rho \leq C(n, \lambda)$, we get

$$\int_{B_1} \rho^q \leq C(n, \lambda, \Lambda).$$

For general f , the L^q estimate (5) follows by homogeneity.

Remark 8.1. There are two restrictions on q . The heat-kernel bound (38) controls the source integral only when $q < 1 + \frac{1}{2n}$ in the kinetic case. Absorbing the coefficient term imposes a second restriction, depending on ellipticity. The first reflects how the auxiliary flow spreads mass; the second comes from estimating the averaged coefficient by its size. It is this second restriction that makes our source exponent grow with the ellipticity ratio.

9. ON EXPONENTS IN THE ALEKSANDROV ESTIMATE

A careful tracking of constants in the argument of the previous section shows that we must take

$$q - 1 \leq c(n)(\lambda/\Lambda)^2.$$

Thus, the exponent in the Aleksandrov estimate (4) produced by this argument goes to infinity with the ellipticity ratio $R := \Lambda/\lambda$.

On the other hand, in the elliptic and parabolic cases, the classical methods tell us much more about the Aleksandrov exponent.

For example, when $n \geq 2$, in the elliptic case we can take $p = n - \epsilon$ for some $\epsilon(n, R) > 0$ (see [7]). It was conjectured by Pucci [16] that the estimate holds for all

$$p > \max \left\{ \frac{n}{2}, \frac{n}{1 + (n-1)/R} \right\}.$$

This threshold is predicted by radial homogeneous examples. It was proven to be sufficient by Astra [14].

In the parabolic case, we can take $p = n + 1 - \epsilon$ for some $\epsilon(n, R) > 0$ ([7]). Small modifications of the elliptic examples show that, for any exponent smaller than n , the Aleksandrov estimate fails at sufficiently large ellipticity ratio. Here, the condition

$$p > \max \left\{ \frac{n+2}{2}, \frac{n}{1 + (n-1)/R} \right\}$$

is necessary (see [6, Theorem 4.8]), while its sufficiency was established by Astra [14].

Finally, in the kinetic case, small modifications of stationary kinetically homogeneous examples show that, for any exponent smaller than $4n$, the Aleksandrov estimate fails at sufficiently large ellipticity ratio. Moreover, under certain restrictions on the variables on which the coefficients depend, Aleksandrov estimates have been proven to hold for the best possible exponents. These results, along with a more thorough discussion of optimal exponents and some explicit conjectures, can be found in [13]. The kinetic Pucci conjecture seems to be an interesting research direction as it presumably requires a novel alternative approach to the kinetic Aleksandrov estimate.

REFERENCES

- [1] F. Abedin and G. Tralli. Harnack inequality for a class of Kolmogorov-Fokker-Planck equations in non-divergence form. *Arch. Ration. Mech. Anal.*, 233(2):867–900, 2019.
- [2] A. D. Aleksandrov. Certain estimates for the Dirichlet problem. *Soviet Mathematics. Doklady*, 1:1151–1154, 1961.
- [3] I. J. Bakel'man. On the theory of quasilinear elliptic equations. *Sibirsk. Mat. Ž.*, 2:179–186, 1961.
- [4] D. Bakry and M. Émery. Diffusions hypercontractives. In *Séminaire de probabilités, XIX, 1983/84*, volume 1123 of *Lecture Notes in Math.*, pages 177–206. Springer, Berlin, 1985.
- [5] F. Baudoin. Bakry-émery meet Villani. *J. Funct. Anal.*, 273(7):2275–2291, 2017.
- [6] H. Dong and S. Kitano. Second order L_p estimates for subsolutions of fully nonlinear equations. *Calc. Var. Partial Differential Equations*, 64(3):Paper No. 76, 20, 2025.
- [7] E. B. Fabes and D. W. Stroock. The L^p -integrability of Green's functions and fundamental solutions for elliptic and parabolic equations. *Duke Math. J.*, 51(4):997–1016, 1984.

- [8] Z. Hao and X. Zhang. Harnack estimates for nondivergence kinetic equations, 2026. arXiv:2609.15534.
- [9] F. Hérau. Short and long time behavior of the Fokker-Planck equation in a confining potential and applications. *J. Funct. Anal.*, 244(1):95–118, 2007.
- [10] L. Hörmander. Hypoelliptic second order differential equations. *Acta Math.*, 119:147–171, 1967.
- [11] N. V. Krylov. On passing to the limit in degenerate Bellman equations. ii. *Mathematics of the USSR-Sbornik*, 35(3):351–362, 1979.
- [12] N. V. Krylov and M. V. Safonov. A certain property of solutions of parabolic equations with measurable coefficients. *Mathematics of the USSR-Izvestiya*, 16(1):151–164, 1981.
- [13] A. Loher, C. Mooney, and C. Mouhot. Döblin–fourier cancellation and kinetic aleksandrov estimates, 2026. arXiv:2610.03083.
- [14] A. Loher, C. Mooney, C. Mouhot, L. Niebel, L. Silvestre, and Y. Zhu. We need ABP. <https://weneedabp.github.io/>. Accessed: 2026-10-06.
- [15] C. Pucci. Limitazioni per soluzioni di equazioni ellittiche. *Ann. Mat. Pura Appl. (4)*, 74:15–30, 1966.
- [16] C. Pucci. Operatori ellittici estremanti. *Ann. Mat. Pura Appl. (4)*, 72:141–170, 1966.
- [17] M. V. Safonov. The Harnack inequality for elliptic equations and Hölder continuity of their solutions. *Journal of Soviet Mathematics*, 21(5):851–863, 1983.
- [18] L. Silvestre. Regularity estimates and open problems in kinetic equations. In *A³N²M: approximation, applications, and analysis of nonlocal, nonlinear models*, volume 165 of *IMA Vol. Math. Appl.*, pages 101–148. Springer, Cham, [2023] ©2023.
- [19] K. Tso. On an Aleksandrov-Bakel’man type maximum principle for second-order parabolic equations. *Comm. Partial Differential Equations*, 10(5):543–553, 1985.
- [20] Y. Zhu. Stationary kinetic krylov-safonov estimates in space dimension one, 2026. arXiv:2608.31158.

ALL SOULS COLLEGE, UNIVERSITY OF OXFORD
 Email address: amelie.loher@all-souls.ox.ac.uk

DEPARTMENT OF MATHEMATICS, UC IRVINE
 Email address: mooneycr@uci.edu

DPMMS, UNIVERSITY OF CAMBRIDGE
 Email address: c.mouhot@dpms.cam.ac.uk

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH
 Email address: lukas.niebel@math.ethz.ch

MATHEMATICS DEPARTMENT, UNIVERSITY OF CHICAGO
 Email address: luiss@uchicago.edu

SSE, THE CHINESE UNIVERSITY OF HONG KONG, SHENZHEN
 Email address: zhuyuzhe@cuhk.edu.cn