

Ques  
approximate value of  $\pi$  ( $x_a$ ) = 3.1428571  
true value " " ( $x_t$ ) = 3.1415926

$$\text{absolute error} = |x_t - x_a|$$
$$= |3.1415926 - 3.1428571|$$
$$= 0.0012645$$

$$\text{relative error} = \frac{x_t - x_a}{x_t}$$

$$= \frac{-0.0012645}{3.1415926}$$


$$= -0.0004025$$

C ~~error~~

$$\text{relative error percentage} = 0.0004025 \times 100\%$$

$$= 0.0402502\%$$

(percentage can't be -ve)



20.

$$f(x) = 3x + \sin x - e^x$$

wirg. Jede Position  
rechts

x	f(x)
0	-1
1	1.12318

$$\text{Hence } x_0 = 0, f_0 = -1 \\ x_1 = 1, f_1 = 1.12318$$

Since  $f_0 \cdot f_1 < 0 \Rightarrow$  root lies between 0 & 1.

$$x_2 = \frac{x_0 f_1 - x_1 f_0}{f_1 - f_0}$$
$$= \frac{0 \times 1.12318 - 1 \times (-1)}{1.12318 + 1}$$
$$= 0.47098$$

$$f(x_2) = 3 \times 0.47098 + \sin(0.47098) -$$
$$e^{0.47098}$$
$$= 0.26516$$

1b

Headline Table:

itno	$x_0$	$f_0$	$x_1$	$f_1$	$x_2$	$f(x_2)$
1	0	-1	1	1.12318	0.47099	0.26516
2	0	-1	0.47099	0.26516	0.37228	0.02953
3	0	-1	0.37228	0.02953	0.36160	0.00295
4	0	-1	0.36160	0.00295	0.36054	0.00029
5	0	-1	0.36054	0.00029	0.36044	0.00003
6	0	-1	0.36044	0.00003	0.36043	0.00002

∴ root of  $f(x)$  upto four decimal  
point = 0.3604.

10

2a

Given

$$f(x) = x^3 + x - 1$$

using secant method,

x	f(x)
0	-1
1	1

$$\text{Now } x_0 = 0, f_0 = -1 \\ x_1 = 1, f_1 = 1$$

$$x_2 = \frac{x_0 f_1 - x_1 f_0}{f_1 - f_0}$$

$$= \frac{0 \times 1 - 1 \times (-1)}{1 - 1}$$

$$= 0.5$$

$$f(x_2) = 0.5^3 + 0.5 - 1 \\ = -0.37500$$



2a

iterative table :

itno	x <sub>0</sub>	f <sub>0</sub>	x <sub>1</sub>	f <sub>1</sub>	x <sub>2</sub>	f(x <sub>2</sub> )
1	0	-1	1	1	0.5	-0.37500
2	1	1	0.5	-0.37500	0.63636	-0.10534
3	0.5	-0.37500	0.63636	-0.10594	0.69005	0.01863
4	0.63636	-0.10594	0.69005	0.01863	0.68202	-0.00074
5	0.69005	0.01863	0.68202	-0.00074	0.68233	0.00000

Since,  $f(x_2) = 0$  which has error less than  $10^{-4}$ , so root of  $f(x) = 0.6823$

(9)

## Algorithm for bisection method

- 1) Start with two points  $x_0$  and  $x_1$  such that  $f(x_0) \neq 0$  and  $f(x_1) \neq 0$
- 2) define  $f(x)$
- 3) choose initial guess  $x_0$  and  $x_1$  such that  $f(x_0) \cdot f(x_1) < 0$
- 4) define tolerable error ( $\epsilon$ )

5) calculate  $x_2 = \frac{x_0 + x_1}{2}$

6) calculate  $f(x_2)$

i) if  $f(x_2) = 0$ , then root =  $x_2$  and  
goto stop.

ii) if  $f_0 \cdot f(x_2) < 0$  set

$$x_0 = x_2, x_1 = x_1$$

$f_0 = f(x_2), f_1 = f_1$  and goto step 3

iii) if  $f_0 \cdot f(x_2) > 0$  set

$$x_0 = x_0, x_1 = x_2$$

$f_0 = f(x_2), f_1 = f_1$  and goto step 8

iv) if  $f(x_2) = 0$ , then root =  $x_2$   
and goto stop.

8. if  $|f(x_0)| < \epsilon$ , goto stop  
else goto step 5.

9. stop.

Given, 39.

$$f(x) = \log x - \cos x$$

x	f(x)	f'(x)
2	1.30929	$\frac{1}{2} + \sin x$
3	2.08860	

using, newton's method:

let, initial guess ( $x_0$ ) = 2.5

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$= 2.5 - \frac{1.71743}{0.99847}$$
$$= 0.77994$$

$$f(x_1) = \log(0.77994) - \cos(0.77994)$$

$$= -0.95949$$

$$f'(x_1) = 1.98539$$

No.	$x_0$	$f(x_0)$	$f'(x_0)$	$x_1$	$f(x_1)$
1	2.5	1.71743	0.39897	0.72344	-0.35943
2	0.77597	-0.95949	1.98533	1.26322	-0.06303
3	1.26322	-0.06909	1.74470	1.30282	-0.00024
4	1.30282	-0.00025	1.73188	1.30294	-0.00002

Since  $|f(x_1)| = 0$   
 which has less error than tolerable  
 error.

(g)

$\therefore \text{root of } f(x) = 1.3029$

Ques.  
given system of equations

$$2x - y - 7z = 3$$

$$5x - 2y + 3z = -1$$

$$-3x + 3y + z = 2$$

given system in matrix form:


$$\begin{bmatrix} 2 & -1 & -7 \\ 5 & -2 & 3 \\ -3 & 9 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$$

$$AX = B \quad \text{--- (1)}$$

where

$$A = \begin{bmatrix} 2 & -1 & -7 \\ 5 & -2 & 3 \\ -3 & 9 & 1 \end{bmatrix}$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$



$$B = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$$

Ans.

A.

[2  
5]

4a

again using LU decomposition method:

$$A = LU$$

using crout's factorization method:

$$\begin{bmatrix} 2 & -1 & -7 \\ 5 & -2 & 3 \\ -3 & 9 & 1 \end{bmatrix} = \begin{bmatrix} L_{11} & 0 & 0 \\ L_{21} & L_{22} & 0 \\ L_{31} & L_{32} & L_{33} \end{bmatrix} \begin{bmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} L_{11} & L_{11}U_{12} & L_{11}U_{13} \\ L_{21} & L_{21}U_{12} + L_{22} & L_{21}U_{13} + L_{22}U_{23} \\ L_{31} & L_{31}U_{12} + L_{32} & L_{31}U_{13} + L_{32}U_{23} + L_{33} \end{bmatrix}$$

Comparing:

$$L_{11} = 2 \rightarrow L_{21} = 5, \quad L_{31} = -3$$

$$L_{11}U_{12} = -1 \Rightarrow U_{12} = -\frac{1}{2}$$

$$L_{11}U_{13} = -7 \Rightarrow U_{13} = -\frac{7}{2}$$

$$L_{21}U_{12} + L_{22} = -2$$

$$5 \times (-\frac{1}{2}) + L_{22} = -2 \Rightarrow L_{22} = \frac{1}{2}$$

$$L_{21}U_{13} + L_{22}U_{23} = 3$$

$$5 \times (-\frac{7}{2}) + \frac{1}{2} \times U_{23} = 3 \Rightarrow U_{23} = \frac{7}{2} + 4$$

-epsing ?

process once.  
→ Disadvantage:  
more

4a

again wing-LU decomposition method:

$$A = LU$$

wing crout's factorization method:

$$\begin{bmatrix} 2 & -1 & -7 \\ 5 & -2 & 3 \\ -3 & 9 & 1 \end{bmatrix} = \begin{bmatrix} L_{11} & 0 & 0 \\ L_{21} & L_{22} & 0 \\ L_{31} & L_{32} & L_{33} \end{bmatrix} \begin{bmatrix} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{bmatrix}$$

$$= \begin{bmatrix} L_{11} & L_{11}U_{12} & L_{11}U_{13} \\ L_{21} & L_{21}U_{12} + L_{22} & L_{21}U_{13} + L_{22}U_{23} \\ L_{31} & L_{31}U_{12} + L_{32} & L_{31}U_{13} + L_{32}U_{23} + L_{33} \end{bmatrix}$$

L11

Comparing:

$$L_{11} = 2 \rightarrow L_{21} = 5, L_{31} = -3$$

$$L_{11}U_{12} = -1 \Rightarrow U_{12} = -1/2$$

$$L_{11}U_{13} = -7 \Rightarrow U_{13} = -7/2$$

$$L_{21}U_{12} + L_{22} = -2$$

$$5 \times (-1/2) + L_{22} = -2 \Rightarrow L_{22} = 1/2$$

$$L_{21}U_{13} + L_{22}U_{23} = 3$$

$$5 \times (-7/2) + 1/2 \times U_{23} = 3 \Rightarrow U_{23} = 7/0 + 41$$

efficiency? Date: NVIDIA  
Page: SPIN

process multiple

once.

→ Disadvantages:

• more memory



# National Academy of science and Technology

Affiliated to Pashchim University

Dhangadhi, Kailali (Nepal)

Accredited by University Grants Commission (UGC), Nepal (2022)

S477

NV22A  
WING A

1a date

by U.P.

Additional answerbook No.

Number of additional answerbooks used

Name of the student

Roll No. (Class) : Exam

Trade/Semester Exam

Engineering Diploma

Programme

Year

\* Students are required to write answer on both sides starting from the first page of this copy.

\* This answerbook must be stitched properly with the original answer book before delivering to the invigilator.

9a

$$L_{31}U_{12} + L_{32} = 9$$
$$-3 \times (-\frac{1}{2}) + L_{32} = 9$$

$$\Rightarrow L_{32} = 15/2$$

$$L_{31}U_{13} + L_{32}U_{23} + L_{33} = 1$$
$$-3 \times (-7/2) + 15/2 \times 76/41 + L_{33} = 1$$

$$L_{33} = -317$$

$$\therefore L = \begin{bmatrix} 2 & 0 & 0 \\ 5 & 1/2 & 0 \\ -3 & 15/2 & -317 \end{bmatrix}$$

~~$$U = \begin{bmatrix} 1 & -1/2 & -7/2 \\ 0 & 1 & 41 \\ 0 & 0 & 1 \end{bmatrix}$$~~

7a

eqn ① can be written as:

$$L(UX) = B$$

again,

$$\begin{aligned} UX &= Z \quad - \textcircled{i} \\ LZ &= B \quad - \textcircled{iii} \end{aligned}$$

Solving \textcircled{iii}

$$\left[ \begin{array}{ccc|c} 2 & 0 & 0 & 7 \\ 5 & 1/2 & 0 & \\ -3 & 15/2 & -317 & \end{array} \right] \left[ \begin{array}{c} z_1 \\ z_2 \\ z_3 \end{array} \right] = \left[ \begin{array}{c} 3 \\ -1 \\ 2 \end{array} \right]$$

$$\Rightarrow \left[ \begin{array}{ccc|c} 2z_1 & & & 3 \\ 5z_1 + 1/2z_2 & & & -1 \\ -3z_1 + 15/2z_2 & -317z_3 & & 2 \end{array} \right]$$

$$z_1 = 3/2$$

$$5 \times 3/2 + \frac{1}{2}z_2 = -1$$

$$z_2 = -17$$

$$-3 \times 3/2 + 15/2 \times (-17) - 317z_3 = 2$$

$$z_3 = -\frac{134}{37}$$

7a

Solving: (ii)

$$\left[ \begin{array}{ccc|c} 1 & -1/2 & -7/2 & x \\ 0 & 1 & 4/2 & y \\ 0 & 0 & 1 & z \end{array} \right] \xrightarrow{x = z} \left[ \begin{array}{c|c|c} x & y & z \\ \hline 3/2 & -17 & -134/137 \end{array} \right]$$

$$\left[ \begin{array}{c|c|c} x - 1/2y - 7/2z & y + 4/2z & z \\ \hline 3/2 & -17 & -134/137 \end{array} \right]$$

$$\therefore z = \frac{-134}{137}$$

$$y + 4/2x \left( \frac{-134}{137} \right) = -17$$

$$y = \frac{3165}{137}$$

$$x - \frac{1}{2} \times \frac{3165}{137} - \frac{7}{2} \times \frac{2}{137} = \frac{3}{2}$$

15

$$x = \frac{2747}{137}$$

$$\therefore \boxed{\begin{array}{l} x = 20.05 \\ y = 23.10 \\ z = -0.97 \end{array}}$$

much a system of equations

$$10x_1 + 37x_2 + 4x_3 = 16 \quad (1)$$

$$3x_1 + 18x_2 + 37x_3 = 37 \quad (2)$$

$$2x_1 + 22x_2 + 10x_3 = -10 \quad (3)$$

solving (1), (2) & (3) for  $x_1, x_2$  &  $x_3$  respectively.

$$x_1 = \frac{1}{10} [15 - 37x_2 - 4x_3] \quad (1)$$

$$x_2 = \frac{1}{10} [37 - 2x_1 - 37x_3] \quad (2)$$

$$x_3 = \frac{1}{10} [-10 - 3x_1 - 2x_2] \quad (3)$$

put,  $x_2 = x_3 = 0$  for initial  
using gauss seidel method:

$$x_1 = \frac{1}{10} [15] = \frac{3}{2} = 1.50$$

$$x_2 = -\frac{1}{10} [37 - 2 \times 1.50 - 0] \\ = -3.40$$

$$x_3 = -\frac{1}{10} [-10 - 3 \times 1.50 - 2 \times (-3.40)] \\ = 0.77 \quad \checkmark$$

4b

Iterative table :

itno	$x_1$ from eqn ①	$x_2$ from eqn ②	$x_3$ from eqn ③
1	1.56	-3.40	0.77
2	2.21	1.89 - 3.02	2.64 - 3.22
3	-0.11	1.64	1.36
4	0.46	-3.20	0.79
5	1.98	-2.98	0.99
4	2.00	-3.00	0.99
5	2.00	-3.00	0.99

$\therefore$ 

$x_1 = 2$
$x_2 = -3$
$x_3 = 0.99$

(6)

5A

Initial equations

$$\begin{aligned} R_1 &+ R_2 + R_3 + R_4 = 7 \\ R_1 &+ 10R_2 + 13 + 9R_4 = 12 \\ R_1 &+ 4R_2 + 6R_3 + R_4 = -5 \\ R_1 &+ 8R_2 + 12R_3 + 4R_4 = -6 \end{aligned}$$

augmented no. coefficient matrix:

$$A = \left[ \begin{array}{cccc|c} 1 & 1 & 1 & 1 & 7 \\ 1 & 10 & 1 & 1 & 12 \\ 1 & 4 & 6 & 1 & -5 \\ 1 & 8 & 12 & 4 & -6 \end{array} \right]$$

using Gaussian elimination method:

$$R_2 \rightarrow 5R_2 - R_1, R_3 \rightarrow 5R_3 - R_1, R_4 \rightarrow 5R_4 -$$

$$\left[ \begin{array}{cccc|c} 1 & 1 & 1 & 1 & 7 \\ 0 & 34 & 1 & 19 & 56 \\ 0 & 4 & 99 & 4 & -29 \\ 0 & 1 & 1 & 19 & -26 \end{array} \right] \quad \text{partial pivoting}$$

$$R_3 \rightarrow \frac{1}{2} R_3 - R_2, R_4 \rightarrow \frac{1}{2} R_4 - R_2$$

$$\left[ \begin{array}{cccc|c} 5 & 1 & 1 & 1 & 4 \\ 0 & 34 & 4 & 19 & 56 \\ 0 & 0 & 485/2 & 15 & -605/2 \\ 0 & 0 & 0 & 285/2 & 165 \end{array} \right]$$

$$\therefore \frac{285x_4}{2} = 165 \quad \text{---(1)}$$

$$x_4 = \frac{22}{19} = 1.15 \text{ #}$$

$$\therefore \frac{485x_3 + 15x_4}{2} = -\frac{605}{2}$$

$$\frac{485x_3}{2} + \frac{15 \times 22}{19} = -\frac{605}{2}$$

$$(6) \quad x_3 = \frac{-2931}{1843} = -1.61 \text{ #}$$

$$34x_2 + 4x_3 + 19x_4 = 56$$

$$x_2 = \frac{2129}{1843} = 1.15 \text{ #}$$

$$5x_1 + x_2 + x_3 + x_4 = 4$$

$$x_1 = \frac{1108}{1843} = 0.60 \text{ #}$$

Qb

Sim. System:

$$\begin{aligned} Q_1 + Q_2 + Q_3 &= 6 \\ Q_2 + Q_3 + Q_1 &= 6 \\ Q_1 - Q_2 + Q_3 &= 0 \end{aligned}$$

Augmented matrix:

$$A = \left[ \begin{array}{ccc|c} 2 & 2 & 1 & : 6 \\ 4 & 2 & 3 & : 6 \\ 1 & -1 & 1 & : 0 \end{array} \right]$$

$$R_2 \rightarrow R_2 - 2R_1, \quad R_3 \rightarrow 2R_3 - R_1$$

$$\left[ \begin{array}{ccc|c} 2 & 2 & 1 & : 6 \\ 0 & -2 & 1 & : -6 \\ 0 & -4 & 1 & : -6 \end{array} \right]$$

$$R_1 \rightarrow R_1 + R_2, \quad R_3 \rightarrow R_3 + 2R_2$$

$$\left[ \begin{array}{ccc|c} 2 & 0 & 2 & : 0 \\ 0 & -2 & 1 & : -6 \\ 0 & 0 & -1 & : 0 \end{array} \right]$$

$$R_2 \rightarrow R_2 + QR_3, \quad R_2 \rightarrow R_2 + R_3$$

National  
Academy  
of  
Technology  
and  
Management  
Additional answer  
Number of additional  
Name of the student  
Roll No. (Class)  
Paper  
\* Students are  
This answer



# National Academy of Science and Technology

(Affiliated to Tribhuvan University)

Dhangadhi, Kailali (Nepal)

Accredited by University Grants Commission (UGC), Nepal (2011)

National University No.

Number of Admissions since last year

3333

Admission's Number

Name of the student

Roll No. (Class)

Exam

Grade Required

Practical

Paper

1910

\* Students are required to write answer on both sides starting from the first page of this copy

\* This answer book must be submitted together with the original answer book before returning to the invigilator

$$\begin{bmatrix} 2 & 0 & 0 : 12 \\ 3 & -2 & 0 : 0 \\ 0 & 0 & -1 : 6 \end{bmatrix}$$

$$R_1 \rightarrow R_1/2, R_2 \rightarrow R_2/(-2), R_3 \rightarrow R_3 + 1$$

$$\begin{bmatrix} 1 & 0 & 0 : 6 \\ 0 & 1 & 0 : 0 \\ 0 & 0 & 1 : -6 \end{bmatrix}$$

∴

$x_1 = 6$
$x_2 = 0$
$x_3 = -6$

⑦

Q8.

Let  $x$  be the cube root of 11.

$$\text{Let } x = \sqrt[3]{11}$$

$$f(x) = x^3 - 11$$

$$f'(x) = 3x^2$$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$\begin{array}{|c|c|} \hline x & f(x) \\ \hline 0 & -11 \\ \hline \end{array} \quad x_1 = 3 - \frac{16}{27} = 2.40$$

1st initial guess ( $x_0$ ) = 3

$$f(x_0) = 3^3 - 11 = 16$$

$$f'(x_0) = 3x^2 = 27$$

iterative table

$x_0$	$f(x_0)$	$f'(x_0)$	$x_1$	$f(x_1)$
3	16	27	2.40741	2.95240
2.40741	2.95240	26.15000	2.29451	1.08004
2.29451	1.08004	15.79493	2.22613	0.03191
2.22613	0.03191	14.86696	2.22398	0.00000

Since  $f(x_2) = 0$

" root of  $f(x) = 2.22398$  #"

neural networks.

6b

Error: Error is the difference between actual value and calculated value.  
i.e.  $|error| = |x_a - x_c|$

These are different types of error arises during calculations like truncation error, inherent errors etc.

a) Round off errors

Round off errors are arises due to roundoff decimal point.

Let we have,  $x = \frac{100}{3} = 33.333\ldots$

If we write  $x = 33.33$  it is rounded.  
 ~~$x = 33.333$~~  is more accurate than  
 ~~$x = 33.33$~~

Solution: Write more number after decimal point.

To get more precise value we must use as more as possible the bit after decimal point.

6b

### b) Truncation Errors.

Truncation Errors are arising due to replacing infinite terms by a finite one.

Example: we have a series:

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

let, put  $x = 2$

$$e^2 = 1 + 2 + \frac{2^2}{2!} = 5 \quad \text{--- (1)}$$

$$e^2 = 1 + 2 + \frac{2^2}{2!} + \frac{2^3}{3!} = 6.33 \quad \text{--- (2)}$$

eqn (2) is more accurate than eqn (1).

Solution: writing more term reduce the truncation error. The more you need accuracy, the more you write the terms.

(7)

$$x^3 - x + 1 = 0$$

$$x = \frac{1}{2}$$

$$x = 0.5$$

3b.

$$f(x) = x^3 + x^2 - 1$$

$x$	$f(x)$
0	-1
1	1

initial guess = 0.5

using fixed point iteration method:

$$x = g(x) = \sqrt[3]{1 - x^2}$$

$$g'(x) = \frac{1}{3} (1 - x^2)^{-\frac{2}{3}} (-2x)$$

$$= -\frac{2x}{3} (1 - x^2)^{\frac{2}{3}}$$

$$g'(0.5) = -0.20637 < 1$$

$$x_1 = g(0.5) = (1 - 0.5^2)^{\frac{1}{3}} = 0.90856$$

~~$$x_2 = (1 - 0.90856^2)^{\frac{1}{3}} = 0.55883$$~~

~~$$x_3 = (1 - 0.55883^2)^{\frac{1}{3}} = 0.88268$$~~

Similarly,

$$x_4 = 0.60448$$

$$x_5 = 0.85935$$

3b

$$x_6 = \frac{(1 - 0.85935^2)^{1/3}}{0.63949}$$

$$x_7 = (1 - 0.63949^2)^{1/3}$$

$$= 0.83992$$

$$= 0.66623$$

~~x<sub>8</sub>~~

~~and so on ...~~