

# Computational Weber electrodynamics with symplectic numerical integrators

Samer

samerweb3@gmail.com

December 8, 2025

## Abstract

This paper presents

## 1 Introduction

The two main equations in Weber's electrodynamics are the *velocity-dependent potential*

$$V = \frac{q_1 q_2}{r} \left( 1 - \frac{\dot{r}^2}{2c^2} \right) \quad (1)$$

and *Weber's force law*

$$\vec{F}_{12} = q_1 q_2 \frac{\hat{r}}{r^2} \left( 1 - \frac{\dot{r}^2}{2c^2} + \frac{r\ddot{r}}{c^2} \right) \quad (2)$$

where  $r^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2$  is the squared distance between two particles with electric charges  $q_1$  and  $q_2$ ,  $\dot{r} = dr/dt$  is the relative radial velocity,  $\ddot{r} = d\dot{r}/dt$  is the relative radial acceleration and  $c$  is the speed of light in vacuum. This setup is described in more detail in Section 7.1, Section 2 and follows conventions used in Assis [1994].

In SI units, (1) and (2) would have a factor of  $1/(4\pi\epsilon_0)$  multiplying the right-hand side, where  $\epsilon_0$  is the permittivity of free space. We choose to omit this factor by using an alternative system of units that better suits our computational purposes, see Section 5.

The force (2) can be derived from the potential (1), see equation (6).

We also have  $\vec{F}_{12} = m_1 \vec{a}_1$  and  $\vec{F}_{21} = m_2 \vec{a}_2$ , where  $m_1$  and  $m_2$  are the *inertial masses* of the particles and  $\vec{a}_1$  and  $\vec{a}_2$  are their accelerations with respect to an *inertial frame of reference*. We discuss frames of reference in more detail in Section 7.1.

This setup satisfies Newton's third law of motion and therefore  $\vec{F}_{21} = -\vec{F}_{12}$  is true in general. Therefore, energy, momentum and angular momentum are always conserved.

This is a significant *computational advantage* of Weber’s electrodynamics compared to Maxwell-Heaviside’s field formalism together with the Lorentz force law. We discuss this advantage in more detail in Section 7.1.

Weber’s force law (2) generalizes Coulomb’s law for electrostatics to include *velocity dependent* (magnetic) and *acceleration dependent* (induction) forces. Helmholtz argued in the 19th century that Weber’s model violates the *principle of conservation of energy*, see Assis [2021b]. Weber replied with great detail to these claims and proved energy conservation extensively in his 6th and 7th memoirs. []

Nowadays, Weber’s electrodynamics is said to be valid only for small velocities (*quasi-static approximation*) compared to  $c$  and is not suitable for describing electromagnetic radiation. But the validity of these statements is being challenged since the 1990’s as part of a renewed interest in Weber’s electrodynamics, see [].

To test the strengths and weaknesses of Weber’s electrodynamics, we provide a computational framework for investigating  $n$ -body systems of charged particles through numerical simulations. This allows us to explore the dynamics of charged particles interacting via Weber’s force law (2) in various configurations and scenarios. Important aspects like energy, momentum and angular momentum conservation can be verified numerically. Furthermore, it paves the way for studying complex systems with oscillatory behavior and wave phenomena within the context of Weber’s electrodynamics.

## 2 Lagrangian and Hamiltonian formulation of Weber’s electrodynamics

The Lagrangian and Hamiltonian formulation of Weber’s electrodynamics was introduced in 1868 by Carl Neumann, see English translation in [Assis, 2021b]. The standard formalism is described in modern notation in Assis [1994]. We repeat some of the main equations and derive the general Hamiltonian and associated canonical equations for  $n$  particles.

Consider a system of two particles with charges  $q_1$  and  $q_2$  and inertial masses  $m_1$  and  $m_2$ . The total kinetic energy of this system is:

$$T = m_1 \frac{\vec{v}_1 \cdot \vec{v}_1}{2} + m_2 \frac{\vec{v}_2 \cdot \vec{v}_2}{2} \quad (3)$$

where  $\vec{v}_1 = (\dot{x}_1, \dot{y}_1, \dot{z}_1)$  and  $\vec{v}_2 = (\dot{x}_2, \dot{y}_2, \dot{z}_2)$  are the velocities of particles 1 and 2 with respect to an *inertial frame of reference*. We discuss frames of reference in more detail in Section 7.1.

To define the Lagrangian, we first introduce an expression nearly identical to (1) but with a positive sign in front of the velocity-dependent term:

$$S = \frac{q_1 q_2}{r} \left( 1 + \frac{\dot{r}^2}{2c^2} \right) \quad (4)$$

The Lagrangian is then defined as:

$$\boxed{L = T - S} \quad (5)$$

Although the following expression is not directly relevant for our goals, it is worth noting that we can apply the Euler-Lagrange equation to obtain the force (2) for particle 1 in the  $x$ -direction:

$$F_{12}^x = \frac{d}{dt} \frac{\partial S}{\partial \dot{x}_1} - \frac{\partial S}{\partial x_1} = q_1 q_2 \frac{(x_1 - x_2)}{r_{12}^3} \left( 1 - \frac{\dot{r}^2}{2c^2} + \frac{r\ddot{r}}{c^2} \right) \quad (6)$$

with analogous equations for particle 2 and for the  $y$  and  $z$  directions. This completes our Lagrangian formulation of Weber's electrodynamics.

## 2.1 Hamiltonian for 2 particles

The Hamiltonian, by definition, is derived from the Lagrangian. For two particles in three spatial dimensions, it is given by

$$H = \sum_{i=1}^2 \left( \dot{x}_i \frac{\partial L}{\partial \dot{x}_i} + \dot{y}_i \frac{\partial L}{\partial \dot{y}_i} + \dot{z}_i \frac{\partial L}{\partial \dot{z}_i} \right) - L \quad (7)$$

This Hamiltonian is a function of individual particle *positions* and *velocities* in three spatial dimensions

$$H = H(x_1, y_1, z_1, x_2, y_2, z_2, \dot{x}_1, \dot{y}_1, \dot{z}_1, \dot{x}_2, \dot{y}_2, \dot{z}_2) \quad (8)$$

Expanding and simplifying (7) yields the Hamiltonian for two charged particles in Weber's electrodynamics

$$H = m_1 \frac{\vec{v}_1 \cdot \vec{v}_1}{2} + m_2 \frac{\vec{v}_2 \cdot \vec{v}_2}{2} + \frac{q_1 q_2}{r} \left( 1 - \frac{\dot{r}^2}{2c^2} \right) \quad (9)$$

or more compactly,

$$\boxed{H = T + V} \quad (10)$$

where  $T$  is given by (3) and  $V$  is the velocity-dependent potential (1).

Before we can apply Hamilton's canonical equations, we perform a *Legendre transform* to express the Hamiltonian in terms of *positions* and *momenta*. For a system of two particles in three spatial dimensions we have

$$H = H(x_1, y_1, z_1, x_2, y_2, z_2, p_{x_1}, p_{y_1}, p_{z_1}, p_{x_2}, p_{y_2}, p_{z_2}) \quad (11)$$

where the velocity  $\dot{x}_1$  in (9) is replaced by  $p_{x_1}/m_1$  and similarly for all other velocity components.

Defining

$$\alpha_x = \frac{q_1 q_2}{c^2} \frac{\dot{r}_{12}(x_1 - x_2)}{r_{12}^2} \quad (12)$$

with analogous definitions for  $\alpha_y$  and  $\alpha_z$ , we apply Hamilton's equations (18) and (19) to obtain

$$\dot{x}_1 = \frac{1}{m_1} (p_{x_1} + \alpha_x) \quad (13)$$

$$\dot{x}_2 = \frac{1}{m_2} (p_{x_2} - \alpha_x) \quad (14)$$

and

$$\dot{p}_{x_1} = \frac{q_1 q_2}{r_{12}^2} \left[ \frac{(x_1 - x_2)}{r_{12}} \left( 1 - \frac{3\dot{r}_{12}^2}{2c^2} \right) + \frac{\dot{r}_{12}(\dot{x}_1 - \dot{x}_2)}{c^2} \right] \quad (15)$$

$$\dot{p}_{x_2} = -\dot{p}_{x_1} \quad (16)$$

with analogous equations for the  $y$  and  $z$  directions. Note that  $\dot{p}_{x_1} + \dot{p}_{x_2} = 0$ , but  $\dot{x}_1 - \dot{x}_2 \neq 0$  in general.

## 2.2 Conserved quantities and symmetries

Weber's electrodynamics for 2 particles conserves energy, momentum and angular momentum. These conserved quantities arise from symmetries in the Hamiltonian (10) according to Noether's theorem.

The conservation of energy follows from the *time-translation symmetry* of the Hamiltonian, meaning that the Hamiltonian has no explicit time dependence.

The conservation of linear momentum arises from the *translational symmetry* of the Hamiltonian, meaning that shifting all particle positions by a constant vector does not change the form of the Hamiltonian.

The conservation of angular momentum is a consequence of the *rotational symmetry* of the Hamiltonian, indicating that rotating the entire system around any axis does not alter the Hamiltonian's structure.

## 2.3 Hamiltonian for $n$ particles

For a system with  $n$  particles, the Hamiltonian (9) generalizes to:

$$H = \sum_{i=1}^n \frac{1}{2} m_i v_i^2 + \sum_{i=1}^n \sum_{j=i+1}^n \frac{q_i q_j}{r_{ij}} \left( 1 - \frac{\dot{r}_{ij}^2}{2c^2} \right) \quad (17)$$

where the double sum is taken over every unique particle pair  $i$  and  $j$ , see Section 3. The conservation laws stated previously still hold for this general case.

For a system of  $n$  particles in one spatial dimension  $x$ , Hamilton's equations of motion are given by

$$\dot{x}_i = \frac{\partial H}{\partial p_{x_i}} \quad (18)$$

$$\dot{p}_{x_i} = -\frac{\partial H}{\partial x_i} \quad (19)$$

for particle  $i \in \{1, 2, \dots, n\}$ . In three dimensions, we also have the analogous equations for  $y_i$  and  $z_i$ .

For a system of  $n$  particles, (18) expands to

$$\dot{x}_i = \frac{1}{m_i} \left( p_{x_i} + \sum_{\substack{j=1 \\ j \neq i}}^n \frac{q_i q_j}{c^2} \frac{\dot{r}_{ij}(x_i - x_j)}{r_{ij}^2} \right) \quad (20)$$

for particle  $i \in \{1, 2, \dots, n\}$ , with analogous expressions for  $\dot{y}_i$  and  $\dot{z}_i$ , and (19) expands to

$$\dot{p}_{x_i} = \sum_{\substack{j=1 \\ j \neq i}}^n \frac{q_i q_j}{r_{ij}^2} \left[ \frac{(x_i - x_j)}{r_{ij}} \left( 1 - \frac{3\dot{r}_{ij}^2}{2c^2} \right) + \frac{\dot{r}_{ij}(\dot{x}_i - \dot{x}_j)}{c^2} \right] \quad (21)$$

for particle  $i \in \{1, 2, \dots, n\}$ , with analogous expressions for  $\dot{p}_{y_i}$  and  $\dot{p}_{z_i}$ .

Equations (20) and (21) together with their analogs for  $y$  and  $z$  directions form a system of  $6n$  first-order ordinary differential equations that describe the time evolution in *phase space* of an  $n$ -particle system for Weber's electrodynamics. For any time  $t$ , the *state* of the system is represented by a point  $\mathbf{z}$  in a  $6n$ -dimensional phase space

$$\mathbf{z} = (x_1, y_1, z_1, \dots, x_n, y_n, z_n, p_{x_1}, p_{y_1}, p_{z_1}, \dots, p_{x_n}, p_{y_n}, p_{z_n}) \in \mathbb{R}^{6n} \quad (22)$$

where every component is a function of time  $t$ . To evolve the system in time using small discrete time steps  $\Delta t$ , we compute


$$x_i^{t+\Delta t} = x_i^t + \Delta t \dot{x}_i^t \quad (23)$$

$$p_{x_i}^{t+\Delta t} = p_{x_i}^t + \Delta t \dot{p}_{x_i}^t \quad (24)$$

and similarly for  $y_i$  and  $z_i$ , where  $\dot{x}_i^t$  and  $\dot{p}_{x_i}^t$  are computed using (20) and (21) at time  $t$ . Repeating this process iteratively allows us to evolve a system of  $n$  charged particles interacting via Weber's electrodynamics over time given some initial positions and momenta.

### 3 Computational complexity of Hamiltonian Weber electrodynamics

To understand the total number of unique interactions, consider an  $n \times n$  matrix where entry  $(i, j)$  represents the interaction between particles  $i$  and  $j$ :



For a system of  $n$  particles, we thus have a total of  $n(n-1)/2$  unique particle pairs, resulting in a computational complexity of order  $\mathcal{O}(n^2)$ .

A naive implementation would compute all  $n(n-1)$  interactions, but by exploiting (12), (13), (14), (15) and (16), we reduce this to  $n(n-1)/2$  interactions.

Furthermore, each evaluation of (20) and (21) only needs to compute  $r_{ij}$  and  $\dot{r}_{ij}$  and their powers once per unique pair, which further reduces the computational load.

### 4 Symplectic numerical integration of non-separable Hamiltonian systems

A numerical method is called *symplectic* if it preserves the *phase space volume* of Hamiltonian systems over time according to *Liouville's theorem*. Standard symplectic methods require the Hamiltonian to be separable into kinetic and potential energy terms that depend only on positions or momenta. The general Hamiltonian for Weber's electrodynamics (17) is non-separable because the velocity-dependent term

$$-\frac{q_1 q_2 \dot{r}_{ij}^2}{2c^2 r_{ij}} \quad (25)$$

is both a function of the positions and momenta. Standard symplectic methods cannot be applied directly. Recent work by [Pihajoki, 2015] and [Tao, 2016] introduced explicit symplectic methods using the clever extended phase space method introduced by [Pihajoki, 2015]. Building on these approaches, [Jayawardana and Ohsawa, 2023] developed a semi-explicit symplectic integrator for non-seperable Hamiltonian systems that also preserves quadratic invariants, see [Ohsawa, 2023]. As of today, this is the most suitable method for our purposes. We describe it briefly here.

We introduce compact vector notation  $\mathbf{q} = (x_1, y_1, z_1, \dots, x_n, y_n, z_n) \in \mathbb{R}^{3n}$  for positions and  $\mathbf{p} = (p_{x_1}, p_{y_1}, p_{z_1}, \dots, p_{x_n}, p_{y_n}, p_{z_n}) \in \mathbb{R}^{3n}$  for momenta, so that  $\mathbf{z} = (\mathbf{q}, \mathbf{p}) \in \mathbb{R}^{6n}$  as in (22). The extended phase space doubles the phase space by introducing another pair of positions and momenta  $\mathbf{x} \in \mathbb{R}^{3n}$  and  $\mathbf{y} \in \mathbb{R}^{3n}$ , giving the state vector in extended phase space

$$\mathbf{Z} = (\mathbf{q}, \mathbf{x}, \mathbf{p}, \mathbf{y}) \in \mathbb{R}^{12n} \quad (26)$$

The condition  $\mathbf{x} = \mathbf{q}$  and  $\mathbf{y} = \mathbf{p}$  is equivalent to a system in the original phase space. If

$$\mathbf{A} = \begin{pmatrix} \mathbf{I} & -\mathbf{I} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I} & -\mathbf{I} \end{pmatrix} \quad (27)$$

where  $\mathbf{I}$  and  $\mathbf{0}$  denote the  $3n \times 3n$  identity and zero matrices, then the condition we are interested in is expressed as

$$\mathbf{AZ} = (\mathbf{q} - \mathbf{x}, \mathbf{p} - \mathbf{y}) = \mathbf{0} \quad (28)$$

We define a *symplectic map* that updates the state vector (26) over a time step  $\Delta t$

$$\Phi_{\Delta t} : \mathbf{Z}_t \mapsto \mathbf{Z}_{t+\Delta t} \quad (29)$$

Using the the update steps (23) and (24) to compute (20) and (21) for  $(\mathbf{q}, \mathbf{p})$  and  $(\mathbf{x}, \mathbf{y})$  separately, we can evolve the system in the extended phase space by defining two separate flows  $\Phi_{\Delta t}^A$  and  $\Phi_{\Delta t}^B$  that we apply in this order

- $\Phi_{\Delta t/2}^A$  evolves  $(\mathbf{x}, \mathbf{p})$  from  $t$  to  $t + \Delta t/2$ , using the values of  $(\mathbf{q}, \mathbf{y})$  at time  $t$
- $\Phi_{\Delta t}^B$  evolves  $(\mathbf{q}, \mathbf{y})$  from  $t$  to  $t + \Delta t$ , using the values of  $(\mathbf{x}, \mathbf{p})$  at time  $t + \Delta t/2$
- $\Phi_{\Delta t/2}^A$  evolves  $(\mathbf{x}, \mathbf{p})$  from  $t + \Delta t/2$  to  $t + \Delta t$ , using the values of  $(\mathbf{q}, \mathbf{y})$  at time  $t + \Delta t$

Our mapping (29) becomes

$$\boxed{\Phi_{\Delta t} = \Phi_{\Delta t/2}^A \circ \Phi_{\Delta t}^B \circ \Phi_{\Delta t/2}^A} \quad (30)$$

This staggered update scheme is referred to as the *leapfrog* method, which is a second-order symplectic integrator.

The key innovation of [Jayawardana and Ohsawa, 2023] is the *symmetric projection method*, which applies a carefully chosen *shift vector*  $\boldsymbol{\mu} \in \mathbb{R}^{6n}$  before and after the explicit evolution step (30) to ensure (28).

The full computation can then be succinctly expressed as a non-linear equation

$$\mathbf{f}(\boldsymbol{\mu}) = \mathbf{A}(\Phi_{\Delta t}(\mathbf{Z}_n + \mathbf{A}^T \boldsymbol{\mu}) + \mathbf{A}^T \boldsymbol{\mu}) = 0 \quad (31)$$

where the solution satisfies (28). One can solve this equation iteratively using Newton's method. Starting from an initial guess  $\boldsymbol{\mu}_{(0)} = 0$ , we compute successive approximations using the *simplified Newton approximations*, see [Jayawardana and Ohsawa, 2023] and [Hairer et al., 2006]

$$\boldsymbol{\mu}_{(k+1)} = \boldsymbol{\mu}_{(k)} - \frac{1}{4} \mathbf{f}(\boldsymbol{\mu}_{(k)}) \quad (32)$$

where  $k$  is the iteration index. We repeat this process until

$$\|\boldsymbol{\mu}_{(k+1)} - \boldsymbol{\mu}_{(k)}\| < \epsilon \quad (33)$$

where  $\epsilon$  is a small tolerance value.

The full specification of this semi-explicit symplectic integrator can be found in Jayawardana and Ohsawa [2023].

## 5 Units

When performing numerical computations we often rely on the 64-bit floating point representation of decimal numbers. This number representation introduces a trade-off between range and precision. To maintain optimal precision, it is advantageous to work with number values near the order of magnitude of 1. In Weber's electrodynamics, we are mostly interested in numerical values that are relatively small when expressed in SI-units. In our numerical computations, we will convert to smaller units to obtain larger numerical values for the same physical quantities.

In the 19th century, Gauss and Weber used an absolute system of units based on the *millimeter*, *milligram*, and *second*. This system is not to be confused with the CGS-system of units, also known as the Gaussian system of units [Assis, 2021a]. (1) and (2) were originally introduced by Weber in this absolute system of units. The units in this system for the potential (1) are  $mg \cdot mm^2/s^2$  and for the force (2) are  $mg \cdot mm/s^2$ .



Thus, the original system of Gauss and Weber, compared to the SI system, is better suited for our computational purposes, though not sufficient. In practice, we need even smaller units like *micrometer*, *microgram* and *microsecond* or even smaller to obtain numerical values near the order of magnitude of 1 for typical values of charge, mass and distance encountered in Weber's electrodynamics.

## 6 Two body problem in Weber's electrodynamics

## 7 Discussion

### 7.1 Details

*Universal material frame of reference* from Relational Mechanics [[Assis, 1999](#)]

## A Notation and expanded definitions

For a system of two particles, we have the following quantities. Particle 1 and particle 2 have electric charges  $q_1, q_2$  and inertial masses  $m_1, m_2$ . In the Gauss-Weber system of units, charge has dimensions of  $\sqrt{mg \cdot mm^3/s^2}$ .

The individual particle positions, velocities and accelerations are

$$\begin{aligned}\vec{r}_1 &= (x_1, y_1, z_1) & \vec{r}_2 &= (x_2, y_2, z_2) \\ \vec{v}_1 &= (\dot{x}_1, \dot{y}_1, \dot{z}_1) & \vec{v}_2 &= (\dot{x}_2, \dot{y}_2, \dot{z}_2) \\ \vec{a}_1 &= (\ddot{x}_1, \ddot{y}_1, \ddot{z}_1) & \vec{a}_2 &= (\ddot{x}_2, \ddot{y}_2, \ddot{z}_2)\end{aligned}$$

The individual particle momenta are

$$\begin{aligned}p_{x_1} &= m_1 \dot{x}_1 & p_{x_2} &= m_2 \dot{x}_2 \\ p_{y_1} &= m_1 \dot{y}_1 & p_{y_2} &= m_2 \dot{y}_2 \\ p_{z_1} &= m_1 \dot{z}_1 & p_{z_2} &= m_2 \dot{z}_2\end{aligned}$$

The relative positions, velocities and accelerations between particle 1 and particle 2 are

$$\begin{aligned}
x &= x_1 - x_2 & \dot{x} &= \dot{x}_1 - \dot{x}_2 & \ddot{x} &= \ddot{x}_1 - \ddot{x}_2 \\
y &= y_1 - y_2 & \dot{y} &= \dot{y}_1 - \dot{y}_2 & \ddot{y} &= \ddot{y}_1 - \ddot{y}_2 \\
z &= z_1 - z_2 & \dot{z} &= \dot{z}_1 - \dot{z}_2 & \ddot{z} &= \ddot{z}_1 - \ddot{z}_2
\end{aligned}$$

The relative position, velocity and acceleration vectors are

$$\vec{r} = (x, y, z)$$

$$\vec{v} = (\dot{x}, \dot{y}, \dot{z})$$

$$\vec{a} = (\ddot{x}, \ddot{y}, \ddot{z})$$

The relative distance is

$$r = |\vec{r}| = \sqrt{x^2 + y^2 + z^2}$$

The relative radial velocity is

$$\dot{r} = \frac{x\dot{x} + y\dot{y} + z\dot{z}}{r} = \frac{\vec{r} \cdot \vec{v}}{r} = \hat{r} \cdot \vec{v}$$

The relative radial acceleration is

$$\begin{aligned}
\ddot{r} &= \frac{x\ddot{x} + y\ddot{y} + z\ddot{z}}{r} + \frac{\dot{x}^2 + \dot{y}^2 + \dot{z}^2}{r} - \frac{(x\dot{x} + y\dot{y} + z\dot{z})^2}{r^3} \\
&= \frac{\vec{r} \cdot \vec{a}}{r} + \frac{\vec{v} \cdot \vec{v}}{r} - \frac{(\vec{r} \cdot \vec{v})^2}{r^3} \\
&= \frac{1}{r} (\vec{r} \cdot \vec{a} + \vec{v} \cdot \vec{v} - \dot{r}^2)
\end{aligned}$$

The relative unit vector is

$$\hat{r} = \frac{\vec{r}}{r}$$

## Acknowledgments

I would like to thank Norman Wildberger, Andre Assis, Ton Kuiper and Koen van Vlaenderen for educating me.

## References

Andre Koch Torres Assis. *Weber's Electrodynamics*. Springer, 1994.

Andre Koch Torres Assis. *Relational Mechanics*. Apeiron, 1999.

- Andre Koch Torres Assis. *Wilhelm Weber's Main Works on Electrodynamics Translated into English: Volume I: Gauss and Weber's Absolute System of Units*. Apeiron, 2021a.
- Andre Koch Torres Assis. *Wilhelm Weber's Main Works on Electrodynamics Translated into English: Volume IV: Conservation of Energy, Weber's Planetary Model of the Atom and the Unification of Electromagnetism and Gravitation*. Apeiron, 2021b.
- Ernst Hairer, Christian Lubich, and Gerhard Wanner. *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. Springer, 2006.
- Buddhika Jayawardana and Tomoki Ohsawa. Semiexplicit symplectic integrators for non-separable hamiltonian systems. *Mathematics of Computation*, 2023.
- Tomoki Ohsawa. Preservation of quadratic invariants by semiexplicit symplectic integrators for non-separable hamiltonian systems. *SIAM Journal on Numerical Analysis*, 61(3):1293–1315, 2023.
- Pauli Pihajoki. Explicit methods in extended phase space for inseparable Hamiltonian problems. *Celestial Mechanics and Dynamical Astronomy*, 2015.
- Molei Tao. Explicit symplectic approximation of nonseparable Hamiltonians: algorithm and long time performance. *Physical Review E*, 2016.