

Does the Brain's E:I Balance Really Shape Long-Range Temporal Correlations? Lessons Learned from 3T MRI

Lydia Sochan¹, Jessica Archibald⁴, Alexander Mark Weber^{1,2,3*}

¹, School of Biomedical Engineering, The University of British Columbia,
Vancouver, BC, Canada.

², BC Children's Hospital Research Institute, The University of British
Columbia, Vancouver, BC, Canada.

³, Pediatrics, The University of British Columbia, Vancouver, BC, Canada.

⁴, Department of Radiology, Weill Cornell Medicine, New York, New York,
USA.

*Corresponding author(s). E-mail(s): aweber@bcchr.ca;

Contributing authors: lydiasochan@gmail.com; jea4025@med.cornell.edu;

Keywords: Hurst exponent, long range temporal correlation, excitatory / inhibitory
balance, criticality, complex systems, visual task, functional magnetic resonance
imaging, magnetic resonance spectroscopy, functional magnetic resonance spectroscopy

¹ School of Biomedical Engineering, The University of British Columbia, Vancouver,
BC, Canada

² BC Children's Hospital Research Institute, The University of British Columbia,
Vancouver, BC, Canada

³ Pediatrics, The University of British Columbia, Vancouver, BC, Canada

⁴ Department of Radiology, Weill Cornell Medicine, New York, New York, USA

* Correspondence: Alexander Mark Weber <aweber@bcchr.ca>

Abstract

A 3T multimodal MR study of healthy adults ($n=18$; 10 female; 21.3 - 53.4 years) was performed to investigate the relationship between fMRI long-range temporal correlations and excitatory/inhibitory balance (via single voxel magnetic resonance spectroscopy measurements of glutamate and GABA). The study objective was to determine if the Hurst exponent (H) — an estimate of self-correlation and signal complexity — of the blood-oxygen-level-dependent signal was correlated with the excitatory-inhibitory (E:I) ratio. E:I has been proposed to serve as a control parameter for brain criticality — the theory that the brain operates near a critical point between order and disorder, optimizing information processing and adaptability — which H is believed to be a measure of. Thus, understanding if H and E:I are indeed correlated would clarify this relationship. Moreover, findings in this domain have implications for neurological and neuropsychiatric conditions with disrupted E:I balance, such as autism, schizophrenia, and Alzheimer’s disease. From a practical perspective, H is easier to accurately measure than E:I ratio at 3T MRI. If H can serve as a proxy for E:I, it may serve as a more practical clinical biomarker for this imbalance and for neuroscience research in general. The study collected functional MRI and magnetic resonance spectroscopy data during rest and movie-watching. H and E:I (glutamate/GABA) were not found to be correlated. H was found to increase with movie-watching compared to rest, while E:I did not change between conditions. This study represents the first attempt to investigate this connection *in vivo* in humans. We conclude that, at 3T and with our particular methodologies, no association was found. We end with lessons learned and suggestions for future research.

1 Introduction

The human brain consists of interconnected networks of neurons whose collective activity gives rise to complex behaviors and cognition. In order to understand how neural systems generate this emergent functionality, neuroscience is increasingly using tools and concepts from the science of complexity. Recent advances have led to the emergence of the Critical Brain Hypothesis, which posits that the brain — across all regions and networks — operates at a ‘critical point’, a state where both order and disorder are balanced to enable the most efficient information processing^{1–14}. In fact, criticality is perhaps the main way that nature produces stable complex systems^{15,16}. When in a critical state, the brain is optimally responsive to both internal and external stimuli while maintaining a balance between stability and instability^{1,2,13,14}.

Currently, the most popular theory proposing how the brain achieves this delicate balance — what is known as the ‘control parameter’ — is the excitation-inhibition (E:I) ratio: the balance of excitatory and inhibitory neural activity, often operationalized as the ratio of the primary excitatory and inhibitory neurotransmitters (i.e. glutamate (Glu) and γ -aminobutyric acid (GABA))^{6,8–12}. This theory essentially relies on the fact that E:I is a primary driver of the signal-to-noise ratio (SNR) of neuronal signalling, and hence could drive transitions between order and disorder^{6,17}. Thus, the E:I theory of criticality posits that altering the E:I ratio would allow for movement between

sub- and super-critical states⁶. When excitation is perfectly balanced with inhibition, the system is at a critical point, but if either excitation or inhibition are favoured, the system can become overly noisy (excess excitation) or overly controlled (excess inhibition), respectively⁶. Thus, the relationship between E:I balance and large-scale brain dynamics offers a promising link between synaptic physiology and macroscopic neural computation, and for elucidating fundamental neural computational principles. Furthermore, understanding this relationship could hold great clinical relevance, as disruptions in E:I homeostasis have been implicated in multiple neuropsychiatric and neurodevelopmental disorders — including autism, schizophrenia, and Alzheimer’s disease — making this an important target for biomarker development and therapeutic intervention^{10,18,19}.

In vivo measurement of brain criticality can be achieved using several available methods, as systems near a critical state often exhibit fractal-like fluctuations or scale-invariance²⁰. Scale-invariance has been observed extensively across various neuronal spatio-temporal scales, from dendritic branching structures²¹ in the spatial domain, to neurotransmitter release²², neuronal firing rates²³, local field potentials²⁴, electroencephalography (EEG)²⁵ and functional magnetic resonance imaging (fMRI) — which measures the blood oxygen level-dependent (BOLD) signal^{26–28} — in the time domain. This fractal structure can be quantified by assessing the signal’s long-range temporal correlations (LRTC), which reflect the persistence or memory in a time series, where future values are statistically related to past values over long extended periods. The Hurst exponent (H) is a useful measure for evaluating LRTC^{28,29}. An H value between 0.5 and 1 indicates a long-range positive correlation, with large values being followed by large values and small values by small ones. Conversely, an H exponent value between 0 and 0.5 indicates a long-range negative correlation, where large values are likely to be followed by small ones and vice versa. An H value of 0.5 is uncorrelated random noise³⁰.

H has emerged as a valuable tool in neuroscience and clinical research. Typically, H values reported in adult brains are above 0.5, with higher H values in grey matter than white matter or cerebrospinal fluid^{31,32}. Some key findings from neuroscience research include: a decrease in H during task performance^{33,34}; an increase in H during movie watching in the visual resting state network³⁵; negative correlations with task novelty and difficulty³⁶; increases with age in the frontal and parietal lobes³¹, and hippocampus³⁷; decreases with age in the insula, and limbic, occipital and temporal lobes³¹; and $H < 0.5$ in preterm infants³⁸. For a more in depth review, see Campbell et al. (2022)²⁸. In terms of clinical findings, abnormal H values have been identified in Alzheimer’s disease (AD)^{39,40}, autism^{41–44}, mild traumatic brain injury⁴⁵, major depressive disorder^{46,47} and schizophrenia^{44,48}. Crucially, these same disorders have been associated with imbalances in E:I^{10,18,19,49–52}.

In addition to its implications for the critical brain hypothesis, establishing a connection between E:I and H could facilitate simpler estimation of excitatory and inhibitory neurotransmitters, as precise measurement of E:I is technically challenging⁵³. Currently, magnetic resonance spectroscopy (MRS) is the only non-invasive method for *in vivo* assessment of the Glu/GABA ratio (excitatory to inhibitory neurotransmitters) in humans^{54,55}. Unfortunately, MRS suffers from limited spatial and temporal resolution,

with single-voxel spectroscopy sizes between 20 and 27 cm³, and scan times between 5-10 minutes^{12,53,54}. In addition, quantification of GABA at 3T necessitates the use of a specialized J-difference spectral-editing sequence (such as MEGA-PRESS), in order to isolate GABA (at 3 ppm, 2.3 ppm, and 1.9 ppm) from overlapping peaks (e.g. creatine and glutamate)⁵⁶. The need for this additional sequence therefore reduces the temporal resolution further. If H could function as a substitute for E:I, it could simplify the estimation of E:I.

Beyond clinical observations suggesting that both H and E:I imbalance are altered in neuropsychiatric conditions, several studies have attempted to show that critical-state dynamics emerge when excitation and inhibition are balanced. Indirect evidence comes from pharmacological manipulation of GABAergic neurotransmission, which has been shown to change the strength of LRTC *in vitro*⁵⁷ and in the beta-frequency range in humans using EEG⁵⁸; and to eliminate the power-law scaling of neuronal avalanches in rats⁵⁹⁻⁶¹. More direct efforts linking H and E:I have mostly relied on computational models or animal studies⁶⁻¹², and the results remain inconsistent, ranging from positive to negative or even U-shaped associations. These discrepancies may reflect differences in model systems (e.g., animal vs. computer simulations), methodological choices (such as how H or E:I are estimated), or may represent counter evidence against the theory that these measures are systematically related. Indeed, other theories for how the brain maintains criticality exist, such as neuromodulator regulation⁶²⁻⁶⁴ or synaptic plasticity⁶⁵, **ikeda2025emergentfunctionsnoisedrivenpontaneous?**, to name a few. Despite the lack of direct evidence, several recent human studies have used fMRI-derived H as a direct proxy for E:I balance^{43,44,66,67}; yet, no study to date has tested the empirical relationship between H and neurochemical estimates of E:I (e.g., Glu/GABA ratio) *in vivo* in humans.

The theory that E:I serves as the primary control parameter governing H — and, by extension, brain criticality — offers a framework with clear, falsifiable predictions that can be directly tested. Specifically, if this framework holds, E:I and H should covary systematically in the human brain across individuals, age, brain regions, and cognitive states. To empirically test this prediction, the present study examined the E:I–Hurst relationship in the human visual cortex during rest and naturalistic movie-watching. Although a single condition would be sufficient to test the correlation between E:I and H, we included two conditions to broaden the dynamic range of neural and neurochemical measures — thereby increasing our sensitivity to detect a correlation — and to provide validation that our metrics behave as expected across qualitatively different brain states. Naturalistic movie-watching is known to reliably engage the visual cortex with complex, temporally structured input, making it a useful condition for probing changes in the temporal dynamics of neural activity^{35,68}. Prior work from our group has shown that H values increased during movie-watching compared to rest³⁵, while work from others have found changes in Glu and GABA with sustained visual stimulation⁶⁹⁻⁷². We hypothesized that H and Glu would increase during movie-watching^{35,69,70}, that GABA would decrease^{71,72}, and that H would be negatively correlated with E:I in both conditions¹¹.

2 Methods

2.1 Participants

Twenty-seven healthy adult participants were recruited to the study. One participant was not scanned due to claustrophobia while in the scanner. After our analysis and performing quality assurance (see below for details), a further eight participants were removed, leaving eighteen final participants, between the ages of 21.3 and 53.4 (mean age $\pm$ sd: 29.6 ± 8.7 years; 8 males). Sex, handedness and date of birth were recorded on the day of the scan.

2.2 Ethics Statement

Written informed consent was obtained from all participants. Ethics approval was granted by the Clinical Research Ethics Board at the University of British Columbia and BC Children’s & Women’s Hospital (H21-02686).

2.3 Scanning Procedure

After two anatomical sequences were acquired, participants were instructed to visually fixate on a cross-hair for 24 minutes (see Table 1 and below for more details). During this period, an fMRI, single-voxel semi Localization by Adiabatic SElective Refocusing (sLASER)⁷³, and single-voxel MEscher-Garwood Point-REsolved SpectroScopy (MEGA-PRESS)⁷⁴ sequences were acquired (see Figure 1 A and Section 2.4). Next, participants were instructed to watch a nature documentary (Our Planet (2019), Episode 3, “Jungles”⁷⁵) for 24 minutes. During this period, another set of fMRI, sLASER, and MEGA-PRESS sequences were acquired. See Figure 1 B for a visual representation of the scanning protocol. Total scan duration was approximately 1 hour of uninterrupted scanning (i.e. subjects were not removed from the scanner). All participants followed the same order of conditions: rest then movie; all participants saw the same movie segment, beginning at the same time during the scan.




Figure 1. Overview of the MRI acquisition protocol. A) fMRI coverage was across the whole-brain (example coverage in red). MRS voxels were placed in the visual cortex (blue). Background image is of a T1w acquisition from a sample subject. Figure was inspired by Ip et al. (2017)⁷⁶. B) fMRI, sLASER, and MEGA-PRESS were acquired first with participants looking at a white cross for 24 minutes. Next, the same sequences were acquired with the participants viewing a nature documentary for an identical period of time.

2.4 Acquisition Details

Scans were performed at BC Children’s Hospital MRI Research Facility on a 3.0 Tesla GE Discovery MR750 scanner (scanner software version: DV26.0_R03) with a Nova Medical 32 channel head coil. Participants changed into scrubs and were screened by an MRI technologist. Participants were given wax earplugs and a fiducial was placed on their left temple. Participants were provided with an audio headset and blanket once lying down on the scanner bed. Since visual stimuli were rear-projected, position and angle of mirror above patient eyes was adjusted for optimal movie viewing.

The following MRI scans were acquired. A 3D T1-weighted sagittal fast spoiled gradient echo (FSPGR) sequence; a 3D T2-weighted sagittal CUBE; a 2D echo-planar imaging (EPI) multi-echo gradient-echo fMRI sequence; a single-voxel MEGA-PRESS sequence with CHESS (CHEMical Shift Selective saturation) water suppression; and a single-voxel sLASER sequence with VAPOR (VARIABLE Power and Optimized Relaxations) water suppression. Details are listed in Table 1.

Sequence	TE (ms)	TR (ms)	Flip Angle	FOV	Slice Thick- ness (mm)	In-Plane Resolu- tion (mm ²)	Other Parameters	Time (mins)
3D T1- weighted FSPGR	2.176	7.216	12	256 × 256	0.9	0.9375 × 0.9375	-	5
3D T2- weighted CUBE	75.242	2,504.090		256 x 256	0.9	0.9375 × 0.9375	-	4
2D EPI	12.2,	1,500	52	64 x	3.6	3.5938 ×	Acceleration	12
Multi-Echo	35.352,			64		3.5938	factor = 6	
fMRI	58.504							
sLASER	35	2,000	90	-	28.0	28.0 x 28.0	Transients = 128*; Spectral width = 5,000 Hz; Data points = 4,096	4
MEGA- PRESS	68	1,800	90	-	28.0	28.0 x 28.0	Transients = 128 ON, 128 OFF; Spectral width = 5,000 Hz; Data points = 4,096	8

Table 1. Summary of MRI acquisition details TE = echo time; TR = repetition time; FOV = field of view; FSPGR = fast spoiled gradient-echo; CUBE = a GE acronym; EPI = echo-planar imaging; fMRI = functional magnetic resonance imaging; sLASER = semi localization by adiabatic selective refocusing; MEGA-PRESS = Mescher-Garwood point-resolved spectroscopy. *Note, the GE semi-LASER sequence averaged the 128 transients into groups of 4, each with 32 transients, before being saved as raw P-file.

For both MRS sequences, the voxel size was set to 2.8 x 2.8 x 2.8 cm³. MRS voxels were rotated and placed in the occipital lobe, aligned along the calcarine fissure, using the high resolution T1w sequence as anatomical guide. Voxel placement was set once at the beginning and copied for the remaining MRS sequences. Glutamate was quantified from semi-LASER acquisitions as it offers the highest quality spectra for unedited metabolites⁷⁷. The GE semi-LASER sequence automatically aligned and averaged every 32 transients, resulting in 4 averaged transients to process. GABA was measured

using MEGA-PRESS, which applies frequency-selective editing pulses to isolate GABA from overlapping signals⁵⁶. This dual-sequence approach is standard in the field and reflects the differing spectral requirements of each metabolite⁷⁸. Finally, blip-up and blip-down spin-echo versions of the fMRI sequence were acquired at the end to estimate the B0 non-uniformity map for fMRI phase distortion correction.

2.5 Image Processing

Our full image processing pipeline has been can be accessed from our Github account page: github.com/WeberLab/EI_Hurst_Analysis

Images were downloaded offline from the scanner in raw Digital Imaging and Communications in Medicine (DICOM) format. DICOM files were then converted to Neuroimaging Informatics Technology Initiative (NIfTI) using Chris Rorden’s `dcm2niix`⁷⁹ (v1.0.20211006) and then to Brain Imaging Data Structure (BIDS)⁸⁰ format using `dcm2bids`⁸¹ (v2.1.6). MRS files were downloaded as raw P files.

2.5.1 Structural Images

The T1w image was corrected for intensity non-uniformity with `N4BiasFieldCorrection`⁸² (`ANTs`⁸³; v2.3.335) to be used as a T1w-reference for the rest of the workflow. The T1w-reference was skull-stripped using a `Nipype`⁸⁴ implementation of `antsBrainExtraction.sh` from `ANTs`; `OASIS30ANTs` was used as a target template. `Fast`⁸⁵ (`FSL`⁸⁶ v.6.0.5.1:57b01774, RRID: SCR_002823) was used for brain tissue segmentation into cerebrospinal fluid (CSF), white matter (WM), and gray matter (GM). Brain surfaces were reconstructed with `recon-all`⁸⁷ (`FreeSurfer`⁸⁷ 7.3.2, RRID: SCR_001847). The previously-estimated brain mask was refined with `Mindboggle`⁸⁸ (RRID:SCR_002438) to reconcile `ANTs`-derived and `FreeSurfer`-derived segmentations of cortical GM. `AntsRegistration`⁸³ (`ANTs` 2.3.3) was used to perform volume-based spatial normalization to two standard spaces: MNI152NLin2009cAsym and MNI152NLin6Asym. Normalization used brain-extracted versions of both T1w reference and T1w template.

2.5.2 fMRI

Using `fMRIPrep`⁸⁹, the shortest echo of the BOLD run was used to generate a reference volume (both skull-stripped and skull-included). Head-motion parameters with respect to the BOLD reference (transformation matrices as well as six corresponding rotation and translation parameters) were estimated before spatiotemporal filtering using `mcflirt`⁹⁰ (`FSL` v6.0.5.1:57b01774). The fieldmap was aligned with rigid registration to the target EPI reference run. Field coefficients were mapped to the reference EPI using the transform. BOLD runs were slice-time corrected to 643 ms (half of slice acquisition range of 0-1290 ms) using `3dTshift` from `AFNI`⁹¹ (RRIS: SCR_005927). To estimate T2* map from preprocessed EPI echoes, a voxel-wise fitting was performed by fitting the maximum number of echoes with reliable echoes in a particular voxel to a monoexponential signal decay model with nonlinear regression. Initial values were *T2/S0 estimates from a log-linear regression fit. This calculated T2 map was then*

used to optimally combine preprocessed BOLD across echoes using the method by Posse et al. (1999)⁹². The generated BOLD reference was then co-registered (6 degrees of freedom) to the T1w reference with **bbregister** (**FreeSurfer**⁸⁷) using boundary-based registration. First, a reference volume and its skull-stripped equivalent were generated with **fMRIPrep**. Confounding time series were calculated from preprocessed BOLD: framewise displacement (FD), DVARS, and three region-wise global signals. **Tedana**⁹³ was then used to denoise the data by decomposing the multi-echo BOLD data via principal component analysis (PCA) and independent component analysis (ICA). The resulting components are automatically analyzed to determine whether they are TE-dependent or -independent. TE-dependent components were classified as BOLD, while TE-independent components were classified as non-BOLD and were discarded as part of data cleaning.

Participants were excluded from further analysis if their mean FD was > 0.15 mm.

2.6 MRS

sLASER data were processed using **Osprey**⁹⁴ (v2.9.5) and fit via the embedded **LCModel** (v6.3-1N) wrapper⁹⁵. Because of substantial overlap among Glu and Gln resonances^{96,97}, in order to avoid errors in spectral assignment — especially since it is controversial whether Glu can reliably be separated from Gln at 3T^{98,99} — we report Glx (Glu+ Gln) as the primary outcome measure. Individual Glu estimates are provided in Supplementary Materials Table 6, alongside all other quantified metabolites. Several Osprey functions were modified to accommodate scanner-specific features of the GE implementation, where individual transients had been averaged by the scanner into four sub-spectra prior to export. These changes primarily affected **GEload.m** to correct water-scaling inconsistencies. RF coil combination was performed using generalized least squares (GLS) weighting, following An et al. (2013)^{100,101}.

MEGA-PRESS data were processed with **Osprey**, and the difference of the ON and OFF sequences were fit the **LCModel** algorithm. Due to the J-editing sequence of MEGA-PRESS, a challenge of GABA quantification is macromolecule quantification¹⁰². As a result, we report GABA+, a measure which co-reports GABA with macromolecules. Macromolecules (MM) are expected to account for approximately 45% of the GABA+ signal¹⁰². While a macromolecule-suppressed estimate of GABA seems ideal, a recent 25-site and multi-vendor study conducted at 3T found that GABA+ showed much lower coefficient of variation than MM-suppressed GABA, meaning that GABA+ is more consistent across research sites and MRI vendors (i.e., Philips, GE, Siemens)¹⁰³. Moreover, GABA+ shows greater reliability for both creatine-referenced and water-suppressed estimates^{103,104}. MM-suppressed GABA and GABA+ estimates are also correlated, albeit weakly- to moderately-so^{102–104}. Consequently, we report GABA+ to allow for easier comparison of our results to other studies as well as reproducibility. All metabolite values, including GABA, are reported in the Supplementary Materials Table 7.

MRS voxels were co-registered to the T1w reference image and segmented by **SPM12**¹⁰⁵ into CSF, GM, and WM. sLASER and MEGA-PRESS data were water-scaled

as well as tissue- and relaxation-corrected by the Gasparovic et al. (2006) method¹⁰⁶. Concentrations are reported in millimoles. Full width half-maximum (FWHM) of the single-Lorentzian fit of the water peak, signal-to-noise (SNR) ratio of the creatine signal, frequency shift, and Cramer-Rao lower bounds (CRLB) were calculated for quality assurance purposes. **Osprey** basis sets were used for linear combination. Metabolites included for both basis sets were: Asc, Asp, Cr, CrCH2, EA, GABA, GPC, GSH, Gln, Glu, Gly, H2O, Lac, NAA, NAAG, PCh, PCr, PE, Ser, Tau, mI, and sI. Excitatory-inhibitory ratio (E:I) was calculated as [Glx from sLASER]/[GABA+ from MEGA-PRESS], a common practice to report E:I using MRS¹⁰⁷.

Participants were excluded from further analysis if any of their MRS scans had a water peak FWHM $> 10^{108}$.

To quantify the inter-subject consistency across the MRS masks, we computed a consensus mask using majority voting and calculated the mean Dice coefficient between each individual mask and the consensus.

2.6.1 Re-analysis using Gannet

Glutamate and GABA were acquired using different MRS sequences and in separate blocks during the same scan session. While acquisitions were closely timed and performed under similar conditions, we cannot fully exclude the possibility of session-related variability affecting E:I ratio estimates. Therefore, we re-ran our analysis using the Glx and GABA+ values from the MEGA-PRESS sequence alone using **Gannet**¹⁰⁹ (v3.3.0).

2.7 Hurst Exponent Calculation

Hurst exponent was calculated from the power spectrum density (PSD) of the BOLD signal. A log-log plot was used, where log power was plotted against log frequency; generally, if a log-log plot results in a linear relationship, it is assumed that the mean slope of this line represents the power-law exponent⁵. A PSD shows the distribution of signal variance ('power') across frequencies. Complex signals are classified into two categories: fractional Gaussian noise (fGn) and fractional Brownian motion (fBm)^{29,110}. The former is a stationary signal (i.e., does not vary over time), while the latter is non-stationary with stationary increments²⁹. Most physiological signals consist of fBm, but fMRI BOLD is typically conceptualized as fGn once motion-corrected; otherwise put, unprocessed BOLD signal is initially fBm which is converted to fGn with appropriate processing¹¹¹. fBm and fGn require distinct H calculation methods²⁹. PSD was estimated using Welch's method¹¹² from the Python **Scipy.Signal** library¹¹³. Data were divided into 8 windows of 50% overlap and averaged within each window, as is standard¹¹⁴. Welch's method was selected based on prior comparisons with other Hurst exponent algorithms, which found it to perform best across all evaluation criteria¹¹⁴. The spectral index, β , was calculated from the full frequency spectrum. The spectral index was then converted to H using the following equation^{29,115}:

$$H = \frac{1 + \beta}{2}$$

Since it cannot be assumed that all fBm is removed from the signal, we employed the ‘extended Hurst’ (H’) concept in this study: for $0 < H < 1$, the signal is understood as fGn, while for $1 < H < 2$, the signal is understood to be fBm^{35,116,117}. More generally, it is assumed that when $0.5 < H < 1.5$, the signal displays 1/f behaviour⁵. H was calculated for all voxels in the brain of each subject. A brain mask was then applied which included only GM and the region of the MRS voxel in the visual cortex. H was averaged across the brain mask area, using only non-zero voxels.

2.8 Statistics

All statistical analyses were performed using R¹¹⁸ (v4.5.1) and RStudio.

A linear mixed-effect model was used to test for the correlation of H and E:I, with H as the dependent variable, Glx/GABA+ as the primary predictor of interest, and Condition (rest vs. movie) as a fixed effect. Mean FD, FWHM(sLASER), FWHM(MEGA-PRESS), Frequency Shift (sLASER), Frequency Shift (MEGA-PRESS) were included as covariates, and subjects were modeled as random effects:

$$\begin{aligned} H \sim & \frac{\text{Glx}}{\text{GABA+}} + \text{Condition} + \text{meanFD} \\ & + \text{FWHM}_{\text{sLASER}} + \text{FWHM}_{\text{MEGA-PRESS}} \\ & + \text{FreqShift}_{\text{sLASER}} + \text{FreqShift}_{\text{MEGA-PRESS}} \\ & + (1|\text{Subject}) \end{aligned}$$

A post-hoc calculation of the power of the model was calculated using `powerSim` from the `simr` package using 1,000 simulations.

Exploratory post hoc analyses were then performed. Difference in the means of H, Glx, and GABA+ between rest and movie conditions were assessed using paired Student’s t-tests¹¹⁹. Correlations between H and Glx, GABA+ and E:I were calculated using Pearson’s method¹²⁰.

3 Results

3.1 Participant Demographics

Twenty-seven participants were originally recruited for the study. Twenty-six of these participants were successfully scanned, but one participant experienced claustrophobia and chose not to continue. Of the remaining 26 participants, 18 were included in the final analysis: three were removed due to low MRS quality (FWHM > 10) and five were removed due to low fMRI quality (mean FD > 0.15 mm). See Figure 2.

The final study sample included 8 males and 10 females between ages 21.3 and 53.4, with a mean age and standard deviation of 29.6 ± 8.7 years.




Figure 2. Flowchart of participant, recruitment, scanning and exclusion. The number of participants at each stage is indicated within each node. Participants were excluded based on MRS FWHM and fMRI Mean FD thresholds, resulting in 18 participants included in the final analysis

3.2 Data Summary

An average of all MRS voxel placements can be seen in Figure 3 A, and a sample of the Osprey sLASER and MEGA-PRESS spectrum fits at rest can be seen in Figure 3 B and C, respectively. The mean consensus-based inter-subject Dice score for MRS voxel placements was 0.77 ± 0.14 (range: [0.35, 0.95]).




Figure 3. MRS data quality A) Average MRS location. Colored overlay of the average voxel location across all nineteen participants, from 0 (black) to 1 (white). Brighter yellow represents voxel locations shared by all participants, while darker purple represents voxel locations unique to participants. MRS voxels were registered to MNI space and averaged. Underlay is T1w MNI152 at 0.5 mm resolution. B) Sample Osprey sLASER Spectrum. Red spectrum indicates overall model fit. Model fits for individual metabolites are shown in blue below overall fit. C) Sample Osprey MEGA-PRESS Spectrum. Yellow spectrum indicates overall model fit. Model fits for individual metabolites are shown in blue below overall fit.

Average water FWHM, creatine SNR, and frequency shift are reported in Supplementary Material Table 4.

Average voxel rotation along the x-axis, along with average tissue fractions are reported in Supplementary Material Table 5.

A sample of the combined grey-matter and MRS voxel mask used to average H values, along with a sample Hurst exponent map, and sample fits for H calculation during rest can be found in Figure 4 A, B, and C, respectively.




Figure 4. fMRI data quality. A) sample fMRI grey-matter mask within the MRS voxel. Background: a sample coronal, sagittal, and axial slice is displayed of the mean fMRI scan from the rest acquisition. Foreground: the greymatter/MRS mask used to calculate mean H. B) Sample H map for whole brain. Coronal, sagittal, and axial views of are shown, colour-coded by H values. Colour-coding shows evident tissue differentiation by H value (i.e., gray matter (red), white matter (light red), and cerebrospinal fluid (white)). C) Sample PSD spectrum. All participants' PSD spectrums during rest are plotted using light grey lines. Mean PSD is plotted in solid red Mean linear regression line is plotted in dashed blue. H for each participant was calculated from the slope of their mean linear regression line.

Figure 5 shows the average Hurst maps during rest and movie watching.




Figure 5. Average Hurst maps for rest and movie conditions Group averaged maps of the Hurst exponent computed to visualize whole-brain patterns across sessions: rest (top) and movie watching (bottom.)

3.3 Linear-mixed effects model: H and E:I

To examine the relationship between H and E:I, we ran a linear mixed-effects model. The model's total explanatory power was substantial (conditional $r^2 = 0.79$), with fixed effects (marginal r^2) equal to 0.22. The model's intercept, corresponding to $E:I = 0$, Condition = Movie, $FWHM_{sLASER} = 0$, $FWHM_{MEGA-PRESS} = 0$, $FreqShift_{sLASER} = 0$, $FreqShift_{MEGA-PRESS} = 0$, and $meanFD = 0$, was 1.3 (95% CI [0.66, 1.95], $t(26) = 3.95$, $p < 0.01$).

Among the fixed effects, only the effect of Condition [Rest] ($\beta = -0.07$, 95% CI [-0.11, -0.02], $t(26) = -2.84$, $p = 0.02$); and Frequency Shift (sLASER) ($\beta = 0.03$, 95% CI [0.01, 0.06], $t(26) = 2.31$, $p = 0.05$) were statistically significant. All other predictors — including E:I ($\beta = 0.07$, 95% CI [-0.05, 0.19], $t(26) = 1.12$, $p = 0.28$); along with $FWHM_{MEGA-PRESS}$ ($\beta = 0.01$, 95% CI [-0.05, 0.07], $t(26) = 0.28$, $p = 0.78$); $FWHM_{sLASER}$ ($\beta = -0.02$, 95% CI [-0.09, 0.05], $t(26) = -0.58$, $p = 0.57$); $FreqShift_{MEGA-PRESS}$ ($\beta = 0.03$, 95% CI [-0.01, 0.06], $t(26) = 1.56$, $p = 0.14$); $FreqShift_{sLASER}$ ($\beta = 0.03$, 95% CI [0.01, 0.06], $t(26) = 2.31$, $p = 0.05$); and $meanFD$ ($\beta = -0.59$, 95% CI [-2.14, 0.96], $t(26) = -0.75$, $p = 0.46$) — were not statistically significant.

3.4 Post-hoc analyses

Diagnostics of our linear-mixed effects model (check for normality) can be found in the Supplementary Materials (Figure 8)

To calculate the post-hoc power of our model, we required an effect size. While *in silico* and animal studies indicate that changes in E:I ratio influence the Hurst exponent¹¹, these models do not provide standardized effect sizes, and translation to human neuroimaging remains uncertain. Therefore, we used several effect sizes, and report here the power of each. The effect sizes we used to calculate power were: the value calculated from our model ($\beta = 0.07$); and β values two and three times that size: 0.14 and 0.21. As a reminder, β represents the increase of H by that amount per every increase of E:I by 1. The power for these effect sizes were 25%, 62.7%, and 92.9%, respectively.

Furthermore, as our spectroscopy quality metrics could be highly correlated with each other (e.g., FWHM measures from the two sequences, or frequency shifts), we checked variance inflation factors (VIFs) to ensure all values were < 5 (generally acceptable). VIFs were: EI = 1.24, Condition = 1.55, FWHMsLASER = 1.76, FWHMMEGA = 1.45, FreqShiftsLASER = 1.15, FreqShiftMEGA = 1.48, and MeanFD = 1.28.

Mean $\pm$ sd of the main metabolites, their Cramer-Rao lower bounds, E:I, and H during rest and movie are reported in Table 2. Boxplots between rest and movie can be seen in Figure 6. Neither Glx nor GABA+ were different between movie and rest conditions. E:I ratio did not change between conditions either. H was found to be greater during movie watching than rest.

	Rest	Movie	p-value
Glx	9.84 ± 1.06	10.04 ± 0.79	0.26
GABA+	5.48 ± 0.77	5.23 ± 0.71	0.17
E:I Ratio	1.82 ± 0.25	1.94 ± 0.25	0.08
H	0.98 ± 0.98	1.05 ± 1.05	< 0.01

Table 2. Summary of main measures Glx and GABA+ values are in millimoles.




Figure 6. Paired comparison of metabolite values during rest (green) and movie (orange) conditions. A) Glx (mM); B) GABA+ (mM); and C) E:I. Paired dots represent the same participant across conditions. Mean difference with 95% confidence intervals, and as Cohen's D are reported at the top of each plot.

H was not found to correlate with Glx, GABA+, or E:I, during rest or movie (Figure 7).




Figure 7. Scatter plots of H vs. metabolites. A) Glx (mM); B) GABA+ (mM); and C) E/I. Rest is in green, while movie is in orange. p-values are reported at the top of each plot.

3.4.1 Data quality

Mean FD was not correlated with H during rest ($r = -0.33$, $p = 0.18$) but was moderately negatively correlated with H during movie watching ($r = -0.49$, $p = 0.04$; see Supplementary Materials Figure 9 A).

Glx and GABA+ were tested for associations with FWHM values during rest and movie watching. No significant correlations were found (see Supplementary Materials Figure 9 B and C). Glx and GABA+ were tested for associations with frequency shift during rest and movie watching. No significant correlations were found (see Supplementary Materials Figure 9 D and E).

Furthermore, a group averaged plot of the frequency shift between rest and movie can be found in the Supplementary Materials (Figure 10).

3.5 Re-analysis using Gannet

The re-analysis of our results using only the MEGA-PRESS sequence and **Gannet** did not result in any significant changes from the results reported above. We have included these results in the Supplementary Material.

4 Discussion

This study provides the first *in vivo* human investigation of the relationship between H and the E:I ratio, measured using fMRI and MRS-derived glutamate and GABA levels, respectively. No association was observed between H and E:I, suggesting that these indices may reflect distinct dimensions of neural dynamics. During movie-watching, H increased relative to rest, consistent with stronger long-range temporal dependencies in BOLD activity during visual stimulation. In contrast, Glx, GABA+, and their ratio (E:I) remained stable across conditions. These results indicate that functional signal complexity, as indexed by H, is sensitive to sensory engagement but does not appear to covary with MRS-derived estimates of excitatory–inhibitory balance under the current experimental conditions.

An increase in H during movie-watching relative to rest aligns with previous work from our lab³⁵, which reported increased H in the visual network (as defined by Yeo’s seven resting-state networks¹²¹) during movie-watching using data from the Human Connectome Project¹²². While this increase in H is consistent with our prior findings, other studies have reported decreases in H during active tasks^{33,34,36,123}. The current results therefore suggest that the naturalistic, passive nature of movie-watching elicits a different effect on H compared to more active tasks. This interpretation is consistent with evidence indicating distinct neural responses and BOLD signal characteristics between conventional active visual tasks and naturalistic passive visual stimuli^{35,124}. Higher H during movie-watching may reflect richer scaling properties that support the continuous integration of visual stimuli³⁵.

No significant changes were observed in either Glx or GABA+ between conditions. Consequently, the E:I ratio remained stable. The experimental design included both resting and movie-watching conditions to induce measurable differences in H, Glx, and GABA+, providing a wider range of values to test for correlations between these metrics. However, the absence of condition-related differences in Glx and GABA+ complicates interpretation of the null findings. Establishing such changes would have provided a foundation for directly examining the correlation between H and the E:I ratio. Without clear evidence that the paradigm elicited detectable neurochemical variation, it remains uncertain whether the lack of association between H and E:I reflects a genuine independence of these measures or limited sensitivity of the MRS protocol to capture subtle metabolic changes.

A recent meta-analysis of studies measuring Glu and Glx⁹⁶ reported a minimal task-induced increase in Glu and Glx within the visual cortex, primarily in the context of pain, learning, or motor paradigms rather than visual stimulation. Additionally, many of the studies included were conducted at 7T, offering increased sensitivity for

detecting changes in Glu and Glx. Nonetheless, several studies have demonstrated changes in Glu and Glx at 3T^{125–127}. Regarding our GABA+ findings, the same meta-analysis⁹⁶ reported no task-dependent change in GABA within the visual cortex. This may be attributable to technical difficulties associated with capturing GABA levels using MRS at 3T due to its low concentration and spectral overlap with more abundant metabolites⁹⁶. Although, several studies have successfully reported changes in GABA at 3T using different paradigms and/or regions of interest^{128–130}.

Another reason changes in Glx or GABA+ may not have been observed could be due to our use of a block-design: collecting 24 minutes of rest data, then collecting 24 minutes of movie-watching data. While a block design has the advantage of a more robust metabolite quantification due to greater signal averaging, brain homeostasis during these long blocks may lead to an erasure of any real metabolic changes^{127,131}. Indeed, Pasanta et al. (2023)⁹⁶ and Archibald et al. (2020)¹³² reported larger metabolite changes during event-related paradigms compared to block designs, underscoring how task timing and structure can strongly influence the detectability of metabolic modulation with MRS.

The concept of E:I balance encompasses multiple biological scales and mechanisms beyond neurotransmitter concentrations, including the relative densities of excitatory and inhibitory neurons, synaptic properties, interneuron subtype interactions, and electrophysiological signatures such as 1/f spectral scaling. While the glutamate-to-GABA ratio measured via MRS provides a noninvasive — albeit indirect — estimate of neurochemical E:I balance in humans, it remains a simplification of the underlying complexity. MRS derived values reflect total glutamate and GABA metabolite pools, predominantly intracellular and metabolically active, rather than direct synaptic neurotransmitter release underlying real-time excitation and inhibition. Moreover, conventional MRS measures such as Glx and GABA+ include contributions from glutamine and macromolecules, respectively, limiting their chemical specificity. Therefore, while these metabolites serve as useful proxies related to excitatory and inhibitory neural systems, caution is warranted when interpreting them as direct indices of neurotransmission. Nonetheless, these metabolic pools provide valuable insight into the overall balance of excitation and inhibition within the brain’s microenvironment and remain widely used in human *in vivo* studies of neural function and dysfunction. Finally, few studies have directly quantified both glutamate and GABA in the same individuals, highlighting the novelty and importance of this approach. Our study acknowledges these limitations and aims to provide a step toward linking neurochemical measures with large-scale brain dynamics related to criticality.

Alternatively, the present findings may in fact indicate that H and E:I are not directly coupled, or that their relationship is more complex than a simple linear association. This would perhaps not be surprising given the large disparity of findings in the literature, especially with regard to the directionality and linearity of the proposed E:I-Hurst relationship^{6–12} (see Table 3). The heterogeneity across these studies underscores the challenges of studying this phenomenon and suggests that any potential E:I-Hurst relationship may depend on experimental context, data acquisition methods, or analysis

strategies. Further work is needed to clarify the true nature of this relationship and the conditions under which it might emerge.

Citation	Study Type	H Data Type	H Calculation Method	E:I Type	E:I-Hurst Relationship
Poil et al. (2012) ⁷	Computational with in-house simulated model	Neuronal avalanche size	Detrended fluctuation analysis (DFA)	Structural: number of E-to-I neurons	Inverse U
Bruining et al. (2020) ¹⁰	Computational with model by Poil et al. (2012); modified in-house	Neuronal oscillation amplitude	DFA	Structural: number of E-to-I synapses	Inverse U
Gao et al. (2017) ¹²	Computational; in vivo in rats and macaques	Local field potential (LFP)	PSD	Estimated from LFP	Positive linear
Lombardi et al. (2017) ⁸	Computational with in-house model	Neuronal avalanche size	PSD	Structural: number of E-to-I neurons	Negative linear
Trakoshis et al. (2020) ¹¹	Computational with simulated data; in vivo in mice	fMRI BOLD signal	Wavelet-based maximum likelihood method	E-to-I synaptic conductance	Positive linear

Table 3. Summary of methods for existing E:I-Hurst studies

Our findings specifically show an absence of a robust association between the Hurst exponent and MRS-derived E:I ratio, challenging the assumption that these measures directly reflect a shared underlying mechanism, such as the brain’s distance from criticality. One possibility is that while both H and E:I may independently vary with neural excitability, their relationship is nonlinear or not detectable at the temporal and spatial scales of fMRI and MRS. Additionally, the E:I balance estimated via glutamate and GABA concentrations may reflect metabolic rather than synaptic activity, limiting its correspondence with network dynamics captured by BOLD signal autocorrelation. These results have important implications for studies that use H as a

proxy for E:I, particularly in neurodevelopmental or neurodegenerative contexts, and suggest caution in interpreting H as a stand-in for neurochemical balance without direct validation. Future research should investigate whether the relationship emerges under task-driven conditions, in specific brain regions, or in populations with pronounced E:I dysregulation.

Finally, it is also possible that while an E:I-Hurst relationship exists, it is not observed within the visual cortex. This theory seems plausible given that MRS studies of disrupted E:I, mostly conducted within the context of autistic adults, have found changes in E:I within other brain regions such as the anterior cingulate cortex, frontal lobe, or temporal lobe⁵³. Moreover, findings with reference to changes in excitatory or inhibitory neurotransmitters within the visual cortex tend to be difficult to capture, perhaps indicating that E:I shows less changes in this region⁹⁶. However, this would suggest that the E:I-H relationship would be region dependent, and therefore not a global theory as it is often portrayed.

As both H and E:I have been known to change with age, we did not seek to limit our sample to any specific age range. Furthermore, other physiological details — such as caffeine, nicotine, phase of the menstrual cycle, gender, and medication use — were not collected, as the core E:I-Hurst theory posits a global relationship that should hold irrespective of specific brain states or transient physiological factors. Additionally, our within-subjects design strengthens this approach: our core analyses assessing changes between resting state and movie-watching were paired, with each participant serving as their own control. This reduces the influence of inter-individual variability, including age-related and other brain-state effects.

4.1 Limitations and Strengths

Beyond the limitations already mentioned (field strength, visual cortex, passive task, block design), another limitation was our small sample size. As this was a pilot study, 26 participants were initially scanned. Once individuals were excluded for poor MRI quality, only 18 participants’ data were analyzed. With this small sample size, it is difficult to make conclusions about a concept as complex as the E:I-Hurst relationship.

It is also important to note that MRS measures reflect total glutamate and GABA metabolite pools, predominantly intracellular and involved in metabolic processes, rather than direct synaptic neurotransmitter release. Furthermore, commonly used measures such as Glx and GABA+ include signals from glutamine and macromolecules, respectively, limiting their chemical specificity. Therefore, while these metabolites serve as useful proxies related to excitatory and inhibitory neural systems, caution is warranted when interpreting them as direct indices of neurotransmission. Nonetheless, these metabolic pools provide valuable insight into the overall balance of excitation and inhibition within the brain’s microenvironment and remain widely used in human *in vivo* studies of neural function and dysfunction. Although voxel placement was guided by anatomical landmarks and performed by trained personnel, some variability in voxel location across subjects and sessions may have introduced noise in metabolite quantification. Future studies may benefit from automated alignment and formal voxel

overlap analysis to reduce this source of variability. Furthermore, our MRS voxels were placed once for the first MRS sequence, and copied for the subsequent MRS runs. Future studies may also wish to acquire high resolution anatomical scans before each MRS run in order to re-align the MRS voxel. We did not explicitly monitor scanner frequency drift following the fMRI runs. Heating-related drift can affect the stability of MRS sequences, potentially introducing variability in glutamate and GABA quantification¹³³. While all MRS data were visually inspected for quality, we cannot rule out subtle effects of drift, which may have contributed to noise in the E:I measurements. We included frequency drift in our linear mixed-effects models in an attempt to control for these confounds. Finally, the order of scanning conditions (rest followed by movie-watching) was not pseudorandomized. This may introduce potential order-related confounds, such as physiological adaptation or scanner drift, which could influence the observed differences between conditions.

An additional avenue for future research involves dynamic fMRS, which seeks to capture time-varying changes in metabolite levels during task performance. While promising, such approaches currently face substantial limitations in signal-to-noise ratio and temporal resolution, particularly for edited GABA measures. Our study was not optimized for dynamic fitting, but future work could explore how fluctuations in neurometabolites relate to moment-to-moment variations in BOLD signal complexity.

We hope that by future researchers can use our reported effect sizes to calculate potential sample sizes. We would also like to list some of the strengths of our study, which include: using sLASER (as opposed to PRESS), which has been shown to have enhanced detection of complex multiplets such as Glx/Glu⁷⁷; using the J-editing sequence MEGA-PRESS for improved GABA detection⁵⁶; using a large MRS voxel size (~22 ml) as per consensus recommendations^{134–136}; measuring H within the same region as our single-voxel MRS; and using multi-echo fMRI for improved motion artifact regression¹³⁷.

4.2 Lessons for Future Researchers

Finally, we hope that we can provide some guidance to future researchers. The following is a non-exhaustive list of suggestions for future work:

- future studies should consider using ultra-high-field 7T MRI, which provides improved spectral resolution and more reliable detection of Glu and GABA compared to conventional 3T MRI;
- examining paradigms that have consistently demonstrated alterations in these metabolites, such as pain studies, may enhance the likelihood of detecting changes in Glu and GABA^{125,126,132}.
- exploring brain regions beyond the visual cortex, such as the anterior cingulate cortex, which is commonly implicated in pain processing^{132,138–140};
- including a more diverse participant sample — beyond healthy controls — may help capture a wider range of H and E:I values, potentially improving sensitivity to metabolite-related changes;
- increasing sample size in order to detect small effect sizes;

- using an event-related design, as opposed to the block design we used, may provide more sensitivity to detecting rapid and transient neural responses due to its ability to isolate specific events within the scan session¹⁴¹. However, see Pasanta et al. (2023)⁹⁶ for a longer discussion and possible downsides to this approach;
- and finally, using a combined fMRI-MRS sequence^{76,142} to measure BOLD and Glu/GABA near-simultaneously.

Together, these considerations may help in overcoming the limitations observed in the present study and contribute to a clearer understanding of the potential relationship between E:I and H.

5 Conclusion

In conclusion, the results do not support a relationship between H and the E:I ratio in the visual cortex either during rest or during movie-watching at 3T in humans. Although a task-related increase in H was observed, no accompanying changes were detected in Glu, GABA, or E:I between movie and rest. Comparing our findings to the broader literature, E:I balance may be too subtle to be detected with conventional 3T MRS methods. With regards to the broader E:I-Hurst relationship, we similarly suggest that either this relationship is insufficiently captured with our methods, or that the relationship between these two variables may be more complex than originally envisaged — perhaps they are not directly related, but rather connected through other mediating variables in a non-linear fashion. To our knowledge, this is the first *in vivo* human study to test for this relationship. It is our hope that as the literature grows, more authors will examine this relationship with respect to other brain regions and using other methods, and will use the lessons learned in this study to inform their own. Hopefully then it will be possible to corroborate findings to probe the complex relationships that may exist with regards to H and E:I in the human brain.

Data availability statement

All code used in this paper is available at github.com/WeberLab/EI_Hurst_Analysis. The raw MRI data used in this paper is available by contacting the author. The manuscript was written in a ‘reproducible manner’: the entire manuscript, including statistics reported, figures, and tables, can be reproduced here: weberlab.github.io/EI_Hurst_Manuscript/

Competing interests

The authors declare they have no known competing interests.

Author contributions

LS performed data curation, formal analysis, investigation, software, visualization, and wrote the original draft. JA helped with validation, visualization, and reviewed and

edited the manuscript. AMW was responsible for conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, validation, visualization, and writing - review & editing.

Acknowledgements

The work presented in this paper was supported in part from funding from the British Columbia Children’s Hospital Foundation, and an NSERC Discovery Grant.

References

1. Deco, G., Jirsa, V. K. & McIntosh, A. R. [Resting brains never rest: Computational insights into potential cognitive architectures](#). *Trends in Neurosciences* **36**, 268–274 (2013).
2. Beggs, J. & Timme, N. Being Critical of Criticality in the Brain. *Frontiers in Physiology* **3**, (2012).
3. Baranger, M. Chaos, Complexity, and Entropy: A physics talk for non-physicists. *New England Complex Systems Institute* (2000).
4. Bassett, D. S. & Gazzaniga, M. S. [Understanding complexity in the human brain](#). *Trends in cognitive sciences* **15**, 200–209 (2011).
5. Zimmern, V. Why Brain Criticality Is Clinically Relevant: A Scoping Review. *Frontiers in Neural Circuits* **14**, (2020).
6. Liang, J., Yang, Z. & Zhou, C. Excitation–Inhibition Balance, Neural Criticality, and Activities in Neuronal Circuits. *The Neuroscientist* 10738584231221766 (2024) doi:[10.1177/10738584231221766](#).
7. Poil, S.-S., Hardstone, R., Mansvelder, H. D. & Linkenkaer-Hansen, K. [Critical-State Dynamics of Avalanches and Oscillations Jointly Emerge from Balanced Excitation/Inhibition in Neuronal Networks](#). *Journal of Neuroscience* **32**, 9817–9823 (2012).
8. Lombardi, F., Herrmann, H. J. & de Arcangelis, L. [Balance of excitation and inhibition determines 1/f power spectrum in neuronal networks](#). *Chaos: An Interdisciplinary Journal of Nonlinear Science* **27**, 047402 (2017).
9. Baumgarten, L. & Bornholdt, S. [Critical excitation-inhibition balance in dense neural networks](#). *Physical Review E* **100**, 010301 (2019).
10. Bruining, H. *et al.* [Measurement of excitation-inhibition ratio in autism spectrum disorder using critical brain dynamics](#). *Scientific Reports* **10**, 9195 (2020).
11. Trakoshis, S. *et al.* [Intrinsic excitation-inhibition imbalance affects medial prefrontal cortex differently in autistic men versus women](#). *eLife* **9**, e55684.
12. Gao, R., Peterson, E. J. & Voytek, B. [Inferring synaptic excitation/inhibition balance from field potentials](#). *NeuroImage* **158**, 70–78 (2017).
13. Tian, Y. *et al.* [Theoretical foundations of studying criticality in the brain](#). *Network Neuroscience* **6**, 1148–1185 (2022).

14. Rubinov, M., Sporns, O., Thivierge, J.-P. & Breakspear, M. [Neurobiologically Realistic Determinants of Self-Organized Criticality in Networks of Spiking Neurons](#). *PLoS Computational Biology* **7**, e1002038 (2011).
15. Langton, C. G. [Computation at the edge of chaos: Phase transitions and emergent computation](#). *Physica D: Nonlinear Phenomena* **42**, 12–37 (1990).
16. Bak, P. *How Nature Works*. (Springer, New York, NY, 1996). doi:[10.1007/978-1-4757-5426-1](#).
17. Rubenstein, J. L. R. & Merzenich, M. M. Model of autism: Increased ratio of excitation/inhibition in key neural systems. *Genes, brain, and behavior* **2**, 255–267 (2003).
18. Lauterborn, J. C. *et al.* [Increased excitatory to inhibitory synaptic ratio in parietal cortex samples from individuals with Alzheimer’s disease](#). *Nature Communications* **12**, 2603 (2021).
19. Sohal, V. S. & Rubenstein, J. L. R. [Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders](#). *Molecular Psychiatry* **24**, 1248–1257 (2019).
20. Muñoz, M. A. [Colloquium : Criticality and dynamical scaling in living systems](#). *Reviews of Modern Physics* **90**, 031001 (2018).
21. Caserta, F. *et al.* [Determination of fractal dimension of physiologically characterized neurons in two and three dimensions](#). *Journal of Neuroscience Methods* **56**, 133–144 (1995).
22. Lowen, S. B., Cash, S. S., Poo, M. & Teich, M. C. [Quantal Neurotransmitter Secretion Rate Exhibits Fractal Behavior](#). *The Journal of Neuroscience* **17**, 5666–5677 (1997).
23. Mazzoni, A. *et al.* [On the Dynamics of the Spontaneous Activity in Neuronal Networks](#). *PLoS ONE* **2**, e439 (2007).
24. Bédard, C. & Destexhe, A. [Macroscopic Models of Local Field Potentials and the Apparent 1/f Noise in Brain Activity](#). *Biophysical Journal* **96**, 2589–2603 (2009).
25. Bullmore, E. T. *et al.* [Fractal analysis of electroencephalographic signals intracerebrally recorded during 35 epileptic seizures: Evaluation of a new method for synoptic visualisation of ictal events](#). *Electroencephalography and Clinical Neurophysiology* **91**, 337–345 (1994).
26. Zarahn, E., Aguirre, G. K. & D’Esposito, M. [Empirical Analyses of BOLD fMRI Statistics](#). *NeuroImage* **5**, 179–197 (1997).
27. Fadili, M. J. & Bullmore, E. T. [Wavelet-Generalized Least Squares: A New BLU Estimator of Linear Regression Models with 1/f Errors](#). *NeuroImage* **15**, 217–232 (2002).
28. Campbell, O. L. & Weber, A. M. [Monofractal analysis of functional magnetic resonance imaging: An introductory review](#). *Human Brain Mapping* **43**, 2693–2706 (2022).
29. Eke, A., Herman, P., Kocsis, L. & Kozak, L. R. [Fractal characterization of complexity in temporal physiological signals](#). *Physiological Measurement* **23**, R1–38 (2002).

30. Hurst, H. E. [Long-Term Storage Capacity of Reservoirs](#). *Transactions of the American Society of Civil Engineers* **116**, 770–799 (1951).
31. Dong, J. *et al.* [Hurst Exponent Analysis of Resting-State fMRI Signal Complexity across the Adult Lifespan](#). *Frontiers in Neuroscience* **12**, (2018).
32. Wink, A.-M., Bullmore, E., Barnes, A., Bernard, F. & Suckling, J. [Monofractal and multifractal dynamics of low frequency endogenous brain oscillations in functional MRI](#). *Human Brain Mapping* **29**, 791–801 (2008).
33. Ciuciu, P., Abry, P. & He, B. J. [Interplay between functional connectivity and scale-free dynamics in intrinsic fMRI networks](#). *NeuroImage* **95**, 248–263 (2014).
34. He, B. J. [Scale-Free Properties of the Functional Magnetic Resonance Imaging Signal during Rest and Task](#). *Journal of Neuroscience* **31**, 13786–13795 (2011).
35. Campbell, O., Vanderwal, T. & Weber, A. M. [Fractal-Based Analysis of fMRI BOLD Signal During Naturalistic Viewing Conditions](#). *Frontiers in Physiology* **12**, (2022).
36. Churchill, N. W. *et al.* [The suppression of scale-free fMRI brain dynamics across three different sources of effort: Aging, task novelty and task difficulty](#). *Scientific Reports* **6**, 30895 (2016).
37. Wink, A. M., Bernard, F., Salvador, R., Bullmore, E. & Suckling, J. [Age and cholinergic effects on hemodynamics and functional coherence of human hippocampus](#). *Neurobiology of Aging* **27**, 1395–1404 (2006).
38. Mella, A. E., Vanderwal, T., Miller, S. P. & Weber, A. M. [Temporal complexity of the BOLD-signal in preterm versus term infants](#). *Cerebral Cortex* **34**, bhae426 (2024).
39. Maxim, V. *et al.* [Fractional Gaussian noise, functional MRI and Alzheimer’s disease](#). *NeuroImage* **25**, 141–158 (2005).
40. Warsi, M. A., Molloy, W. & Noseworthy, M. D. [Correlating brain blood oxygenation level dependent \(BOLD\) fractal dimension mapping with magnetic resonance spectroscopy \(MRS\) in Alzheimer’s disease](#). *Magnetic Resonance Materials in Physics, Biology and Medicine* **25**, 335–344 (2012).
41. Dona, O., Hall, G. B. & Noseworthy, M. D. [Temporal fractal analysis of the rs-BOLD signal identifies brain abnormalities in autism spectrum disorder](#). *PLOS ONE* **12**, e0190081 (2017).
42. Lai, M.-C. *et al.* [A Shift to Randomness of Brain Oscillations in People with Autism](#). *Biological Psychiatry* **68**, 1092–1099 (2010).
43. Linke, A. C. *et al.* [Altered Development of the Hurst Exponent in the Medial Prefrontal Cortex in Preschoolers With Autism](#). *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* S2451902224002714 (2024) doi:10.1016/j.bpsc.2024.09.003.
44. Uscătescu, L. C. *et al.* [Using the Excitation/Inhibition Ratio to Optimize the Classification of Autism and Schizophrenia](#). 2022.05.24.22275531 (2023) doi:10.1101/2022.05.24.22275531.
45. Dona, O., Noseworthy, M. D., DeMatteo, C. & Connolly, J. F. [Fractal Analysis of Brain Blood Oxygenation Level Dependent \(BOLD\) Signals from Children with Mild Traumatic Brain Injury \(mTBI\)](#). *PLOS ONE* **12**, e0169647 (2017).

46. Wei, M. *et al.* Identifying major depressive disorder using Hurst exponent of resting-state brain networks. *Psychiatry Research: Neuroimaging* **214**, 306–312 (2013).
47. Jing, B. *et al.* Identifying current and remitted major depressive disorder with the Hurst exponent: A comparative study on two automated anatomical labeling atlases. *Oncotarget* **8**, 90452–90464 (2017).
48. Sokunbi, M. O. *et al.* Nonlinear Complexity Analysis of Brain fMRI Signals in Schizophrenia. *PLoS ONE* **9**, e95146 (2014).
49. Vico Varela, E., Etter, G. & Williams, S. Excitatory-inhibitory imbalance in Alzheimer’s disease and therapeutic significance. *Neurobiology of Disease* **127**, 605–615 (2019).
50. Uzunova, G., Pallanti, S. & Hollander, E. Excitatory/inhibitory imbalance in autism spectrum disorders: Implications for interventions and therapeutics. *The World Journal of Biological Psychiatry* **17**, 174–186 (2016).
51. Page, C. E. & Coutellier, L. Prefrontal excitatory/inhibitory balance in stress and emotional disorders: Evidence for over-inhibition. *Neuroscience & Biobehavioral Reviews* **105**, 39–51 (2019).
52. Kang, E. *et al.* Interplay between a Mental Disorder Risk Gene and Developmental Polarity Switch of GABA Action Leads to Excitation-Inhibition Imbalance. *Cell Reports* **28**, 1419–1428.e3 (2019).
53. Ajram, L. A. *et al.* The contribution of [1H] magnetic resonance spectroscopy to the study of excitation-inhibition in autism. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **89**, 236–244 (2019).
54. Stanley, J. A. & Raz, N. Functional Magnetic Resonance Spectroscopy: The ‘New’ MRS for Cognitive Neuroscience and Psychiatry Research. *Frontiers in Psychiatry* **9**, 76 (2018).
55. Harris, A. D., Saleh, M. G. & Edden, R. A. E. Edited 1 H magnetic resonance spectroscopy in vivo: Methods and metabolites. *Magnetic Resonance in Medicine* **77**, 1377–1389 (2017).
56. Peek, A. L. *et al.* A comprehensive guide to MEGA-PRESS for GABA measurement. *Analytical Biochemistry* **669**, 115113 (2023).
57. Poil, S.-S. *et al.* Fast network oscillations in vitro exhibit a slow decay of temporal auto-correlations: Stability of fast network oscillations. *European Journal of Neuroscience* **34**, 394–403 (2011).
58. Monto, S., Vanhatalo, S., Holmes, M. D. & Palva, J. M. Epileptogenic Neocortical Networks Are Revealed by Abnormal Temporal Dynamics in Seizure-Free Subdural EEG. *Cerebral Cortex* **17**, 1386–1393 (2007).
59. Beggs, J. M. & Plenz, D. Neuronal Avalanches in Neocortical Circuits. *The Journal of Neuroscience* **23**, 11167–11177 (2003).
60. Gireesh, E. D. & Plenz, D. Neuronal avalanches organize as nested theta- and beta/gamma-oscillations during development of cortical layer 2/3. *Proceedings of the National Academy of Sciences* **105**, 7576–7581 (2008).

61. Yang, H., Shew, W. L., Roy, R. & Plenz, D. [Maximal Variability of Phase Synchrony in Cortical Networks with Neuronal Avalanches](#). *The Journal of Neuroscience* **32**, 1061–1072 (2012).
62. O’Byrne, J. & Jerbi, K. [How critical is brain criticality?](#) *Trends in Neurosciences* **45**, 820–837 (2022).
63. Plenz, D. *et al.* [Self-Organized Criticality in the Brain](#). *Frontiers in Physics* **9**, 639389 (2021).
64. Avrămiea, A.-E. Regulation of critical brain dynamics and its functional implications. HFAAA72020230307 (Vrije Universiteit Amsterdam, 2023). doi:[10.5463/thesis.144](#).
65. Zeraati, R., Priesemann, V. & Levina, A. [Self-Organization Toward Criticality by Synaptic Plasticity](#). *Frontiers in Physics* **9**, 619661 (2021).
66. Xie, K. *et al.* Pharmacoresistant temporal lobe epilepsy gradually perturbs the cortex-wide excitation-inhibition balance. (2024) doi:[10.1101/2024.04.22.590555](#).
67. Fotiadis, P. *et al.* [Myelination and excitation-inhibition balance synergistically shape structure-function coupling across the human cortex](#). *Nature Communications* **14**, 6115 (2023).
68. Vanderwal, T., Eilbott, J. & Castellanos, F. X. [Movies in the magnet: Naturalistic paradigms in developmental functional neuroimaging](#). *Developmental Cognitive Neuroscience* **36**, 100600 (2019).
69. Lin, Y., Stephenson, M. C., Xin, L., Napolitano, A. & Morris, P. G. [Investigating the Metabolic Changes due to Visual Stimulation using Functional Proton Magnetic Resonance Spectroscopy at 7 T](#). *Journal of Cerebral Blood Flow & Metabolism* **32**, 1484–1495 (2012).
70. Martínez-Maestro, M., Labadie, C. & Möller, H. E. [Dynamic metabolic changes in human visual cortex in regions with positive and negative blood oxygenation level-dependent response](#). *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism* **39**, 2295–2307 (2019).
71. Saucedo, A., Guo, F., Pappas, I. & Wang, D. Changes in GABA and Glutamate levels with peripheral and central visual stimulation using functional magnetic resonance spectroscopy at 7T. in *2024 ISMRM & ISMRT Annual Meeting 1922* (Toronto, ON, Canada). doi:[10.58530/2024/1922](#).
72. Kurcyus, K. *et al.* [Opposite Dynamics of GABA and Glutamate Levels in the Occipital Cortex during Visual Processing](#). *The Journal of Neuroscience* **38**, 9967–9976 (2018).
73. Oz, G. & Tkáč, I. [Short-echo, single-shot, full-intensity proton magnetic resonance spectroscopy for neurochemical profiling at 4 T: Validation in the cerebellum and brainstem](#). *Magnetic Resonance in Medicine* **65**, 901–910 (2011).
74. Mescher, M., Merkle, H., Kirsch, J., Garwood, M. & Gruetter, R. [Simultaneous in vivo spectral editing and water suppression](#). *NMR in biomedicine* **11**, 266–272 (1998).
75. Cordey, H. *Jungles. Our Planet* (2019).

76. Ip, I. B. *et al.* Combined fMRI-MRS acquires simultaneous glutamate and BOLD-fMRI signals in the human brain. *NeuroImage* **155**, 113–119 (2017).
77. Wilson, M. *et al.* Methodological consensus on clinical proton MRS of the brain: Review and recommendations. *Magnetic Resonance in Medicine* **82**, 527–550 (2019).
78. Van Veenendaal, T. M. *et al.* Glutamate quantification by PRESS or MEGA-PRESS: Validation, repeatability, and concordance. *Magnetic Resonance Imaging* **48**, 107–114 (2018).
79. Li, X., Morgan, P. S., Ashburner, J., Smith, J. & Rorden, C. The first step for neuroimaging data analysis: DICOM to NIfTI conversion. *Journal of Neuroscience Methods* **264**, 47–56 (2016).
80. Gorgolewski, K. J. *et al.* The brain imaging data structure, a format for organizing and describing outputs of neuroimaging experiments. *Scientific Data* **3**, 160044 (2016).
81. Boré, A., Guay, S., Bedetti, C., Meisler, S. & GuenTher, N. Dcm2Bids. (2023) doi:10.5281/zenodo.8436509.
82. Tustison, N. J. *et al.* N4ITK: Improved N3 Bias Correction. *IEEE Transactions on Medical Imaging* **29**, 1310–1320 (2010).
83. Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis* **12**, 26–41 (2008).
84. Gorgolewski, K. J. *et al.* Nipype: A flexible, lightweight and extensible neuroimaging data processing framework in Python. 0.12.0-Rc1. (2016) doi:10.5281/zenodo.50186.
85. Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical Imaging* **20**, 45–57 (2001).
86. Smith, S. M. *et al.* Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage* **23 Suppl 1**, S208–219 (2004).
87. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage* **9**, 179–194 (1999).
88. Klein, A. *et al.* Mindboggling morphometry of human brains. *PLoS computational biology* **13**, e1005350 (2017).
89. Esteban, O. *et al.* fMRIPrep: A robust preprocessing pipeline for functional MRI. *Nature Methods* **16**, 111–116 (2019).
90. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage* **17**, 825–841 (2002).
91. Cox, R. W. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages. *Computers and Biomedical Research* **29**, 162–173 (1996).
92. Posse, S. *et al.* Enhancement of BOLD-contrast sensitivity by single-shot multi-echo functional MR imaging. *Magnetic Resonance in Medicine* **42**, 87–97 (1999).

93. DuPre, E. *et al.* [TE-dependent analysis of multi-echo fMRI with *tedana*](#). *Journal of Open Source Software* **6**, 3669 (2021).
94. Oeltzschner, G. *et al.* [Osprey: Open-Source Processing, Reconstruction & Estimation of Magnetic Resonance Spectroscopy Data](#). *Journal of neuroscience methods* **343**, 108827 (2020).
95. Provencher, S. W. [Automatic quantitation of localized in vivo ¹H spectra with LCModel](#). *NMR in biomedicine* **14**, 260–264 (2001).
96. Pasanta, D. *et al.* [Functional MRS studies of GABA and glutamate/Glx – A systematic review and meta-analysis](#). *Neuroscience & Biobehavioral Reviews* **144**, 104940 (2023).
97. Ramadan, S., Lin, A. & Stanwell, P. [Glutamate and Glutamine: A Review of In Vivo MRS in the Human Brain](#). *NMR in biomedicine* **26**, 10.1002/nbm.3045 (2013).
98. Zöllner, H. J. *et al.* [Comparison of different linear-combination modeling algorithms for short-TE proton spectra](#). *NMR in biomedicine* **34**, e4482 (2021).
99. Zöllner, H. J. *et al.* [Comparison of linear combination modeling strategies for edited magnetic resonance spectroscopy at 3 T](#). *NMR in biomedicine* **35**, e4618 (2022).
100. An, L., Willem Van Der Veen, J., Li, S., Thomasson, D. M. & Shen, J. [Combination of multichannel single-voxel MRS signals using generalized least squares](#). *Journal of Magnetic Resonance Imaging* **37**, 1445–1450 (2013).
101. Bouchard, A. E. & Mikkelsen, M. [Noise decorrelation coil combination optimizes SNR of edited ¹H MRS data](#). *Magnetic Resonance Imaging* **122**, 110452 (2025).
102. Harris, A. D., Puts, N. A. J., Barker, P. B. & Edden, R. A. E. [Spectral-Editing Measurements of GABA in the Human Brain with and without Macromolecule Suppression](#). *Magnetic resonance in medicine* **74**, 1523–1529 (2015).
103. Mikkelsen, M. *et al.* [Big GABA II: Water-referenced edited MR spectroscopy at 25 research sites](#). *NeuroImage* **191**, 537–548 (2019).
104. Mikkelsen, M. *et al.* [Big GABA: Edited MR spectroscopy at 24 research sites](#). *NeuroImage* **159**, 32–45 (2017).
105. Friston, K., Ashburner, J., Kiebel, S., Nichols, T. & Penny, W. *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. (Elsevier Ltd., 2007).
106. Gasparovic, C. *et al.* [Use of tissue water as a concentration reference for proton spectroscopic imaging](#). *Magnetic Resonance in Medicine* **55**, 1219–1226 (2006).
107. Rideaux, R. [No balance between glutamate+glutamine and GABA+ in visual or motor cortices of the human brain: A magnetic resonance spectroscopy study](#). *NeuroImage* **237**, 118191 (2021).
108. Juchem, C. *et al.* [B₀ shimming for in vivo magnetic resonance spectroscopy: Experts' consensus recommendations](#). *NMR in Biomedicine* **34**, e4350 (2021).
109. Edden, R. A. E., Puts, N. A. J., Harris, A. D., Barker, P. B. & Evans, C. J. [Gannet: A batch-processing tool for the quantitative analysis of gamma-aminobutyric acid-edited MR spectroscopy spectra](#). *Journal of Magnetic Resonance Imaging* **40**, 1445–1452 (2014).

110. Duff, E. P. *et al.* The power of spectral density analysis for mapping endogenous BOLD signal fluctuations. *Human Brain Mapping* **29**, 778–790 (2008).
111. Bullmore, E. *et al.* Wavelets and functional magnetic resonance imaging of the human brain. *NeuroImage* **23**, S234–S249 (2004).
112. Welch, P. The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms. *IEEE Transactions on Audio and Electroacoustics* **15**, 70–73 (1967).
113. Virtanen, P. *et al.* SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods* **17**, 261–272 (2020).
114. Rubin, D., Fekete, T. & Mujica-Parodi, L. R. Optimizing Complexity Measures for fMRI Data: Algorithm, Artifact, and Sensitivity. *PLoS ONE* **8**, e63448 (2013).
115. Schaefer, A., Brach, J. S., Perera, S. & Sejdić, E. A comparative analysis of spectral exponent estimation techniques for $1/f\beta$ processes with applications to the analysis of stride interval time series. *Journal of neuroscience methods* **222**, 118–130 (2014).
116. Hartmann, A. *et al.* Real-time fractal signal processing in the time domain. *Physica A: Statistical Mechanics and its Applications* **392**, 89–102 (2013).
117. Eke, A. *et al.* Physiological time series: Distinguishing fractal noises from motions. *Pflügers Archiv - European Journal of Physiology* **439**, 403–415 (2000).
118. Team, R. C. R: A language and environment for statistical computing. (2021).
119. Student. The Probable Error of a Mean. *Biometrika* **6**, 1–25 (1908).
120. Freedman, D., Pisani, R. & Purves, R. *Statistics*. vol. 4 (W. W. Norton & Company, New York, 2007).
121. Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology* **106**, 1125–1165 (2011).
122. Van Essen, D. C. *et al.* The Human Connectome Project: A data acquisition perspective. *NeuroImage* **62**, 2222–2231 (2012).
123. Barnes, A., Bullmore, E. T. & Suckling, J. Endogenous Human Brain Dynamics Recover Slowly Following Cognitive Effort. *PLoS ONE* **4**, e6626 (2009).
124. Hasson, U., Malach, R. & Heeger, D. J. Reliability of cortical activity during natural stimulation. *Trends in Cognitive Sciences* **14**, 40–48 (2010).
125. Gutzeit, A. *et al.* Differential NMR spectroscopy reactions of anterior/posterior and right/left insular subdivisions due to acute dental pain. *European Radiology* **23**, 450–460 (2013).
126. Cleve, M., Gussew, A., Wagner, G., Bär, K.-J. & Reichenbach, J. R. Assessment of intra- and inter-regional interrelations between GABA+, Glx and BOLD during pain perception in the human brain - A combined ^1H fMRS and fMRI study. *Neuroscience* **365**, 125–136 (2017).
127. Apšvalka, D., Gadie, A., Clemence, M. & Mullins, P. G. Event-related dynamics of glutamate and BOLD effects measured using functional magnetic resonance spectroscopy (fMRS) at 3T in a repetition suppression paradigm. *NeuroImage* **118**, 292–300 (2015).

128. Floyer-Lea, A., Wylezinska, M., Kincses, T. & Matthews, P. M. [Rapid Modulation of GABA Concentration in Human Sensorimotor Cortex During Motor Learning](#). *Journal of Neurophysiology* **95**, 1639–1644 (2006).
129. Sampaio-Baptista, C. *et al.* [Changes in functional connectivity and GABA levels with long-term motor learning](#). *NeuroImage* **106**, 15–20 (2015).
130. Stagg, C. J., Bachtier, V. & Johansen-Berg, H. [The Role of GABA in Human Motor Learning](#). *Current Biology* **21**, 480–484 (2011).
131. Mangia, S., Giove, F. & Dinuzzo, M. [Metabolic pathways and activity-dependent modulation of glutamate concentration in the human brain](#). *Neurochemical Research* **37**, 2554–2561 (2012).
132. Archibald, J. *et al.* [Metabolite activity in the anterior cingulate cortex during a painful stimulus using functional MRS](#). *Scientific Reports* **10**, 19218 (2020).
133. Hui, S. C. N. *et al.* [Frequency drift in MR spectroscopy at 3T](#). *NeuroImage* **241**, 118430 (2021).
134. Peek, A. L. *et al.* [Brain GABA and glutamate levels across pain conditions: A systematic literature review and meta-analysis of 1H-MRS studies using the MRS-Q quality assessment tool](#). *NeuroImage* **210**, 116532 (2020).
135. Choi, I.-Y. *et al.* [Spectral editing in 1H magnetic resonance spectroscopy: Experts' consensus recommendations](#). *NMR in Biomedicine* **34**, e4411 (2021).
136. Lin, A. *et al.* [Minimum Reporting Standards for in vivo Magnetic Resonance Spectroscopy \(MRSinMRS\): Experts' consensus recommendations](#). *NMR in biomedicine* **34**, e4484 (2021).
137. Kundu, P. *et al.* [Multi-echo fMRI: A review of applications in fMRI denoising and analysis of BOLD signals](#). *NeuroImage* **154**, 59–80 (2017).
138. Mullins, P. G., Rowland, L. M., Jung, R. E. & Sibbitt, W. L. [A novel technique to study the brain's response to pain: Proton magnetic resonance spectroscopy](#). *NeuroImage* **26**, 642–646 (2005).
139. Gussew, A. *et al.* [Time-resolved functional 1H MR spectroscopic detection of glutamate concentration changes in the brain during acute heat pain stimulation](#). *NeuroImage* **49**, 1895–1902 (2010).
140. Gutzeit, A. *et al.* [Insula-specific responses induced by dental pain. A proton magnetic resonance spectroscopy study](#). *European Radiology* **21**, 807–815 (2011).
141. Mullins, P. G. [Towards a theory of functional magnetic resonance spectroscopy \(fMRS\): A meta-analysis and discussion of using MRS to measure changes in neurotransmitters in real time](#). *Scandinavian Journal of Psychology* **59**, 91–103 (2018).
142. Dwyer, G. E. *et al.* [Simultaneous Measurement of the BOLD Effect and Metabolic Changes in Response to Visual Stimulation Using the MEGA-PRESS Sequence at 3 T](#). *Frontiers in Human Neuroscience* **15**, 644079 (2021).

Supplementary Material

MRS quality

Measure	Rest sLASER	Rest MEGA-PRESS	Movie sLASER	Movie MEGA-PRESS
water FWHM (Hz)	8.47 ± 0.66	7.86 ± 0.8	8.32 ± 0.72	7.95 ± 0.82
Cr SNR	147.61 ± 20.46	101.05 ± 21.42	140.6 ± 26	98.75 ± 20.01
Frequency Shift (ppm)	-3.08 ± 0.81	-3.98 ± 0.94	-3.12 ± 0.76	-3.85 ± 1.05

Table 4. Summary of spectroscopy quality measures. FWHM = full-width half-maximum; Cr = creatine; SNR = signal-to-noise ratio.

Measure	Amount
Voxel Rotation (degrees)	-39 ± 8.13
GM Tissue Fraction	0.64 ± 0.04
WM Tissue Fraction	0.24 ± 0.04
CSF Tissue Fraction	0.12 ± 0.03

Table 5. Summary of spectroscopy voxel placement values GM = grey matter; WM = white matter; CSF = cerebrospinal fluid.

Linear-mixed effects model diagnostics




Figure 8. Linear-mixed effects model tests for normality. Model diagnostics for the linear mixed effects model predicting Hurst exponent from the EI ratio. A) Residuals vs. fitted values showing homoscedasticity. B) Histogram of residuals. C) Q-Q plot indicating normally distributed residuals.

Metabolites

sLASER

	Rest Conc	Rest CRLB	Movie Conc	Movie CRLB
Asc	0.34 ± 0.29	239.72 ± 364.52	0.34 ± 0.24	194.39 ± 307.8
Asp	3.49 ± 0.44	9.56 ± 1.34	3.39 ± 0.43	9.72 ± 1.81
Cr	2.04 ± 0.62	16.33 ± 7.65	2.1 ± 0.49	15.22 ± 5.25

	Rest Conc	Rest CRLB	Movie Conc	Movie CRLB
GABA	0.69 ± 0.41	52.78 ± 30.69	0.67 ± 0.32	49.06 ± 28.51
GPC	0.92 ± 0.11	12.11 ± 4.83	0.9 ± 0.11	13.56 ± 4.23
GSH	1.3 ± 0.18	7.61 ± 1.38	1.32 ± 0.13	7.44 ± 1.04
Gln	2.82 ± 0.58	10.11 ± 1.78	2.93 ± 0.47	9.72 ± 1.02
Glu	7.02 ± 0.71	4.33 ± 0.49	7.11 ± 0.62	4.17 ± 0.51
mI	4.77 ± 0.34	3 ± 0	4.8 ± 0.29	3 ± 0
Lac	0.32 ± 0.13	62.78 ± 38.94	0.28 ± 0.16	182.33 ± 311.77
NAA	11.88 ± 1.11	1.94 ± 0.24	11.83 ± 1.03	1.83 ± 0.38
NAAG	0 ± 0	999 ± 0	0.07 ± 0.22	890.78 ± 314.99
PCh	0.16 ± 0.12	292.5 ± 394.55	0.18 ± 0.1	179.89 ± 299.51
PCr	5.68 ± 0.84	5.83 ± 1.04	5.7 ± 0.69	6.11 ± 1.23
PE	1.67 ± 0.49	31.28 ± 11.85	1.71 ± 0.53	29.5 ± 8.58
sI	0.13 ± 0.12	198.61 ± 369.4	0.13 ± 0.11	127.56 ± 244.45
Tau	1.63 ± 0.35	15.44 ± 3.2	1.65 ± 0.3	15.06 ± 3.23
CrCH2	0.26 ± 0.28	433.94 ± 469.27	0.24 ± 0.2	285.11 ± 397.57
tCho	1.07 ± 0.09	3.72 ± 0.46	1.07 ± 0.09	3.56 ± 0.62
tCr	7.72 ± 0.41	1.44 ± 0.51	7.8 ± 0.37	1.39 ± 0.5
tNAA	11.88 ± 1.11	1.94 ± 0.24	11.91 ± 1.04	1.83 ± 0.38
Glx	9.84 ± 1.06	4.39 ± 0.5	10.04 ± 0.79	4.17 ± 0.51
Lip13a	18.63 ± 7.85	76.06 ± 230.43	20 ± 7.47	21.78 ± 6.48
Lip13b	0.07 ± 0.3	944.39 ± 231.7	0 ± 0	999 ± 0
Lip09	4.91 ± 2.05	25 ± 9.44	5.22 ± 2.04	27 ± 10.96
MM09	3.1 ± 2.15	112.67 ± 226.41	3.03 ± 1.5	103.56 ± 227.35
Lip20	3.64 ± 1.34	15.5 ± 4.26	4.05 ± 1	14.44 ± 3.07
MM20	3.88 ± 3.43	242.11 ± 383.63	3.41 ± 2.16	143 ± 262.51
MM12	0.95 ± 0.93	316.28 ± 389.01	0.99 ± 0.97	333.78 ± 425.72
MM14	4.45 ± 2.14	34.39 ± 21.5	5.11 ± 2.63	35.67 ± 25.59
MM17	2.56 ± 1.84	246.39 ± 414.02	2.69 ± 1.1	39.89 ± 31.22

Table 6. All metabolite concentrations and Cramer-Rao lower bounds from semi-LASER. Concentration values are in millimoles; Cramer-Rao lower bounds are in percentages. Conc = concentration; CRLB = Cramer-Rao lower bounds.

MEGA-PRESS

	Rest Conc	Rest CRLB	Movie Conc	Movie CRLB
GABA	1.64 ± 0.36	16.61 ± 5.51	1.55 ± 0.43	18.5 ± 5.86
GSH	2.41 ± 0.6	4.72 ± 0.57	2.17 ± 0.62	5.06 ± 0.94
Gln	2.59 ± 0.95	74.56 ± 230.78	2.4 ± 0.56	24.11 ± 9.54
Glu	12.41 ± 1.76	4.06 ± 0.64	12.11 ± 1.65	4.33 ± 0.77
NAA	16.47 ± 1.49	9.94 ± 2.21	16.29 ± 2.01	10.39 ± 1.5
NAAG	1.8 ± 0.57	16.61 ± 5.51	2.02 ± 1.07	18.5 ± 5.86

	Rest Conc	Rest CRLB	Movie Conc	Movie CRLB
tNAA	18.27 ± 1.41	4.72 ± 0.57	18.31 ± 1.65	5.06 ± 0.94
Glx	15 ± 1.51	74.56 ± 230.78	14.51 ± 1.57	24.11 ± 9.54
MM09	3.84 ± 0.77	4.06 ± 0.64	3.69 ± 0.49	4.33 ± 0.77
GABAplus	5.48 ± 0.77	9.94 ± 2.21	5.23 ± 0.71	10.39 ± 1.5

Table 7. All metabolite concentrations and Cramer-Rao lower bounds from J-edited MEGA-PRESS. Concentration values are in millimoles; Cramer-Rao lower bounds are in percentages. Conc = concentration; CRLB = Cramer-Rao lower bounds.

Data quality





Figure 9. Data quality correlations. A) Correlation plots of H vs. mean FD (mm) for rest (green) and movie (orange). B) and C) Correlation plots of Glx (B) and GABA+ (C) vs. water FWHM (Hz) for rest (green) and movie (orange). D) and E) Correlation plots of Glx (B) and GABA+ (C) vs. frequency shift (ppm) for rest (green) and movie (orange).

A



B




Figure 10. Frequency shift between rest and movie. Overlay of all participants' frequency drift (ppm) plots during rest (blue) and movie (red) for A) semi-LASER and B) MEGA-PRESS. Note, the GE semi-LASER sequence averaged all transients (128) into groups of 4, each with 32 transients, before being saved as raw P-file.

Gannet Results

An example fit result from Gannet of a subject at rest is shown in Figure 11.




Figure 11. Gannet Fit. Sample Gannet fit result from a subject at rest. Glx (3.75ppm) and GABA (3) data (blue) and model fit (red) are shown top left. Water and creatine reference signals are shown bottom left. Table on right shows GABA+ and Glx values before tissue-correction.

To examine the relationship between H and E:I, we ran a linear mixed-effects model. The model's total explanatory power was substantial (conditional $r^2 = 0.64$), with fixed effects (marginal r^2) equal to 0.23. The model's intercept, corresponding to $E:I = 0$, Condition = Movie, $FWHM_{MEGA-PRESS} = 0$, $FreqShift_{MEGA-PRESS} = 0$, and $meanFD = 0$, was 1.2 (95% CI [0.66, 1.7], $t(26) = 4.64$, $p < 0.01$). Within this model, only the effect of Condition [Rest] was statistically significant (beta = -0.08, 95% CI [-0.13, -0.03], $t(26) = -3.32$, $p < 0.01$). The effect of E:I was statistically non-significant (beta = -0.07, 95% CI [-0.21, 0.06], $t(26) = -1.04$, $p = 0.31$); along with $FWHM_{MEGA-PRESS}$ (beta = 0.03, 95% CI [-0.03, 0.08], $t(26) = 0.96$, $p = 0.35$); $FreqShift_{MEGA-PRESS}$ (beta = 0.01, 95% CI [-0.02, 0.05], $t(26) = 0.63$, $p = 0.53$); and $meanFD$ (beta = -0.76, 95% CI [-2.34, 0.82], $t(26) = -0.95$, $p = 0.35$).

Mean $\pm$ sd of metabolites and E:I when analyzed using Gannet, during rest and movie, are reported in Table 8. Boxplots between rest and movie can be seen in Figure 12. Neither Glx nor GABA+ were different between movie and rest conditions. E:I ratio did not change between conditions either.

	Rest	Movie	p-value
Glx	10.84 $\pm$ 1.06	10.72 $\pm$ 0.68	0.43
GABA+	3.59 $\pm$ 0.38	3.53 $\pm$ 0.36	0.33
E:I Ratio	1.82 $\pm$ 0.25	1.94 $\pm$ 0.25	0.08

Table 8. Summary of main metabolite results using Gannet




Figure 12. Paired comparison of metabolite values analyzed using Gannet during rest (green) and movie (orange) conditions. A) Glx (mM); B) GABA+ (mM); and C) E:I. Paired dots represent the same participant across conditions. Mean difference with 95% confidence intervals, and as Cohen's D are reported at the top of each plot.

H was not found to correlate with Glx (Rest: $r = 0$, $p = 0.99$; Movie: $r = -0.27$, $p = 0.28$), GABA+ (Rest: $r = 0.02$, $p = 0.93$; Movie: $r = 0.11$, $p = 0.67$), or E:I (Rest: $r = -0.04$, $p = 0.89$; Movie: $r = -0.32$, $p = 0.2$) (Figure 13).




Figure 13. Scatter plots of H vs. metabolites when analyzed using Gannet. A) Glx (mM); B) GABA+ (mM); and C) E:I. Rest is in green, while movie is in orange. P-values are reported at the top of each plot.