

EGU24-3916

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Seismic Attenuation Imaging in the Western Part of the North Anatolian Fault Zone

Wei-Mou Zhu^{1,2}, Luca De Siena³, Lian-Feng Zhao^{1,4}, David G. Cornwell⁵, Xiao-Bi Xie⁶, Simona Gabrielli⁷, Aqeel Abbas⁸, Xi He¹, Lei Zhang⁹, Panayiota Sketsiou⁵, Stella Lamest¹⁰, and Zhen-Xing Yao¹

¹Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China. (zhuwm@mail.iggcas.ac.cn)

²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, China.

³Department of Physics and Astronomy "Augusto Righi", University of Bologna, Bologna, Italy.

⁴Heilongjiang Mohe Observatory of Geophysics, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China.

⁵Department Geology and Geophysics, University of Aberdeen, Aberdeen, UK.

⁶Institute of Geophysics and Planetary Physics, University of California at Santa Cruz, CA95064, USA.

⁷Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Roma, Roma, Italy.

⁸Earth and Environmental Sciences Programme, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

⁹School of Earth and Space Sciences, Institute of Theoretical and Applied Geophysics, Peking University, Beijing, China.

¹⁰Institute of Geosciences, Johannes Gutenberg University, Mainz, Germany.

Colossal and devastating earthquakes are typically associated with the slip and rupture of fault zones. Fault zone imaging is challenging yet crucial to understand fault structure and behavior, and consequently hazard assessment and mitigation. Seismic attenuation imaging provides constraints on the fault zone structure that are independent of seismic velocity imaging. Here, we image the S-wave total attenuation (Q_s) structure of the western part of the North Anatolian Fault Zone (NAFZ) using data recorded by the DANA (Dense Array for North Anatolia) array. The area of interest is divided into three distinct regions by the northern and southern segments of the NAFZ, which extends from north to south: the Istanbul Zone to the north, the Armutlu Block in between, and the Sakarya Terrane to the south, respectively. The Armutlu Block exhibits much higher attenuation compared to the other two regions. The anomaly body has an attention value of 0.0008 with a notable 3D distribution pattern: It extends from $30.2^\circ E$ to $30.6^\circ E$ and around roughly $40.6^\circ N$ following the northern strand of the NAF and shows a west-east trend, dipping deeper into the crust to the east from depths of 5 to 15 km. Combining previous geological, geodetic, micro-seismicity, and other geophysical observations, we inferred that the high values are a sign of fluid pathways. Micro-seismicity and strain distributions around the Armutlu Block are in line with the assumption fluids migrate through cracks and increased permeability attributed to the background stress, particularly residual stress after the 1999 M7.4 Izmit earthquake and M7.2 Düzce earthquake.

This research is supported by the National Natural Science Foundation of China (U2139206, 41974061, 41974054) and the Special Fund of China Seismic Experimental Site (2019CSES0103).

The first author has also been financially supported by the China Scholarship Council (202204910302).