
Disjunctive Programming/Lift and Project Methods for IP

Endre Boros 26:711:653: Discrete Optimization

Spring 2019



 Consider the polyhedron P defined by the inequalities

$$\mathbf{P} \; = \; \left\{ (x_1, \, x_2) \in \mathbb{R}^2 \; \left| \begin{array}{rrr} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & \leq 1 \\ x_1, & x_2 & \leq 1 \\ x_1, & x_2 & \geq 0 \end{array} \right. \right\}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_1 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$


Then we have

$$\mathbf{P}_{\mathbf{I}} \subseteq \mathbf{P}^{(x_1)} = \operatorname{conv}((\mathbf{Q}_0 \cap \mathbf{P}) \cup (\mathbf{Q}_1 \cap \mathbf{P})) \subseteq \mathbf{P}$$

Conside

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_2 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$

$$\mathbf{P}_{\mathbf{I}} \ = \ \left(\mathbf{P}^{\left(x_{1}\right)}\right)^{\left(x_{2}\right)} \ = \ \mathrm{conv}\left(\left(\mathbf{Q}_{0} \cap \mathbf{P}^{\left(x_{1}\right)}\right) \cup \left(\mathbf{Q}_{1} \cap \mathbf{P}^{\left(x_{1}\right)}\right)\right)$$



 Consider the polyhedron P defined by the inequalities

$$\mathbf{P} \; = \; \left\{ (x_1, \, x_2) \in \mathbb{R}^2 \; \left| \begin{array}{ccc} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & & \leq 1 \\ x_2 & \leq 1 \\ x_1, & x_2 & \geq 0 \end{array} \right. \right\}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_1 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$


Then we have

$$\mathbf{P}_{\mathbf{I}} \subseteq \mathbf{P}^{(x_1)} = \operatorname{conv}((\mathbf{Q}_0 \cap \mathbf{P}) \cup (\mathbf{Q}_1 \cap \mathbf{P})) \subseteq \mathbf{P}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_2 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$

$$\mathbf{P}_{\mathbf{I}} \; = \; \left(\mathbf{P}^{\left(x_{1}\right)}\right)^{\left(x_{2}\right)} \; = \; \mathrm{conv}\left(\left(\mathbf{Q}_{0} \cap \mathbf{P}^{\left(x_{1}\right)}\right) \cup \left(\mathbf{Q}_{1} \cap \mathbf{P}^{\left(x_{1}\right)}\right)\right)$$



Consider the polyhedron P defined by the inequalities

$$\mathbf{P} \; = \; \left\{ (x_1, \, x_2) \in \mathbb{R}^2 \; \middle| \; \begin{array}{ccc} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & \leq 1 \\ x_2 & \leq 1 \\ x_1, & x_2 & \geq 0 \end{array} \right\}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_1 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$


Then we have

$$\mathbf{P_I} \;\subseteq\; \mathbf{P}^{\left(x_1\right)} \;=\; \mathrm{conv}\left((\mathbf{Q}_0 \cap \mathbf{P}) \cup (\mathbf{Q}_1 \cap \mathbf{P})\right) \;\subseteq\; \mathbf{P}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_2 = 0\}$$
 and $\mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$

$$\mathbf{P_{I}} \ = \ \left(\mathbf{P}^{\left(x_{1}\right)}\right)^{\left(x_{2}\right)} \ = \ \mathrm{conv}\left(\left(\mathbf{Q}_{0} \cap \mathbf{P}^{\left(x_{1}\right)}\right) \cup \left(\mathbf{Q}_{1} \cap \mathbf{P}^{\left(x_{1}\right)}\right)\right)$$



Consider the polyhedron P defined by the inequalities

$$\mathbf{P} \, = \, \left\{ (x_1, \, x_2) \in \mathbb{R}^2 \, \left| \begin{array}{rrr} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & \leq 1 \\ x_2 & \leq 1 \\ x_1, & x_2 & \geq 0 \end{array} \right. \right\}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_1 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$


Then we have

$$\mathbf{P_I} \subseteq \mathbf{P}^{(x_1)} = \operatorname{conv}\left((\mathbf{Q}_0 \cap \mathbf{P}) \cup (\mathbf{Q}_1 \cap \mathbf{P})\right) \subseteq \mathbf{P}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_2 = 0\}$$
 and $\mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$

$$\mathbf{P}_{\mathbf{I}} \ = \ \left(\mathbf{P}^{\left(x_{1}\right)}\right)^{\left(x_{2}\right)} \ = \ \mathrm{conv}\left(\left(\mathbf{Q}_{0} \cap \mathbf{P}^{\left(x_{1}\right)}\right) \cup \left(\mathbf{Q}_{1} \cap \mathbf{P}^{\left(x_{1}\right)}\right)\right)$$



Consider the polyhedron P defined by the inequalities

$$\mathbf{P} \; = \; \left\{ (x_1, \, x_2) \in \mathbb{R}^2 \; \left| \begin{array}{ccc} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & & \leq 1 \\ x_1 & & \leq 1 \\ x_1, & x_2 & \geq 0 \end{array} \right. \right\}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_1 = 0\} \quad \text{and} \quad \mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$$

Then we have

$$\mathbf{P_I} \subseteq \mathbf{P}^{(x_1)} = \operatorname{conv}((\mathbf{Q}_0 \cap \mathbf{P}) \cup (\mathbf{Q}_1 \cap \mathbf{P})) \subseteq \mathbf{P}$$

Consider

$$\mathbf{Q}_0 = \{(x_1, x_2) \mid x_2 = 0\}$$
 and $\mathbf{Q}_1 = \{(x_1, x_2) \mid x_2 = 1\}$

$$\mathbf{P_I} \ = \ \left(\mathbf{P}^{\left(x_1\right)}\right)^{\left(x_2\right)} \ = \ \mathrm{conv}\left(\left(\mathbf{Q}_0 \cap \mathbf{P}^{\left(x_1\right)}\right) \cup \left(\mathbf{Q}_1 \cap \mathbf{P}^{\left(x_1\right)}\right)\right)$$

Consider the polyhedron defined by

$$\begin{array}{ccccc} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & & \leq 1 \\ x_1 & & \leq 1 \\ x_2 & & \leq 1 \\ x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

Let us multiply all inequalities by $t_1(x) = x_1$ and $t_2(x) = 1 - x_1$...

Consider the polyhedron defined by

$$\begin{array}{cccc} x_1 & +x_2 & \geq \frac{1}{2} \\ x_1 & +x_2 & \leq \frac{3}{2} \\ x_1 & & \leq 1 \\ x_1 & & \leq 1 \\ x_2 & & \leq 1 \\ x_1 & & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

Let us multiply all inequalities by $t_1(x) = x_1$ and $t_2(x) = 1 - x_1$...

Let us multiply all inequalities by $t_1(x) = x_1$ and $t_2(x) = 1 - x_1$...

Substitute $x_1^2 = x_1, x_1(1 - x_1) = 0 \dots$

Let us multiply all inequalities by $t_1(x) = x_1$ and $t_2(x) = 1 - x_1 \dots$

Substitute $x_1^2 = x_1, x_1(1 - x_1) = 0 \dots$

Substitute $x_1^2 = x_1, x_1(1 - x_1) = 0 \dots$

Substitute $y_{21} = x_2t_1(x) = x_1x_2$, and $y_{22} = x_2t_2(x) = x_2(1-x_1)$, and eliminate redundant inequalities ...

Substitute $x_1^2 = x_1, x_1(1 - x_1) = 0 ...$

Substitute $y_{21} = x_2t_1(x) = x_1x_2$, and $y_{22} = x_2t_2(x) = x_2(1-x_1)$, and eliminate redundant inequalities ...

Substitute $y_{21} = x_2t_1(x) = x_1x_2$, and $y_{22} = x_2t_2(x) = x_2(1-x_1)$, and eliminate redundant inequalities ...

$$y_{21} \quad + y_{22} \quad = x_2$$

$$x_1 + 2x_2 \ge 1$$

$$x_1 + 2x_2 \le 2$$

Substitute $y_{21} = x_2t_1(x) = x_1x_2$, and $y_{22} = x_2t_2(x) = x_2(1-x_1)$, and eliminate redundant inequalities ...

$$y_{21} + y_{22} = x_2$$

$$x_1 + 2x_2 \ge 1$$

١

$$x_1 + 2x_2 \le 2$$

Substitute $y_{21} = x_2t_1(x) = x_1x_2$, and $y_{22} = x_2t_2(x) = x_2(1-x_1)$, and eliminate redundant inequalities ...

$$y_{21} + y_{22} = x_2$$

 $x_1 + 2x_2 > 1$

 $x_1+2x_2 \leq 2$