


Hilbert Bases

Endre Boros
26:711:653: Discrete Optimization

Spring 2019


Hilbert Bases of Pointed Cones

- $\mathbf{C} = \text{cone} \left(\begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right)$
- $H^*(\mathbf{C}) = \left\{ \begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right\}$
- Note the determinant values of the 2×2 matrices formed by the consecutive vectors.



Hilbert Bases of Pointed Cones

- ▶ $\mathbf{C} = \text{cone} \left(\begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right)$
- ▶ $H^*(\mathbf{C}) = \left\{ \begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right\}$
- ▶ Note the determinant values of the 2×2 matrices formed by the consecutive vectors.



Hilbert Bases of Pointed Cones

- ▶ $\mathbf{C} = \text{cone} \left(\begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right)$
- ▶ $H^*(\mathbf{C}) = \left\{ \begin{pmatrix} 3 \\ 7 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 5 \\ 2 \end{pmatrix} \right\}$
- ▶ Note the determinant values of the 2×2 matrices formed by the consecutive vectors.

