
Iona Ecological Conservation



STRM

Utilizing Earth Observations to Understand Landscape Patterns and Assist in Wildlife Management in Iona National Park, Angola

Synopsis

Following the end of the Angolan Civil War in 2002, the human and livestock population in Iona National Park drastically increased. Since 2017, an on-going drought has increased competition between wildlife, humans, and livestock for the park's scarce resources. In collaboration with African Parks, we used Earth observation data to aid in wildlife preservation efforts. We developed a land use/land cover (LULC) classification model using Google Earth Engine, to produce a high-resolution LULC time series analysis between 1990 and 2023. The results provided African Parks with quantitative data on landscape change over the past thirty years. They will use the findings to understand the park's history prior to their involvement and to track changes into the future. Remote sensing and machine learning are useful resources to improve conservation strategies in arid, data-scarce areas where ground-based surveys are challenging.

Objectives


- Create a 2023 LULC map with vegetation details for Iona National Park
- ▶ **Generate** and analyze a time-series of LULC between 1990-2024 that analyzes change over time
- ▶ Produce a short video highlighting the importance of the project, methods, and results

Project Partner


African Parks

Study Area

Iona National Park, Namibe Preserve, & Associated Areas




Methodology



Team Members







Wei Lun Tay



Lukka Wolff




Emmanuel N. Aklie


This material is based upon work supported by NASA through contract 80LARC23FA024. Any mention of a commercial product, service, or activity in this material does not constitute NASA endorsement. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Aeronautics and Space Administration and partner organizations


Earth Observations



Results







Conclusions

- The random forest model was highly successful in detecting fine scale landcover types, doing so with over 90% overall accuracy across the time series.
- Overall diversity decreased across the study period, which indicates a potential threat to local flora and fauna.
- The land cover change analysis was difficult to interpret and would benefit from the inclusion of additional dates to the time series.

Acknowledgements

- Dr. Marguerite Madden, Megan Rich (DEVELOP, Georgia Athens)
- Dr. Sergio Bernardes (University of Georgia, Center for Geospatial Research)
- Daniel van de Vyver, Pedro Monterroso, Priya Tekriwal, Evan Trotzuk (African Parks)



