

Find Laplace Inverse

$$(ii) \quad F(s) = \frac{3s+4}{s^2+9} = \frac{3s}{s^2+9} + \frac{4}{s^2+9}$$

$$L^{-1}\{F(s)\} = 3 \cos 3t + \frac{4}{3} \sin 3t$$

$$(iv) \quad F(s) = \frac{2s+2}{s^2+2s+10} = \frac{2(s+1)}{(s+1)^2+9}$$

$$L^{-1}\{F(s)\} = 2 e^{-t} L^{-1} \left\{ \frac{s}{s^2+9} \right\} = 2 e^{-t} \cos 3t$$

$$(v) \quad F(s) = \frac{2s+3}{s^2+2s+2} = \frac{2s+2+1}{(s+1)^2+1} = \frac{2(s+1)+1}{(s+1)^2+1} = 2 \frac{(s+1)}{(s+1)^2+1} + \frac{1}{(s+1)^2+1}$$

$$L^{-1}\{F(s)\} = 2 e^{-t} L^{-1} \left\{ \frac{s}{s^2+1} \right\} + e^{-t} L^{-1} \left\{ \frac{1}{s^2+1} \right\} = 2 e^{-t} \cos t + e^{-t} \sin t$$

$$(vii) \quad F(s) = \frac{s}{(2s+1)^2} = \frac{1}{4} \frac{s + \frac{1}{2} - \frac{1}{2}}{\left(s + \frac{1}{2}\right)^2} = \frac{1}{4} \left[\frac{1}{s + \frac{1}{2}} - \frac{1}{2} \cdot \frac{1}{\left(s + \frac{1}{2}\right)^2} \right]$$

$$L^{-1}\{F(s)\} = \frac{1}{4} L^{-1} \left\{ \frac{1}{s + \frac{1}{2}} \right\} - \frac{1}{8} e^{-\frac{t}{2}} L^{-1} \left\{ \frac{1}{s^2} \right\} = \frac{1}{4} e^{-\frac{t}{2}} - \frac{1}{8} e^{-\frac{t}{2}} t$$

$$(ix) \quad F(s) = \frac{3s+1}{(s+1)^4} = \frac{3(s-1)-2}{(s+1)^4} = \frac{3}{(s+1)^3} - \frac{2}{(s+1)^4}$$

$$L^{-1}\{F(s)\} = 3e^{-t} L^{-1} \left\{ \frac{1}{s^3} \right\} - 2e^{-t} L^{-1} \left\{ \frac{1}{s^4} \right\} = 3e^{-t} \frac{t^2}{2!} - 2e^{-t} \frac{t^3}{3!} = \frac{3}{2} e^{-t} t^2 - \frac{1}{3} e^{-t} t^3$$

Partial Fractions Cases

Case I: Factors are linear and distinct,

$$F(s) = \frac{P(s)}{(s+a)(s+b)}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+a} + \frac{B}{s+b}$$

Case II: Factors are linear and repeated,

$$F(s) = \frac{P(s)}{(s+a)(s+b)^n}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+a} + \frac{B_1}{s+b} + \frac{B_2}{(s+b)^2} + \dots + \frac{B_n}{(s+b)^n}$$

Case III: Factors are quadratic and distinct,

$$F(s) = \frac{P(s)}{(s^2+as+b)(s^2+cs+d)}$$

By partial fraction expansion,

$$F(s) = \frac{As+B}{s^2+as+b} + \frac{Cs+D}{s^2+cs+d}$$

Case IV: Factors are quadratic and repeated,

$$F(s) = \frac{P(s)}{(s^2+as+b)(s^2+cs+d)^n}$$

By partial fraction expansion,

$$F(s) = \frac{As+B}{s^2+as+b} + \frac{C_1s+D_1}{s^2+cs+d} + \frac{C_2s+D_2}{(s^2+cs+d)^2} + \dots + \frac{C_ns+D_n}{(s^2+cs+d)^n}$$

$$(i) F(s) = \frac{s+2}{s(s+1)(s+3)}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s} + \frac{B}{s+1} + \frac{C}{s+3}$$

$$s+2 = A(s+1)(s+3) + Bs(s+3) + Cs(s+1) \quad \dots (1)$$

Putting $s = 0$ in Eq. (1),

$$2 = 3A$$

$$A = \frac{2}{3}$$

Putting $s = -1$ in Eq. (1),

$$1 = B(-1)(2)$$

$$B = -\frac{1}{2}$$

Putting $s = -3$ in Eq. (1),

$$-1 = C(-3)(-2)$$

$$C = -\frac{1}{6}$$

$$F(s) = \frac{2}{3} \cdot \frac{1}{s} - \frac{1}{2} \cdot \frac{1}{s+1} - \frac{1}{6} \cdot \frac{1}{s+3}$$

$$L^{-1}\{F(s)\} = \frac{2}{3} L^{-1}\left\{\frac{1}{s}\right\} - \frac{1}{2} L^{-1}\left\{\frac{1}{s+1}\right\} - \frac{1}{6} L^{-1}\left\{\frac{1}{s+3}\right\}$$

$$= \frac{2}{3} - \frac{1}{2} e^{-t} - \frac{1}{6} e^{-3t}$$

$$(ii) F(s) = \frac{s+2}{s^2(s+3)}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s+3}$$

$$s+2 = As(s+3) + B(s+3) + Cs^2 \quad \dots (1)$$

Putting $s = 0$ in Eq. (1),

$$2 = 3B$$

$$B = \frac{2}{3}$$

Putting $s = -3$ in Eq. (1),

$$-1 = 9C$$

$$C = -\frac{1}{9}$$

Equating the coefficients of s^2 ,

$$0 = A + C$$

$$A = \frac{1}{9}$$

$$F(s) = \frac{1}{9} \cdot \frac{1}{s} + \frac{2}{3} \cdot \frac{1}{s^2} - \frac{1}{9} \cdot \frac{1}{s+3}$$

$$L^{-1}\{F(s)\} = \frac{1}{9} L^{-1}\left\{\frac{1}{s}\right\} + \frac{2}{3} L^{-1}\left\{\frac{1}{s^2}\right\} - \frac{1}{9} L^{-1}\left\{\frac{1}{s+3}\right\} = \frac{1}{9} + \frac{2}{3}t - \frac{1}{9}e^{-3t}$$

$$(iii) F(s) = \frac{5s^2 - 15s - 11}{(s+1)(s-2)^2}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+1} + \frac{B}{s-2} + \frac{C}{(s-2)^2}$$

$$5s^2 - 15s - 11 = A(s-2)^2 + B(s+1)(s-2) + C(s+1) \quad \dots (1)$$

Putting $s = -1$ in Eq. (1),

$$9 = 9A$$

$$A = 1$$

Putting $s = 2$ in Eq. (1),

$$-21 = 3C$$

$$C = -7$$

Equating the coefficients of s^2 ,

$$5 = A + B$$

$$B = 4$$

$$F(s) = \frac{1}{s+1} + \frac{4}{s-2} - \frac{7}{(s-2)^2}$$

$$L^{-1}\{F(s)\} = L^{-1}\left\{\frac{1}{s+1}\right\} + 4L^{-1}\left\{\frac{1}{s-2}\right\} - 7L^{-1}\left\{\frac{1}{(s-2)^2}\right\}$$

$$= e^{-t} + 4e^{2t} - 7te^{2t}$$

$$(iv) F(s) = \frac{s+2}{(s+3)(s+1)^3}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+3} + \frac{B}{s+1} + \frac{C}{(s+1)^2} + \frac{D}{(s+1)^3}$$

$$s+2 = A(s+1)^3 + B(s+3)(s+1)^2 + C(s+3)(s+1) + D(s+3) \quad \dots (1)$$

Putting $s = -3$ in Eq. (1),

$$-1 = -8A$$

$$A = \frac{1}{8}$$

Putting $s = -1$ in Eq. (1),

$$1 = 2D$$

$$D = \frac{1}{2}$$

Equating the coefficients of s^3 ,

$$0 = A + B$$

$$B = -\frac{1}{8}$$

Equating the coefficients of s^2 ,

$$0 = 3A + 5B + C$$

$$C = -\frac{3}{8} + \frac{5}{8} = \frac{1}{4}$$

$$F(s) = \frac{1}{8} \cdot \frac{1}{s+3} - \frac{1}{8} \cdot \frac{1}{s+1} + \frac{1}{4} \cdot \frac{1}{(s+1)^2} + \frac{1}{2} \cdot \frac{1}{(s+1)^3}$$

$$L^{-1}\{F(s)\} = \frac{1}{8}L^{-1}\left\{\frac{1}{s+3}\right\} - \frac{1}{8}L^{-1}\left\{\frac{1}{s+1}\right\} + \frac{1}{4}L^{-1}\left\{\frac{1}{(s+1)^2}\right\} + \frac{1}{2}L^{-1}\left\{\frac{1}{(s+1)^3}\right\}$$

$$= \frac{1}{8}e^{-3t} - \frac{1}{8}e^{-t} + \frac{1}{4}te^{-t} + \frac{1}{2} \cdot \frac{t^2}{2} \cdot e^{-t}$$

$$= \frac{1}{8}[e^{-3t} + (2t^2 + 2t - 1)e^{-t}]$$

$$(v) F(s) = \frac{s^3 + 6s^2 + 14s}{(s+2)^4}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+2} + \frac{B}{(s+2)^2} + \frac{C}{(s+2)^3} + \frac{D}{(s+2)^4}$$

$$\begin{aligned} s^3 + 6s^2 + 14s &= A(s+2)^3 + B(s+2)^2 + C(s+2) + D \\ &= As^3 + (6A+B)s^2 + (12A+4B+C)s + (8A+4B+2C+D) \quad \dots (1) \end{aligned}$$

Equating the coefficients of s^3 ,

$$A = 1$$

Equating the coefficients of s^2 ,

$$6 = 6A + B$$

$$B = 0$$

Equating the coefficients of s ,

$$14 = 12A + 4B + C$$

$$C = 14 - 12 - 0 = 2$$

Equating the coefficients of s^0 ,

$$0 = 8A + 4B + 2C + D$$

$$D = -8 - 0 - 4 = -12$$

$$F(s) = \frac{1}{s+2} + \frac{2}{(s+2)^3} - \frac{12}{(s+2)^4}$$

$$\begin{aligned} L^{-1}\{F(s)\} &= L^{-1}\left\{\frac{1}{s+2}\right\} + 2L^{-1}\left\{\frac{1}{(s+2)^3}\right\} - 12L^{-1}\left\{\frac{1}{(s+2)^4}\right\} \\ &= e^{-2t} + 2 \cdot \frac{t^2}{2} \cdot e^{-2t} - 12 \cdot \frac{t^3}{6} \cdot e^{-2t} = e^{-2t} (1 + t^2 - 2t^3) \end{aligned}$$

$$(vi) F(s) = \frac{3s+1}{(s+1)(s^2+2)}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s+1} + \frac{Bs+C}{s^2+2}$$

$$3s+1 = A(s^2+2) + (Bs+C)(s+1) \quad \dots (1)$$

Putting $s = -1$ in Eq. (1),

$$-2 = 3A$$

$$A = -\frac{2}{3}$$

Equating the coefficients of s^2 ,

$$0 = A + B$$

$$B = \frac{2}{3}$$

Equating the coefficients of s^0 ,

$$1 = 2A + C$$

$$C = 1 + \frac{4}{3} = \frac{7}{3}$$

$$F(s) = -\frac{2}{3} \cdot \frac{1}{s+1} + \frac{2}{3} \cdot \frac{s}{s^2+2} + \frac{7}{3} \cdot \frac{1}{s^2+2}$$

$$\begin{aligned} L^{-1}\{F(s)\} &= -\frac{2}{3} L^{-1}\left\{\frac{1}{s+1}\right\} + \frac{2}{3} L^{-1}\left\{\frac{s}{s^2+2}\right\} + \frac{7}{3} L^{-1}\left\{\frac{1}{s^2+2}\right\} \\ &= -\frac{2}{3} e^{-t} + \frac{2}{3} \cos \sqrt{2}t + \frac{7}{3\sqrt{2}} \sin \sqrt{2}t \end{aligned}$$

$$(vii) F(s) = \frac{s+4}{s(s-1)(s^2+4)}$$

By partial fraction expansion,

$$F(s) = \frac{A}{s} + \frac{B}{s-1} + \frac{Cs+D}{s^2+4}$$

$$s+4 = A(s-1)(s^2+4) + Bs(s^2+4) + (Cs+D)s(s-1) \quad \dots (1)$$

Putting $s = 0$ in Eq. (1),

$$4 = -4A$$

$$A = -1$$

Putting $s = 1$ in Eq. (1),

$$5 = 5B$$

$$B = 1$$

Equating the coefficients of s^3 ,

$$0 = A + B + C$$

$$C = 1 - 1 = 0$$

Equating the coefficients of s ,

$$1 = 4A + 4B - D$$

$$D = -4 + 4 - 1 = -1$$

$$F(s) = -\frac{1}{s} + \frac{1}{s-1} - \frac{1}{s^2+4}$$

$$L^{-1}\{F(s)\} = -L^{-1}\left\{\frac{1}{s}\right\} + L^{-1}\left\{\frac{1}{s-1}\right\} - L^{-1}\left\{\frac{1}{s^2+4}\right\} = -1 + e^t - \frac{1}{2} \sin 2t$$

$$(viii) F(s) = \frac{s}{(s^2+1)(s^2+4)} = \frac{s}{3} \left[\frac{s^2+4-s^2-1}{(s^2+1)(s^2+4)} \right] = \frac{1}{3} \left[\frac{s}{s^2+1} - \frac{s}{s^2+4} \right]$$

$$L^{-1}\{F(s)\} = \frac{1}{3} \left[L^{-1}\left\{\frac{s}{s^2+1}\right\} - L^{-1}\left\{\frac{s}{s^2+4}\right\} \right] = \frac{1}{3} [\cos t - \cos 2t]$$

$$(ix) F(s) = \frac{s^2}{(s^2 + a^2)(s^2 + b^2)}$$

$$\text{Let } s^2 = x$$

$$G(x) = \frac{x}{(x + a^2)(x + b^2)}$$

By partial fraction expansion,

$$G(x) = \frac{A}{x + a^2} + \frac{B}{x + b^2}$$

$$x = A(x + b^2) + B(x + a^2) \quad \dots (1)$$

Putting $x = -a^2$ in Eq. (1),

$$-a^2 = A(-a^2 + b^2)$$

$$A = \frac{a^2}{a^2 - b^2}$$

Putting $x = -b^2$ in Eq. (1),

$$-b^2 = B(-b^2 + a^2)$$

$$B = -\frac{b^2}{a^2 - b^2}$$

$$G(x) = \frac{a^2}{a^2 - b^2} \cdot \frac{1}{x + a^2} - \frac{b^2}{a^2 - b^2} \cdot \frac{1}{x + b^2}$$

$$F(s) = \frac{a^2}{a^2 - b^2} \cdot \frac{1}{s^2 + a^2} - \frac{b^2}{a^2 - b^2} \cdot \frac{1}{s^2 + b^2}$$

$$L^{-1}\{F(s)\} = \frac{a^2}{a^2 - b^2} L^{-1}\left\{\frac{1}{s^2 + a^2}\right\} - \frac{b^2}{a^2 - b^2} L^{-1}\left\{\frac{1}{s^2 + b^2}\right\}$$

$$= \frac{a^2}{a^2 - b^2} \frac{1}{a} \sin at - \frac{b^2}{a^2 - b^2} \frac{1}{b} \sin bt$$

$$= \frac{1}{a^2 - b^2} (a \sin at - b \sin bt)$$

$$(x) F(s) = \frac{s^2 + 2s + 3}{(s^2 + 2s + 5)(s^2 + 2s + 2)}$$

$$\text{Let } s^2 + 2s = x$$

$$G(x) = \frac{x + 3}{(x + 5)(x + 2)}$$

By partial fraction expansion,

$$G(x) = \frac{A}{x + 5} + \frac{B}{x + 2}$$

$$x + 3 = A(x + 2) + B(x + 5) \quad \dots (1)$$

Putting $x = -5$ in Eq. (1),

$$-2 = -3A$$

$$A = \frac{2}{3}$$

Putting $x = -2$ in Eq. (1),

$$1 = 3B$$

$$B = \frac{1}{3}$$

$$G(x) = \frac{2}{3} \cdot \frac{1}{x+5} + \frac{1}{3} \cdot \frac{1}{x+2}$$

$$F(s) = \frac{2}{3} \cdot \frac{1}{(s^2 + 2s + 5)} + \frac{1}{3} \cdot \frac{1}{(s^2 + 2s + 2)} = \frac{2}{3} \cdot \frac{1}{(s+1)^2 + 4} + \frac{1}{3} \cdot \frac{1}{(s+1)^2 + 1}$$

$$L^{-1}\{F(s)\} = \frac{2}{3} L^{-1}\left\{\frac{1}{(s+1)^2 + 4}\right\} + \frac{1}{3} L^{-1}\left\{\frac{1}{(s+1)^2 + 1}\right\}$$

$$= \frac{2}{3} e^{-t} \cdot \frac{1}{2} \sin 2t + \frac{1}{3} e^{-t} \sin t = \frac{1}{3} e^{-t} (\sin 2t + \sin t)$$

$$\begin{aligned} \text{(xi)} \quad F(s) &= \frac{s+2}{(s^2 + 4s + 8)(s^2 + 4s + 13)} = \frac{s+2}{5} \left[\frac{s^2 + 4s + 13 - s^2 - 4s - 8}{(s^2 + 4s + 8)(s^2 + 4s + 13)} \right] \\ &= \frac{1}{5} \left[\frac{s+2}{s^2 + 4s + 8} - \frac{s+2}{s^2 + 4s + 13} \right] = \frac{1}{5} \left[\frac{s+2}{(s+2)^2 + 4} - \frac{s+2}{(s+2)^2 + 9} \right] \end{aligned}$$

$$L^{-1}\{F(s)\} = \frac{1}{5} \left[L^{-1}\left\{\frac{s+2}{(s+2)^2 + 4}\right\} - L^{-1}\left\{\frac{s+2}{(s+2)^2 + 9}\right\} \right]$$

$$= \frac{1}{5} \left[e^{-2t} L^{-1}\left\{\frac{s}{s^2 + 4}\right\} - e^{-2t} L^{-1}\left\{\frac{s}{s^2 + 9}\right\} \right]$$

$$= \frac{1}{5} (e^{-2t} \cos 2t - e^{-2t} \cos 3t) = \frac{e^{-2t}}{5} (\cos 2t - \cos 3t)$$