

Published in final edited form as:

Virology. 2012 February 5; 423(1): 58–67. doi:10.1016/j.virol.2011.11.018.

Genetic and Phenotypic Characterization of Sylvatic Dengue Virus Type 4 Strains

S.L. Rossi^{1,2,3}, F. Nasar^{1,2,3}, J. Cardoso⁴, S.V. Mayer¹, R.B Tesh^{1,2,3}, K.A. Hanley⁵, S.C. Weaver^{1,2,3}, and Nikos Vasilakis^{1,2,3,*}

¹Center for Biodefense and Emerging Infectious Diseases and Department of Pathology, University of Texas Medical Branch, Galveston, TX 77555-0610

²Institute of Human Infections and Immunity, University of Texas Medical Branch, Galveston, TX 77555-0610

³Center for Tropical Diseases, University of Texas Medical Branch, Galveston, TX 77555-0610

⁴Institute of Health & Community Medicine, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia

⁵Department of Biology, New Mexico State University, Las Cruces, NM 88005

Abstract

Four serotypes of dengue virus (DENV 1-4) currently circulate between humans and domestic/peridomestic *Aedes* mosquitoes, resulting in 100 million infections per year. All four serotypes emerged, independently, from sylvatic progenitors transmitted among non-human primates by arboreal *Aedes* mosquitoes. This study investigated the genetic and phenotypic changes associated with emergence of human DENV-4 from its sylvatic ancestors. Analysis of complete genomes of 3 sylvatic and 4 human strains revealed high conservation of both the 5'- and 3'-untranslated regions but considerable divergence within the open reading frame. Additionally, the two ecotypes did not differ significantly in replication dynamics in cultured human liver (Huh-7), monkey kidney (Vero) or mosquito (C6/36) cells, although significant inter-strain variation within ecotypes was detected. These findings are in partial agreement with previous studies of DENV-2, where human strains produced a larger number of progeny than sylvatic strains in human liver cells but not in monkey or mosquito cells.

Keywords

Dengue virus (DENV); sylvatic DENV; human DENV; phylogenetic and phenotypic analysis

Introduction

There is growing concern about the potential emergence of new pathogens, particularly arthropod-borne viruses (arboviruses), from animal reservoirs into humans (Weaver and

© 2011 Elsevier Inc. All rights reserved.

*Corresponding author: Center for Biodefense and Emerging Infectious Diseases and Department of Pathology, Institute for Human Infection and Immunity, and Center for Tropical Diseases, University of Texas Medical Branch, Galveston, TX 77555-0610. Tel: +1 409 266.6919; Fax: +1 409 266.6810 nivasila@utmb.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Reisen, 2010; Wilder-Smith and Gubler, 2008). To gain insight into this process of emergence, it is particularly instructive to study viruses that have completed the trajectory from an enzootic into a human reservoir, such as the four serotypes of dengue virus (DENV-1-4, genus *Flavivirus*, family *Flaviviridae*). These viruses originated in a sylvatic cycle between nonhuman primates, and possibly other enzootic hosts, and arboreal *Aedes* (*Ae.*) mosquitoes. Each serotype emerged independently into a human transmission cycle, wherein humans now serve as the exclusive reservoir and amplification hosts for the endemic/epidemic lineages (Vasilakis et al., 2011). In this human cycle, DENV-1,-2,-3, and -4 are transmitted by domestic and peridomestic *Aedes* mosquitoes, primarily *Ae. aegypti aegypti* and *Ae. albopictus* (Halstead, 1997; Halstead et al., 1964; Rosen et al., 1954; Sabin, 1952; Simmons et al., 1931). The human DENV cycle is presently found in nearly all urban and peri-urban environments throughout the tropics and subtropics. In recent decades, DENV transmission among humans has intensified due to increased travel, uncontrolled urbanization and lack of effective and sustainable vector control programs (Guzman et al., 2010). By current estimates, DENV infects approximately 100 million people each year in over 100 countries.

Unlike the ancestors of many other human viruses, the ancestral sylvatic cycle of DENV remains extant and has been documented in two foci: one in West Africa involving arboreal *Aedes* spp. (e.g. *Ae. fuscifer*, *Ae. luteocephalus*) and primates including patas monkeys (*Erythrocebus patas*), African green monkeys (*Chlorocebus sabaeus*), and Guinea baboons (*Papio papio*) (Cordellier et al., 1983; Diallo et al., 2003; Diallo et al., 2005; Hervy et al., 1984; Rodhain, 1991; Saluzzo et al., 1986a; Vasilakis et al., 2008c) and the other in peninsular Malaysia involving *Ae. niveus s.l.* and primates including cynomolgus macaques (*Macaca fascicularis*), pig-tailed macaques (*Macaca nemestrina*) and silvered leaf monkeys (*Presbytis cristata*) (Rudnick, 1986). The continued circulation of sylvatic DENV provides an opportunity to perform comparative studies to elucidate the virus attributes that promote arboviral emergence. However these sylvatic viruses also pose a considerable threat, because they may retain the capacity to re-emerge even as efforts to control circulation of human dengue intensify (Vasilakis et al., 2011), in a manner analogous to urban yellow fever.

Although sylvatic and human DENV strains show substantial genetic differences, our previous studies of DENV-2 demonstrated that such differences do not constitute an adaptive barrier to emergence into the human transmission cycle. Specifically sylvatic DENV-2 showed no detectable deficit relative to human DENV-2 in replication kinetics in cultured human or *Ae. albopictus* cells (Vasilakis et al., 2008b), in replication in two proxy measures for human infection, monocyte-derived dendritic cells and SCID mice engrafted with human hepatoma cells (Vasilakis et al., 2007b), or in infection of *Ae. aegypti* and *Ae. albopictus* *in vivo* (Hanley and Vasilakis, unpublished data). Furthermore, some sylvatic *Aedes* species sympatric with sylvatic DENV are capable of transmitting these viruses and may act as bridge vectors when they move between forest and human habitations. For example in West Africa a highly susceptible vector of sylvatic DENV, the forest-dwelling *Ae. fuscifer*, (Diallo et al., 2005), disperses into villages (Diallo et al., 2003), while in Southeast Asia, *Ae. albopictus* disperses from the forest into surrounding agricultural settlements (Smith, 1956). Both species bite humans, resulting in potential transmission of sylvatic DENV to humans. Indeed, several reports have now documented spillover of sylvatic DENV, resulting in infection of individual humans or small outbreaks (Cardosa et al., 2009; Carey et al., 1971; Franco et al., 2011; Monlun et al., 1992; Saluzzo et al., 1986a; Saluzzo et al., 1986b; Vasilakis et al., 2008c). Currently it is not possible to distinguish sylvatic and human DENV infections with antibody-based assays, and thus sylvatic DENV infections may frequently be misclassified as human DENV. Nonetheless two recent reports

from Southeast Asia (Cardosa et al., 2009) and West Africa (Franco et al., 2011) reveal that sylvatic DENV infections can result in severe disease.

Collectively, these data indicate that DENV has evolved as an ecological generalist capable of utilizing a broad range of *Aedes* vectors and primate hosts (including humans) and suggest that the public health impact of sylvatic dengue spillover may be substantially greater than is currently appreciated. However to date the vast majority of both experimental (Cox et al., 2011; Mota and Rico-Hesse, 2009; Mota and Rico-Hesse, 2011; Vasilakis et al., 2009; Vasilakis et al., 2008a; Vasilakis et al., 2008b; Vasilakis et al., 2007b) and phylogenetic (Vasilakis et al., 2007a) studies of sylvatic DENV have focused exclusively on DENV-2, and it is critical to extend these efforts to encompass the full range of genetic and phenotypic diversity within sylvatic DENV. Thus, we investigated the genetic relationships of sylvatic and human DENV-4 using complete genome sequences from each ecotype. Previous studies have generated phylogenies of DENV-4 that include a only a single sylvatic DENV-4 isolate or a single gene from the all 3 known sylvatic DENV-4 isolates (AbuBakar et al., 2002; Wang et al., 2000); in this study we utilized the complete genome sequences of the three sylvatic DENV-4 isolates as well as 59 human DENV-4 sequences from GenBank that represent the complete genotypic diversity known for this serotype. To better link genetic and phenotypic variation, we also measured the replication kinetics of a subset of human versus sylvatic isolates in both mammalian and mosquito cells in culture.

Results and Discussion

Phylogenetic analyses

Complete genome sequences from 59 human DENV-4 isolates that span the genetic, geographic and temporal range of DENV-4 diversity (Chen and Vasilakis, 2011; Holmes and Twiddy, 2003; Vasilakis and Weaver, 2008; Villabona-Arenas and Zanotto, 2011; Weaver and Vasilakis, 2009) as well as the only 3 known sylvatic DENV-4 strains (Rudnick, 1986) were used for phylogenetic analysis. Figure 1 shows a representative phylogenetic tree derived from Bayesian analysis; several consensus trees obtained based on Maximum-likelihood (ML) and Bayesian analyses exhibited similar topologies.

All 3 sylvatic DENV-4 strains were genetically distinct from and basal to human DENV-4 strains. Among the sylvatic strains, P73-1120 and P75-514 were more closely related to each other than to the P75-215 strain, an observation that reflects their history. Strain P73-1120 was isolated from a sentinel silver leaf monkey (*Presbytis cristata*) in the Gunong Besut forest reserve in Malaysia in 1973 (Table 1). Two years later an aliquot of the original serum sample collected from the sentinel monkey was used for experimental infections of silver leaf monkeys, which led to the isolation of P75-514 strain (Rudnick, 1986). Our sequences indicated that, during this single passage, several mutations occurred, which were primarily synonymous and resulted in a 0.1% nucleotide divergence from the parent strain P73-1120 (Table 1). Strain P75-215 was isolated in 1975 from a pool of *Ae. niveus s. l.* mosquitoes collected in the forest canopy of the Gunong Besut forest reserve in late 1974 (Rudnick, 1986).

As shown in previous studies (AbuBakar et al., 2002; Bennett et al., 2003; Foster et al., 2003; Klungthong et al., 2004; Vasilakis and Weaver, 2008; Villabona-Arenas and Zanotto, 2011; Weaver and Vasilakis, 2009), human DENV-4 strains clustered into three major genotypes. Genotype I includes strains from the Philippines, Thailand, Vietnam, Myanmar, Malaysia, Sri Lanka, and India (Cecilia et al., 2011; Dash et al., 2011). The India G11337 strain, which was isolated in 1961 and is therefore one of the oldest DENV-4 strains sampled, is genetically distinct and basal to other isolates of this clade. Genotype II is composed of two distinct sub-lineages: IIa, including strains from Malaysia, Thailand and

Taiwan and I Ib, comprising strains from the Caribbean and the Americas (Bennett et al., 2003; Foster et al., 2003). Genotype III includes Thai strains sampled between 1997 – 2001 that are distinct from the Thai isolates of genotype II (Klungthong et al., 2004).

Genetic analysis

The DENV genome is comprised of approximately 10.7 kilobases (kb) of single stranded RNA of positive polarity. A single open reading frame (ORF) of 10,164 nucleotides (Table 2) is flanked by untranslated regions (UTRs) at the 5' and 3' ends. The 5'-UTR is 101 nt long and is capped with a type I 5'cap (Cleaves and Dubin, 1979; Lindenbach and Rice, 2003), while the 402 nt 3'-UTR lacks the classical polyadenylation site (Wengler et al., 1978) (Table 2).

5'- and 3'-UTRs—The sequences of both the 5' and 3' UTRs are highly conserved among the four DENV serotypes (Markoff, 2003). In the DENV-4 strains analyzed, the sequence of both UTRs was highly conserved, and the 5' UTR had a higher sequence identity between the strains (ranging from 95.2 – 98.0%) than the 3' UTR (88.7–95.1%) (Table 2; sequence data is presented in Supplemental data figures 1 and 2). The 5' cyclization sequence (5'CS) located within the coding region of the capsid gene (nt 35–42 after the start codon) (Hahn et al., 1987) was present in all strains analyzed. The flavivirus 3' UTR comprises three sub-regions: the Variable Region (VR), core, and 3' terminus (Markoff, 2003). Several previously identified conserved regions within the core and 3' terminus (RCS2, CS2 and CS1 and CPS (Hahn et al., 1987)) were present in both the sylvatic and human strains (Supplemental data 3c,d). The highly conserved 3' terminal dinucleotide of the plus-strand (UC-3') was present in all DENV-4 sequences for which the 3' terminal sequence was available, and was complementary to the 5'-terminal dinucleotides (5'AG), purportedly enabling cyclization of the genome during the early stages of replication (Markoff, 2003; Rice et al., 1985; Wengler and Wengler, 1981). Relative to the sylvatic strains, the VR of human genotypes I – III exhibited 3, 13 or 15 nt deletions, respectively (supplemental data 2). A similar accrual of deletions in human strains relative to sylvatic strains has been documented in DENV-2 strains (Vasilakis et al., 2008b), and other investigators (Leitmeyer et al., 1999; Shurtleff et al., 2001) have suggested that these deletions occurred during the evolution of human strains from sylvatic progenitors. Proutski and colleagues suggested that the VR region may act as a spacer to protect conserved and structurally important distal regions (Proutski et al., 1999) but the precise effects of these particular deletions in human strains have not been determined.

The predicted RNA secondary structures of representative sylvatic (P73-1120) and human (IndiaG11337) DENV-4 strains were extremely similar at the 5' and 3' UTRs (Supplemental data figures 3a–d) and were also similar to previously predicted structures for these regions (Markoff, 2003; Rauscher et al., 1997; Romero et al., 2006; Thurner et al., 2004; Yu et al., 2008). As in previous predictions of the 5' UTR in isolation (Brinton and Dispoto, 1988), the RNA secondary structure of each strain contains a long stem loop (SLA) with a shorter side loop (SLB) closest to the start codon; alterations to this sequence may affect RNA-dependent RNA polymerase binding (Filomatori et al., 2011) and viral replication (Cahour et al., 1995; Filomatori et al., 2006). There has been disagreement in the prediction of the DENV-4 CS2, with some studies showing a dumbbell structure and others showing a single turret (Romero et al., 2006)(Gritsun and Gould, 2006; Proutski et al., 1999). The current study found a dumbbell structure at this region for both the sylvatic and human strain. CS2 is particularly important because it is the site of the delta30 mutation that has been used for generation of dengue virus vaccines (Durbin and Whitehead, 2010). Moreover this 30 nt deletion confers attenuation on some but not all DENV strains (Blaney et al., 2004a; Blaney et al., 2004b; Durbin et al. 2011). Thus experimental analysis of the structure of this region

in sylvatic DENV-4 would be particularly valuable once a reverse genetics system is established.

Open Reading Frame (ORF)—To identify sequence differences between sylvatic DENV-4 and each of the genotypes of human DENV-4, we generated both multiple ORF sequence alignments that included DENV-4 strains from all genotypes as well as human and sylvatic transmission cycles (Figure 1 and Material and Methods section). Consistent differences between the ancestral sylvatic and derived human strains were inferred to be mutations in the human strains. These comparisons revealed a large number of nucleotide differences, most of which were nonsynonymous (Table 3).

Structural proteins—The DENV capsid (C) is a small, positively charged protein (Trent, 1977) containing membrane-associated alpha-helical structures (Jones et al., 2003; Markoff et al., 1997) that form the inner core of the virion. There are 6 positions that show differences between the sylvatic and human strains with most nonsynonymous mutations located outside of predicted alpha-helical regions (Table 3).

The prM protein is the immature precursor to the membrane (M) protein found in the virions, and forms a scaffold over the viral envelope (E) protein to prevent premature fusion during virion maturation (Li et al., 2008). Seven non-synonymous differences between the sylvatic and human DENV4 strains were detected, 2 of which are found within the mature M protein (Table 3).

The E protein comprises the majority of the surface area on the mature virion and is responsible for receptor binding (Chen et al., 1996; Johnson et al., 1994; Rey et al., 1995) and host cell surface fusion (Modis et al., 2004). Of the 29 amino acid changes (Table 3), no changes occurred to the 12 conserved disulfide bridge-forming cysteine residues, to the two N-linked glycosylation sites in the E protein at residues N67 and N153 (Johnson et al., 1994; Zhao et al., 1986), or to proposed heparan sulfate binding sites (Chen et al., 1997; Modis et al., 2005). Given strong antibody neutralization of both sylvatic and human strains of the same serotype by antibodies raised against either ecotype (Vasilakis et al., 2008a) it was not surprising that highly immunogenic residues 104, 106, 107 (located in the fusion loop) and 126, 226 and 231 (Crill and Chang, 2004) remain unchanged between sylvatic and human strains. Interestingly, mutations were observed within the DIII region at positions 310 and 364, which are potentially significant since they are located within the binding domains of several well-characterized monoclonal antibodies including 4E11 (Thullier et al., 2001) and 9D12 (Crill et al., 2009).

Nonstructural proteins—The functions of NS1 are not completely known. Like the E protein, NS1 contains 12 disulfide bridge-forming cysteine residues and encodes N-linked glycosylation sites at N130 and N207 (Pryor and Wright, 1994). Of the 26 non-synonymous differences between the sylvatic and human DENV-4 strains, none involved critical cysteines or glycosylation sites. However, one mutation is found in all human genotypes at position 131 (N131S) within the variable residue in the NXT glycosylation signal at this site (Table 3). Furthermore recent evidence suggests that epitope variation within NS1 (Chen et al., 2010; Masrinoul et al., 2011) may influence progression to severe dengue disease by a mechanism of molecular mimicry to human endothelial cell antigens (Liu et al., 2011).

NS2A is a small, hydrophobic protein that in several flaviviruses plays a role in modulating the type I interferon (IFN) response (Munoz-Jordan et al., 2003). Human DENV-4 strains differed from sylvatic strains at 26 amino acids in NS2A (Table 3). The observed high level of variability is characteristic of proteins involved in immune suppression (Franzosa and Xia, 2011; Obbard et al., 2009). NS2B is another small, hydrophobic protein that acts as a

cofactor for the enzymatic activity of NS3. Six nonsynonymous mutations were detected (Table 3), none of which occurred at critical residues) in NS2B required for NS3 function (Niyomrattanakit et al., 2004). NS3, along with NS2B, has multiple functions including protease and helicase. Nineteen non-synonymous differences were noted (Table 3).

NS4A and NS4B are small, multifunctional, hydrophobic proteins that have recently been shown to play a role in modulating the host's type I IFN responses (Jones et al., 2005; Munoz-Jordan et al., 2005; Munoz-Jordan et al., 2003). Other functions of NS4A have not been fully elucidated, but it associates with host cell membranes (Jiang et al., 2009; Miller et al., 2007) and is required for viral replication (Tajima et al., 2011). Nine differences were observed (Table 3) in human NS4A DENV proteins. NS4B, which has also been shown to act as a helper for the helicase function of NS3 (Umareddy et al., 2006) contained 11 differences between sylvatic and human strains (Table 3). A previous study examining the evolutionary pressures on sylvatic DENV-2 identified 13 sites in the NS4B gene under weak positive selective pressure (Vasilakis et al., 2007a). This observation was striking in that no such sites were identified in human DENV-2, and relatively few positively selected sites were identified in any of the other genes in either of the two ecotypes. Four of the observed NS4B mutations in this current study of DENV-4 (positions 11, 18, 20 and 198) were adjacent to sites under positive selection in sylvatic DENV-2 (Vasilakis et al., 2007a).

NS5 acts as the RNA-dependent RNA-polymerase (RdRp) and is responsible for RNA cap processing. Recently, anti-IFN activity of NS5 has also been reported (Ashour et al., 2009). Forty-eight non-synonymous differences were found in NS5 (Table 3), half of which were seen in all genotypes. The N-terminus region has shown to be responsible for methyltransferase activity, and residues critical for this process (positions 14, 18, 29, 61, 146, 150 181 and 217) were unchanged between sylvatic and human strains (Egloff et al., 2002). Five non-synonymous mutations occur within the interdomain region between the N-terminal methyl transferase and the C-terminal RdRp (positions 320–405) which contains two nuclear localization signals (NLS) (Pryor et al., 2007), however all critical residues were unaffected.

Phenotypic analyses

Replication kinetics of four human DENV-4 strains (Genotype I IndiaG11337 and H241, as well as Genotype II Haiti73 and INH6412, Table 1) and the 3 available sylvatic strains (P73-1120, P75-215 and P75-514, Table 1) were characterized in primate kidney and human hepatoma (Vero and Huh-7, respectively) and *Ae. albopictus* mosquito (C6/36) cells. Cultured cells are admittedly an imperfect model of dengue replication *in vivo*; notably Vero cells and C6/36 cells are deficient in the IFN (Mosca and Pitha, 1986) and RNA interference response (Brackney et al., 2010; Scott et al., 2010), respectively. However replication dynamics in these cells is sometimes correlated with replication *in vivo* (Hanley et al., 2003) and therefore they offer a useful first screen for phenotypic variation among virus strains.

Titers of most viruses belonging to both human and sylvatic genotypes increased in both C6/36 and Vero cells until day 4 post-infection (p.i.) (Figure 2a and 2b respectively), and in Huh-7 cells until day 5 p.i. (Figure 2c), after which titers stabilized or declined in all cell types. Huh-7 cells exhibited pronounced cytopathic effect (CPE) by day 4 p.i., which became severe by 6 p.i. in response to infection with all DENV-4 strains (data not shown). Vero cells also exhibited CPE following infection by all strains, but it was less severe than that observed in Huh-7 cells; C6/36 cells exhibited no CPE.

Overall, ecotype (human or sylvatic) had no impact on patterns of DENV-4 multistep replication kinetics in any of the 3 cells lines tested ($P > 0.25$ for all comparisons) (Figure 2d–f). However significant inter-strain variation was detected within ecotypes during

infection of the 3 cell types. The patterns of replication kinetics of individual strains in each cell line are denoted in Figure 2a–c. In C6/36 cells, both time (DF = 7, $F = 4209.7$, $P < 0.0001$) and virus (DF = 6, $F = 132.3$, $P < 0.0001$) as well as the interaction between time and virus (DF = 42, $F = 6.5$, $P < 0.0001$) had a significant effect on viral replication dynamics. Notably, the human India and H241 strains replicated to significantly lower titers than all other strains (Figure 2a). In Vero cells, time (DF = 7, $F = 5276.97$, $P < 0.0001$) and virus (DF = 6, $F = 115.1$, $P < 0.0001$) as well as the interaction between time and virus (DF = 42, $F = 8.7$, $P < 0.0001$) had significant effects on replication dynamics. Contrary to its performance in mosquito cells, strain H241 replicated to significantly higher levels than other strains in Vero cells, while the India strain replicated to significantly lower levels than all other strains (Figure 2b). In Huh-7 cells, both time (DF = 7, $F = 1009.8$, $P < 0.0001$) and virus (DF = 6, $F = 29.2$, $P < 0.0001$) as well as the interaction between time and virus (DF = 42, $F = 4.6$, $P < 0.0001$), had a significant effect on viral replication dynamics. Similar to the pattern observed in Vero cells, H241 replicated to significantly higher levels than most other strains and IndiaG11337 replicated to significantly lower levels than all other strains (Figure 2c).

In conclusion, our phylogenetic analyses of a diverse array of complete DENV-4 genome sequences demonstrated that sylvatic DENV-4 isolates are basal to and evolutionarily distinct from human DENV-4 isolates, supporting previous classifications of DENV-4 into two discrete ecotypes. This evolutionary and ecologic divergence is further corroborated by our genetic analyses, which revealed considerable differences at both the nucleotide and protein levels between human and sylvatic strains. While all sequence and structural elements previously identified as critical for virus replication (i.e. UTR secondary structures, disulfide bridges, and glycosylation sites) were fully conserved in both ecotypes, nonetheless, we identified several amino acids that may have played a role in the emergence of human DENV-4 strains from sylvatic progenitors. Human and sylvatic DENV-4 strains replicated with similar dynamics in green monkey kidney cells, human hepatoma cells and mosquito cells. These studies in cultured cells should be complemented by testing the phenotypes of sylvatic versus human DENV-4 in peridomestic mosquitoes *in vivo* and in models for human replication. However the evidence to date suggests that sylvatic strains of DENV-4 in Asia retain the potential to emerge into human transmission cycles.

Materials and Methods

Viruses and Sequencing

To avoid mutations resulting from adaptation to cell cultures, low passage DENV-4 isolates of the following strains were obtained from the World Reference Center for Emerging Viruses and Arboviruses (WRCEVA) at the University of Texas Medical Branch: P75-514, P73-1120 INH6412, JF262781, Haiti73, JF262782, India G11337, and JF262783 (Table 1). All strains were amplified once in cultured C6/36 *Ae. albopictus* cells to achieve sufficiently high titers for amplification and sequencing. We have previously shown that DENV accumulates relatively few mutations during passage in C6/36 cells (Vasilakis et al., 2009). Initial overlapping cDNA fragments and amplicons of viruses were generated using primer pairs specific to DENV P75-215 or H241 in a one-step (Roche Diagnostics, Indianapolis) or two-step (New England Biolabs, MA) reverse transcriptase polymerase chain reactions (RT-PCR) in 8–13 overlapping cDNA fragments spanning the DENV genome. Two-step RT-PCR was generated from a cDNA template using a specific antisense primer (10666: 5'-AGAACCTGTTGGATCAACAACACC) and M-MuLV reverse transcriptase (New England Biolabs, Ipswich) as per manufacture's instructions. The PCR reaction was performed using the high-fidelity Phusion polymerase (Finnzymes, Lafayette, CO) for 35x amplification cycles at an annealing temperature approximately 5C below the lowest T_m .

Amplified sequences were purified (Qiagen, Valencia, CA) and automated sequencing with specific sequencing primers for both strands provided consensus sequences.

Nucleotide Sequence Accession Numbers

The following DENV-4 genome sequences were determined in this study: INH6412, JF262781; Haiti73, JF262782; India G11337, JF262783; and the sylvatic strains P75-514 and P73-1120, accession numbers JF262779 and JF262780 respectively. The GenBank accession numbers for the genome sequences of DENV viruses used in the phylogenetic analyses are summarized in Supplemental Data 4.

Phylogenetic analyses

Sequences were aligned using the ClustalW multiple sequence alignment function of MacVector® version 8.0 (Accelrys, Cary, NC) with default gap penalties. Phylogenetic analyses of the aligned genomic sequences were estimated with maximum likelihood and distance/neighbor joining using the PAUP* program version 4.10 (D.L. Swofford, Illinois Natural History Survey, Champaign) under the general time-reversible model of nucleotide substitution. The parameter values used for the substitution type, optimal base composition, proportion of invariable sites, as well as the shape parameter of the Γ distribution of rate variation among sites were estimated from the data and are available upon request. We also utilized Bayesian analysis (MrBayes v3.1.0) where 4 MCMC tree searches of 4 million generations each were run simultaneously sampling 1 in 100 trees and computing a 50% majority-rule consensus tree out of the last 9,800 sampled trees, where the initial 10% of trees removed as burn in. Bootstrapping obtained by the Bayesian analysis was used to place confidence values on grouping within the consensus tree (Felsenstein, 1985). Character evolution was traced using MacClade 4.08 (Sinauer Associates, Sunderland, MA).

Viral replication kinetics

Comparative multi-step growth curves of all available sylvatic (P73-1120, P75-215 and P75-514) and selected human (Haiti73, IndiaG11337, INH6412 and H241) DENV-4 strains were generated in triplicate on African green monkey kidney (Vero), human hepatoma (Huh-7) and *Ae. albopictus* (C6/36) cells. Huh-7 cells (clone JTC-39) were obtained from the Japanese Health Sciences Foundation, Osaka. All 3 cell lines were plated in 12-well plates at 3.6×10^5 , 3.6×10^5 , and 7.5×10^5 cells per well respectively, and infected with a MOI of 0.01 per cell in triplicate. Infected dishes containing Huh-7 and Vero cells were incubated for one hour with periodic gentle rocking at 37°C, whereas dishes containing mosquito C6/36 cells were incubated at 28°C. Viral inocula were removed and cell monolayers were washed thrice with PBS to remove unadsorbed virus. Two ml of complete cell media (MEM supplemented with 5% FBS, 2mM L-Glutamine, 1% nonessential amino acids (NEAA), and 50mg/ml penicillin/streptomycin) were then added and dishes were incubated at 28°C or 37°C for the mosquito or mammalian cell lines respectively. Virus from individual dishes was harvested daily through day 7 pi, clarified by low speed centrifugation, and assayed in C6/36 cells to determine virus titer by focus forming immunoassay (FFA). Virus yield at each timepoint is recorded as FFU/cell, represented as the ratio of the total amount of virus present in the sample by the number of cells originally infected.

Focus Forming Assays and Immunostaining

Ten-fold serial dilutions of virus in MEM supplemented with 2% FBS and antibiotics (Invitrogen, Carlsbad, CA), were added in duplicate to confluent C6/36 cell monolayers attached to 24-well Costar® (Corning, NY) plates, and incubated for one hour with periodic gentle rocking to facilitate virus adsorption at 28°C. Wells were then overlaid with 1 ml of 0.8% methylcellulose (Sigma-Aldrich, St. Louis) diluted in warm OptiMem (Invitrogen)

supplemented with 2% FBS, antibiotics and 1% (w/v) L-glutamine and incubated undisturbed for 4 days at 28°C. Methylcellulose overlay was aspirated and cell monolayer rinsed once with phosphate buffered saline (PBS), pH 7.4 (Invitrogen) followed by fixation with a mixture of ice-cold acetone and methanol (1:1) solution and allowed to incubate for 30 minutes at room temperature (RT). Fixation solution was aspirated and plates were allowed to air dry. Plates were washed thrice with PBS supplemented with 3% FBS, followed by hour-long incubation with a dengue-specific hyperimmune mouse ascitic fluid. Plates were washed thrice followed by hour-long incubation with a secondary antibody conjugated to horseradish peroxidase (KPL, Gaithersburg, MD). Detection proceeded with the addition of aminoethylcarbazole (AEC) substrate (ENZO Life sciences, Farmingdale, CT) prepared according to vendor instructions.

Statistical Analyses

To compare multistep replication dynamics, repeated measures ANOVA was first used to compare the replication of each of the seven strains tested, with each of the three cell lines analyzed separately. When a significant effect of virus was detected, the differences among the strains were compared using a Tukey-Kramer post-hoc test. Subsequently, the replication of the two ecotypes (sylvatic and human) was compared in each cell line separately using the mean value for each virus strain at each time point, resulting in $N = 3$ for the sylvatic ecotype and $N = 4$ for the human ecotype.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Naomi Forester and Thomas Monath for helpful comments on the manuscript, and The Japanese Health Sciences Foundation, Osaka, for providing the human hepatoma Huh-7 cells (clone JTC-39). SR is supported by 2T32AI007536-11A1 postdoctoral training grant; SW is supported for dengue research by NIH grant RO1 AI069145; KH is supported by NIH grants RO1 AI069145 and P20RR016480 and a Woods Hole Marine Biological Laboratory Karush Library Fellowship; RT is supported by NIH contract HHSN272201000040I/HHSN27200004/D04; NV was supported by start-up funds provided by the Department of Pathology, UTMB.

References Cited

- AbuBakar S, Wong PF, Chan YF. Emergence of dengue virus type 4 genotype IIA in Malaysia. *J Gen Virol.* 2002; 83(Pt 10):2437–2442. [PubMed: 12237425]
- Ashour J, Laurent-Rolle M, Shi PY, Garcia-Sastre A. NS5 of dengue virus mediates STAT2 binding and degradation. *J Virol.* 2009; 83(11):5408–5418. [PubMed: 19279106]
- Bennett SN, Holmes EC, Chirivella M, Rodriguez DM, Beltran M, Vorndam V, Gubler DJ, McMillan WO. Selection-driven evolution of emergent dengue virus. *Mol Biol Evol.* 2003; 20(10):1650–1658. [PubMed: 12832629]
- Blaney JE Jr, Hanson CT, Firestone CY, Hanley KA, Murphy BR, Whitehead SS. Genetically modified, live attenuated dengue virus type 3 vaccine candidates. *Am J Trop Med Hyg.* 2004a; 71(6):811–821. [PubMed: 15642976]
- Blaney JE Jr, Hanson CT, Hanley KA, Murphy BR, Whitehead SS. Vaccine candidates derived from a novel infectious cDNA clone of an American genotype dengue virus type 2. *BMC Infect Dis.* 2004b; 4(1):39. [PubMed: 15461822]
- Brackney DE, Scott JC, Sagawa F, Woodward JE, Miller NA, Schilkey FD, Mudge J, Wilusz J, Olson KE, Blair CD, Ebel GD. C6/36 *Aedes albopictus* cells have a dysfunctional antiviral RNA interference response. *PLoS Negl Trop Dis.* 2010; 4(10):e856. [PubMed: 21049065]
- Brinton MA, Disposito JH. Sequence and secondary structure analysis of the 5'-terminal region of flavivirus genome RNA. *Virology.* 1988; 162(2):290–299. [PubMed: 2829420]

- Cahour A, Pletnev A, Vazille-Falcoz M, Rosen L, Lai CJ. Growth-restricted dengue virus mutants containing deletions in the 5' noncoding region of the RNA genome. *Virology*. 1995; 207(1):68–76. [PubMed: 7871753]
- Cardosa J, Ooi MH, Tio PH, Perera D, Holmes EC, Bibi K, Abdul Manap Z. Dengue virus serotype 2 from a sylvatic lineage isolated from a patient with dengue hemorrhagic Fever. *PLoS Negl Trop Dis*. 2009; 3(4):e423. [PubMed: 19399166]
- Carey DE, Causey OR, Reddy S, Cooke AR. Dengue viruses from febrile patients in Nigeria, 1964–68. *Lancet*. 1971; 1(7690):105–106. [PubMed: 4099602]
- Cecilia D, Kakade MB, Bhagat AB, Vallentyne J, Singh A, Patil JA, Todkar SM, Varghese SB, Shah PS. Detection of dengue-4 virus in Pune, western india after an absence of 30 years--its association with two severe cases. *Virology Journal*. 2011; 8(1):46. [PubMed: 21284832]
- Chen R, Vasilakis N. Dengue - Quo tu et quo vadis. *Viruses*. 2011 In Press.
- Chen Y, Maguire T, Hileman RE, Fromm JR, Esko JD, Linhardt RJ, Marks RM. Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. *Nat Med*. 1997; 3(8):866–871. [PubMed: 9256277]
- Chen Y, Maguire T, Marks RM. Demonstration of binding of dengue virus envelope protein to target cells. *J Virol*. 1996; 70(12):8765–8772. [PubMed: 8971005]
- Chen Y, Pan Y, Guo Y, Qiu L, Ding X, Che X. Comprehensive mapping of immunodominant and conserved serotype- and group-specific B-cell epitopes of nonstructural protein 1 from dengue virus type 1. *Virology*. 2010; 398(2):290–298. [PubMed: 20079511]
- Cleaves GR, Dubin DT. Methylation status of intracellular dengue type 2 40 S RNA. *Virology*. 1979; 96(1):159–165. [PubMed: 111410]
- Cordellier R, Bouchite B, Roche JC, Monteny N, Diaco B, Akoliba P. Circulation selvatique du virus Dengue 2, en 1980, dans les savanes sub-soudaniennes de Cote d'Ivoire. *Cah ORSTOM ser Ent Med et Parasitol*. 1983; 21:165–179.
- Cox J, Brown HE, Rico-Hesse R. Variation in vector competence for dengue viruses does not depend on mosquito midgut binding affinity. *PLoS Negl Trop Dis*. 2011; 5(5):e1172. [PubMed: 21610852]
- Crill WD, Chang GJ. Localization and characterization of flavivirus envelope glycoprotein cross-reactive epitopes. *J Virol*. 2004; 78(24):13975–13986. [PubMed: 15564505]
- Crill WD, Hughes HR, Delorey MJ, Chang GJ. Humoral immune responses of dengue fever patients using epitope-specific serotype-2 virus-like particle antigens. *PLoS One*. 2009; 4(4):e4991. [PubMed: 19337372]
- Dash PK, Sharma S, Srivastava A, Santhosh SR, Parida MM, Neeraja M, Subbalaxmi MV, Lakshmi V, Rao PV. Emergence of dengue virus type 4 (genotype I) in India. *Epidemiology and Infection*. 2011; 139(6):867–861. [PubMed: 20822577]
- Diallo M, Ba Y, Sall AA, Diop OM, Ndione JA, Mondo M, Girault L, Mathiot C. Amplification of the sylvatic cycle of dengue virus type 2, Senegal, 1999–2000: entomologic findings and epidemiologic considerations. *Emerg Infect Dis*. 2003; 9(3):362–367. [PubMed: 12643833]
- Diallo M, Sall AA, Moncayo AC, Ba Y, Fernandez Z, Ortiz D, Coffey LL, Mathiot C, Tesh RB, Weaver SC. Potential role of sylvatic and domestic african mosquito species in dengue emergence. *Am J Trop Med Hyg*. 2005; 73 (2):445–449. [PubMed: 16103619]
- Durbin AP, Whitehead SS. Dengue vaccine candidates in development. *Curr Top Microbiol Immunol*. 2010; 338:129–143. [PubMed: 19802583]
- Durbin AP, Whitehead SS, Shaffer D, Elwood D, Wanionek K, Thumar B, Blaney JE, Murphy BR, Schmidt AC. A single dose of the DENV-1 candidate vaccine rDEN1Delta30 is strongly immunogenic and induces resistance to a second dose in a randomized trial. *PLoS Negl Trop Dis*. 2011; 5(8):e1267. [PubMed: 21829748]
- Egloff MP, Benarroch D, Selisko B, Romette JL, Canard B. An RNA cap (nucleoside-2'-O-)-methyltransferase in the flavivirus RNA polymerase NS5: crystal structure and functional characterization. *Embo J*. 2002; 21(11):2757–2768. [PubMed: 12032088]
- Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*. 1985; 39(4):783–791.

- Filomatori CV, Iglesias NG, Villordo SM, Alvarez DE, Gamarnik AV. RNA sequences and structures required for the recruitment and activity of the dengue virus polymerase. *J Biol Chem*. 2011; 286(9):6929–6939. [PubMed: 21183683]
- Filomatori CV, Lodeiro MF, Alvarez DE, Samsa MM, Pietrasanta L, Gamarnik AV. A 5' RNA element promotes dengue virus RNA synthesis on a circular genome. *Genes Dev*. 2006; 20(16): 2238–2249. [PubMed: 16882970]
- Foster JE, Bennett SN, Vaughan H, Vorndam V, McMillan WO, Carrington CV. Molecular evolution and phylogeny of dengue type 4 virus in the Caribbean. *Virology*. 2003; 306(1):126–134. [PubMed: 12620805]
- Franco L, Palacios G, Martinez JA, Vazquez A, Savji N, de Ory F, Sanchez-Seco MP, Martin D, WIL, Tenorio A. First report of sylvatic DENV-2-associated dengue hemorrhagic fever in West Africa. *PLoS NTD*. 2011; 5(8):e1251.
- Franzosa EA, Xia Y. Structural principles within the human-virus protein-protein interaction network. *Proc Natl Acad Sci U S A*. 2011; 108(26):10538–10543. [PubMed: 21680884]
- Gritsun TS, Gould EA. Direct repeats in the 3' untranslated regions of mosquito-borne flaviviruses: possible implications for virus transmission. *J Gen Virol*. 2006; 87(Pt 11):3297–3305. [PubMed: 17030864]
- Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler DJ, Hunsperger E, Kroeger A, Margolis HS, Martinez E, Nathan MB, Pelegriño JL, Simmons C, Yoksan S, Peeling RW. Dengue: a continuing global threat. *Nat Rev Microbiol*. 2010; 8(12 Suppl):S7–S16. [PubMed: 21079655]
- Hahn CS, Hahn YS, Rice CM, Lee E, Dalgarno L, Strauss EG, Strauss JH. Conserved elements in the 3' untranslated region of flavivirus RNAs and potential cyclization sequences. *J Mol Biol*. 1987; 198(1):33–41. [PubMed: 2828633]
- Halstead SB. Epidemiology of dengue and dengue hemorrhagic fever. In: Gubler, DJ.; Kuno, G., editors. *Dengue and dengue hemorrhagic fever*. CABI International; Oxon: 1997. p. 23–44.
- Halstead SB, Sukhavachana P, Nisalak A. In Vitro Recovery Of Dengue Viruses From Naturally Infected Human Beings And Arthropods. *Nature*. 1964; 202:931–932. [PubMed: 14190111]
- Hanley KA, Manlucu LR, Gilmore LE, Blaney JE Jr, Hanson CT, Murphy BR, Whitehead SS. A trade-off in replication in mosquito versus mammalian systems conferred by a point mutation in the NS4B protein of dengue virus type 4. *Virology*. 2003; 312(1):222–232. [PubMed: 12890635]
- Hervy JP, Legros F, Roche JC, Monteny N, Diaco B. Circulation du Dengue 2 dans plusieurs milieux boisés des savanes soudanaises de la région de Bobo-Dioulasso (Burkina Faso). *Considerations entomologiques et épidémiologiques*. *Cah ORSTOM ser Ent Med et Parasitol*. 1984; 22:135–143.
- Holmes EC, Twiddy SS. The origin, emergence and evolutionary genetics of dengue virus. *Infect Genet Evol*. 2003; 3(1):19–28. [PubMed: 12797969]
- Jiang L, Yao H, Duan X, Lu X, Liu Y. Polypyrimidine tract-binding protein influences negative strand RNA synthesis of dengue virus. *Biochem Biophys Res Commun*. 2009; 385(2):187–192. [PubMed: 19450550]
- Johnson AJ, Guirakhoo F, Roehrig JT. The envelope glycoproteins of dengue 1 and dengue 2 viruses grown in mosquito cells differ in their utilization of potential glycosylation sites. *Virology*. 1994; 203(2):241–249. [PubMed: 8053148]
- Jones CT, Ma L, Burgner JW, Groesch TD, Post CB, Kuhn RJ. Flavivirus capsid is a dimeric alpha-helical protein. *J Virol*. 2003; 77(12):7143–7149. [PubMed: 12768036]
- Jones M, Davidson A, Hibbert L, Gruenwald P, Schlaak J, Ball S, Foster GR, Jacobs M. Dengue virus inhibits alpha interferon signaling by reducing STAT2 expression. *J Virol*. 2005; 79(9):5414–5420. [PubMed: 15827155]
- Klungthong C, Zhang C, Mammen MP Jr, Ubol S, Holmes EC. The molecular epidemiology of dengue virus serotype 4 in Bangkok, Thailand. *Virology*. 2004; 329(1):168–179. [PubMed: 15476884]
- Leitmeyer KC, Vaughn DW, Watts DM, Salas R, Villalobos I, de C, Ramos C, Rico-Hesse R. Dengue virus structural differences that correlate with pathogenesis. *J Virol*. 1999; 73(6):4738–4747. [PubMed: 10233934]

- Li L, Lok SM, Yu IM, Zhang Y, Kuhn RJ, Chen J, Rossmann MG. The flavivirus precursor membrane-envelope protein complex: structure and maturation. *Science*. 2008; 319(5871):1830–1834. [PubMed: 18369147]
- Lindenbach, BD.; Rice, CM. Molecular Biology of Flaviviruses. In: Chambers, TJ.; Monath, TP., editors. *The Flaviviruses: Structure, Replication and Evolution*. Elsevier Academic Press; 2003.
- Liu JJ, Chiu CY, Chen YC, Wu HC. Molecular mimicry of human endothelial cell antigen by autoantibodies to nonstructural protein 1 of dengue virus. *J Biol Chem*. 2011; 286(11):9726–9736. [PubMed: 21233208]
- Markoff, L. 5'- and 3'-Noncoding Regions in Flavivirus RNA. In: Monath, TP.; Chambers, TJ., editors. *The Flaviviruses: Structure, Replication and Evolution*. Vol. 59. Elsevier Academic Press; San Diego: 2003. Vol. *Advances in Virus Research*
- Markoff L, Falgout B, Chang A. A conserved internal hydrophobic domain mediates the stable membrane integration of the dengue virus capsid protein. *Virology*. 1997; 233(1):105–117. [PubMed: 9201220]
- Masrinoul P, Diata MO, Pambudi S, Limkittikul K, Ikuta K, Kurosu T. Highly conserved region 141168 of the NS1 protein is a new common epitope region of dengue virus. *Jpn J Infect Dis*. 2011; 64(2):109–115. [PubMed: 21519123]
- Miller S, Kastner S, Krijnse-Locker J, Buhler S, Bartenschlager R. The non-structural protein 4A of dengue virus is an integral membrane protein inducing membrane alterations in a 2K-regulated manner. *J Biol Chem*. 2007; 282(12):8873–8882. [PubMed: 17276984]
- Modis Y, Ogata S, Clements D, Harrison SC. Structure of the dengue virus envelope protein after membrane fusion. *Nature*. 2004; 427(6972):313–319. [PubMed: 14737159]
- Modis Y, Ogata S, Clements D, Harrison SC. Variable surface epitopes in the crystal structure of dengue virus type 3 envelope glycoprotein. *J Virol*. 2005; 79(2):1223–1231. [PubMed: 15613349]
- Monlun E, Zeller H, Traore-Lamizana M, Hervy JP, Adam F, Mondo M, Digoutte JP. Caracteres cliniques et epidemiologiques de la dengue 2 au Senegal. *Med Mal Infect*. 1992; 22:718–721.
- Mosca JD, Pitha PM. Transcriptional and posttranscriptional regulation of exogenous human beta interferon gene in simian cells defective in interferon synthesis. *Mol Cell Biol*. 1986; 6(6):2279–2283. [PubMed: 3785197]
- Mota J, Rico-Hesse R. Humanized mice show clinical signs of dengue fever according to infecting virus genotype. *J Virol*. 2009; 83(17):8638–8645. [PubMed: 19535452]
- Mota J, Rico-Hesse R. Dengue virus tropism in humanized mice recapitulates human dengue fever. *PLoS One*. 2011; 6(6):e20762. [PubMed: 21695193]
- Munoz-Jordan JL, Laurent-Rolle M, Ashour J, Martinez-Sobrido L, Ashok M, Lipkin WI, Garcia-Sastre A. Inhibition of alpha/beta interferon signaling by the NS4B protein of flaviviruses. *J Virol*. 2005; 79(13):8004–8013. [PubMed: 15956546]
- Munoz-Jordan JL, Sanchez-Burgos GG, Laurent-Rolle M, Garcia-Sastre A. Inhibition of interferon signaling by dengue virus. *Proc Natl Acad Sci U S A*. 2003; 100(24):14333–14338. [PubMed: 14612562]
- Niyomrattanakit P, Winoyanu wattikun P, Chanprapaph S, Angsuthanasombat C, Panyim S, Katzenmeier G. Identification of residues in the dengue virus type 2 NS2B cofactor that are critical for NS3 protease activation. *J Virol*. 2004; 78(24):13708–13716. [PubMed: 15564480]
- Obbard DJ, Gordon KH, Buck AH, Jiggins FM. The evolution of RNAi as a defence against viruses and transposable elements. *Philos Trans R Soc Lond B Biol Sci*. 2009; 364(1513):99–115. [PubMed: 18926973]
- Proutski V, Gritsun TS, Gould EA, Holmes EC. Biological consequences of deletions within the 3'-untranslated region of flaviviruses may be due to rearrangements of RNA secondary structure. *Virus Res*. 1999; 64(2):107–123. [PubMed: 10518708]
- Pryor MJ, Rawlinson SM, Butcher RE, Barton CL, Waterhouse TA, Vasudevan SG, Bardin PG, Wright PJ, Jans DA, Davidson AD. Nuclear localization of dengue virus nonstructural protein 5 through its importin alpha/beta-recognized nuclear localization sequences is integral to viral infection. *Traffic*. 2007; 8(7):795–807. [PubMed: 17537211]
- Pryor MJ, Wright PJ. Glycosylation mutants of dengue virus NS1 protein. *J Gen Virol*. 1994; 75 (Pt 5):1183–1187. [PubMed: 8176380]

- Rauscher S, Flamm C, Mandl CW, Heinz FX, Stadler PF. Secondary structure of the 3'-noncoding region of flavivirus genomes: comparative analysis of base pairing probabilities. *Rna*. 1997; 3(7): 779–791. [PubMed: 9214660]
- Rey FA, Heinz FX, Mandl C, Kunz C, Harrison SC. The envelope glycoprotein from tick-borne encephalitis virus at 2 Å resolution. *Nature*. 1995; 375(6529):291–298. [PubMed: 7753193]
- Rice CM, Lenches EM, Eddy SR, Shin SJ, Sheets RL, Strauss JH. Nucleotide sequence of yellow fever virus: implications for flavivirus gene expression and evolution. *Science*. 1985; 229(4715): 726–733. [PubMed: 4023707]
- Rodhain F. The role of monkeys in the biology of dengue and yellow fever. *Comp Immunol Microbiol Infect Dis*. 1991; 14(1):9–19. [PubMed: 1905605]
- Romero TA, Tumban E, Jun J, Lott WB, Hanley KA. Secondary structure of dengue virus type 4 3' untranslated region: impact of deletion and substitution mutations. *J Gen Virol*. 2006; 87(Pt 11): 3291–3296. [PubMed: 17030863]
- Rosen L, Rozeboom LE, Sweet BH, Sabin AB. The transmission of dengue by *Aedes polynesiensis* Marks. *Am J Trop Med Hyg*. 1954; 3(5):878–882. [PubMed: 13197723]
- Rudnick A. Dengue virus ecology in Malaysia. *Inst Med Res Malays Bull*. 1986; 23:51–152.
- Sabin AB. Research on dengue during World War II. *Am J Trop Med Hyg*. 1952; 1(1):30–50. [PubMed: 14903434]
- Saluzzo JF, Cornet M, Adam C, Eyraud M, Digoutte JP. Dengue 2 in eastern Senegal: serologic survey in simian and human populations. 1974–85. *Bull Soc Pathol Exot Filiales*. 1986a; 79(3):313–322. [PubMed: 3769119]
- Saluzzo JF, Cornet M, Castagnet P, Rey C, Digoutte JP. Isolation of dengue 2 and dengue 4 viruses from patients in Senegal. *Trans R Soc Trop Med Hyg*. 1986b; 80(1):5. [PubMed: 3726996]
- Scott JC, Brackney DE, Campbell CL, Bondu-Hawkins V, Hjelle B, Ebel GD, Olson KE, Blair CD. Comparison of dengue virus type 2-specific small RNAs from RNA interference-competent and -incompetent mosquito cells. *PLoS Negl Trop Dis*. 2010; 4(10):e848. [PubMed: 21049014]
- Shurtleff AC, Beasley DW, Chen JJ, Ni H, Suderman MT, Wang H, Xu R, Wang E, Weaver SC, Watts DM, Russell KL, Barrett AD. Genetic variation in the 3' non-coding region of dengue viruses. *Virology*. 2001; 281(1):75–87. [PubMed: 11222098]
- Simmons JS, St John JH, Reynolds FHK. Experimental studies of dengue. *Philippine J Sci*. 1931; 44:1–252.
- Smith CE. The history of dengue in tropical Asia and its probable relationship to the mosquito *Aedes aegypti*. *J Trop Med Hyg*. 1956; 59(10):243–251. [PubMed: 13368255]
- Tajima S, Takasaki T, Kurane I. Restoration of replication-defective dengue type 1 virus bearing mutations in the N-terminal cytoplasmic portion of NS4A by additional mutations in NS4B. *Arch Virol*. 2011; 156(1):63–69. [PubMed: 20882304]
- Thullier P, Demangel C, Bedouelle H, Megret F, Jouan A, Deubel V, Mazie JC, Lafaye P. Mapping of a dengue virus neutralizing epitope critical for the infectivity of all serotypes: insight into the neutralization mechanism. *J Gen Virol*. 2001; 82(Pt 8):1885–1892. [PubMed: 11457994]
- Turner C, Witwer C, Hofacker IL, Stadler PF. Conserved RNA secondary structures in Flaviviridae genomes. *J Gen Virol*. 2004; 85(Pt 5):1113–1124. [PubMed: 15105528]
- Trent DW. Antigenic characterization of flavivirus structural proteins separated by isoelectric focusing. *J Virol*. 1977; 22(3):608–618. [PubMed: 69040]
- Umareddy I, Chao A, Sampath A, Gu F, Vasudevan SG. Dengue virus NS4B interacts with NS3 and dissociates it from single-stranded RNA. *J Gen Virol*. 2006; 87(Pt 9):2605–2614. [PubMed: 16894199]
- Vasilakis N, Cardosa J, Hanley KA, Holmes EC, Weaver SC. Fever from the forest: prospects for the continued emergence of sylvatic dengue virus and its impact on public health. *Nat Rev Microbiol*. 2011; 9(7):532–541. [PubMed: 21666708]
- Vasilakis N, Deardorff ER, Kenney JL, Rossi SL, Hanley KA, Weaver SC. Mosquitoes put the brake on arbovirus evolution: experimental evolution reveals slower mutation accumulation in mosquito than vertebrate cells. *PLoS Pathog*. 2009; 5(6):e1000467. [PubMed: 19503824]

- Vasilakis N, Durbin A, Travassos da Rosa APA, Munoz-Jordan JL, Tesh RB, Weaver SC. Antigenic relationships between sylvatic and endemic dengue viruses. *Am J Trop Med Hyg.* 2008a; 79(1): 128–132. [PubMed: 18606776]
- Vasilakis N, Fokam EB, Hanson CT, Weinberg E, Sall AA, Whitehead SS, Hanley KA, Weaver SC. Genetic and phenotypic characterization of sylvatic dengue virus type 2 strains. *Virology.* 2008b; 377(2):296–307. [PubMed: 18570968]
- Vasilakis N, Holmes EC, Fokam EB, Faye O, Diallo M, Sall AA, Weaver SC. Evolutionary processes among sylvatic Dengue-2 viruses. *Journal of Virology.* 2007a; 81:9591–9595. [PubMed: 17553878]
- Vasilakis N, Shell EJ, Fokam EB, Mason PW, Hanley KA, Estes DM, Weaver SC. Potential of ancestral sylvatic dengue-2 viruses to re-emerge. *Virology.* 2007b; 358(2):402–412. [PubMed: 17014880]
- Vasilakis N, Tesh RB, Weaver SC. Sylvatic dengue virus type 2 activity in humans, Nigeria, 1966. *Emerg Infect Dis.* 2008c; 14(3):502–504. [PubMed: 18325274]
- Vasilakis N, Weaver SC. The History and Evolution of Human Dengue Emergence. *Adv Vir Res.* 2008; 72:1–76.
- Villabona-Arenas CJ, Zanotto PM. Evolutionary history of Dengue virus type 4: Insights into genotype phylodynamics. *Infect Genet Evol.* 2011; 11(5):878–885. [PubMed: 21335103]
- Wang E, Ni H, Xu R, Barrett AD, Watowich SJ, Gubler DJ, Weaver SC. Evolutionary relationships of endemic/epidemic and sylvatic dengue viruses. *J Virol.* 2000; 74(7):3227–3234. [PubMed: 10708439]
- Weaver SC, Reisen WK. Present and future arboviral threats. *Antiviral Res.* 2010; 85(2):328–345. [PubMed: 19857523]
- Weaver SC, Vasilakis N. Molecular evolution of dengue viruses: contributions of phylogenetics to understanding the history and epidemiology of the preeminent arboviral disease. *Infect Genet Evol.* 2009; 9(4):523–540. [PubMed: 19460319]
- Wengler G, Wengler G. Terminal sequences of the genome and replicative-from RNA of the flavivirus West Nile virus: absence of poly(A) and possible role in RNA replication. *Virology.* 1981; 113(2): 544–555. [PubMed: 7269253]
- Wengler G, Wengler G, Gross HJ. Studies on virus-specific nucleic acids synthesized in vertebrate and mosquito cells infected with flaviviruses. *Virology.* 1978; 89(2):423–437. [PubMed: 568848]
- Wilder-Smith A, Gubler DJ. Geographic expansion of dengue: the impact of international travel. *Med Clin North Am.* 2008; 92(6):1377–1390. x. [PubMed: 19061757]
- Yu L, Nomaguchi M, Padmanabhan R, Markoff L. Specific requirements for elements of the 5' and 3' terminal regions in flavivirus RNA synthesis and viral replication. *Virology.* 2008; 374(1):170–185. [PubMed: 18234265]
- Zhao B, Mackow E, Buckler-White A, Markoff L, Chanock RM, Lai CJ, Makino Y. Cloning full-length dengue type 4 viral DNA sequences: analysis of genes coding for structural proteins. *Virology.* 1986; 155(1):77–88. [PubMed: 3022479]





Figure 1. Phylogenetic analysis of sylvatic DENV-4 isolates

Phylogenetic tree derived from complete genome nucleotide sequences of sylvatic and human DENV-4 strains using Bayesian analysis. Numbers of posterior probability values for monophyletic groups to the right. Asterisks indicate DENV strains used to evaluate replication kinetics. Virus strains are coded by abbreviated country of collection/strain name/year of collection. The scale shows a genetic distance of 0.02 or 2% nucleotide sequence divergence






Figure 2. Comparative replication curves of DENV-4 strains

a–c: Daily virus output from days 0–6 following infection at MOI 0.01 by sylvatic (P73-1120, P75-215 and P75-514) and human (Haiti73, IndiaG11337, INH6412 and H241) DENV-4 strains in mosquito cell line C6/36 (*Ae. albopictus*) and vertebrate cell lines Vero (African green monkey kidney) and Huh-7 (human hepatoma) (a–c, respectively). **d–f:** Mean virus output of human (squares) and sylvatic (diamonds) DENV-4 strains in C6/36, Vero, and Huh-7 cells (d–f, respectively); means derived from 3 strains of sylvatic and four human DENV-4 shown in panels a–c. The limit of detection of the assay is 0.9 log₁₀ ffu/mL.

Table 1

Characterization of genetic sequence of selected dengue-4 sylvatic strains.

Virus ^a	Ecotype ^b	Location	Host ^c	Year	GenBank Accession No.	Genome Length	% nt Identity ^c	% aa Identity ^c	ORF (nt)	5'UTR (nt)	3'UTR (nt)
H241	Human	The Philippines	Human	1956	AY947539	10643 ^d	--	--	10164	101	282 ^d
IndiaG11337	Human	India	Human	1961	JF262783	10659	95.1%	99.2%	10164	101	394
INH6412	Human	Venezuela	Human	1995	JF262781	10649	94.3%	99.1%	10164	101	384
Haiti73	Human	Haiti	Human	1994	JF262782	10649	94.3%	99.1%	10164	101	384
P73-1120	Sylvatic	Malaysia	<i>A. niveus</i> s.l.	1975	EF457906	10667	86.9%	98.3%	10164	101	402
P75-514	Sylvatic	Malaysia	<i>Presbytis cristata</i> (experimental infection)	1975	JF262779	10667	87.0%	98.3%	10164	101	402
P75-215	Sylvatic	Malaysia	<i>Presbytis cristata</i> (Sentinel monkey)	1973	JF262780	10666	86.9%	98.3%	10164	101	401

^aLow passage DENV isolates were obtained from the UTMB World Reference Center for Emerging Viruses and Arboviruses and amplified once on C6/36 mosquito cells to achieve sufficiently high titers for further evaluation

^bHuman indicates isolates associated with the human transmission cycle or associated with peridomestic transmission; sylvatic indicates sentinel monkey or canopy-dwelling mosquito isolate; see also Figure 1 for phylogenetic groupings.

^cSource of virus isolation.

^cDenotes % identity compared to endemic H241 reference strain

^dPartial sequence; the 3' terminus of the 3'UTR is missing

Table 2

Nucleotide identity (%) between the consensus sequence of the 5'-UTR (above the diagonal) and 3'-UTR (below the diagonal) of human DENV-4 genotypes I–III and sylvatic DENV-4

5' – UTR	Genotype I	Genotype II	Genotype III	Sylvatic
Genotype I	100.0	96.2	97.1	98.0
Genotype II	93.1	100.0	95.2	96.2
Genotype III	93.8	95.1	100.0	97.0
Sylvatic	91.9	88.7	88.9	100.0

Table 3

Summary of consistent amino acid differences between sylvatic and human DENV-4 genotypes

Gene	Site ^a	Type ^b	AA substitution	Comment
C	8	I – III	A → V	T → T; PR 1995 strains
	34	III	T → I	
	34	II	T → S	
	74	III	N → T	
	81	I – II	T → I	
	102	III	M → V	
	102	I – II	M → I	
	111	I – III	I → V	
prM	5	I – III	S → T	
	41	I – III	V → L	
	70	I – III	S → L	
	76	I – III	T → M	
	83	III	S → N	
M (mature)	6	I – III	A → T	
	17	I – III	T → A	
E	4	III	V → I	I → I: G11337;0348;0087; 0485; BID-V3361
	19	I – III	T → A	
	46	I – II	I → T	
	132	I – III	V → I	
	148	I – III	A → T	
	154	I – III	S → D	
	162	I – III	T → A	
	174	III	K → E	
	182	III	T → S	
	202	III	K → E	
	203	I – III	G → K	
	233	III	Y → H	

Gene	Site ^a	Type ^b	AA substitution	Comment
NS1	265	I - II	T → A	S → S: G11337;0348;0087; 0485; BID-V3361
	310	III	K → R	
	329	I - II	T → A	
	335	I - III	I → V	
	340	I - III	K → R	
	342	I - III	M → V	
	351	I	I → V	
	355	I - III	I → T	
	357	II	F → L	
	364	I - III	I → V	
	382	I - III	A → V	
	384	II	D → N	
	455	III	I → V	
	461	I - II	I → F	
	461	III	I → L	
	474	III	S → P	
	478	I - III	S → T	
	9	III	S → T	
NS1	22	I - II	I → V	L → L: G11337;0348;0087; 0485; BID-V3361
	22	III	I → T	
	33	III	K → Q	
	57	I - III	V → I	
	90	I - III	T → A	
	98	III	T → V	
	103	I - III	V → A	
	104	I - III	I → L	
	105	I - III	A → T	
	129	III	R → K	
	131	I - III	N → S	
	152	II	F → S	
	153	I	L → F	

Gene	Site ^a	Type ^b	AA substitution	Comment
	156	I – III	D → E	
	166	III	N → S	
	171	I – III	L → F	
	188	I – III	V → I	
	192	I – III	R → K	
	246	I – II	T → S	
	251	I – III	S → F	
	265	II	M → Y	
	275	I – III	M → I	
	292	I – III	G → D	
	338	I – III	S → L	
	341	III	K → R	
	NS2A	4	III	P → S
		4	I – II	P → T
		19	III	M → I
		26	III	R → K
		38	III	A → T
		38	II	A → I
		40	III	L → F
		54	I – III	V → L
		57	I – II	I → L
		65	III	F → L
		65	I – II	F → S
		66	III	G → S
		68	III	M → V
		68	I – II	M → I
		72	III	V → T
		78	III	A → I
		113	II	F → L
		127	I – III	A → S
		135	I – II	M → I
				M → M: G11337;0348;0087; 0485; BID-V3361
				M → M: G11337;0348;0087; 0485; BID-V3361

Gene	Site ^a	Type ^b	AA substitution	Comment
	137	I – III	Y → T	K→K; G11337;0348;0087; 0485; BID- V3361
	138	II	H → Q	
	142	I – III	A → T	
	144	I – III	L → V	
	155	I – III	K → R	
	156	I – III	T → S	
	161	III	I → T	
	161	I – II	I → V	
	169	I – III	V → A	
	173	III	V → A	
	197	I – II	I → L	
	217	I – II	K → R	
NS2B	55	III	K → R	A→A; G11337;0348;0087; 0485; BID- V3361
	73	I – III	V → I	
	94	I – III	L → M	
	116	III	V → I	
	119	II – III	A → T	
	127	III	K → R	
NS3	12	I – II	T → A	T→T; G11337;0348;0087; 0485; BID- V3361
	13	I – II	A → T	
	17	II	M → A	
	17	I, III	M → T	
	19	I – II	T → S	
	31	I – II	L → F	
	40	I – III	V → I	
	42	I – III	V → M	
	87	I – III	E → D	
	100	I – III	V → I	
	142	II	K → R	
	170	I – III	K → R	

Gene	Site ^a	Type ^b	AA substitution	Comment
NS4A	190	I – III	V → I	S → S: G11337;0348;0087; 0485; BID-V3361
	249	I – III	A → S	
	292	I	S → C	
	321	II	T → A	
	322	II – III	I → T	
	381	I	K → R	K → K: G11337;0348;0087; 0485;BID-V3361
	518	III	R → K	
	538	I – III	R → K	
	20	I	R → K	
	58	I – III	I → V	R → R: G11337;0348;0087; 0485;BID-V3361
NS4B	83	III	M → V	
	87	II	A → T	
	97	I – III	I → V	
	140	I – II	A → T	
	141	I – III	L → I	
	146	I – III	S → G	
	148	III	I → V	
	11	III	T → A	A → A: G11337;0348;0087; 0485;BID-V3361
	18	I – III	E → V	
	20	I – III	P → T	
	87	I – III	S → G	
	93	I – III	L → M	
	175	I	A → V	
	187	I	F → L	F → F: G11337;0348;0087; 0485;BID-V3361
	198	I – II	A → I	
	198	III	A → V	
	199	I	L → M	
NS5	205	I – III	G → N	L → L: G11337;0348;0087; 0485;BID-V3361
	240	I – III	T → A	
	42	I – III	L → S	

Gene	Site ^a	Type ^b	AA substitution	Comment
	45	I	K → R	K → K: G11337;0348;0087; 0485;BID-V3361
	70	II	I → V	I → I: G11337;0348;0087; 0485;BID-V3361
	90	I – III	V → M	
	100	I – III	R → K	
	102	I – III	F → Y	
	163	I – III	I → V	
	165	I – III	R → K	
	199	I	K → R	K → K: G11337;0348;0087; BID-V3361
	200	I – III	Y → H	
	203	I, III	N → S	
	205	I – II	I → V	
	233	I – III	V → T	
	234	I	T → I	T → T: G11337;0348;0087; BID-V3361
	235	I – III	K → R	
	275	I – III	V → I	
	284	I	Q → R	Q → Q: 0348;0087;0485; BID-V3361
	320	I – III	I → V	
	367	III	P → L	
	371	I – II	V → M	
	372	I	V → I	V → V: 0348;0087;0485; BID-V3361
	386	I – II	R → K	
	419	I – III	A → T	
	432	I – III	K → E	
	500	I – III	D → E	
	520	II	D → E	
	523	III	K → R	
	525	I – III	E → D	
	529	I	M → I	
	547	II	L → Q	
	559	I – III	G → H	
	561	III	K → R	

Gene	Site ^a	Type ^b	AA substitution	Comment
	562	I	I → T	I → E: G11337;0348;0087; 0485;BID-V3361
	585	II	K → R	
	642	I – III	K → R	
	648	II	K → R	
	675	II	S → G	
	717	I – III	L → F	
	749	I – II	K → R	
	785	I – III	V → T	
	788	I – III	V → F	
	831	I – III	Y → H	
	854	I – III	T → S	
	864	I – III	Q → H	
	885	I – III	A → V	
	887	III	K → R	
	892	III	P → L	
	893	II – III	L → S	L → F: G11337;0348;0087; 0485;BID-V3361; 27949

NOTES

Amino acid to the left represents amino acid in sylvatic viruses and to the right on the human viruses. Arrows represent directionality of mutation

^aIndicated amino acid location from the start of the gene

^bI, II, III: represents genotype I, II and III of dengue virus serotype 4.