

Notes for General Topology

Zhao Wenchuan

April 19, 2021

Chapter 1

Definitions

1.1 Basic Definitions

Definition 1.1.1 (Topology and Topological Space). Let X be a set, and let a family $\mathcal{T} \subseteq \mathcal{P}(X)$. \mathcal{T} is called a topology on X iff

- (i) $\emptyset, X \in \mathcal{T}$;
- (ii) \mathcal{T} is closed under arbitrary union;
- (iii) \mathcal{T} is closed under finite intersection.

The pair (X, \mathcal{T}) is called a *topological space*. The elements of \mathcal{T} are called *open sets* in (X, \mathcal{T}) .

Definition 1.1.2 (Metrizible Topology). Let (X, \mathcal{T}) be a topological space.

1.2 Untitled

Definition 1.2.1 (Cover). Let (X, \mathcal{T}) be a topological space, and let $U \subseteq X$, then a family $\mathcal{C} \subseteq \mathcal{P}(X)$ is called a *cover* of U iff the union of \mathcal{C} is a superset of U . That is,

$$U \subseteq \bigcup \mathcal{C}.$$

If $\mathcal{C} \subseteq \mathcal{T}$, then we call \mathcal{C} an *open cover* of U .

Let $\mathcal{C}' \subseteq \mathcal{C}$, iff the union of \mathcal{C}' is still a superset of U , then we call \mathcal{C}' a subcover of \mathcal{C} .

Definition 1.2.2 (Basis). Let (X, \mathcal{T}) be a topological space, let $U \subseteq X$, and let $\mathcal{B} \subseteq \mathcal{P}(X)$ be a cover of X . We call \mathcal{B} a *base* of (X, \mathcal{T}) iff $\mathcal{B} \subseteq \mathcal{T}$ and the union of \mathcal{B} is exactly U itself. That is,

$$\mathcal{B} \subseteq \mathcal{T}, \text{ and } U = \bigcup \mathcal{B}.$$

Definition 1.2.3 (Subspace Topology). Let (X, \mathcal{T}) be a topological space and let $A \subseteq X$. The *subspace topology* \mathcal{T}_A on A is defined to be the family of the intersections of open sets in (X, \mathcal{T}) and A . That is,

$$\mathcal{T}_A = \{U \cap A : U \in \mathcal{T}\}.$$

Definition 1.2.4 (Quotient Topology). Let (X, \mathcal{T}) be a topological space and let \sim be an equivalence relation on X . The *quotient topology* is a topology on $\mathcal{P}(X/\sim)$; it is defined as

$$\mathcal{T}_{X/\sim} = \{U \in \mathcal{P}(X/\sim) : \{x \in X : [x] \in U\} \in \mathcal{T}_X\}.$$

Definition 1.2.5 (Continuous Functions). Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A function $f : X \rightarrow Y$ is said to be *continuous* iff for all open subset U of Y , the preimage $f^{-1}[U]$ is open in X . That is,

$$\forall U \in \mathcal{T}_Y, \quad f^{-1}[U] \in \mathcal{T}_X.$$

Definition 1.2.6 (Homeomorphisms). Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A bijection $f : X \rightarrow Y$ is called a *homeomorphism* iff it is continuous and its inverse is also continuous.

Definition 1.2.7 (Homeomorphic). Two topological spaces (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) are said to be *homeomorphic* or *topologically equivalent*, denoted $X \cong Y$, iff there is an homeomorphism between them.

Definition 1.2.8 (Compactness). A topological space (X, \mathcal{T}) is said to be *compact* iff every open cover of X has a finite subcover. That is,

$$\forall \mathcal{C} \subseteq \mathcal{T} : \bigcup \mathcal{C} = X, \quad \exists \mathcal{S} \subseteq \mathcal{C} : \bigcup \mathcal{S} = X, \quad |\mathcal{S}| < \aleph_0.$$

Definition 1.2.9 (Connectedness). Let (X, \mathcal{T}) be a topological space. (X, \mathcal{T}) is said to be *connected* iff X is not empty and it is not the union of any disjoint open sets. That is,

$$\forall U, V \in \mathcal{T} : X = U \cup V, \quad U \cap V \neq \emptyset.$$

Definition 1.2.10 (Path-connectedness). Let (X, \mathcal{T}) be a topological space.

- (i) A map $\gamma : [0, 1] \rightarrow X$ is called a *path* in X iff it is continuous. If $\gamma(0) = x$ and $\gamma(1) = y$, we say that γ is path from x to y in X .
- (ii) X is said to be *path-connected* iff for all $x, y \in X$ there is a path from x to y in X .

Definition 1.2.11 (T_0 space). A topological space (X, \mathcal{T}) is said to be T_0 or *Kolmogorov*, iff all distinct points $x, y \in X$ are *topologically distinguishable*. That is

$$\forall x, y \in X : x \neq y, \exists N \in \mathcal{T}, \quad x \in N \wedge y \notin N.$$

Definition 1.2.12 (R_0 space).