

Notes for General Topology

Zhao Wenchuan

April 26, 2021

Contents

1	Topological Spaces	2
1.1	Topological Spaces	2
1.2	Cover and Basis	3
1.3	Continuity	4
1.4	Interiors and Closures	5
1.5	Boundaries	7
1.6	Limit Points	9
1.7	Subspaces	9
2	Creating New Spaces	10
2.1	Quotient Spaces	10
2.2	Product Spaces	10
2.3	Homeomorphisms	10
3	Topological Properties	11
3.1	Cardinal Functions	11
3.2	Separation Axioms	11
3.3	Countability Axioms	15
3.4	Metrizability	15
3.5	Compactness	15
3.6	Connectedness	15

Chapter 1

Topological Spaces

1.1 Topological Spaces

Definition 1.1.1 (topology). Let X be a set, and let a family $\mathcal{T} \subseteq \mathcal{P}(X)$. \mathcal{T} is called a topology on X iff

- (i) $\emptyset, X \in \mathcal{T}$;
- (ii) \mathcal{T} is closed under arbitrary union;
- (iii) \mathcal{T} is closed under finite intersection.

Definition 1.1.2 (topological spaces). Let X be any set, and let \mathcal{T} be a topology on X , then the pair (X, \mathcal{T}) is called a *topological space*. All subsets of X in \mathcal{T} are called *open sets* in (X, \mathcal{T}) .

Definition 1.1.3 (closed sets). Let (X, \mathcal{T}) be a topological space. A subset V of X is said to be *closed* iff there is an open set U in X such that

$$V = X \setminus U.$$

Proposition 1.1.1. Let X be a set, and let \mathcal{C} be the family of all closed sets in X . Then

- (i) $\emptyset, X \in \mathcal{C}$;
- (ii) \mathcal{C} is closed under arbitrary intersection;
- (iii) \mathcal{C} is closed under finite union.

Definition 1.1.4 (finer and coarser topology). Let X be any set, and let $\mathcal{T}, \mathcal{T}'$ be topologies on X . \mathcal{T} is said to be *finer* than \mathcal{T}' iff $\mathcal{T} \supseteq \mathcal{T}'$; respectively, \mathcal{T} is said to be *coarser* than \mathcal{T}' iff $\mathcal{T} \subseteq \mathcal{T}'$.

Definition 1.1.5 (neighbourhood). Given (X, \mathcal{T}) as a topological space and a point $x \in X$, a subset $N \subseteq X$ is called a *neighbourhood* iff it contains an open set U containing x .

Proposition 1.1.2. Given (X, \mathcal{T}) as a topological space and $U \subseteq X$, U is open iff for all $x \in U$, there is a neighbourhood N of x contained in U .

Proof. If U is open, then U itself is a neighbourhood of x contained in U .

Conversely, if for all $x \in U$, there is a neighbourhood N_x of x contained in U , then there is a open neighbourhood $U_x \ni x$ contained in N_x . Then we have

$$U \supseteq \bigcup_{x \in U} U_x.$$

Suppose U is not open, then U is a proper superset in the relation above. Then there exists $y \in U$ which is not in any U_x . This implies that such a y does not have any neighbourhood N_y in U , for such an N_y must contains an open $U_y \ni y$. For if it does, then there must be a U_x contains y . This is a contradiction. Thus,

$$U = \bigcup_{x \in U} U_x$$

is open. □

1.2 Cover and Basis

Definition 1.2.1 (cover). Let (X, \mathcal{T}) be a topological space, and let $U \subseteq X$, then a family $\mathcal{C} \subseteq \mathcal{P}(X)$ is called a *cover* of U iff the union of all sets in \mathcal{C} is a superset of U . That is,

$$U \subseteq \bigcup \mathcal{C}.$$

If $\mathcal{C} \subseteq \mathcal{T}$, then we call \mathcal{C} an *open cover* of U .

Let $\mathcal{S} \subseteq \mathcal{C}$, iff the union of \mathcal{S} is still a superset of U , then we call \mathcal{S} a *subcover* of \mathcal{C} .

Definition 1.2.2 (basis). Let (X, \mathcal{T}) be a topological space, let $U \subseteq X$, and let \mathcal{B} be a open cover of X . We call \mathcal{B} a *base* of X iff the union of \mathcal{B} is precisely U itself, i.e.,

$$U = \bigcup \mathcal{B}.$$

Definition 1.2.3 (synthetic basis). Let (X, \mathcal{T}) be a topological space, and let \mathcal{B} be a base of X . \mathcal{B} is said to be *synthetic* iff for any $A, B \in \mathcal{B}$,

$$A \cap B = \bigcup_{i=1}^n B_i, \quad B_i \in \mathcal{B}.$$

1.3 Continuity

Definition 1.3.1 (continuous maps). Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A map $f : X \rightarrow Y$ is said to be *continuous* iff for any open set U in Y , its preimage in X under f is open.

Proposition 1.3.1. Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A map $f : X \rightarrow Y$ is continuous at $x \in X$ iff for any neighbourhood N_y of $f(x)$, there is a neighbourhood N_x of x , such that $f[N_x] \subseteq N_y$.

Proof. Let N_y be a neighbourhood of $f(x)$. Clearly, there exists an open set U_y contains y .

By Definition 1.3.1, f is continuous at x iff $x \in f^{-1}[U_y] \in \mathcal{T}_X$. Clearly, $f^{-1}[U_y]$ is a neighbourhood of x . We have $f[f^{-1}[U_y]] = U_y \subseteq N_y$.

By Proposition 1.1.2, there U_x must contains at least one neighbourhood N_x of x , thus, $f[N_x] \subseteq U_y$. \square

Proposition 1.3.2. Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be metrizable spaces. A map $f : X \rightarrow Y$ is continuous at $p \in X$ iff for any $\varepsilon > 0$, there is a $\delta > 0$, such that for all $x \in B_X(p, \delta)$, $f(x) \in B_Y(f(p), \varepsilon)$, where B_X is defined by any metrics ρ_X induces \mathcal{T}_X , and B_Y is defined by any metrics ρ_Y induces \mathcal{T}_Y .

Proof. Clearly, for all $\varepsilon > 0$, $B_Y(f(x), \varepsilon)$ is an open neighbourhood of $f(x)$.

f is not necessarily be injective, so $f^{-1}[B_Y(f(x), \varepsilon)] = U \in x$. By Definition 1.3.1, U is open, so for some $\delta > 0$, $B_X(x, \delta) \subseteq U$. Thus, By Proposition 1.3.1, f is continuous iff $f[B_X(x, \delta)] \subseteq B_Y(f(x), \varepsilon)$. This satisfies the conditions we have. \square

Proposition 1.3.3. Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A function $f : X \rightarrow Y$ is said to be continuous iff for any closed set V in Y , its preimage in X under f is closed.

Proof. Let U_Y be any open set in Y , let U_X be the preimage of U_Y under f . By Definition 1.3.1, U_X is open in X . Let

$$V_X = f^{-1}[Y \setminus U_Y] = X \setminus U_X,$$

Then V_X is closed. □

1.4 Interiors and Closures

Definition 1.4.1 (interiors). The *interior* of a set A , denoted A° , is defined to be the union of all open subsets of A .

Definition 1.4.2 (closure). The *closure* of a set A , denoted \overline{A} , is defined to be the intersection of all closed supersets of A .

Definition 1.4.3 (dense sets). Let (X, \mathcal{T}) be a topological space, and let $A \subseteq X$. A is said to be *dense*, iff $\overline{A} = X$.

Definition 1.4.4 (nowhere dense sets). A set A is said to be *nowhere dense* iff the interior of its closure is empty.

Proposition 1.4.1 (properties of interiors). Let (X, \mathcal{T}) be any topological space and $A, B \subseteq X$.

- (i) (Intensive) $A^\circ \subseteq A$.
- (ii) A is open iff $A = A^\circ$.
- (iii) (Idempotence) $(A^\circ)^\circ = A^\circ$.
- (iv) $(A \cap B)^\circ = A^\circ \cap B^\circ$.
- (v) $A \subseteq B \implies A^\circ \subseteq B^\circ$.
- (vi) If B is open, then $B \subseteq A$ iff $B \subseteq A^\circ$.

Proof.

- (i) By Definition 1.4.1, naturally, $A^\circ \subseteq A$.
- (ii) By Definition 1.1.2, A° is the union of open sets hence it is open. A is open iff it is the union of all open subsets of A . Thus $A = A^\circ$.
- (iii) A° is open, thus $(A^\circ)^\circ = A^\circ$.

(iv) By Definition 1.4.1, we have

$$\begin{aligned}
(A \cap B)^\circ &= \left\{ \bigcup U : U \in \mathcal{T} \wedge U \subseteq A \cap B \right\} \\
&= \left\{ \bigcup U : (U \in \mathcal{T} \wedge U \subseteq A) \wedge (U \in \mathcal{T} \wedge U \subseteq B) \right\} \\
&= \left\{ \bigcup U : U \in \mathcal{T} \wedge U \subseteq A \right\} \cap \left\{ \bigcup U : U \in \mathcal{T} \wedge U \subseteq B \right\} \\
&= A^\circ \cap B^\circ.
\end{aligned}$$

(v) Clearly, $A^\circ \subseteq A$, thus,

$$A \subseteq B \implies A^\circ \subseteq B^\circ$$

Suppose $A^\circ \not\subseteq B^\circ$, then $A^\circ \setminus B^\circ$ is not empty (\emptyset is the subset of any set, so A° is not empty).

Then there exists $x \in A^\circ$ with $x \in \partial B$ ($x \in B$ but $x \notin B^\circ$). Then there exists neighbourhood $N_x \ni x$, and $N_x \cap \partial B \neq \emptyset$. But this is impossible, for $A^\circ \subseteq B$ implies that $A^\circ \cap \partial B = \emptyset$ (This is a straight consequence of $A^\circ \cap \partial A = \emptyset$. See Proposition 1.5.1), so such N_x does not exist. Thus,

$$A^\circ \subseteq B^\circ.$$

(vi) If B is open, then $B = B^\circ$. Then $B \subseteq A$ iff $B^\circ \subseteq A^\circ$.

□

Proposition 1.4.2 (properties of closures). Let (X, \mathcal{T}) be a topological space, and let $A, B \subseteq X$.

- (i) \overline{A} is closed.
- (ii) A is closed iff $A = \overline{A}$.
- (iii) $A \subseteq B$ implies $\overline{A} \subseteq \overline{B}$.
- (iv) If A is closed, then $A \supseteq B$ iff $A \supseteq \overline{B}$

Proof.

- (i) By Definition 1.4.2, \overline{A} is the intersection of closed sets. By Proposition 1.1.1, \overline{A} is closed.
- (ii) Proposition 1.1.1 implies that any closed set is the intersection of closed sets, this is precisely what Definition 1.4.2 says.

(iii) $A \subseteq B$ iff $X \setminus A \supseteq X \setminus B$. Then we have

$$X \setminus (X \setminus A)^\circ \subseteq X \setminus (X \setminus B)^\circ$$

Clearly, $(X \setminus A)^\circ$ is the union of all open set disjoint from A , then, by De Morgan's laws, $X \setminus (X \setminus A)^\circ$ is the intersection of all closed sets containing A . By Definition 1.4.2, we have $(X \setminus A)^\circ = \overline{A}$. Thus

$$\overline{A} \subseteq \overline{B}.$$

(iv) If A is closed, then $A = \overline{A}$. Suppose $B \subseteq A$, then we have

$$\overline{B} \subseteq \overline{A} \iff \overline{B} \subseteq A.$$

□

1.5 Boundaries

Definition 1.5.1 (boundaries). Let A be any set, the *boundary* of A , denoted ∂A , is defined to be the complement of the interior of A in the closure of A ; i.e.,

$$\partial A = \overline{A} \setminus A^\circ.$$

Proposition 1.5.1 (properties of boundaries). Let (X, \mathcal{T}) be a topological space, and let $A \subseteq X$.

- (i) ∂A is closed.
- (ii) $A^\circ \cap \partial A = \emptyset$.
- (iii) $\overline{A} = A^\circ \cup \partial A$.
- (iv) A is closed iff $\partial A \subseteq A$.
- (v) ∂A is nowhere dense.
- (vi) $\partial \overline{A} \subseteq \partial A \subseteq \partial A^\circ$.
- (vii) $\partial A = \partial(X \setminus A)$.
- (viii) A is dense iff $\partial A = X \setminus A^\circ$.

Proof.

(i) \overline{A} is closed, and $X \setminus A^\circ$ is also closed. Thus

$$\partial A = \overline{A} \setminus A^\circ = \overline{A} \cap (X \setminus A)$$

is closed.

(ii) By Definition 1.5.1, we have

$$\partial A = \overline{A} \setminus A^\circ \iff \partial A \cap A^\circ = \overline{A} \setminus A^\circ \cap A^\circ = \overline{A} \cap \emptyset = \emptyset.$$

(iii) We have

$$\begin{aligned} \partial A = \overline{A} \setminus A^\circ &\iff \partial A \cup A^\circ = \overline{A} \setminus A^\circ \cup A^\circ = \overline{A} \cap (X \setminus A^\circ \cup A^\circ) \\ &\iff \partial A \cup A^\circ = \overline{A} \cap X|_{\text{for } A^\circ} \subseteq X = \overline{A}. \end{aligned}$$

(iv) As A is closed, $A = \overline{A}$ (this can be straightly proved by Definition 1.4.2).

By Definition 1.5.1, it is clear that $\partial A \subseteq \overline{A}$, thus $\partial A \subseteq A$.

(v) By Definition 1.4.4, ∂A is nowhere dense iff $\overline{\partial A}^\circ$ is empty. We have

$$\begin{aligned} \overline{\partial A}^\circ &= \overline{\overline{A} \setminus A^\circ}^\circ \\ &= (\overline{A} \setminus A^\circ) \cup (\overline{A} \setminus A^\circ) \setminus (\overline{A} \setminus A^\circ) \\ &= \emptyset. \end{aligned}$$

(vi) $\overline{A} \supseteq A^\circ$ implies $\overline{A}^\circ \supseteq (A^\circ)^\circ = A^\circ$, then we have,

$$\partial \overline{A} = \overline{\overline{A}} \setminus \overline{A}^\circ \subseteq \overline{A} \setminus A^\circ = \partial A.$$

$A^\circ \subseteq A$ implies $\overline{A}^\circ \subseteq \overline{A}$, then we have,

$$\partial A^\circ = \overline{A^\circ} \setminus (A^\circ)^\circ \supseteq \overline{A} \setminus A^\circ.$$

(vii) We have

$$\begin{aligned} \partial(X \setminus A) &= \overline{X \setminus A} \setminus (X \setminus A)^\circ \\ &= X \setminus A^\circ \setminus (X \setminus \overline{A}) \\ &= X \setminus A^\circ \cap \overline{A} \\ &= \overline{A} \setminus A^\circ \\ &= \partial A. \end{aligned}$$

(viii) By Definition 1.4.3, A is dense in X iff $\overline{A} = X$. Then we have,

$$\begin{aligned} \overline{A} = X &\iff \overline{A} \setminus A^\circ = X \setminus A^\circ \\ &\iff \partial A = X \setminus A^\circ. \end{aligned}$$

□

1.6 Limit Points

Definition 1.6.1 (limit points). Let (X, \mathcal{T}_X) be a topological space, and let $A \subseteq X$. A point $x \in X$ is called a *limit point* of A iff for all neighbourhood N_x of x , $N_x \setminus \{x\}$ intersects A .

Proposition 1.6.1. Let A be any set, and let x be a limit point of A , then x is an element of the closure of A .

Proof. If A is empty, then this is vacuously true. So, suppose A is not empty.

By Definition 1.6.1, for all neighbourhood N_x of x , $N_x \setminus \{x\} \cap A$ is not empty. Naturally, $N_x \cap A$ is not empty.

Assume that $x \notin \overline{A}$, then $X \setminus \overline{A}$ is a neighbourhood of x , by Definition 1.1.5, and is disjoint from A . This is contradicted to the conditions. \square

Note 1.6.1. In this proof, the proposition also holds for $N_x \cap A^\circ = \emptyset$. Because if it is true, then

$$N_x \cap \partial A \supseteq (N_x \cap A) \setminus (N_x \cap A^\circ) = N_x \cap A.$$

This implies that $A \subseteq \partial A$. In this case, $\overline{A} = \partial A$, for

Assume that $x \notin \partial A$, then we have the same conclusion.

Then $A^\circ = A \setminus \partial A = \emptyset$.

Proposition 1.6.2. A set is closed iff it contains all its limit point.

Proof. Let A be a set. By proposition 1.6.1, for every limit point of A , it is also an element of the closure \overline{A} . And A is closed iff $A = \overline{A}$. \square

Definition 1.6.2 (convergent sequences). Let (X, \mathcal{T}_X) be a topological space. A sequence $\{x_n\}$ in X is said to be *convergence* in X iff there is an open set U contains all but finite terms of $\{x_n\}$.

1.7 Subspaces

Definition 1.7.1 (subspace topology). Let (X, \mathcal{T}) be a topological space and let $A \subseteq X$. The *subspace topology* \mathcal{T}_A on A is defined to be the family of the intersections of open sets in (X, \mathcal{T}) and A . That is,

$$\mathcal{T}_A = \{U \cap A : U \in \mathcal{T}\}.$$

Chapter 2

Creating New Spaces

2.1 Quotient Spaces

Definition 2.1.1 (quotient topology). Let (X, \mathcal{T}) be a topological space and let \sim be an equivalence relation on X . The *quotient topology* is a topology on $\mathcal{P}(X/\sim)$; it is defined as

$$\mathcal{T}_{X/\sim} = \{U \in \mathcal{P}(X/\sim) : \{x \in X : [x] \in U\} \in \mathcal{T}_X\}.$$

2.2 Product Spaces

Definition 2.2.1 (product topologies).

2.3 Homeomorphisms

Definition 2.3.1 (homeomorphisms). Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A bijection $f : X \rightarrow Y$ is called a *homeomorphism* iff it is continuous and its inverse is also continuous.

Definition 2.3.2 (homeomorphic). Two topological spaces (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) are said to be *homeomorphic* or *topologically equivalent*, denoted $X \cong Y$, iff there is an homeomorphism between them.

Chapter 3

Topological Properties

3.1 Cardinal Functions

3.2 Separation Axioms

Definition 3.2.1 (topologically indistinguishable). Let (X, \mathcal{T}) be a topological space. Two points $x, y \in X$ are said to be *topologically indistinguishable* iff they share all their neighbourhoods. That is, let \mathcal{N}_x be the family of all neighbourhoods of x and let \mathcal{N}_y be the family of all neighbourhoods of y , we have

$$\mathcal{N}_x = \mathcal{N}_y.$$

Respectively, x, y are said to be *topologically distinguishable* iff they are not topologically indistinguishable; i.e.,

$$\mathcal{N}_x \neq \mathcal{N}_y.$$

Definition 3.2.2 (separated sets). Let (X, \mathcal{T}) be a topological space, and let $A, B \in \mathcal{P}(X)$.

- (i) A and B are said to be *separated* iff each is disjoint from other's closure.
- (ii) A and B are said to be *separated by neighbourhoods* iff there are neighbourhoods N_A of A and N_B of B such that N_A and N_B are disjoint.
- (iii) A and B are said to be *separated by closed neighbourhoods* iff there are closed neighbourhoods \overline{N}_A of A and \overline{N}_B of B such that \overline{N}_A and \overline{N}_B are disjoint.

- (iv) A and B are said to be *separated by a continuous function* iff there is a continuous function $f : X \rightarrow \mathbb{R}$, such that $f[A] = \{0\}$ and $f[B] = \{1\}$.
- (v) A and B are said to be *precisely separated by a continuous function* iff there is a continuous function $f : X \rightarrow \mathbb{R}$, such that $f^{-1}[\{0\}] = A$ and $f^{-1}[\{1\}] = B$

Definition 3.2.3 (T_0 spaces). A topological space (X, \mathcal{T}) is said to be T_0 or *Kolmogorov*, iff all distinct points $x, y \in X$ are *topologically distinguishable*.

Example 3.2.1 (non- T_0 sets). The a set X with the discrete topology is T_0 iff $|X| \in \{0, 1\}$ (vacuously true).

Definition 3.2.4 (R_0 spaces). A topological space (X, \mathcal{T}) is said to be R_0 iff any two topologically distinguishable points in X are separated.

Example 3.2.2 (R_0 but not T_0). Let X be any set, let $U \subsetneq X$ with $|X \setminus U| > 1$, let $V \subsetneq U$ with $|V| = 1$, and let

$$\mathcal{T} = \mathcal{P}(X) \setminus \mathcal{P}(X \setminus U) \setminus \mathcal{P}(V) \cup \{\emptyset, X\}.$$

For any two distinct points $x, y \in U$, the family \mathcal{N}_x of neighbourhoods of x and the family \mathcal{N}_y of neighbourhoods of y are different; and for all such x and y , $x \notin \overline{\{y\}} = \{y\}$ (i.e., they are separated; but, be caution, they are not necessarily be separated by neighbourhoods; for if $y \in V$, the smallest neighbourhood of y is X). Thus (X, \mathcal{T}) is R_0 . But X is not T_0 , because for two distinct points $x, y \in X \setminus U \cup V$, the families of their neighbourhoods are the same.

Example 3.2.3 (T_0 but not R_0). Let $(\mathbb{R}_{\geq 0}, \mathcal{T})$ be a topological space with

$$\mathcal{T} = \{U \subseteq \mathbb{R} : \forall i \in \mathbb{R}_{\geq 0}, U_i = [0, i)\},$$

Then for all $x, y \in (\mathbb{R}_{\geq 0}, \mathcal{T})$, if $x \neq y$, then there are $|y - x|$ neighbourhoods N_x of x do not contain y . Thus, it is T_0 .

On the other hand, it is not R_0 , because for all $x, y \in (\mathbb{R}_{\geq 0}, \mathcal{T})$ with $x < y$, $x \in [0, y]$

Proposition 3.2.1 (alternative definitions of R_0 spaces). Let (X, \mathcal{T}) be R_0 , then the following conditions are equivalent.

- (i) The closure of all singletons in X are not T_0 subspace.
- (ii) For any two points $x, y \in X$, $x \in \overline{\{y\}}$ iff $y \in \overline{\{x\}}$.

(iii) Every open set is the union of closed sets.

Proof.

- (i) By Definition 3.2.4, if y and x are topologically distinguishable, by Definition 3.2.4, x and y are separated; i.e., $x \notin \overline{\{y\}}$ and $y \notin \overline{\{x\}}$.
- (ii) By Definition 3.2.4, for all $x, y \in X$, x, y are not separated only if they are topologically indistinguishable. By Definition 3.2.1, they share all their neighbourhoods, thus they have the same closure; i.e., $\overline{\{x\}} = \overline{\{y\}}$.
- (iii) (Remained as a problem!)

□

Definition 3.2.5 (T_1 spaces). A topological space (X, \mathcal{T}) is said to be T_1 or *Fréchet* iff it is T_0 and R_0 .

Proposition 3.2.2 (alternative definitions of T_1 spaces). Let (X, \mathcal{T}) be T_1 , then the following conditions are equivalent.

- (i) All singletons in X are closed.
- (ii) Every subset of X is the intersection of all open sets containing it.
- (iii) Every cofinite subset of X is open.

Proof.

□

Proposition 3.2.3. All singletons in a T_1 space are closed, That is, if a topological space (X, \mathcal{T}) is T_1 , then

$$\forall x \in (X, \mathcal{T}) : \exists U \in \mathcal{T} : \{x\} = X \setminus U.$$

Definition 3.2.6 (R_1 spaces). A topological space (X, \mathcal{T}) is said to be R_1 iff any two topologically distinguishable points in X are separated by neighbourhoods.

Example 3.2.4 (R_0 but not R_1). (Remained as a problem!)

Definition 3.2.7 (T_2 spaces). A topological space (X, \mathcal{T}) is said to be T_2 or *Hausdorff* or *separated* iff any two distinct points in (X, \mathcal{T}) are separated by neighbourhoods.

Example 3.2.5 (T_2 but not T_1). Let X be a nonempty set, and let $\mathcal{U} = \mathcal{P}(X \setminus \{x \in X\})$. Then the topological space (X, \mathcal{T}) with

$$\mathcal{T} = \mathcal{U} \cup \{X\}$$

is T_1 . But it is T_2 iff $|X| = 1$ (This is vacuously true). As $|X| > 1$, $\{x\}$ is not open.

Proposition 3.2.4. All metric spaces are Hausdorff.

Definition 3.2.8 ($T_{2^{1/2}}$ spaces). A topological space (X, \mathcal{T}) is said to be $T_{2^{1/2}}$ or *Urysohn* iff two distinct points in X are separated by closed neighbourhoods.

Example 3.2.6 (T_2 but not $T_{2^{1/2}}$).¹ (Remained as a problem)

Definition 3.2.9 (T_3 spaces). A topological space (X, \mathcal{T}) is said to be T_3 or *regular* iff it is T_0 and given any point $x \in (X, \mathcal{T})$ and closed set $V \subseteq X$ with $x \notin V$ are separated by neighbourhoods.

Definition 3.2.10 ($T_{3^{1/2}}$ spaces). A topological space (X, \mathcal{T}) is said to be $T_{3^{1/2}}$, or *Tychonoff* or, *completely T_3* , or *completely regular*, iff it is T_0 and given any point x and closed set $V \subseteq X$ with $x \notin V$, they are separated by a continuous function.

Definition 3.2.11 (T_4 spaces). A topological space (X, \mathcal{T}) is said to be T_4 or *normal* iff it is Hausdorff and any two disjoint closed subsets of X are separated by neighbourhoods.

Proposition 3.2.5 (Urysohn's lemma). A topological space is normal iff any two disjoint closed sets are separated by a continuous function.

Definition 3.2.12 (T_5 spaces). A topological space (X, \mathcal{T}) is said to be T_5 or *completely T_4* iff it is T_1 any two separated sets are separated by neighbourhoods.

Proposition 3.2.6. Every subspace of a T_5 space is normal.

Definition 3.2.13 (T_6 spaces). A topological space (X, \mathcal{T}) is said to be T_6 , or *perfectly T_4* or *perfectly normal* iff it is T_1 and any two disjoint closed sets are precisely separated by a continuous function.

Proposition 3.2.7 (Tietze extension theorem). Let (X, \mathcal{T}) be normal topological space, and let $f : A \rightarrow (\mathbb{R}, \mathcal{T}')$ be a continuous map where A is a closed

¹ See [MathPlanet](#).

subset of X and \mathcal{T}' is the standard topology (induced by Euclidean metric). Then there exists a continuous map

$$F : (X, \mathcal{T}) \rightarrow (\mathbb{R}, \mathcal{T}'),$$

such that

$$\forall x \in A : f(x) = g(x).$$

3.3 Countability Axioms

3.4 Metrizable

3.5 Compactness

Definition 3.5.1 (compactness). A topological space (X, \mathcal{T}) is said to be *compact* iff every open cover of X has a finite subcover. That is,

$$\forall \mathcal{C} \subseteq \mathcal{T} : \bigcup \mathcal{C} = X : \exists \mathcal{S} \subseteq \mathcal{C} : \bigcup \mathcal{S} = X : |\mathcal{S}| < \aleph_0.$$

3.6 Connectedness

Definition 3.6.1 (connectedness). Let (X, \mathcal{T}) be a topological space. (X, \mathcal{T}) is said to be *connected* iff X is not empty and it is not the union of any disjoint open sets. That is,

$$\forall U, V \in \mathcal{T} : X = U \cup V : U \cap V \neq \emptyset.$$

Definition 3.6.2 (path-connectedness). Let (X, \mathcal{T}) be a topological space.

- (i) A map $\gamma : [0, 1] \rightarrow X$ is called a *path* in X iff it is continuous. If $\gamma(0) = x$ and $\gamma(1) = y$, we say that γ is path from x to y in X .
- (ii) X is said to be *path-connected* iff for all $x, y \in X$ there is a path from x to y in X .