The Large Deviation Principle (LDP)

Graph Limit Theory

LDP for Random Graphs

Questions & Answers

## Large Deviations for Random Graphs Honours Year Project Introductory Talk

NG Say-Yao Mark<sup>1</sup> A/P SUN Rongfeng<sup>2</sup>

<sup>1</sup>Student, <sup>2</sup>Supervisor Department of Mathematics National University of Singapore (NUS)

October 5, 2016

LDP for Random Graphs

Questions & Answers

- 1 Introduction
- 2 The Large Deviation Principle (LDP)
- **3** Graph Limit Theory
- 4 LDP for Random Graphs
- **6** Questions & Answers

The Large Deviation Principle (LDP)

Graph Lim Theory

LDP for Random Graphs

Questions &

The Large Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions &

#### General information about this talk.

Introduce fundamentals and motivation for chosen topic.

Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions &

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory

The Large Deviation Principle (LDP)

Graph Limit Theory

LDP for Random Graphs

Questions &

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.

Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions &

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.
- Assuming some knowledge in:

The Large Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions & Answers

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.
- Assuming some knowledge in:
  - Probability theory: probability measure, expected value, almost sure convergence.

The Large Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions &

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.
- Assuming some knowledge in:
  - Probability theory: probability measure, expected value, almost sure convergence.
  - Analysis: *limits, metric spaces, measurable functions.*

The Large Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions & Answers

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.
- Assuming some knowledge in:
  - Probability theory: probability measure, expected value, almost sure convergence.
  - Analysis: *limits, metric spaces, measurable functions.*
  - Graph Theory: edge, vertex, graph homomorphisms.

General information about this talk.

- Introduce fundamentals and motivation for chosen topic.
- Large deviation theory for random graph models.
- Assuming some knowledge in:
  - Probability theory: probability measure, expected value, almost sure convergence.
  - Analysis: limits, metric spaces, measurable functions.
  - Graph Theory: edge, vertex, graph homomorphisms.
- Slides available at:

http://mollymr305.github.io

LDP for Random

Questions &

## Large Deviations Theory

Let's begin with soft introduction to large deviations theory.

LDP for Random Graphs

Questions &

## Large Deviations Theory

Let's begin with *soft* introduction to large deviations theory.

Large deviation theory is a branch of probability theory which, roughly speaking, concerns the study of two things:

Questions &

## Large Deviations Theory

Let's begin with *soft* introduction to large deviations theory.

Large deviation theory is a branch of probability theory which, roughly speaking, concerns the study of two things:

Probabilities of rare events.

Questions & Answers

## Large Deviations Theory

Let's begin with *soft* introduction to large deviations theory.

Large deviation theory is a branch of probability theory which, roughly speaking, concerns the study of two things:

- Probabilities of rare events.
- Conditional probabilities of events, given that a rare event has occurred.

Graph Limi

LDP for Random Graphs

Questions & Answers

## Large Deviations Theory

Let's begin with *soft* introduction to large deviations theory.

Large deviation theory is a branch of probability theory which, roughly speaking, concerns the study of two things:

- Probabilities of rare events.
- Conditional probabilities of events, given that a rare event has occurred.

The theory of large deviations is also applied in many fields such as statistical physics, statistical hypothesis testing, etc.

LDP for Random Graphs

Questions &

#### A Coin Tossing Example

Let  $X_1, \ldots, X_n$  be IID random variables with law

$$P(X_i = 0) = P(X_i = 1) = \frac{1}{2}.$$

Questions &

#### A Coin Tossing Example

Let  $X_1, \ldots, X_n$  be IID random variables with law

$$P(X_i = 0) = P(X_i = 1) = \frac{1}{2}.$$

Denote their sum by  $S_n = X_1 + \cdots + X_n$ .

Questions &

#### A Coin Tossing Example

Let  $X_1, \ldots, X_n$  be IID random variables with law

$$P(X_i = 0) = P(X_i = 1) = \frac{1}{2}.$$

Denote their sum by  $S_n = X_1 + \cdots + X_n$ .

The Strong Law of Large Numbers (LLN) tells us that

$$\frac{S_n}{n} \xrightarrow{a.s.} \frac{1}{2}$$
, as  $n \longrightarrow \infty$ .

Questions &

#### A Coin Tossing Example

Let  $X_1, \ldots, X_n$  be IID random variables with law

$$P(X_i = 0) = P(X_i = 1) = \frac{1}{2}.$$

Denote their sum by  $S_n = X_1 + \cdots + X_n$ .

The Strong Law of Large Numbers (LLN) tells us that

$$\frac{S_n}{n} \xrightarrow{a.s.} \frac{1}{2}$$
, as  $n \longrightarrow \infty$ .

i.e. "The *empirical mean* converges to  $\frac{1}{2}$  with probability 1."

LDP for Random Graphs

Questions &

#### A Coin Tossing Example

We can then ask about the *asymptotic behaviour* of the probabilities rare events, for example

$$\mathbf{P}\left(\frac{S_n}{n} \ge \frac{2}{3}\right)$$

LDP for Random Graphs

Questions &

## A Coin Tossing Example

We can then ask about the *asymptotic behaviour* of the probabilities rare events, for example

$$\mathbf{P}\left(\frac{S_n}{n} \ge \frac{2}{3}\right) = \mathbf{P}\left(S_n \ge \frac{2}{3}n\right)$$

LDP for Random Graphs

Questions & Answers

## A Coin Tossing Example

We can then ask about the *asymptotic behaviour* of the probabilities rare events, for example

$$\mathbf{P}\left(\frac{S_n}{n} \geq \frac{2}{3}\right) = \mathbf{P}\left(S_n \geq \frac{2}{3}n\right), \text{ as } n \longrightarrow \infty.$$

LDP for Random Graphs

Questions &

## A Coin Tossing Example

It turns out that,

$$\lim_{n\to\infty}\frac{1}{n}\log\mathbf{P}\left(S_n\geq\frac{2}{3}n\right)$$

LDP for Random Graphs

Questions &

#### A Coin Tossing Example

It turns out that,

$$\lim_{n\to\infty}\frac{1}{n}\log\mathbf{P}\left(S_n\geq\frac{2}{3}n\right)\,=-\left(\log2+\frac{2}{3}\log\frac{2}{3}-\frac{1}{3}\log\frac{1}{3}\right)$$

LDP for Random Graphs

Questions &

#### A Coin Tossing Example

It turns out that,

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbf{P} \left( S_n \ge \frac{2}{3} n \right) = -\left( \log 2 + \frac{2}{3} \log \frac{2}{3} - \frac{1}{3} \log \frac{1}{3} \right)$$
$$= -\log \left( \frac{2^{5/3}}{3} \right).$$

(LDP)

LDP for Random Graphs

Questions &

#### A Coin Tossing Example

It turns out that,

$$\begin{split} \lim_{n\to\infty} \frac{1}{n} \log \mathbf{P} \left( S_n \geq \frac{2}{3} n \right) &= -\left( \log 2 + \frac{2}{3} \log \frac{2}{3} - \frac{1}{3} \log \frac{1}{3} \right) \\ &= -\log \left( \frac{2^{5/3}}{3} \right). \end{split}$$

In other words: the probability of the rare event

Questions &

#### A Coin Tossing Example

It turns out that,

$$\begin{split} \lim_{n\to\infty} \frac{1}{n} \log \mathbf{P} \left( S_n \geq \frac{2}{3} n \right) &= -\left( \log 2 + \frac{2}{3} \log \frac{2}{3} - \frac{1}{3} \log \frac{1}{3} \right) \\ &= -\log \left( \frac{2^{5/3}}{3} \right). \end{split}$$

In other words: the probability of the rare event

$$\mathbf{P}\left(S_n\geq \frac{2}{3}n\right),\,$$

It turns out that,

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbf{P} \left( S_n \ge \frac{2}{3} n \right) = -\left( \log 2 + \frac{2}{3} \log \frac{2}{3} - \frac{1}{3} \log \frac{1}{3} \right)$$
$$= -\log \left( \frac{2^{5/3}}{3} \right).$$

A Coin Tossing Example

In other words: the probability of the rare event

$$\mathbf{P}\left(S_n\geq\frac{2}{3}n\right),\,$$

decays exponentially as  $n \longrightarrow \infty$ , with rate

Questions &

#### A Coin Tossing Example

It turns out that,

$$\begin{split} \lim_{n\to\infty} \frac{1}{n} \log \mathbf{P} \left( S_n \geq \frac{2}{3} n \right) &= -\left( \log 2 + \frac{2}{3} \log \frac{2}{3} - \frac{1}{3} \log \frac{1}{3} \right) \\ &= -\log \left( \frac{2^{5/3}}{3} \right). \end{split}$$

In other words: the probability of the rare event

$$\mathbf{P}\left(S_n\geq\frac{2}{3}n\right),\,$$

decays exponentially as  $n \longrightarrow \infty$ , with rate

$$\frac{2^{5/3}}{3}$$

The Large Deviation Principle (LDP)

Graph Lin Theory

LDP for Random Graphs

Questions &

## A Coin Tossing Example

More generally, for any  $\alpha>\frac{1}{2}$  we have

LDP for Random Graphs

Questions & Answers

#### A Coin Tossing Example

More generally, for any  $\alpha>\frac{1}{2}$  we have

$$\lim_{n\to\infty} \mathbf{P}(S_n \geq \alpha n) = -I(\alpha),$$

where  $I: \mathbb{R} \to \overline{\mathbb{R}}$ , called the *rate function* is defined by

$$I(z) = egin{cases} \log 2 + z \log z + (1-z) \log (1-z) & \text{if } z \in [0,1], \\ +\infty & \text{otherwise.} \end{cases}$$

Questions & Answers

## A Coin Tossing Example

More generally, for any  $\alpha>\frac{1}{2}$  we have

$$\lim_{n\to\infty} \mathbf{P}(S_n \ge \alpha n) = -I(\alpha),$$

where  $I: \mathbb{R} \to \overline{\mathbb{R}}$ , called the *rate function* is defined by

$$I(z) = egin{cases} \log 2 + z \log z + (1-z) \log (1-z) & \text{if } z \in [0,1], \\ +\infty & \text{otherwise.} \end{cases}$$

Remark 1. We set  $0 \log 0 = 0$  by convention.

Questions &

#### A Coin Tossing Example

More generally, for any  $\alpha>\frac{1}{2}$  we have

$$\lim_{n\to\infty} \mathbf{P}(S_n \ge \alpha n) = -I(\alpha),$$

where  $I: \mathbb{R} \to \overline{\mathbb{R}}$ , called the *rate function* is defined by

$$I(z) = egin{cases} \log 2 + z \log z + (1-z) \log (1-z) & \text{ if } z \in [0,1], \\ +\infty & \text{ otherwise.} \end{cases}$$

Remark 1. We set  $0 \log 0 = 0$  by convention.

Remark 2. Some properties of this rate function include:

- $I(z) \geq 0$  for all  $z \in \mathbb{R}$ .
- I(z) = 0 if and only if  $z = \frac{1}{2}$ .

LDP for Random Graphs

Questions &

# Cramer's Theorem for the Empirical Average

A further generalisation of the Coin Tossing Example.

Questions &

# Cramer's Theorem for the Empirical Average

A further generalisation of the Coin Tossing Example.

Theorem (Cramer)

Let  $\{X_i\}_{i\in\mathbb{N}}$  be IID  $\mathbb{R}$ -valued random variables such that

$$\varphi(t) = \mathbf{E}e^{tX_i} < \infty.$$

Let  $S_n = \sum_{i=1}^n X_i$ . Then for any  $\alpha > \mathbf{E}X_i$ ,

$$\lim_{n\to\infty}\frac{1}{n}\log\mathbf{P}\left(S_{n}\geq\alpha n\right)=-I(\alpha),$$

where

$$I(z) = \sup_{t \in \mathbb{R}} \{zt - \log \varphi(t)\}.$$

LDP for Random

Questions &

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

LDP for Random Graphs

Questions &

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

Definition (Good Rate Function)

A function  $I: \mathcal{X} \to [0, \infty]$  is said to be a *good rate function* if

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

Let  $(\mathcal{X},d)$  be a Polish space (i.e. separable and complete).

Definition (Good Rate Function)

A function  $I: \mathcal{X} \to [0, \infty]$  is said to be a *good rate function* if D1  $I \not\equiv \infty$ .

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

Definition (Good Rate Function)

A function  $I: \mathcal{X} \to [0, \infty]$  is said to be a *good rate function* if D1  $I \not\equiv \infty$ .

D2 / is lower semi-continuous.

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the *general* large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

Definition (Good Rate Function)

A function  $I: \mathcal{X} \to [0, \infty]$  is said to be a *good rate function* if D1  $I \not\equiv \infty$ .

D2 I is lower semi-continuous. (iff  $\liminf_{n\to\infty} I(x_n) \ge I(x)$  for any  $x_n \to x$  in  $\mathcal{X}$ .)

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the  $\emph{general}$  large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

## Definition (Good Rate Function)

A function  $I:\mathcal{X}\to [0,\infty]$  is said to be a *good rate function* if

D1  $I \not\equiv \infty$ .

D2 I is lower semi-continuous. (iff  $\liminf_{n\to\infty} I(x_n) \ge I(x)$  for any  $x_n \to x$  in  $\mathcal{X}$ .)

D3 I has compact level sets.

LDP for Random Graphs

Questions &

## The Large Deviation Principle

 $\star$  Starting point for the  $\emph{general}$  large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

## Definition (Good Rate Function)

A function  $I:\mathcal{X}\to [0,\infty]$  is said to be a *good rate function* if

- D1  $I \not\equiv \infty$ .
- D2 I is lower semi-continuous. (iff  $\liminf_{n\to\infty} I(x_n) \ge I(x)$  for any  $x_n \to x$  in  $\mathcal{X}$ .)
- D3 I has compact level sets. (i.e.  $I^{-1}([-\infty, c])$  is compact for every  $c \in \mathbb{R}$ )

LDP for Random Graphs

Questions & Answers

## The Large Deviation Principle

 $\star$  Starting point for the  $\emph{general}$  large deviations theory.

Let  $(\mathcal{X}, d)$  be a Polish space (i.e. separable and complete).

## Definition (Good Rate Function)

A function  $I:\mathcal{X}\to [0,\infty]$  is said to be a *good rate function* if

- D1  $I \not\equiv \infty$ .
- D2 I is lower semi-continuous. (iff  $\liminf_{n\to\infty} I(x_n) \ge I(x)$  for any  $x_n \to x$  in  $\mathcal{X}$ .)
- D3 I has compact level sets. (i.e.  $I^{-1}([-\infty, c])$  is compact for every  $c \in \mathbb{R}$ )

For any subset  $S \subseteq \mathcal{X}$  we also define  $I(S) = \inf\{I(s) : s \in S\}$ .

Graph Limit

LDP for Random Graphs

Questions &

## The Large Deviation Principle

## Definition (LDP)

A sequence of probability measures  $\{\mathbf{P}_n\}_{n\in\mathbb{N}}$  on the space  $\mathcal{X}$  is said to satisfy the Large Deviation Principle (LDP) with rate n and good rate function I if

LDP for Random Graphs

Questions &

## The Large Deviation Principle

## Definition (LDP)

A sequence of probability measures  $\{\mathbf{P}_n\}_{n\in\mathbb{N}}$  on the space  $\mathcal{X}$  is said to satisfy the Large Deviation Principle (LDP) with rate n and good rate function I if

L1 *I* is a good rate function.

LDP for Random Graphs

Questions &

## The Large Deviation Principle

## Definition (LDP)

A sequence of probability measures  $\{\mathbf{P}_n\}_{n\in\mathbb{N}}$  on the space  $\mathcal{X}$  is said to satisfy the Large Deviation Principle (LDP) with rate n and good rate function I if

- L1 *I* is a good rate function.
- L2 For any *closed* subset  $C \subseteq \mathcal{X}$ ,

$$\limsup_{n\to\infty}\frac{1}{n}\log\mathbf{P}_n(C)\leq -I(C).$$

Graph Limit

LDP for Random Graphs

Questions &

## The Large Deviation Principle

## Definition (LDP)

A sequence of probability measures  $\{\mathbf{P}_n\}_{n\in\mathbb{N}}$  on the space  $\mathcal{X}$  is said to satisfy the Large Deviation Principle (LDP) with rate n and good rate function I if

- L1 *I* is a good rate function.
- L2 For any *closed* subset  $C \subseteq \mathcal{X}$ ,

$$\limsup_{n\to\infty}\frac{1}{n}\log\mathbf{P}_n(C)\leq -I(C).$$

L3 For any *open* subset  $O \subseteq \mathcal{X}$ ,

$$\liminf_{n\to\infty}\frac{1}{n}\log \mathbf{P}_n(O)\geq -I(O).$$

LDP for Random

Questions &

## Graph Limit Theory

We first define what is means for a graph sequences to have a limit.

LDP for Random

Questions &

## Graph Limit Theory

We first define what is means for a graph sequences to have a limit. For a graph G, let V(G) and E(G) denote the vertex and edge set respectively.

LDP for Random Graphs

Questions &

## Graph Limit Theory

We first define what is means for a graph sequences to have a limit. For a graph G, let V(G) and E(G) denote the vertex and edge set respectively.

• Let  $\{G_n\}_{n\in\mathbb{N}}$  be a sequence of simple graphs whose number of vertices tend to infinity.

LDP for Random Graphs

Questions & Answers

## Graph Limit Theory

We first define what is means for a graph sequences to have a limit. For a graph G, let V(G) and E(G) denote the vertex and edge set respectively.

- Let  $\{G_n\}_{n\in\mathbb{N}}$  be a sequence of simple graphs whose number of vertices tend to infinity.
- For a fixed simple graph H, let hom $(H, G_n)$  denote the number of homomorphisms from H into  $G_n$ .

LDP for Random Graphs

Questions & Answers

## **Graph Limit Theory**

We first define what is means for a graph sequences to have a limit. For a graph G, let V(G) and E(G) denote the vertex and edge set respectively.

- Let  $\{G_n\}_{n\in\mathbb{N}}$  be a sequence of simple graphs whose number of vertices tend to infinity.
- For a fixed simple graph H, let hom $(H, G_n)$  denote the number of homomorphisms from H into  $G_n$ .
- The probability that a random map  $\phi:V(H)\to V(G_n)$  is a homomorphism is

$$t(H, G_n) = \frac{\mathsf{hom}(H, G_n)}{|V(G_n)|^{|V(H)|}}.$$

LDP for Random Graphs

Questions & Answers

## **Graph Limit Theory**

Lovász and Szegedy proved the following:

## **Theorem**

Suppose  $t(H, G_n)$  tends to a limit t(H) for every H. Then there exists a symmetric measurable function

$$f: [0,1] \times [0,1] \to [0,1]$$

such that

$$\lim_{n\to\infty} t(H,G_n) = t(H,f) \stackrel{\text{def.}}{=} \int_{[0,1]^k} \prod_{\{i,j\}\in E(H)} f(x_i,x_j) dx_1 \cdots dx_k.$$

Conversely, every symmetric measurable function of the form  $f:[0,1]\times[0,1]\to[0,1]$  corresponds to the limit of an appropriate graph sequence.

LDP for Random Graphs

Questions & Answers

## Graph Limit Theory

• The function f is the 'natural limit object' of  $\{G_n\}_{n\in\mathbb{N}}$ .

LDP for Random Graphs

Questions &

## Graph Limit Theory

- The function f is the 'natural limit object' of  $\{G_n\}_{n\in\mathbb{N}}$ .
- This gives us a 'working definition' for a 'graph limit'.

LDP for Random Graphs

Questions (

## **Graph Limit Theory**

- The function f is the 'natural limit object' of  $\{G_n\}_{n\in\mathbb{N}}$ .
- This gives us a 'working definition' for a 'graph limit'.

### Definition

A sequence of simple graphs  $\{G_n\}_{n\in\mathbb{N}}$  whose number of vertices tend to infinity is said to converge to a symmetric measurable function  $f:[0,1]\times[0,1]\to[0,1]$  if for every finite simple graph H,

$$\lim_{n\to\infty} t(H,G_n) = t(H,f).$$

LDP for Random Graphs

Questions & Answers

## **Graph Limit Theory**

- The function f is the 'natural limit object' of  $\{G_n\}_{n\in\mathbb{N}}$ .
- This gives us a 'working definition' for a 'graph limit'.

### Definition

A sequence of simple graphs  $\{G_n\}_{n\in\mathbb{N}}$  whose number of vertices tend to infinity is said to converge to a symmetric measurable function  $f:[0,1]\times[0,1]\to[0,1]$  if for every finite simple graph H,

$$\lim_{n\to\infty}t(H,G_n)=t(H,f).$$

The function f is also referred to as a graphon.

LDP for Random Graphs

Questions & Answers

## **Graph Limit Theory**

- The function f is the 'natural limit object' of  $\{G_n\}_{n\in\mathbb{N}}$ .
- This gives us a 'working definition'\* for a 'graph limit'.

### Definition

A sequence of simple graphs  $\{G_n\}_{n\in\mathbb{N}}$  whose number of vertices tend to infinity is said to converge to a symmetric measurable function  $f:[0,1]\times[0,1]\to[0,1]$  if for every finite simple graph H,

$$\lim_{n\to\infty}t(H,G_n)=t(H,f).$$

The function f is also referred to as a graphon.

LDP for Random

Questions &

## Erdős-Rényi Random Graphs

■ The Erdős-Rényi random graph  $G_{n,p}$  is a graph vertex set  $\{1,\ldots,n\}$ , and the law  $\mathbf{P}(\{i,j\})$  is an edge  $\mathbf{P}(\{i,j\})$  independently for all  $1 \leq i < j \leq n$ .

LDP for Random Graphs

Questions &

## Erdős-Rényi Random Graphs

- The Erdős-Rényi random graph  $G_{n,p}$  is a graph vertex set  $\{1,\ldots,n\}$ , and the law  $\mathbf{P}(\{i,j\})$  is an edge p independently for all  $1 \le i < j \le n$ .
- For a given H, note that  $t(H, G_{n,p})$  is a random variable.

LDP for Random Graphs

Questions &

## Erdős-Rényi Random Graphs

- The Erdős-Rényi random graph  $G_{n,p}$  is a graph vertex set  $\{1,\ldots,n\}$ , and the law  $\mathbf{P}(\{i,j\})$  is an edge p0 independently for all  $1 \le i < j \le n$ .
- For a given H, note that  $t(H, G_{n,p})$  is a random variable.
- It can be shown that

$$t(H, G_{n,p}) \xrightarrow{a.s.} p^{|E(H)|}, \text{ as } n \longrightarrow \infty.$$

LDP for Random Graphs

Questions & Answers

## Erdős-Rényi Random Graphs

- The Erdős-Rényi random graph  $G_{n,p}$  is a graph vertex set  $\{1,\ldots,n\}$ , and the law  $\mathbf{P}(\{i,j\})$  is an edge p0 independently for all  $1 \le i < j \le n$ .
- For a given H, note that  $t(H, G_{n,p})$  is a random variable.
- It can be shown that

$$t(H, G_{n,p}) \xrightarrow{a.s.} p^{|E(H)|}, \text{ as } n \longrightarrow \infty.$$

• Also, note that the constant graphon f(x, y) = p gives

$$t(H,f) = \int_{[0,1]^k} \prod_{\{i,j\} \in E(H)} f(x_i,x_j) dx_1 \cdots dx_k = \rho^{|E(H)|}.$$

LDP for Random Graphs

Questions & Answers

## Erdős-Rényi Random Graphs

- The Erdős-Rényi random graph  $G_{n,p}$  is a graph vertex set  $\{1,\ldots,n\}$ , and the law  $\mathbf{P}(\{i,j\})$  is an edge p0 independently for all  $1 \le i < j \le n$ .
- For a given H, note that  $t(H, G_{n,p})$  is a random variable.
- It can be shown that

$$t(H, G_{n,p}) \stackrel{a.s.}{\longrightarrow} p^{|E(H)|}, \text{ as } n \longrightarrow \infty.$$

• Also, note that the constant graphon f(x, y) = p gives

$$t(H,f) = \int_{[0,1]^k} \prod_{\{i,j\} \in E(H)} f(x_i, x_j) dx_1 \cdots dx_k = p^{|E(H)|}.$$

• In other words, the graph sequence  $\{G_{n,p}\}_{n\in\mathbb{N}}$  converges to  $f\equiv p$  with probability 1.

LDP for Random Graphs

Questions &

# Large Deviation Principle for Random Graphs: Research

General information about this field of research:

LDP for Random Graphs

Questions & Answers

# Large Deviation Principle for Random Graphs: Research

### General information about this field of research:

 Large deviation theory for the Erdős-Rényi random graph was developed not too long ago by Chatterjee & Varadhan (2011).

LDP for Random Graphs

Questions & Answers

# Large Deviation Principle for Random Graphs: Research

### General information about this field of research:

- Large deviation theory for the Erdős-Rényi random graph was developed not too long ago by Chatterjee & Varadhan (2011).
- Other developments include large deviation theory for the Exponential Random Graph Model (ERGM) by Chatterjee & Diaconis (2013).

LDP for Random Graphs

Questions & Answers

# Large Deviation Principle for Random Graphs: Research

### General information about this field of research:

- Large deviation theory for the Erdős-Rényi random graph was developed not too long ago by Chatterjee & Varadhan (2011).
- Other developments include large deviation theory for the Exponential Random Graph Model (ERGM) by Chatterjee & Diaconis (2013).
- However, less has been established with respect to sparse random graphs.

LDP for Random Graphs

Questions &

# Large Deviation Principle for Random Graphs: Research

Some interesting work regarding the Erdős-Rényi random graph:

LDP for Random Graphs

Questions & Answers

# Large Deviation Principle for Random Graphs: Research

Some interesting work regarding the Erdős-Rényi random graph:

• For an Erdős-Rényi random graph  $G_{n,p}$ , let  $T_{n,p}$  denote the number of *triangles*.

# Large Deviation Principle for Random Graphs: Research

Some interesting work regarding the Erdős-Rényi random graph:

- For an Erdős-Rényi random graph  $G_{n,p}$ , let  $T_{n,p}$  denote the number of triangles.
- It can be shown that  $\mathbf{E}T_{n,p}=\binom{n}{3}p^3$ .
- Consider the *rare* event  $E_n = \{T_{n,p} \ge (1+\delta)\mathbf{E}T_{n,p}\}$ , for some  $\delta > 0$ .

# Large Deviation Principle for Random Graphs: Research

Some interesting work regarding the Erdős-Rényi random graph:

- For an Erdős-Rényi random graph  $G_{n,p}$ , let  $T_{n,p}$  denote the number of triangles.
- It can be shown that  $\mathbf{E}T_{n,p}=\binom{n}{3}p^3$ .
- Consider the *rare* event  $E_n = \{T_{n,p} \ge (1+\delta)\mathbf{E}T_{n,p}\}$ , for some  $\delta > 0$ .
  - ? What is the asymptotic behaviour of  $P(E_n)$  as  $n \to \infty$ .

# Large Deviation Principle for Random Graphs: Research

Some interesting work regarding the Erdős-Rényi random graph:

- For an Erdős-Rényi random graph  $G_{n,p}$ , let  $T_{n,p}$  denote the number of triangles.
- It can be shown that  $\mathbf{E}T_{n,p}=\binom{n}{3}p^3$ .
- Consider the *rare* event  $E_n = \{T_{n,p} \geq (1+\delta)\mathbf{E}T_{n,p}\}$ , for some  $\delta > 0$ .
- ? What is the asymptotic behaviour of  $P(E_n)$  as  $n \to \infty$ .
- ? Can we characterize the conditional distribution of  $G_{n,p}$ given that the rare event  $E_n$  has occurred.

Deviation Principle (LDP)

Graph Limi Theory

LDP for Random Graphs

Questions &

LDP for Random Graphs

Questions & Answers

## Goals for Honours Year Project

- Better understand results concerning large deviations for:
  - Dense random graphs.
  - Sparse random graphs.

LDP for Random Graphs

Questions & Answers

## Goals for Honours Year Project

- Better understand results concerning large deviations for:
  - Dense random graphs.
  - Sparse random graphs.
- On a side note: I hope to spend more focus on sparse random graph models.

LDP for Random Graphs

Questions & Answers

## Goals for Honours Year Project

- Better understand results concerning large deviations for:
  - Dense random graphs.
  - Sparse random graphs.
- On a side note: I hope to spend more focus on sparse random graph models.
- ? Try to answer some open questions in this field.

LDP for Random Graphs

Questions & Answers

## Questions & Answers

Thank you very much for your attention.

## **Selected References**

- 1 Frank den Hollander. *Large deviations*. Fields Institute monographs, ISSN 1069-5273; 14.
- 2 László Lovász and Balázs Szegedy. Limits of dense graph sequences. J. Combin. Theory Ser. B, 96 no. 6, 933-957.
- 3 Sourav Chatterjee. *An introduction to large deviations for random graphs.* To appear in Bull. Amer. Math. Soc.