

## Reading Note

Terms marked with *color* mean they are not yet verified.

The Vlasov equation is studied as a governing equation describing the collisionless plasma physics, in which the anisotropic velocity distribution contributes a significant influence to the dynamics of the system. When coupled with Maxwell equations as the electromagnetism field governing rule, Vlasov is capable to decide the dynamic scenario for particle-field interaction, named Vlasov-Maxwell system (VM). Furthermore, under the assumption that the electrostatic force dominates the interaction, magnetic force is omitted and then comes Vlasov-Poisson problem (VP).

$$(VP) \quad \begin{cases} \partial_t f + a(v) \cdot \nabla_x f + \mu \nabla_x \phi \cdot \nabla_v f = 0 \\ \Delta \phi = \rho(f) := \int_{\mathbb{R}^3} f(t, x, v) dv \end{cases} \quad (1)$$

where,  $\mu \in \{+, -\}$ ,  $a(v) \in \{v, \hat{v}\}$ , and  $\hat{v} := v/\sqrt{1 + |v|^2}$ . The sign of  $\mu$  indicates different physical scenario, "+" for the plasma physics case and "-" for the stellar dynamics case.

Phase space distribution  $f(t, \mathbf{x}, \mathbf{v}) \geq 0$  ( $x \in \mathbb{R}_x^3, v \in \mathbb{R}_v^3, t \geq 0$ ) with initial datum  $f_0(\mathbf{x}, \mathbf{v}) = f(0, \mathbf{x}, \mathbf{v})$  determines the particle density at  $(t, \mathbf{x}, \mathbf{v})$ , *i.e.*, the number of particles per unit volume in phase space.

Some quantities have physical meaning are introduced as follows,

$$\begin{aligned} E(t, x) &= -\nabla_x \phi(t, x), \quad \phi = \frac{1}{4\pi} \frac{1}{|x|} * \rho \\ \rho(t, x) &= \int f(t, x, v) dv, \quad j(t, x) = \int a(v) f(t, x, v) dv \end{aligned} \quad (2)$$

The more realistic equations consider relativistic effect when  $a(v) = \hat{v}$  and limit the max velocity the particle can reach, transforming the Vlasov-Poisson to relativistic

Vlasov-Poisson (RVP) and the Vlasov-Maxwell to relativistic Vlasov-Maxwell (RVM).

$$(VM) \begin{cases} f_t + a(v) \cdot \nabla_x f + \mu(E + a(v) \times B) \cdot \nabla_v f = 0 \\ E_t = \nabla \times B - j \\ B_t = -\nabla \times E \\ \nabla \cdot E = \rho, \quad \nabla \cdot B = 0 \end{cases} \quad (3)$$

For classical solutions, it is well known that the existence and uniqueness result of the Vlasov-Poisson system solution have been presented by Iordanskii <sup>①</sup> in dimension 1, Ukai et al. (1978) in dimension 2, Bardos et al. (1985) in dimensions 3 for small data. The case of (nearly) symmetric data has been treated by Batt (1977), Wollman (1980), Horst et al. (1981), Schaeffer (1987). Schaeffer (1987) treated the relativistic case of symmetric data in one paper.

Glasse et al. (1985) stated the conservative characteristic of mass and energy,  $\|\rho(t, \cdot)\|$  has a bound relying on  $\|f_0\|_\infty$  and  $\mathcal{E}_0$  in plasma case and proved the existence of global classical spherically symmetric solutions to the Cauchy problem with compact support for the 3D RVP system in the plasma physics case. It turns out that, for the case of stellar dynamics, the "small" data with  $40M^{2/3}\|f_0\|_\infty^{1/3}$  will sustain the solution globally while the "large" data for which  $-\infty < \mathcal{E}_0 < 0$ , the solution blows up in finite time.

Lions et al. (1991), based on the representation formula built by the characteristic method considering the source term, proves the propagation of moments in  $v$  higher than 3. More precisely, if  $|v|^m f_0 \in L^1(\mathbb{R}^6)$  for all  $m < m_0$ , with  $m_0 > 3$ , then we build a solution of Vlasov-Poisson equations satisfying  $|v|^m f(t, \mathbf{x}, \mathbf{v}) \in L^1(\mathbb{R}^6)$  for all  $m < m_0$ , with  $m_0 > 3$  for any  $t > 0$ . Moreover, for  $m_0 > 6$ , *Sobolev injections* deduces that  $E \in L^\infty([0, T] \times \mathbb{R}^3)$  for any  $t > 0$ , and, following Horst <sup>②</sup>,  $f$  is a smooth function if  $f_0(x, v)$  is smooth. This improves a recent result of Pfaffelmoser <sup>③</sup> who has shown

① The paper [16] listed in the reference of Lions et al. (1991) is missing, referred as Iordanskii, S.V.: The Cauchy problem for the kinetic equation of plasma. Transl., II. Ser., Am. Math. Soc. 35, 351-363 (1964)

② The paper [14] listed in the reference of Lions et al. (1991) is missing, referred as Horst, E.: Global strong solutions of Vlasov's Equation. Necessary and sufficient conditions for their existence. Partial differential equations. Banach Cent. Publ. 19, 143-153 (1987)

③ The paper [19] listed in the reference of Lions et al. (1991) is missing, referred as Pfaffelmoser Global classical

independently of our work the existence of a classical solution when  $f_0$  is, roughly speaking, Lipschitz continuous in the space  $L_v^1(L_x^\infty)$ .

The existence of global solution for the 3D RVM system with finite energy initial data has been investigated extensively by many authors. Glassey et al. (1986) deduced the result that the classical solution can be globally extended with the strong assumption that  $f$  has compact support in  $v$  for all the time. Staffilani et al. (2001) presents a new perspective with Fourier analysis to the study of 3D RVM system and acquired the same result.

Continuation criterion for the global existence of the Vlasov-Maxwell system may be a more relaxed condition than assuming the compact support in  $v$  assumption. Robert et al. (1989) proved that, if the initial data decay at rate  $|v|^{-7}$  as  $|v| \rightarrow \infty$  and the imposed assumption that bound “ $\iint_{\mathbb{R}^3} |v| f(t, x, v) dv \leq \text{Constant}$ ” holds, then the solution extends globally. Luk et al. (2016), Kunze (2015), Pallard (2015) and Patel (2018) have developed recent improvements on the continuation criterion. To release the limitation of the compact support and get rid of the assumptions on the solution itself, Wang (2018) studied the propagation of regularity and the long time behavior of the 3D RVM system for suitably small initial data.

Chen et al. (2019) recently indicated that with the space torus topology  $\mathbb{T}^3$ , the weak solution of Vlasov-Poisson system exists globally and the velocity moment of order  $> 3$  can propagate, provided that the initial datum  $0 < f_0 \in L^1 \cap L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$ .

## Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system

The most important result in Lions et al. (1991) is the theorem of moment propagation,

**Theorem 0.1:** Let  $f_0 \geq 0, f_0 \in L^1 \cap L^\infty(\mathbb{R}^3 \times \mathbb{R}^3)$ . We assume that

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} |v|^m f_0(x, v) dx dv < +\infty \quad \text{if } m < m_0, \quad (4)$$

where  $3 < m_0$ . Then, there exists a solution  $f \in C(\mathbb{R}^+; L^p(\mathbb{R}^3 \times \mathbb{R}^3)) \cap L^\infty(\mathbb{R}^+; L^\infty(\mathbb{R}^3 \times \mathbb{R}^3))$  (for all  $1 \leq p < +\infty$ ) of Vlasov-Poisson system satis-

---

(footnote from last page) solutions of the Vlasov-Poisson system in three dimensions for general initial data.  
(Preprint)

fying

$$\sup_{t \in [0, T]} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |v|^m f(t, x, v) dx dv < +\infty \quad (5)$$

In the first step of the proof, we prove some general estimates on  $E$ . Then, we conclude assuming that the time interval  $(0, T)$  on which we solve the Vlasov-Poisson system is small enough and that  $E$  is bounded in  $L^{3/2}(\mathbb{R}_x^3)$ , in fact which is false in general but holds true for bounded domains of  $\mathbb{R}_x^3$ . We relax this assumption on  $E$  in a fourth and final step.

Due to the solution lying in general functional space  $\mathcal{D}'$ ,  $E \cdot \nabla_v f = \text{div}_v(Ef)$  <sup>①</sup>

proof (i) *General estimates*. Simplify the proof by handling smooth solutions first, *i.e.*,  $C^\infty$  with compact support. The *standard approximation method* (see[5,6]) would be enough to say the estimates are uniform and expand the function space, so don't worry. Treat the  $E \cdot \nabla_v f$  on the RHS as a source term safely in Vlasov-Poisson equation (??), trace the characteristic and then integrate in  $v$  to acquire  $\rho(t, x)$ .

$$\rho(t, x) = -\text{div}_x \int_0^t \int_{\mathbb{R}^3} [Ef(t-s, x-vs, v)] dv ds + \int_{\mathbb{R}^3} f_0(x-vt, v) dv \quad (6)$$

**Definition 0.1:** The supreme  $k$ -order moment in  $|v|$  the solution has on the time interval  $[0, t]$  is defined as follows:

$$M_k(t) = \sup_{0 \leq s \leq t} \int |v|^k f(s) dx dv \quad (7)$$

The above equation (6) is then *transformed* to inequality concerning  $E$

$$\|E(t, \cdot)\|_{m+3} \leq C + C \left\| \int_0^t \int_{\mathbb{R}^3} Ef(t-s, x-vs, v) dv ds \right\|_{m+3} \quad (8)$$

and *deduces* the following inequality between the derivative of  $M_k(t)$  and its power

---

① There seems to be a typo in the paper where  $E \cdot \nabla_x f = \text{div}_v(Ef)$

$$\frac{k+2}{k+3}.$$

$$\begin{aligned} \frac{d}{dt} M_k(t) &\leq \left| \frac{d}{dt} \int |v|^k f(t) dx dv \right| \\ &\leq C \|E(t)\|_{3+k} M_k(t)^{\frac{k+2}{k+3}}, \end{aligned} \quad (9)$$

with which one can imply by Gronwall lemma that,

$$M_k(t) \leq C \left\{ M_k(0) + \left( t \sup_{s \in (0,t)} \|E(s)\|_{3+k} \right)^{k+3} \right\} \quad (10)$$

confirming the bound of  $M_k(t)$ , *i.e.* the supreme  $k$ -order moment in  $|v|$  the solution has on the time interval  $[0, t]$ , exists and restricted by the supreme  $E$  during the time  $(0, t)$

(ii) *Small time estimates.* Based on the above  $\|E(t, \cdot)\|_{m+3}$  estimate (8) and *its integration on the time interval*  $(0, t_0)$  for any  $r > 3/2$  and  $t_0 \leq t \leq T$ .

$$\left\| \int_0^{t_0} s ds \sup_{\tau \in (0,T)} \left( \int_{\mathbb{R}^3} |E|^r(\tau, y) \frac{dy}{s^3} \right)^{1/r} \left( \int_{\mathbb{R}^3} f(t-s, x-vs, v) dv \right)^{1/r'} \right\|_{m+3} \|f\|_{\infty}^{(r'-1)/r'} \leq C t_0^\gamma (1 + M_m(t)^\delta), \quad (11)$$

where  $r'$  is the conjugate exponent of  $r$ ,  $1/r + 1/r' = 1$ ,  $1 \leq r' < 3$  and  $C$  only depends upon the initial data,  $\gamma = 2 - 3/r > 0$  and  $\delta = 3(k+1)/(m+3)^2 > 0$ ,  $m_0 > k > m$ . Here comes the limitation for  $m_0 > 3$ .

On the other hand,  $|E(t, \cdot)|$  is bounded by choosing  $t_0 = t$  in (8),

$$\|E(t)\|_{m+3} \leq C (1 + t^\gamma M_m(t))^\delta.$$

While in the first step, we introduced (9), the derivative of  $M_k(t)$  is bounded by the product of  $\|E(t, \cdot)\|_{3+k}$  and  $M_k(t)^{\frac{k+2}{k+3}}$ . For  $k = m$ , the above two equations jointly give

$$\frac{d}{dt} M_m(t) \leq C \left( M_m(t)^{\frac{m+2}{m+3}} + t^\gamma M_m(t)^{\delta + \frac{m+2}{m+3}} \right),$$

by which we acquire a bound imposed by  $M_k(t)$  itself on a small time interval  $[0, t_0]$  exclusive of the appearance  $\|E(t, \cdot)\|_{3+k}$ .

In the following proof, step (iii) firstly proves the boundedness of the  $M_m(t)$  on any time interval  $(0, T)$  in the case  $E(t, \cdot) \in L^{3/2}$ , and then step (iv) expanded the range to the general case.

(iii) *The case when  $E \in L^{3/2}$ .* This is a stronger limitation on the functional space than the  $E \in L^{3/2, \infty}(\mathbb{R}^3)$ <sup>①</sup>, relaxed by the author in the next step.

The estimate  $\|E(t, \cdot)\|_{3+m}$  in (8) combined with the bound acquired from step ii(11) can give a bound of the derivative of  $M_m(t)$  restricted by  $M_m(t)$  itself on any time interval  $(0, T)$ .

$$\frac{d}{dt} M_m(t) \leq C (1 + M_m(t) |\log M_m(t)|) \quad (12)$$

makes it possible to say  $M_m(t)$  is bounded on any interval  $(0, T)$ .

(iv) *The general case.* When  $E \notin L^{3/2}(\mathbb{R}^3)$ , similar to the step (i) treating  $E \cdot \nabla_v f$  as a source term, the  $E$  is decomposed into two parts:  $E = \mathbf{E}_1 + \mathbf{F}$

$$E_1 = \frac{\alpha}{4\pi} \left( \chi_R(x) \nabla \frac{1}{|x|} \right) * \rho, \quad (13)$$

where  $0 \leq \chi_R \leq 1$  is smooth such that  $\chi_R(x) = 1$  if  $|x| \leq R$  and  $\chi_R(x) = 0$  if  $|x| \geq 2R$ . But I am a little confused by the numerator why it is  $\alpha$  *rather than 1*. □

Treat the  $\mathbf{E}_1 \cdot \nabla_v f$  as source term on the RHS of the Vlasov-Poisson equation and trace the characteristic again like step (i), then it yields similar equation with (8).

$$\|E(t)\|_{m+3} \leq C + C \left\| \int_0^t \int_{\mathbb{R}^3} \left( \frac{\partial y}{\partial V} \right) E_1 f(t-s, X(s), V(s)) ds dv \right\|_{m+3} \quad (14)$$

According to the *standard approximation method*, it is said that it can keep the desired estimate, but I have not yet understood it.

## On symmetric solutions of the relativistic Vlasov-Poisson system

Glasse et al. (1985) discusses about the existence problem of spherically sym-

① I don't really understand the definition of weak type lebesgue space

metric solutions to the Cauchy problem for the 3D relativistic Vlasov-Poisson (RVP) system.

**Proposition 0.1:** Let  $f$  be a classical solution of (RVP) on some time interval  $(0, T)$  with nonnegative data  $f_0 \in C_0^1(\mathbb{R}^6)$ . Then the following properties hold:

1. If  $f_0$  vanished for  $|x| > k$ , then  $f(x, v, t)$  vanished for  $|x| > t + k$  (casuality).
2. The total mass is conserved, *i.e.*,  $\iint_{\mathbb{R}^6} f d\mathbf{v} d\mathbf{x} = \text{constant} \equiv M$ .
3. The total energy is conserved, *i.e.*,

$$\int_{\mathbb{R}^3} \left\langle \int_{\mathbb{R}^3} \sqrt{1 + |\mathbf{v}|^2} f d\mathbf{v} + \frac{1}{2} \gamma |\mathbf{E}|^2 \right\rangle d\mathbf{x} = \text{constant} = \mathcal{E}_0$$

4. If  $\gamma = +1$ , then there exists a constant  $C$  (depending on  $\|f_0\|_\infty$  and  $\mathcal{E}_0$ ) such that  $\|\rho(t)\|_{4/3} \leq C$  for  $0 \leq t < T$ .

The paper mainly talks about spherically symmetric solutions, *i.e.*, the radial ones. Here is the simplified version of characteristic trace equation.

$$\frac{dR}{ds} = \frac{U \cos A}{\sqrt{1 + U^2}} \quad (15)$$

$$\frac{dU}{ds} = \gamma \frac{\cos A}{R^2} M(R, s) \quad (16)$$

$$\frac{dA}{ds} = - \left( \gamma \frac{M(R, s)}{R^2 U} + \frac{U}{R \sqrt{1 + U^2}} \right) \sin A \quad (17)$$

#### 0.0.1 Velocity Bound in both cases

**Lemma 0.1:** There exists a constant  $C_1$  such that for  $r \geq 0$  and  $0 \leq t < T$

$$|E(x, t)| = \frac{M(r, t)}{r^2} \leq \begin{cases} \min(Mr^{-2}, 100M^{1/3} \|\hat{f}\|_\infty^{2/3} P^2(t)) & \text{if } \gamma = -1 \\ \min(Mr^{-2}, C_1 P^{5/3}(t)) & \text{if } \gamma = +1 \end{cases}$$

**Definition 0.2:** The highest speed the solution  $f$  has on the time interval  $[0, t]$ .

$$P(t) = \sup\{U(s, 0, r, u, \alpha) : 0 \leq s \leq t, (r, u, \alpha) \in \text{support } f\}$$

**Lemma 0.2:** For all  $z \geq 1$ ,

$$\xi^{-1}(z) \leq \left[ (z + A^{-1}B^{-1})^2 - 1 \right]^{1/2}$$

Lemma 0.3: For all  $t \in [0, T_0]$

$$U(t) \leq U(0) + \xi^{-1}(\sqrt{1 + U^2(0)}), \text{ when } \gamma = -1$$

#### 0.0.1.1 The stellar dynamics case

Theorem 0.2: Let  $f$  be a classical solution of (RVP) on some time interval  $[0, T)$  with  $\gamma = -1$  and smooth, nonnegative, spherically symmetric data  $f$  which has compact support and vanishes for  $(r, u, \alpha) \notin (0, \infty) \times (0, \pi)$ . If  $40M^{2/3} \|f^\circ\|_\infty^{1/3} < 1$ , then  $P(t)$  is uniformly bounded on  $[0, T)$ , and hence (RVP) possesses a global classical solution.

#### 0.0.1.2 The plasma physics case

Definition 0.3: To control the derivative of characteristic  $R(s)$ , here comes a function:

$$G(r, t) = - \int_r^\infty \min(M\lambda^{-2}, C_1 P^{5/3}(t)) d\lambda, r \geq 0 \text{ and } t \geq 0 \quad (18)$$

Lemma 0.4: Assume either  $\dot{R} \geq 0$  on  $[t_1, t_2]$  or  $\dot{R} \leq 0$  on  $[t_1, t_2]$ . Assume  $\gamma = +1$ . Then

$$|\sqrt{1 + U^2}(t_2) - \sqrt{1 + U^2}(t_1)| \leq |G(R(t_2), t_2) - G(R(t_1), t_2)| \quad (19)$$

Remark 0.1: There exists a positive constant  $C_2$

$$|G(r_1, t) - G(r_2, t)| \leq C_2 P^{5/6}(t) \text{ for all } r_1 \geq 0, r_2 \geq 0, \text{ and } t \geq 0 \quad (20)$$

Lemma 0.5: Assume  $\gamma = +1$ .  $\dot{R}$  can be zero for at most one value of  $s$ . If  $\dot{R}(t_1) = 0$ , then  $R$  has an absolute minimum at  $t_1$ .

Theorem 0.3: Let  $f$  be a classical solution of (RVP) on some time interval  $[0, T)$  with  $\gamma = +1$  and smooth, nonnegative, spherically symmetric data  $f_0$  which has compact support and vanishes for  $(r, u, \alpha) \notin (0, \infty) \times (0, \infty) \times (0, \pi)$ . Then  $P(t)$  is uniformly bounded on  $[0, T)$ , and hence (RVP) possesses a global classical solution.

proof According to Lemma 1.6 and its following remark,

$$\begin{aligned} \sqrt{1 + U^2}(t_2) &\leq \sqrt{1 + U^2}(t_1) + |G(R(t_2), t_2) - G(R(t_1), t_2)| \\ &\leq \sqrt{1 + U^2}(t_1) + C_2 P^{5/6}(t_2) \end{aligned} \quad (21)$$



Either  $\dot{R}$  never vanishes or vanishes at one value,

$$\sqrt{1 + U^2(t)} \leq \sqrt{1 + U^2(0)} + 2C_2 P^{5/6}(t) \quad (22)$$

holds for  $t \in [0, T)$  as long as we apply Eq. 21 at most twice. Naturally,

$$P(t) \leq \sqrt{1 + P^2(t)} \leq \sqrt{1 + P^2(0)} + 2C_2 P^{5/6}(t) \quad (23)$$

induces that  $P(t)$  has upper bound.  $\square$

### 0.0.2 Blow-up of Radial Solutions in the case of stellar dynamics system

**Theorem 0.4:** Let  $f_0$  be smooth, nonnegative, radial and of compact support on  $\mathbb{R}^6$ . Let  $f(\mathbf{x}, \mathbf{v}, t)$  be a classical solution of (RVP) on an interval  $0 < t < T$  for which  $-\infty < \mathcal{E}_0 < 0$ . Then  $T < \infty$ .

proof The "dilation identity" is used below, and here is its derivation.

$$\begin{aligned} \frac{d}{dt} \iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f d\mathbf{v} d\mathbf{x} &= \iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} [-\hat{\mathbf{v}} \cdot \nabla_{\mathbf{x}} f + \mathbf{E} \cdot \nabla_{\mathbf{v}} f] d\mathbf{v} d\mathbf{x} = \dots \text{(a lot omitted)} \\ &= \iint_{\mathbb{R}^6} \frac{|\mathbf{v}|^2 f}{\sqrt{1 + |\mathbf{v}|^2}} d\mathbf{v} d\mathbf{x} - \int_{\mathbb{R}^3} \rho \mathbf{x} \cdot \mathbf{E} d\mathbf{x} = \dots \text{(a lot omitted)} \\ &= \iint_{\mathbb{R}^6} \frac{|\mathbf{v}|^2 f}{\sqrt{1 + |\mathbf{v}|^2}} d\mathbf{v} d\mathbf{x} - \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 d\mathbf{x} \\ &= \mathcal{E}_0 - \iint_{\mathbb{R}^6} \frac{1}{\sqrt{1 + |\mathbf{v}|^2}} f d\mathbf{v} d\mathbf{x} \end{aligned} \quad (24)$$

Clearly, we have a coarse upper bound estimation of  $\iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f d\mathbf{v} d\mathbf{x}$ , which is a part of  $\frac{d}{dt} \iint_{\mathbb{R}^6} r^2 \sqrt{1 + |\mathbf{v}|^2} f d\mathbf{v} d\mathbf{x}$ .

$$\iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f d\mathbf{v} d\mathbf{x} \leq \iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f_0 d\mathbf{v} d\mathbf{x} + \mathcal{E}_0 t$$

$$\begin{aligned}
\frac{d}{dt} \iint_{\mathbb{R}^6} r^2 \sqrt{1 + |\mathbf{v}|^2} f d\mathbf{v} d\mathbf{x} &= 2 \iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f d\mathbf{v} d\mathbf{x} - \int_{\mathbb{R}^3} r^2 \mathbf{E} \cdot \mathbf{j} d\mathbf{x} \\
(\mathbf{j} &= \int_{\mathbb{R}^3} \hat{\mathbf{v}} f d\mathbf{v} \text{ and } \left| \int_{\mathbb{R}^3} r^2 \mathbf{E} \cdot \mathbf{j} d\mathbf{x} \right| \leq M^2) \\
&\leq 2 \left( \iint_{\mathbb{R}^6} \mathbf{x} \cdot \mathbf{v} f_0 d\mathbf{v} d\mathbf{x} + \mathcal{E}_0 t \right) + M^2 \quad (25) \\
&\leq \text{Constant} + 2\mathcal{E}_0 t \\
\Rightarrow 0 &\leq \iint_{\mathbb{R}^6} r^2 \sqrt{1 + |\mathbf{v}|^2} f d\mathbf{v} d\mathbf{x} \leq C + Ct + \mathcal{E}_0 t^2
\end{aligned}$$

However, since we assume  $f_0$  is a "large" enough initial data, *i.e.*, satisfying the hypothesis that  $\mathcal{E}_0 < 0$ . There must be some time the solution blows up.  $\square$

## List of Equations

Equation 1 .....	1
Equation 2 .....	1
Equation 3 .....	2
Equation 4 .....	3
Equation 5 .....	4
Equation 6 .....	4
Equation 7 .....	4
Equation 8 .....	4
Equation 9 .....	5
Equation 10 .....	5
Equation 11 .....	5
Equation 12 .....	6
Equation 13 .....	6
Equation 14 .....	6
Equation 15 .....	7
Equation 16 .....	7
Equation 17 .....	7
Equation 18 .....	8
Equation 19 .....	8
Equation 20 .....	8
Equation 21 .....	8
Equation 22 .....	9
Equation 23 .....	9
Equation 24 .....	9
Equation 25 .....	10

## Bibliography

- Lions P L, Perthame B. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system.[J/OL]. *Inventiones Mathematicae*, 1991, 105:415. DOI: 10.1007/BF01232273.
- UKAI S, OKABE T. On classical solutions in the large in time of two-dimensional vlasov's equation [J]. *Osaka Journal of Mathematics*, 1978, 15(2):245-261.
- BARDOS C, DEGOND P. Global existence for the vlasov-poisson equation in 3 space variables with small initial data[C]//*Annales de l'Institut Henri Poincare (C) Non Linear Analysis: volume 2*. Elsevier, 1985: 101-118.
- BATT J. Global symmetric solutions of the initial value problem of stellar dynamics[J]. *Journal of Differential Equations*, 1977, 25(3):342-364.
- WOLLMAN S. Global-in-time solutions of the two-dimensional vlasov-poisson systems[J]. *Communications on Pure and Applied Mathematics*, 1980, 33(2):173-197.
- HORST E, NEUNZERT H. On the classical solutions of the initial value problem for the unmodified non-linear vlasov equation i general theory[J]. *Mathematical Methods in the Applied Sciences*, 1981, 3(1):229-248.
- SCHAEFFER J. Global existence for the poisson-vlasov system with nearly symmetric data[J]. *Journal of differential equations*, 1987, 69(1):111-148.
- GLASSEY R T, SCHAEFFER J. On symmetric solutions of the relativistic vlasov-poisson system [J/OL]. 101(4):459-473[2020-03-13]. <https://doi.org/10.1007/BF01210740>.
- GLASSEY R T, STRAUSS W A. Singularity formation in a collisionless plasma could occur only at high velocities[J]. *Archive for rational mechanics and analysis*, 1986, 92(1):59-90.
- STAFFILANI G, KLAINERMAN S. A new approach to study the vlasov-maxwell system[J/OL]. *Communications on Pure and Applied Analysis - COMMUN PURE APPL ANAL*, 2001, 1: 103-125. DOI: 10.3934/cpaa.2002.1.103.
- ROBERT T G, WALTER A S. Large velocities in the relativistic vlasov-maxwell equations[J]. *Journal of the Faculty of Science, the University of Tokyo. Sect. 1 A, Mathematics = 東京大学理学部紀要. 第 1 類 A, 数学*, 1989, 36(3):615-627.
- Luk J, Strain R M. Strichartz Estimates and Moment Bounds for the Relativistic Vlasov-Maxwell System[J/OL]. *Archive for Rational Mechanics and Analysis*, 2016, 219(1):445-552. DOI: 10.1007/s00205-015-0899-1.

- KUNZE M. Yet another criterion for global existence in the 3d relativistic vlasov - maxwell system[J/OL]. Journal of Differential Equations, 2015, 259(9):4413 - 4442. <http://www.sciencedirect.com/science/article/pii/S0022039615003009>. DOI: <https://doi.org/10.1016/j.jde.2015.06.003>.
- PALLARD C. A refined existence criterion for the relativistic vlasov-maxwell system[J/OL]. Communications in Mathematical Sciences, 2015, 13:347-354. DOI: 10.4310/CMS.2015.v13.n2.a4.
- PATEL N. Three new results on continuation criteria for the 3d relativistic vlasov - maxwell system[J/OL]. Journal of Differential Equations, 2018, 264(3):1841 - 1885. <http://www.sciencedirect.com/science/article/pii/S0022039617305442>. DOI: <https://doi.org/10.1016/j.jde.2017.10.008>.
- WANG X. Propagation of regularity and long time behavior of the 3d massive relativistic transport equation II: Vlasov-maxwell system[J/OL]. [2019-12-13]. <http://arxiv.org/abs/1804.06566>.
- CHEN Z, CHEN J. Moments propagation for weak solutions of the vlasov-poisson system in the three-dimensional torus[J/OL]. 472(1):728-737[2020-03-14]. <http://www.sciencedirect.com/science/article/pii/S0022247X18309958>. DOI: 10.1016/j.jmaa.2018.11.049.

综合论文训练记录表

学生姓名		学号		班级	
论文题目					
主要内容以及进度安排	<div>指导教师签字：_____</div> <div>考核组组长签字：_____</div> <div>年 月 日</div>				
中期考核意见	<div>考核组组长签字：_____</div> <div>年 月 日</div>				

指导教师评语	<div>指导教师签字：_____</div> <div>年      月      日</div>
评阅教师评语	<div>评阅教师签字：_____</div> <div>年      月      日</div>
答辩小组评语	<div>答辩小组组长签字：_____</div> <div>年      月      日</div>

总成绩：\_\_\_\_\_

教学负责人签字：\_\_\_\_\_

年      月      日