

Harmonic Analysis Notes

July 25, 2026

Contents

I	Foundations of Harmonic Analysis	4
1	Interpolation Theorems	4
1.1	The Riesz–Thorin Interpolation Theorem	4
1.2	Sublinear Operators and Weak L^p Spaces	5
1.3	The Marcinkiewicz Interpolation Theorem	5
1.4	A Standard Application Pattern	6
2	Fourier Series, Summability Kernels, and Approximate Identities	7
2.1	Convolution and the Physical–Frequency Dictionary	7
2.2	The Dirichlet and Fejér Kernels	7
2.3	Approximate Identities	8
2.4	Convergence of Approximate Identities	9
3	The Hardy–Littlewood Maximal Function	10
3.1	Motivation: A Generalized Newton–Leibniz Formula	10
3.2	Definition and Main Estimate	10
3.3	Proof of the Weak-Type Estimate	10
3.4	The Lebesgue Differentiation Theorem	11
3.5	Lebesgue Points	12
4	Ergodic Theory	13
4.1	Measure-Preserving Dynamical Systems	13
4.2	Ergodicity	13
4.3	The von Neumann Mean Ergodic Theorem	14
4.4	Birkhoff’s Pointwise Ergodic Theorem	15
4.5	The Maximal Ergodic Estimate	15
4.6	Structural Summary	18
II	Weighted Estimates and Singular Integrals	18
5	Weighted Maximal Function Theory	18
5.1	Review and Motivation	18
5.2	Muckenhoupt’s Theorem	19
5.3	The Elementary Implications	20

5.4	Dyadic Cubes and the Dyadic Maximal Operator	21
5.5	A Dyadic Calderón–Zygmund Decomposition	21
5.6	Weighted Boundedness of the Dyadic Maximal Operator	22
6	The Hilbert Transform	24
6.1	Singular Kernels and Singular Integrals	24
6.2	Definition of the Hilbert Transform	24
6.3	The Fourier Multiplier	25
6.4	L^p Boundedness	26
7	The Fourier Transform on Euclidean Space	27
7.1	Convolution and the Multiplication Formula	28
7.2	Basic Examples and Decay	28
7.3	The Gaussian	29
7.4	Fourier Inversion on the Schwartz Space	29
7.5	Plancherel’s Theorem	30
7.6	The Hausdorff–Young Theorem	30
III	Randomization and Pointwise Fourier Convergence	31
8	Probability and Randomization	31
8.1	Basic probability theory	31
8.2	The Borel–Cantelli lemmas	31
8.3	Kolmogorov’s zero–one law	32
8.4	Sub-Gaussian bounds	32
8.5	Khinchine’s inequality	33
9	Fourier Partial Sums and Divergence	34
9.1	Partial sums	34
9.2	Norm convergence and uniform boundedness	34
9.3	Kolmogorov’s divergence theorem	35
9.4	The Banach principle	35
9.5	A measure-theoretic corollary and the construction	37
9.6	A refined almost-everywhere divergence construction	39
10	Poisson Integrals and Boundary Values	40
10.1	The Hilbert transform and the Poisson kernel	40
10.2	Almost-everywhere convergence of Poisson integrals	41
IV	Calderón–Zygmund Theory and BMO	42
11	Homogeneous Singular Integrals	42
11.1	Dini continuity and the Hörmander condition	42
11.2	Translation invariance and dilation structure	43
11.3	Fourier multipliers of truncated kernels	44

12 Maximal Singular Integrals	45
12.1 Calderón–Zygmund decomposition and the good part	45
12.2 Cancellation estimate for the bad part	46
12.3 Consequences of the cancellation estimate	46
12.4 Geometric proof of the cancellation estimate	46
13 BMO and Endpoint Estimates	47
13.1 Averages over nested cubes grow only linearly	47
13.2 Equivalent norm with arbitrary constants	47
13.3 Where BMO sits	48
13.4 Calderón–Zygmund operators map L^∞ to BMO	48
13.5 Local and far-field decomposition	48
14 Calderón–Zygmund Proof Architecture	49
 V Multipliers and Littlewood–Paley Theory	 50
15 Fourier Multipliers	50
15.1 The Mikhlin multiplier theorem	50
15.2 Reduction to Calderón–Zygmund kernels	50
15.3 Dyadic kernel estimates	51
15.4 Proof of the Mikhlin theorem	52
16 Littlewood–Paley Theory	53
16.1 Dimension $d = 1$	53
16.2 Higher dimensions	53
17 Applications of Littlewood–Paley Theory	55
17.1 Homogeneous Sobolev spaces and Sobolev embedding	55
17.2 Hölder spaces and their Littlewood–Paley characterization	55
 VI Oscillatory Integrals and Fourier Restriction	 57
18 Oscillatory Integrals and Stationary Phase	57
18.1 Fourier transforms of surface measures	57
18.2 Stationary phase	58
18.3 Fourier decay of surface measures	61
19 Fourier Restriction Theory	61
19.1 Critical points and the restriction problem	61
19.2 The L^2 restriction theorem	62
19.3 Restriction–extension duality	63
19.4 Necessary exponents, Stein’s conjecture, and Kakeya	64
19.5 A refined stationary-phase lemma	65

Part I

Foundations of Harmonic Analysis

1 Interpolation Theorems

The notes proceed from interpolation to Fourier summation and approximate identities, then to maximal functions and differentiation, and finally to ergodic theory. Throughout, we use the notation

$$e(x) = e^{2\pi i x}, \quad \mathbb{T} = \mathbb{R}/\mathbb{Z}, \quad \widehat{f}(n) = \int_0^1 f(x) e^{-2\pi i n x} dx,$$

and $|E|$ denotes Lebesgue measure. For a measure-preserving transformation T , we write

$$A_m f = \frac{1}{m} \sum_{k=0}^{m-1} f \circ T^k.$$

1.1 The Riesz–Thorin Interpolation Theorem

Theorem 1.1 (Riesz–Thorin). *Let $1 \leq p_1, p_2, q_1, q_2 \leq \infty$, let (X, μ) and $(\tilde{X}, \tilde{\mu})$ be measure spaces, and let T be a linear operator such that*

$$T : L^{p_1}(X) \longrightarrow L^{q_1}(\tilde{X}), \quad T : L^{p_2}(X) \longrightarrow L^{q_2}(\tilde{X})$$

is bounded with norms M_1 and M_2 , respectively. If

$$\left(\frac{1}{p}, \frac{1}{q}\right) = (1-\theta) \left(\frac{1}{p_1}, \frac{1}{q_1}\right) + \theta \left(\frac{1}{p_2}, \frac{1}{q_2}\right), \quad 0 \leq \theta \leq 1,$$

then $T : L^p(X) \rightarrow L^q(\tilde{X})$ is bounded and

$$\|T\|_{L^p \rightarrow L^q} \leq M_1^{1-\theta} M_2^\theta.$$

Proof sketch via the three-lines lemma. For simple functions f and g , consider

$$\Phi(z) = \int_{\tilde{X}} T(f_z) g_z d\tilde{\mu},$$

where

$$f_z = |f|^{p \left(\frac{1-z}{p_1} + \frac{z}{p_2} \right)} \frac{f}{|f|}, \quad g_z = |g|^{q' \left(\frac{1-z}{q'_1} + \frac{z}{q'_2} \right)} \frac{g}{|g|}.$$

The function Φ is holomorphic in the strip $0 < \operatorname{Re} z < 1$, bounded and continuous on its closure, and the endpoint assumptions give

$$|\Phi(iy)| \leq M_1, \quad |\Phi(1+iy)| \leq M_2.$$

Hadamard's three-lines lemma yields

$$|\Phi(\theta)| \leq M_1^{1-\theta} M_2^\theta.$$

Taking the supremum over g and using duality proves the claim. $\square$

Remark 1.2. Linearity is essential in the Riesz–Thorin theorem. The interpolated indices lie on

the line segment joining $(1/p_1, 1/q_1)$ and $(1/p_2, 1/q_2)$.

1.2 Sublinear Operators and Weak L^p Spaces

Definition 1.3. An operator T , defined on a dense subclass $\mathcal{D} \subseteq L^p(X)$, is *sublinear* if

$$|T(f_1 + f_2)| \leq |Tf_1| + |Tf_2|, \quad |T(cf)| = |c| |Tf|.$$

Definition 1.4. For $1 \leq p < \infty$, the weak L^p quasi-norm is

$$\|f\|_{L^{p,\infty}(X,\mu)} = \sup_{\lambda>0} \lambda \mu(\{x \in X : |f(x)| > \lambda\})^{1/p}.$$

We write $L^{\infty,\infty} = L^\infty$.

This is only a quasi-norm: its triangle inequality holds up to a constant. Chebyshev's inequality gives the continuous embedding $L^p \subseteq L^{p,\infty}$:

$$\mu\{|f| > \lambda\} \leq \int_{\{|f|>\lambda\}} \frac{|f|^p}{\lambda^p} d\mu \leq \frac{\|f\|_{L^p}^p}{\lambda^p},$$

and hence

$$\|f\|_{L^{p,\infty}} \leq \|f\|_{L^p}.$$

The converse is false. For example,

$$|x|^{-d/p} \in L^{p,\infty}(\mathbb{R}^d) \setminus L^p(\mathbb{R}^d).$$

The notation $T : L^p(X) \rightarrow L^{q,\infty}(\tilde{X})$ means that

$$\|Tf\|_{L^{q,\infty}} \leq C \|f\|_{L^p};$$

this is called a weak-type estimate.

1.3 The Marcinkiewicz Interpolation Theorem

Theorem 1.5 (Marcinkiewicz). *Let T be sublinear, let $1 \leq p_1, p_2, q_1, q_2 \leq \infty$, assume $p_j \leq q_j$ and $q_1 \neq q_2$, and suppose*

$$T : L^{p_1} \longrightarrow L^{q_1,\infty}, \quad T : L^{p_2} \longrightarrow L^{q_2,\infty}$$

are weak-type bounds. If

$$\left(\frac{1}{p}, \frac{1}{q}\right) = (1-\theta) \left(\frac{1}{p_1}, \frac{1}{q_1}\right) + \theta \left(\frac{1}{p_2}, \frac{1}{q_2}\right), \quad 0 < \theta < 1,$$

then $T : L^p \rightarrow L^q$ is strongly bounded.

Remark 1.6. In contrast with Riesz–Thorin, Marcinkiewicz interpolation applies to sublinear operators and assumes only weak-type endpoint bounds. The price is that the endpoints are excluded and the interpolation segment cannot be horizontal.

We give the proof in the most frequently used diagonal case $p_j = q_j$. Suppose

$$p_1 < p < p_2 < \infty, \quad \|Tf\|_{L^{p_j,\infty}} \leq A_j \|f\|_{p_j}.$$

For every measurable g , the layer-cake formula gives

$$\|g\|_{L^p}^p = p \int_0^\infty \lambda^{p-1} \mu\{|g| > \lambda\} d\lambda.$$

For a fixed $\lambda > 0$, split f at height λ :

$$f_1^\lambda = f \mathbf{1}_{\{|f| > \lambda\}}, \quad f_2^\lambda = f \mathbf{1}_{\{|f| \leq \lambda\}}.$$

On $\{|f| > \lambda\}$,

$$|f|^{p_1} = |f|^p |f|^{p_1-p} \leq |f|^p \lambda^{p_1-p},$$

so $f_1^\lambda \in L^{p_1}$. Similarly,

$$|f|^{p_2} \leq |f|^p \lambda^{p_2-p} \quad \text{on } \{|f| \leq \lambda\},$$

and therefore $f_2^\lambda \in L^{p_2}$.

Sublinearity and the weak-type estimates imply

$$\begin{aligned} \mu\{|Tf| > 2\lambda\} &\leq \mu\{|Tf_1^\lambda| > \lambda\} + \mu\{|Tf_2^\lambda| > \lambda\} \\ &\leq \frac{A_1^{p_1} \|f_1^\lambda\|_{p_1}^{p_1}}{\lambda^{p_1}} + \frac{A_2^{p_2} \|f_2^\lambda\|_{p_2}^{p_2}}{\lambda^{p_2}}. \end{aligned}$$

Consequently, by Fubini's theorem,

$$\begin{aligned} \|Tf\|_p^p &\leq 2^p p \left[A_1^{p_1} \int_0^\infty \lambda^{p-p_1-1} \int_{\{|f| > \lambda\}} |f|^{p_1} d\mu d\lambda \right. \\ &\quad \left. + A_2^{p_2} \int_0^\infty \lambda^{p-p_2-1} \int_{\{|f| \leq \lambda\}} |f|^{p_2} d\mu d\lambda \right]. \end{aligned}$$

Changing the order of integration gives

$$\begin{aligned} \int_0^\infty \lambda^{p-p_1-1} \int_{\{|f| > \lambda\}} |f|^{p_1} d\mu d\lambda &= \frac{\|f\|_p^p}{p-p_1}, \\ \int_0^\infty \lambda^{p-p_2-1} \int_{\{|f| \leq \lambda\}} |f|^{p_2} d\mu d\lambda &= \frac{\|f\|_p^p}{p_2-p}. \end{aligned}$$

Thus

$$\|Tf\|_p \leq 2 \left[p \left(\frac{A_1^{p_1}}{p-p_1} + \frac{A_2^{p_2}}{p_2-p} \right) \right]^{1/p} \|f\|_p.$$

The constant blows up as $p \downarrow p_1$ or $p \uparrow p_2$, which explains why the endpoints are excluded.

If $p_2 = \infty$, split instead at height $\lambda/(2A_2)$. Then

$$\|Tf_2^{\lambda/(2A_2)}\|_\infty \leq A_2 \frac{\lambda}{2A_2} = \frac{\lambda}{2},$$

so the second distribution-function term disappears, and the rest of the argument is unchanged.

1.4 A Standard Application Pattern

Maximal operators often fail to map L^1 to L^1 . The standard route to strong L^p bounds is:

1. prove $T : L^\infty \rightarrow L^\infty$;

2. prove the weak-type bound $T : L^1 \rightarrow L^{1,\infty}$, usually by a covering lemma;
3. interpolate to obtain strong type (p, p) for $1 < p < \infty$.

2 Fourier Series, Summability Kernels, and Approximate Identities

2.1 Convolution and the Physical–Frequency Dictionary

If $f, g \in L^1(\mathbb{T})$, then

$$\widehat{f * g}(n) = \widehat{f}(n)\widehat{g}(n).$$

The following correspondences are fundamental:

	Physical side	Frequency side
1	$f * g$	$\widehat{f}\widehat{g}$
2	f'	multiplication by $2\pi in$
3	multiplication by $1 - e^{2\pi ix}$	$\Delta \widehat{f}(n) = \widehat{f}(n) - \widehat{f}(n-1)$
4	regularity	decay
5	decay	regularity

For example, if $f \in C^k(\mathbb{T})$, repeated integration by parts and periodicity give

$$\widehat{f}(n) = \frac{\widehat{f^{(k)}}(n)}{(2\pi in)^k}, \quad |\widehat{f}(n)| \leq \frac{\|f^{(k)}\|_1}{(2\pi|n|)^k}.$$

Thus smoothness forces polynomial decay of Fourier coefficients. Conversely, suitable decay or summability assumptions on Fourier coefficients imply regularity. Multiplication by $1 - e^{2\pi ix}$, for example, corresponds to discrete differentiation of the Fourier coefficients.

2.2 The Dirichlet and Fejér Kernels

The N -th symmetric Fourier partial sum is

$$S_N f = f * D_N, \quad D_N(x) = \sum_{n=-N}^N e(n x).$$

On the frequency side, S_N is multiplication by $\mathbf{1}_{[-N, N]}$, a sharp cutoff. Accordingly, D_N oscillates, changes sign, and satisfies

$$\|D_N\|_{L^1(\mathbb{T})} \sim \log N.$$

This unboundedness is the source of the failure of $S_N f \rightarrow f$ in general in $C(\mathbb{T})$ and $L^1(\mathbb{T})$.

Cesàro averaging replaces the sharp cutoff by a smoother one:

$$\sigma_N f = \frac{1}{N} \sum_{k=0}^{N-1} S_k f = f * K_N, \quad K_N = \frac{1}{N} \sum_{k=0}^{N-1} D_k.$$

To compute the closed form, set $z = e(x)$. A geometric sum gives

$$D_k = \frac{z^{k+\frac{1}{2}} - z^{-k-\frac{1}{2}}}{z^{\frac{1}{2}} - z^{-\frac{1}{2}}}.$$

Hence

$$\begin{aligned}
K_N(z^{\frac{1}{2}} - z^{-\frac{1}{2}}) &= \frac{1}{N} \sum_{k=0}^{N-1} (z^{k+\frac{1}{2}} - z^{-k-\frac{1}{2}}) \\
&= \frac{1}{N} \left(\frac{z^N - 1}{z^{\frac{1}{2}} - z^{-\frac{1}{2}}} - \frac{1 - z^{-N}}{z^{\frac{1}{2}} - z^{-\frac{1}{2}}} \right) \\
&= \frac{1}{N} \frac{(z^{N/2} - z^{-N/2})^2}{z^{\frac{1}{2}} - z^{-\frac{1}{2}}}.
\end{aligned}$$

Here we used

$$\sum_{k=0}^{N-1} z^{k+\frac{1}{2}} = \frac{z^{\frac{1}{2}}(z^N - 1)}{z - 1} = \frac{z^N - 1}{z^{\frac{1}{2}} - z^{-\frac{1}{2}}}.$$

Therefore

$$K_N(x) = \frac{1}{N} \left(\frac{z^{N/2} - z^{-N/2}}{z^{1/2} - z^{-1/2}} \right)^2 = \frac{1}{N} \frac{\sin^2(\pi Nx)}{\sin^2(\pi x)}.$$

For $|x| \leq 1/2$,

$$0 \leq K_N(x) \lesssim \min \left\{ \frac{1}{Nx^2}, N \right\} = \frac{1}{N} \min \left\{ \frac{1}{x^2}, N^2 \right\}.$$

Indeed,

$$\sin^2(\pi Nx) \leq \min\{1, (\pi Nx)^2\}, \quad \sin^2(\pi x) \geq 4x^2$$

in this range.

On the frequency side, the Fejér kernel has a triangular cutoff:

$$K_N(x) = \sum_{|k| < N} \left(1 - \frac{|k|}{N} \right) e(kx).$$

This follows by counting:

$$\begin{aligned}
\frac{1}{N} \sum_{k=0}^{N-1} \sum_{|j| \leq k} e(jx) &= \frac{1}{N} \sum_{|j| \leq N-1} \#\{k : |j| \leq k \leq N-1\} e(jx) \\
&= \sum_{|j| < N} \left(1 - \frac{|j|}{N} \right) e(jx).
\end{aligned}$$

2.3 Approximate Identities

Definition 2.1. A sequence $\{\Phi_N\}_{N=1}^{\infty} \subseteq L^{\infty}(\mathbb{T})$ is an *approximate identity* if

1. $\int_0^1 \Phi_N(x) dx = 1$;
2. $\sup_N \int_0^1 |\Phi_N(x)| dx < \infty$;
3. for every $\delta > 0$,

$$\lim_{N \rightarrow \infty} \int_{|x| > \delta} |\Phi_N(x)| dx = 0.$$

Example 2.2. The Fejér kernels form an approximate identity. Their zeroth Fourier coefficient

gives $\int K_N = 1$. Since $K_N \geq 0$, $\int |K_N| = 1$. Finally,

$$\int_{|x|>\delta} |K_N(x)| dx \leq \frac{C}{N} \int_{\delta}^{1/2} \frac{dx}{x^2} \leq \frac{C}{N\delta} \rightarrow 0.$$

By contrast, the Dirichlet kernels violate the second condition because $\|D_N\|_1 \sim \log N$.

2.4 Convergence of Approximate Identities

Proposition 2.3. *Let $\{\Phi_N\}$ be an approximate identity.*

1. *If $f \in C(\mathbb{T})$, then $\|\Phi_N * f - f\|_{L^\infty} \rightarrow 0$.*
2. *If $f \in L^p(\mathbb{T})$, $1 \leq p < \infty$, then $\|\Phi_N * f - f\|_{L^p} \rightarrow 0$.*
3. *If $\mu \in M(\mathbb{T})$, then $\mu * \Phi_N \rightarrow \mu$ in the weak-* topology.*

Proof. For the first assertion, normalization gives

$$\Phi_N * f(x) - f(x) = \int_{\mathbb{T}} \Phi_N(y)(f(x-y) - f(x)) dy.$$

Split this integral into $|y| \leq \delta$ and $|y| > \delta$. Uniform continuity of f gives, for sufficiently small δ ,

$$\int_{|y| \leq \delta} |\Phi_N(y)| |f(x-y) - f(x)| dy \leq \varepsilon \|\Phi_N\|_1 \leq C\varepsilon.$$

The remaining part is bounded by

$$2\|f\|_\infty \int_{|y|>\delta} |\Phi_N(y)| dy,$$

which tends to zero.

For $1 \leq p < \infty$, Minkowski's integral inequality gives

$$\|\Phi_N * f - f\|_{L^p} \leq \int_{\mathbb{T}} |\Phi_N(y)| \|f(\cdot - y) - f\|_{L^p} dy.$$

Translations are continuous in L^p :

$$\|f(\cdot - y) - f\|_{L^p} \rightarrow 0 \quad \text{as } y \rightarrow 0.$$

Splitting again at δ proves the second assertion.

For the third assertion, define $\tilde{\Phi}_N(y) = \Phi_N(-y)$. Fubini's theorem gives, for $f \in C(\mathbb{T})$,

$$\int_{\mathbb{T}} f d(\mu * \Phi_N) = \int_{\mathbb{T}} (\tilde{\Phi}_N * f) d\mu \rightarrow \int_{\mathbb{T}} f d\mu,$$

using the uniform convergence from the first assertion and the finiteness of $|\mu|(\mathbb{T})$. $\square$

Corollary 2.4 (Fejér's theorem). *If $f \in C(\mathbb{T})$, then $\sigma_N f \rightarrow f$ uniformly. If $f \in L^p(\mathbb{T})$, $1 \leq p < \infty$, then $\sigma_N f \rightarrow f$ in L^p . In particular, trigonometric polynomials are dense in both $C(\mathbb{T})$ and $L^p(\mathbb{T})$.*

3 The Hardy–Littlewood Maximal Function

3.1 Motivation: A Generalized Newton–Leibniz Formula

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable and define

$$F(x) = \int_a^x f(y) dy.$$

Formally,

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{1}{|I|} \int_I f, \quad I = (x, x+h).$$

This raises the question whether shrinking averages converge almost everywhere to $f(x)$. In $\mathbb{R}^d$, the corresponding question is whether

$$\lim_{\substack{|B| \rightarrow 0 \\ x \in B}} \frac{1}{|B|} \int_B f(y) dy = f(x) \quad \text{for almost every } x.$$

The standard tool for controlling such families of averages is the maximal function.

3.2 Definition and Main Estimate

Definition 3.1. For $f \in L^1(\mathbb{R}^d)$, the uncentered Hardy–Littlewood maximal function is

$$Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy,$$

where the supremum is over all open balls containing x .

Theorem 3.2. If $f \in L^1(\mathbb{R}^d)$, then:

- (i) Mf is measurable;
- (ii) $Mf(x) < \infty$ for almost every x ;
- (iii) for every $\alpha > 0$,

$$|\{x : Mf(x) > \alpha\}| \leq \frac{3^d}{\alpha} \|f\|_{L^1}.$$

The last inequality is the weak-type $(1, 1)$ estimate. In general $Mf \notin L^1$ unless $f = 0$, so weak type is the best possible endpoint result.

3.3 Proof of the Weak-Type Estimate

For $\alpha > 0$, set

$$E_\alpha = \{x \in \mathbb{R}^d : Mf(x) > \alpha\}.$$

If $x \in E_\alpha$, there is an open ball $B \ni x$ such that

$$\frac{1}{|B|} \int_B |f| > \alpha.$$

The same ball works for every $x' \in B$, so $B \subseteq E_\alpha$. Thus E_α is open, Mf is lower semicontinuous, and in particular it is measurable.

Lemma 3.3 (Finite Vitali-type covering lemma). *Let $\mathcal{B} = \{B_1, \dots, B_n\}$ be a finite family of open balls. There is a pairwise disjoint subfamily $B_{i_1}, \dots, B_{i_k}$ such that*

$$\left| \bigcup_{i=1}^n B_i \right| \leq 3^d \sum_{p=1}^k |B_{i_p}|.$$

Proof. Choose a ball of largest radius and discard all balls intersecting it. Repeat the procedure with the remaining balls. The selected balls are pairwise disjoint. Every discarded ball B_j intersects a selected ball B_{i_p} whose radius is at least r_j , and hence

$$B_j \subseteq 3B_{i_p}.$$

Therefore

$$\left| \bigcup_i B_i \right| \leq \left| \bigcup_p 3B_{i_p} \right| \leq \sum_p |3B_{i_p}| = 3^d \sum_p |B_{i_p}|.$$

□

For each $x \in E_\alpha$, choose a ball $B_x \ni x$ satisfying

$$|B_x| < \frac{1}{\alpha} \int_{B_x} |f|.$$

Let $K \subseteq E_\alpha$ be compact. By compactness, finitely many of the B_x cover K . Apply the covering lemma and retain a pairwise disjoint subfamily $\{B_{i_p}\}$. Then

$$|K| \leq 3^d \sum_p |B_{i_p}| \leq \frac{3^d}{\alpha} \sum_p \int_{B_{i_p}} |f| \leq \frac{3^d}{\alpha} \|f\|_{L^1}.$$

Inner regularity of Lebesgue measure now yields

$$|E_\alpha| \leq \frac{3^d}{\alpha} \|f\|_{L^1}.$$

Finally,

$$|\{Mf = \infty\}| \leq |\{Mf > \alpha\}| \leq \frac{3^d}{\alpha} \|f\|_1 \rightarrow 0$$

as $\alpha \rightarrow \infty$, proving almost-everywhere finiteness.

Corollary 3.4. *For $1 < p \leq \infty$,*

$$\|Mf\|_{L^p} \leq C_{d,p} \|f\|_{L^p}.$$

Proof. The L^∞ estimate is immediate, and the weak $(1, 1)$ estimate was proved above. Apply Marcinkiewicz interpolation. □

3.4 The Lebesgue Differentiation Theorem

Theorem 3.5 (Lebesgue differentiation theorem). *For $f \in L^1(\mathbb{R}^d)$,*

$$\lim_{\substack{|B| \rightarrow 0 \\ x \in B}} \frac{1}{|B|} \int_B f(y) dy = f(x) \quad \text{for almost every } x.$$

Proof. For $\alpha > 0$, let

$$E_\alpha = \left\{ x : \limsup_{\substack{|B| \rightarrow 0 \\ x \in B}} \left| \frac{1}{|B|} \int_B f - f(x) \right| > \alpha \right\}.$$

It suffices to prove $|E_\alpha| = 0$.

Choose $g \in C_c^\infty(\mathbb{R}^d)$ such that $\|g - f\|_1 < \varepsilon$. By the triangle inequality,

$$\begin{aligned} \left| \frac{1}{|B|} \int_B f - f(x) \right| &\leq \frac{1}{|B|} \int_B |f - g| + \left| \frac{1}{|B|} \int_B g - g(x) \right| + |g(x) - f(x)| \\ &\leq (f - g)^*(x) + \left| \frac{1}{|B|} \int_B g - g(x) \right| + |g(x) - f(x)|. \end{aligned}$$

As $|B| \rightarrow 0$, the middle term tends to zero by continuity of g . Therefore

$$E_\alpha \subseteq \left\{ (f - g)^* > \frac{\alpha}{2} \right\} \cup \left\{ |g - f| > \frac{\alpha}{2} \right\}.$$

The weak $(1, 1)$ estimate and Chebyshev's inequality give

$$|E_\alpha| \leq \frac{2 \cdot 3^d}{\alpha} \|f - g\|_1 + \frac{2}{\alpha} \|f - g\|_1 \leq \frac{2(3^d + 1)}{\alpha} \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, $|E_\alpha| = 0$. □

3.5 Lebesgue Points

The differentiation theorem implies

$$Mf(x) \geq |f(x)| \quad \text{for almost every } x.$$

It also extends to $f \in L_{\text{loc}}^1(\mathbb{R}^d)$ by localization: on $B(0, R)$, replace f by $f \mathbf{1}_{B(0, 2R)}$.

Definition 3.6. A point x is a *Lebesgue point* of f if

$$\lim_{\substack{|B| \rightarrow 0 \\ x \in B}} \frac{1}{|B|} \int_B |f(y) - f(x)| dy = 0.$$

Corollary 3.7. If $f \in L_{\text{loc}}^1(\mathbb{R}^d)$, then almost every point of $\mathbb{R}^d$ is a Lebesgue point of f .

Proof. For each $q \in \mathbb{Q}$, apply the differentiation theorem locally to $|f - q|$, and let N_q be the corresponding null exceptional set. The union $N = \bigcup_{q \in \mathbb{Q}} N_q$ is still null. If $x \notin N$ and $\varepsilon > 0$, choose $q \in \mathbb{Q}$ such that $|f(x) - q| < \varepsilon$. Then

$$\begin{aligned} \limsup_{\substack{|B| \rightarrow 0 \\ x \in B}} \frac{1}{|B|} \int_B |f - f(x)| &\leq \limsup_{\substack{|B| \rightarrow 0 \\ x \in B}} \frac{1}{|B|} \int_B |f - q| + \varepsilon \\ &= |f(x) - q| + \varepsilon < 2\varepsilon. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ proves the result. □

4 Ergodic Theory

4.1 Measure-Preserving Dynamical Systems

Let $(X, \mathcal{A}, \mu)$ be a probability space. Suppose $T : X \rightarrow X$ is measurable and invertible, T^{-1} is measurable, and T preserves measure:

$$\mu(T^{-1}A) = \mu(A) \quad \text{for all } A \in \mathcal{A}, \quad \text{equivalently} \quad T_*\mu = \mu.$$

Then (X, T) is a measure-preserving $\mathbb{Z}$ -system. Its two-sided orbits are

$$\dots, T^{-2}x, T^{-1}x, x, Tx, T^2x, \dots$$

The associated Koopman operator is

$$U_T : L^2(X) \longrightarrow L^2(X), \quad U_T f = f \circ T.$$

Measure preservation implies

$$\int_X f \circ T \, d\mu = \int_X f \, d\mu$$

and hence

$$\int_X |f \circ T|^2 \, d\mu = \int_X |f|^2 \, d\mu.$$

Thus U_T is an isometry; if T is invertible, U_T is unitary.

Example 4.1. For a torus rotation, take

$$\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d, \quad T(x) = x + \alpha \pmod{\mathbb{Z}^d}.$$

This is a model of structured dynamics.

Example 4.2. For a Bernoulli system, take

$$X = \{0, 1\}^{\mathbb{Z}}$$

with a product probability measure and let T be the shift. If $T((x_n)_{n \in \mathbb{Z}}) = (y_n)_{n \in \mathbb{Z}}$, then $y_{n+1} = x_n$. This is a model of random dynamics.

4.2 Ergodicity

Definition 4.3. The system (X, T) is *ergodic* if every measurable invariant set is trivial:

$$TA = A \text{ a.e.} \implies \mu(A) \in \{0, 1\}.$$

Equivalently, every invariant L^2 function is constant:

$$U_T f = f \implies f \text{ is constant almost everywhere.}$$

Indeed, if the set formulation holds and $U_T f = f$, every superlevel set $\{f > c\}$ is invariant and has measure zero or one. This forces f to be constant. Conversely, apply the functional statement to $\mathbf{1}_A$.

4.3 The von Neumann Mean Ergodic Theorem

Theorem 4.4 (von Neumann mean ergodic theorem). *Let (X, T) be an ergodic measure-preserving probability system and let $f \in L^2(X)$. Then*

$$\underbrace{\frac{1}{N} \sum_{n=0}^{N-1} f \circ T^n}_{\text{time average}} \xrightarrow{L^2} \underbrace{\mathbb{E}[f] \mathbf{1}}_{\text{space average}}.$$

Proof. Write

$$A_N f = \frac{1}{N} \sum_{n=0}^{N-1} U_T^n f.$$

Since U_T is an isometry, $\|A_N\|_{L^2 \rightarrow L^2} \leq 1$. Let

$$C = \overline{\text{conv}}\{f \circ T^n : n \geq 0\} \subseteq L^2.$$

This is a nonempty closed convex set. By the nearest-point property in Hilbert spaces, C contains a unique element g of minimal norm.

The operator U_T maps the orbit into itself and therefore maps C into C . Since it is an isometry, $U_T g$ has the same minimal norm as g . Uniqueness gives $U_T g = g$, and ergodicity implies that $g \equiv c$ is constant.

The functional $h \mapsto \int_X h d\mu$ is continuous on L^2 because $\mu(X) = 1$. It has the value $\int f d\mu$ on every element of the orbit, hence on all of C . In particular,

$$c = \mathbb{E}[c] = \mathbb{E}[f].$$

It remains to show that $A_N f \rightarrow g$. The key estimate is

$$\|A_N(U_T f) - A_N f\|_2 = \frac{1}{N} \|f \circ T^N - f\|_2 \leq \frac{2}{N} \|f\|_2.$$

For a finite convex combination

$$h = \sum_{n=0}^M a_n U_T^n f, \quad a_n \geq 0, \quad \sum_{n=0}^M a_n = 1,$$

iteration gives

$$\|A_N h - A_N f\|_2 \leq \frac{2M \|f\|_2}{N}.$$

Given $\varepsilon > 0$, choose such an h with $\|h - g\|_2 < \varepsilon$. Because $A_N g = g$,

$$\|A_N f - g\|_2 \leq \|A_N f - A_N h\|_2 + \|A_N(h - g)\|_2 \leq \frac{2M \|f\|_2}{N} + \varepsilon.$$

Taking the limit superior and then letting $\varepsilon \rightarrow 0$ proves the theorem. □

Remark 4.5. There is also a standard orthogonal-decomposition proof:

$$L^2 = \ker(U_T - I) \oplus \overline{\text{ran}(U_T - I)}.$$

On $\ker(U_T - I)$, the averages are unchanged. If $f = U_T g - g$, then

$$A_N f = \frac{U_T^N g - g}{N} \longrightarrow 0.$$

The result follows by density and the uniform bound $\|A_N\| \leq 1$.

4.4 Birkhoff's Pointwise Ergodic Theorem

Theorem 4.6 (Birkhoff). *Let (X, T) be an ergodic measure-preserving probability system and let $f \in L^1(X)$. Then*

$$\frac{1}{N} \sum_{n=0}^{N-1} f \circ T^n(x) \longrightarrow \mathbb{E}[f] \quad \text{for almost every } x.$$

The proof has two components. First, convergence is easy on a dense class. Functions of the form

$$f = h + (g - g \circ T), \quad h \circ T = h, \quad g \in L^\infty,$$

are dense in the relevant L^2 space, and

$$A_N f = h + \frac{g - g \circ T^N}{N} \longrightarrow h$$

pointwise. Second, the maximal inequality in the next subsection shows that the class of functions with almost-everywhere convergent averages is closed in L^1 .

To see the closure argument, suppose $f_k \rightarrow f$ in L^1 and each $A_N f_k$ converges almost everywhere. Define the oscillation

$$\Omega f(x) = \limsup_N A_N f(x) - \liminf_N A_N f(x) \geq 0.$$

The oscillation is subadditive, and $\Omega f_k = 0$, so

$$\Omega f \leq \Omega(f - f_k) \leq 2(f - f_k)^*.$$

The maximal inequality gives

$$\mu\{\Omega f > \alpha\} \leq \mu\{(f - f_k)^* > \alpha/2\} \leq \frac{2A}{\alpha} \|f - f_k\|_1 \longrightarrow 0.$$

Thus $\Omega f = 0$ almost everywhere. The value of the limit is identified as $\mathbb{E}[f]$ using the von Neumann theorem and an approximation argument.

4.5 The Maximal Ergodic Estimate

Define

$$M_T f(x) = \sup_{N \geq 1} \frac{1}{N} \sum_{n=0}^{N-1} |f \circ T^n(x)|.$$

Theorem 4.7 (Maximal ergodic weak-type estimate). *Let (X, μ) be a σ -finite measure space, let T be measure preserving, and let $f \in L^1(X)$. Then $M_T f(x) < \infty$ for almost every x , and*

there is a universal constant $A > 0$ such that

$$\mu\{x : M_T f(x) > \alpha\} \leq \frac{A}{\alpha} \|f\|_{L^1} \quad (\alpha > 0).$$

It is enough to consider $f \geq 0$.

Step 1: The Case $X = \mathbb{Z}$

Let $X = \mathbb{Z}$ with counting measure, $T(n) = n + 1$, and $f \geq 0$. Embed the discrete problem into $\mathbb{R}$ by defining

$$\tilde{f}(x) = f(n) \quad \text{when } n \leq x < n + 1.$$

If $E \subseteq \mathbb{Z}$, define

$$\tilde{E} = \bigcup_{n \in E} [n, n + 1).$$

Then

$$|\tilde{E}| = \#E, \quad \int_{\mathbb{R}} \tilde{f}(x) dx = \sum_{n \in \mathbb{Z}} f(n) = \|f\|_{\ell^1}.$$

Moreover,

$$\sum_{k=0}^{N-1} f(n+k) = \int_0^N \tilde{f}(n+t) dt.$$

If $x \in [n, n+1)$, then $[n, n+N] \subseteq [x-1, x+N]$, and therefore

$$\int_0^N \tilde{f}(n+t) dt \leq \int_{-1}^N \tilde{f}(x+t) dt.$$

Since $N \geq 1$, this yields

$$\frac{1}{N} \sum_{k=0}^{N-1} f(n+k) \leq \frac{N+1}{N} \frac{1}{N+1} \int_{-1}^N \tilde{f}(x+t) dt \leq 2 \tilde{f}^{*,\text{HL}}(x),$$

where $\tilde{f}^{*,\text{HL}}$ is the one-dimensional uncentered Hardy–Littlewood maximal function. Taking the supremum over N ,

$$\boxed{M_T f(n) \leq 2 \tilde{f}^{*,\text{HL}}(x) \quad \text{for every } x \in [n, n+1)}.$$

Consequently,

$$\begin{aligned} \#\{n : M_T f(n) > \alpha\} &\leq \left| \left\{ x : \tilde{f}^{*,\text{HL}}(x) > \frac{\alpha}{2} \right\} \right| \\ &\leq \frac{6}{\alpha} \|f\|_{\ell^1(\mathbb{Z})}. \end{aligned}$$

Thus the discrete estimate holds with $A = 6$.

Step 2: Transference from $\mathbb{Z}$ to X

For each N , define the truncated maximal function

$$M_{T,N} f(x) = \sup_{1 \leq m \leq N} \frac{1}{m} \sum_{k=0}^{m-1} f \circ T^k(x).$$

Since $M_{T,N}f \nearrow M_Tf$, it suffices to prove

$$\mu\{M_{T,N}f > \alpha\} \leq \frac{A}{\alpha} \|f\|_{L^1}$$

with A independent of N .

Fix N and write $M_Tf = M_{T,N}f$ temporarily. On $\mathbb{Z}$, set

$$M(F)(n) = \sup_{m \geq 1} \frac{1}{m} \sum_{k=0}^{m-1} F(n+k).$$

Encode each orbit in $X \times \mathbb{Z}$ by

$$F(x, n) = \begin{cases} f \circ T^n(x), & n \geq 0, \\ 0, & n < 0. \end{cases}$$

Then

$$A_m f(T^n x) = \frac{1}{m} \sum_{k=0}^{m-1} F(x, n+k).$$

Choose a positive integer a , let $B = a + N$, and truncate once more:

$$F_B(x, n) = \begin{cases} F(x, n), & n < B, \\ 0, & n \geq B. \end{cases}$$

If $m \leq N$ and $0 \leq n < a$, then $n+k < B$ for $0 \leq k < m$. Hence

$$A_m f(T^n x) \leq M(F_B(x, \cdot))(n),$$

and therefore

$$\boxed{M_T f(T^n x) \leq M(F_B(x, \cdot))(n) \quad \text{for } 0 \leq n < a.}$$

Let $E_\alpha = \{x : M_T f(x) > \alpha\}$. Measure preservation gives

$$\mu(E_\alpha) = \mu\{x : M_T f(T^n x) > \alpha\}$$

for every n . Summing over $0 \leq n < a$, applying Fubini–Tonelli, and then using the discrete estimate,

$$\begin{aligned} a\mu(E_\alpha) &= \int_X \#\{0 \leq n < a : M_T f(T^n x) > \alpha\} d\mu \\ &\leq \int_X \#\{n \in \mathbb{Z} : M(F_B(x, \cdot))(n) > \alpha\} d\mu \\ &\leq \int_X \frac{A}{\alpha} \|F_B(x, \cdot)\|_{\ell^1(\mathbb{Z})} d\mu \\ &= \frac{A}{\alpha} \int_X \sum_{k=0}^{B-1} f \circ T^k(x) d\mu \\ &= \frac{A}{\alpha} (a + N) \|f\|_{L^1}. \end{aligned}$$

Dividing by a and letting $a \rightarrow \infty$ gives

$$\mu(E_\alpha) \leq \frac{A}{\alpha} \|f\|_{L^1}.$$

The constant $A = 6$ is independent of N . Letting $N \rightarrow \infty$ proves the full maximal estimate.

4.6 Structural Summary

The main implications can be summarized as follows:

- The Vitali covering lemma implies the Hardy–Littlewood weak type $(1, 1)$ estimate, which implies Lebesgue differentiation.
- The Hardy–Littlewood weak estimate and Marcinkiewicz interpolation imply strong L^p bounds.
- The discrete maximal estimate and transference imply the maximal ergodic estimate.
- The maximal ergodic estimate and von Neumann’s L^2 convergence imply Birkhoff’s pointwise theorem.

The common principle is

maximal inequality + convergence on a dense subclass $\implies$ almost-everywhere convergence
--

Part II

Weighted Estimates and Singular Integrals

5 Weighted Maximal Function Theory

5.1 Review and Motivation

Recall the uncentered Hardy–Littlewood maximal function

$$Mf(x) = \sup_{x \in B} \frac{1}{|B|} \int_B |f(y)| dy.$$

It satisfies

$$|\{Mf > \lambda\}| \leq \frac{C}{\lambda} \|f\|_{L^1}, \quad \|Mf\|_\infty \leq \|f\|_\infty,$$

and Marcinkiewicz interpolation consequently gives

$$\|Mf\|_p \leq C_p \|f\|_p, \quad 1 < p \leq \infty.$$

The argument extends in several directions: balls may be replaced by cubes, Lebesgue measure may be replaced by a doubling measure, and $\mathbb{R}^d$ may be replaced by a suitable metric space.

Definition 5.1. A measure μ on $\mathbb{R}^d$ is *doubling* if there is a constant C such that

$$\mu(2B) \leq C\mu(B)$$

for every ball B , where $2B$ is the concentric ball with twice the radius.

For a doubling measure, one may define

$$M_\mu f(x) = \sup_{x \in B} \frac{1}{\mu(B)} \int_B |f(y)| d\mu(y).$$

In this section, however, M always denotes the maximal operator whose averages are taken with respect to Lebesgue measure. We ask when this operator is bounded on a weighted space. Let

$$d\mu(x) = w(x) dx, \quad w \geq 0,$$

and use the notation

$$w(E) = \int_E w(x) dx, \quad f_B = \frac{1}{|B|} \int_B f(x) dx.$$

The problem is to characterize those w for which

$$M : L^p(w) \longrightarrow L^p(w).$$

5.2 Muckenhoupt's Theorem

Theorem 5.2 (Muckenhoupt). *Let $1 < p < \infty$ and $w \geq 0$. The following statements are equivalent:*

1. *There is $C > 0$ such that*

$$\|Mf\|_{L^p(w)} \leq C \|f\|_{L^p(w)}.$$

2. *There is $C > 0$ such that*

$$w(\{Mf > \lambda\}) \leq C \frac{\|f\|_{L^p(w)}^p}{\lambda^p}.$$

3. *There is $C > 0$ such that, for every ball B and every $f \geq 0$,*

$$\left(\frac{f(B)}{|B|} \right)^p \leq \frac{C}{w(B)} \int_B f^p w dx, \quad f(B) = \int_B f dx.$$

4. *The A_p characteristic*

$$[w]_{A_p} := \sup_B \left(\frac{1}{|B|} \int_B w dx \right) \left(\frac{1}{|B|} \int_B w^{-p'/p} dx \right)^{p/p'}$$

is finite, where $1/p + 1/p' = 1$.

Definition 5.3. We write $w \in A_p$ if $[w]_{A_p} < \infty$.

Applying the third condition to $f = \mathbf{1}_E$, where $E \subseteq B$, gives

$$\left[\left(\frac{|E|}{|B|} \right)^p \leq C \frac{w(E)}{w(B)} \right].$$

In particular, every A_p weight is doubling. Indeed, take $E = \frac{1}{2}B$ to obtain

$$2^{-dp} \leq C \frac{w(\frac{1}{2}B)}{w(B)},$$

which is equivalent, after rescaling, to $w(2B) \leq C'w(B)$.

5.3 The Elementary Implications

Strong Type Implies Weak Type

Chebyshev's inequality gives

$$\lambda^p w(\{Mf > \lambda\}) \leq \int_{\{Mf > \lambda\}} (Mf)^p w \, dx \leq \|Mf\|_{L^p(w)}^p \leq C \|f\|_{L^p(w)}^p.$$

Weak Type Implies the Local Averaging Inequality

Fix a ball B , and replace f by $f\mathbf{1}_B$ if necessary. For every $x \in B$,

$$Mf(x) \geq \frac{1}{|B|} \int_B |f|.$$

If $\lambda < f(B)/|B|$, then $B \subseteq \{Mf > \lambda\}$, and hence

$$w(B) \leq w(\{Mf > \lambda\}) \leq C \frac{\|f\|_{L^p(w)}^p}{\lambda^p}.$$

Letting $\lambda \uparrow f(B)/|B|$ yields

$$\left(\frac{f(B)}{|B|} \right)^p \leq \frac{C}{w(B)} \int_B f^p w \, dx.$$

The Local Averaging Inequality Implies A_p

Take $f = w^{-p'/p}$. More precisely, first use

$$f = (w + \delta)^{-p'/p} \mathbf{1}_B$$

and then let $\delta \downarrow 0$. Since $p'/p = p' - 1$,

$$f^p w = w^{-p'} w = w^{1-p'} = w^{-p'/p} = f.$$

The local averaging inequality gives

$$\left(\frac{1}{|B|} \int_B w^{-p'/p} \right)^p \leq \frac{C}{w(B)} \int_B w^{-p'/p}.$$

Multiplying by $w(B)$ and dividing by the final integral,

$$|B|^{-p} \left(\int_B w^{-p'/p} \right)^{p-1} w(B) \leq C.$$

Since $p - 1 = p/p'$, this is precisely the A_p condition.

A_p Implies the Local Averaging Inequality

Hölder's inequality gives

$$f(B) = \int_B f w^{1/p} w^{-1/p} \leq \left(\int_B f^p w \right)^{1/p} \left(\int_B w^{-p'/p} \right)^{1/p'}.$$

Raising to the p -th power and dividing by $|B|^p$,

$$\left(\frac{f(B)}{|B|}\right)^p \leq \frac{1}{|B|^p} \left(\int_B f^p w\right) \left(\int_B w^{-p'/p}\right)^{p/p'}.$$

Because $1 + p/p' = p$, the A_p condition implies

$$\frac{w(B)}{|B|^p} \left(\int_B w^{-p'/p}\right)^{p/p'} \leq [w]_{A_p},$$

which proves the desired inequality.

The difficult implication is $w \in A_p \Rightarrow M : L^p(w) \rightarrow L^p(w)$. We prove it first for the dyadic maximal operator. We shall use the dual weight

$$\sigma = w^{-p'/p}.$$

Directly interchanging the two factors in the A_p condition shows that

$$w \in A_p \implies \sigma \in A_{p'}, \quad [\sigma]_{A_{p'}} = [w]_{A_p}^{p'/p}.$$

5.4 Dyadic Cubes and the Dyadic Maximal Operator

Let

$$\mathcal{D} = \left\{ 2^k \ell + [0, 2^k)^d : (\ell, k) \in \mathbb{Z}^d \times \mathbb{Z} \right\}$$

be the standard dyadic grid, and define

$$M_{\text{dyad}} f(x) = \sup_{\substack{Q \ni x \\ Q \in \mathcal{D}}} \frac{1}{|Q|} \int_Q |f(y)| dy.$$

If $Q_1, Q_2 \in \mathcal{D}$ intersect, then one contains the other. The dyadic parent $\pi_{\mathcal{D}}(Q)$ is the smallest dyadic cube strictly containing Q , and

$$|\pi_{\mathcal{D}}(Q)| = 2^d |Q|.$$

5.5 A Dyadic Calderón–Zygmund Decomposition

Lemma 5.4. *Let $f \in L^1(\mathbb{R}^d)$ and $\lambda > 0$. There is a pairwise disjoint family $\mathcal{B}$ of dyadic cubes such that*

(1)

$$\lambda < \frac{1}{|Q|} \int_Q |f| \leq 2^d \lambda \quad (Q \in \mathcal{B});$$

(2)

$$\sum_{Q \in \mathcal{B}} |Q| \leq \frac{1}{\lambda} \|f\|_{L^1};$$

(3)

$$\{M_{\text{dyad}} f > \lambda\} = \bigcup_{Q \in \mathcal{B}} Q.$$

Moreover, if

$$g = f \mathbf{1}_{\left(\bigcup_{Q \in \mathcal{B}} Q\right)^c}, \quad b = f - g = \sum_{Q \in \mathcal{B}} f \mathbf{1}_Q,$$

then $|g| \leq \lambda$ almost everywhere.

Proof. Let $\mathcal{B}$ be the family of maximal dyadic cubes satisfying

$$\frac{1}{|Q|} \int_Q |f| > \lambda.$$

Such maximal cubes exist because

$$\frac{1}{|Q|} \int_Q |f| \leq \frac{\|f\|_1}{|Q|} \rightarrow 0$$

as the side length tends to infinity. Maximality and the nesting property imply that the cubes in $\mathcal{B}$ are pairwise disjoint.

For $Q \in \mathcal{B}$, its parent does not satisfy the defining inequality. Therefore

$$\lambda \geq \frac{1}{|\pi_{\mathcal{D}}(Q)|} \int_{\pi_{\mathcal{D}}(Q)} |f| \geq \frac{1}{2^d |Q|} \int_Q |f|,$$

which proves the upper bound in (1).

Every bad cube lies in the superlevel set in (3). Conversely, if $M_{\text{dyad}} f(x) > \lambda$, then a dyadic cube containing x satisfies the bad-average condition and is contained in a maximal such cube. This proves (3). Finally, disjointness and the lower bound in (1) give

$$\sum_{Q \in \mathcal{B}} |Q| \leq \frac{1}{\lambda} \sum_{Q \in \mathcal{B}} \int_Q |f| \leq \frac{1}{\lambda} \|f\|_1.$$

Outside the union of the bad cubes, every dyadic average containing x is at most λ . The Lebesgue differentiation theorem then gives $|f(x)| \leq \lambda$ almost everywhere there. $\square$

5.6 Weighted Boundedness of the Dyadic Maximal Operator

Theorem 5.5. *If $1 < p < \infty$ and $w \in A_p$, then*

$$M_{\text{dyad}} : L^p(w) \longrightarrow L^p(w)$$

is bounded.

Proof. Choose $c_0 > 2^{d+1} + 1$. For each $k \in \mathbb{Z}$, let $\mathcal{B}_k$ be the family of bad cubes from the preceding decomposition at height c_0^k , and set

$$\mathcal{B} = \bigcup_{k \in \mathbb{Z}} \mathcal{B}_k.$$

For $Q \in \mathcal{B}$, define

$$E(Q) = Q \setminus \bigcup_{\substack{Q' \in \mathcal{B} \\ Q' \subsetneq Q}} Q'.$$

We first claim that

$$|E(Q)| \geq \frac{1}{2} |Q|.$$

Suppose $Q \in \mathcal{B}_k$. For $j \leq k$, the cube Q lies in a maximal cube of $\mathcal{B}_j$, so no member of $\mathcal{B}_j$ is strictly contained in Q . For $j \geq k+1$, disjointness within $\mathcal{B}_j$ and the Calderón–Zygmund bounds give

$$\sum_{\substack{Q' \in \mathcal{B}_j \\ Q' \subseteq Q}} |Q'| \leq \frac{1}{c_0^j} \int_Q |f| \leq \frac{2^d c_0^k}{c_0^j} |Q|.$$

Summing over $j \geq k+1$,

$$\left| \bigcup_{\substack{Q' \in \mathcal{B} \\ Q' \subsetneq Q}} Q' \right| \leq 2^d |Q| \sum_{j \geq k+1} c_0^{k-j} = 2^d |Q| \frac{c_0^{-1}}{1 - c_0^{-1}} \leq \frac{1}{2} |Q|.$$

This proves the claim. The sets $E(Q)$ are pairwise disjoint.

If $x \in E(Q)$ with $Q \in \mathcal{B}_k$, then x belongs to no cube in $\mathcal{B}_{k+1}$, and therefore

$$M_{\text{dyad}} f(x) \leq c_0^{k+1}.$$

Consequently,

$$\begin{aligned} \int |M_{\text{dyad}} f|^p w \, dx &\leq \sum_k \sum_{Q \in \mathcal{B}_k} c_0^{(k+1)p} w(E(Q)) \\ &\leq C_p \sum_k \sum_{Q \in \mathcal{B}_k} \left(\frac{1}{|Q|} \int_Q |f| \right)^p w(E(Q)). \end{aligned}$$

Introduce $\sigma = w^{-p'/p}$. We rewrite

$$\frac{1}{|Q|} \int_Q |f| = \left(\frac{1}{\sigma(Q)} \int_Q |f| \sigma^{-1} \sigma \, dy \right) \frac{\sigma(Q)}{|Q|}.$$

Thus the preceding sum equals

$$\begin{aligned} \sum_{Q \in \mathcal{B}} \left(\frac{1}{\sigma(Q)} \int_Q |f| \sigma^{-1} \sigma \right)^p \sigma(E(Q)) \\ \times \underbrace{\frac{w(E(Q))}{\sigma(E(Q))} \left(\frac{\sigma(Q)}{|Q|} \right)^p}_I. \end{aligned}$$

Since $\sigma \in A_{p'}$, the set estimate for $A_{p'}$ weights and $|E(Q)| \geq |Q|/2$ imply

$$\sigma(E(Q)) \geq c \sigma(Q).$$

Also $w(E(Q)) \leq w(Q)$, so

$$\begin{aligned} I &\leq C \frac{w(Q)}{\sigma(Q)} \frac{\sigma(Q)^p}{|Q|^p} \\ &= C \left(\frac{1}{|Q|} \int_Q w \right) \left(\frac{1}{|Q|} \int_Q \sigma \right)^{p-1} \\ &\leq C[w]_{A_p}. \end{aligned}$$

For $y \in E(Q) \subseteq Q$,

$$\frac{1}{\sigma(Q)} \int_Q |f\sigma^{-1}| \sigma \leq M_\sigma(f\sigma^{-1})(y).$$

Using the disjointness of the $E(Q)$ and the $L^p(\sigma)$ boundedness of M_σ , which holds because σ is doubling, we obtain

$$\begin{aligned} \int |M_{\text{dyad}} f|^p w &\leq C_{p,w} \sum_{Q \in \mathcal{B}} \int_{E(Q)} (M_\sigma(f\sigma^{-1}))^p \sigma \, dy \\ &\leq C_{p,w} \int_{\mathbb{R}^d} (M_\sigma(f\sigma^{-1}))^p \sigma \, dy \\ &\leq C_{p,w} \int_{\mathbb{R}^d} |f\sigma^{-1}|^p \sigma \, dy. \end{aligned}$$

Finally,

$$\sigma^{1-p} = w^{-\frac{p'}{p}(1-p)} = w,$$

and hence

$$\int |M_{\text{dyad}} f|^p w \leq C_{p,w} \int |f|^p w.$$

□

6 The Hilbert Transform

6.1 Singular Kernels and Singular Integrals

Definition 6.1. A kernel

$$K : \mathbb{R}^d \times \mathbb{R}^d \setminus \{(x, x) : x \in \mathbb{R}^d\} \longrightarrow \mathbb{C}$$

is singular if

$$|K(x, y)| \sim |x - y|^{-d} \quad \text{as } |x - y| \rightarrow 0.$$

A singular integral is formally an operator of the form

$$Tf(x) = \int K(x, y)f(y) \, dy.$$

Such operators occur throughout partial differential equations, ergodic theory, and signal processing.

6.2 Definition of the Hilbert Transform

For the Hilbert transform, the convolution kernel is

$$h(x) = \frac{1}{\pi x}, \quad K(x, y) = h(x - y).$$

Since $1/x \notin L^1_{\text{loc}}(\mathbb{R})$, the integral must be interpreted as a principal value.

Recall that the Schwartz space $\mathcal{S}(\mathbb{R})$ is the Fréchet space defined by the seminorms

$$\|f\|_{n,m} = \sup_{x \in \mathbb{R}} |x^m f^{(n)}(x)|.$$

Define the principal-value distribution w_0 by

$$\langle w_0, f \rangle = \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon \leq |x| \leq 1} \frac{f(x)}{x} dx + \frac{1}{\pi} \int_{|x| \geq 1} \frac{f(x)}{x} dx.$$

This is well defined because

$$\int_{\varepsilon \leq |x| \leq 1} \frac{f(x)}{x} dx = \int_{\varepsilon \leq |x| \leq 1} \frac{f(x) - f(0)}{x} dx,$$

and

$$\left| \frac{f(x) - f(0)}{x} \right| \leq \sup_x |f'(x)|.$$

The integral at infinity converges because Schwartz functions decay rapidly.

Definition 6.2. For $f \in L^p(\mathbb{R})$, $1 \leq p < \infty$, define the truncated Hilbert transform by

$$H^{(\varepsilon)} f(x) = \frac{1}{\pi} \int_{|y| \geq \varepsilon} \frac{f(x-y)}{y} dy = \frac{1}{\pi} \int_{|x-y| \geq \varepsilon} \frac{f(y)}{x-y} dy.$$

The truncated integral is well defined by Hölder's inequality. For $f \in \mathcal{S}(\mathbb{R})$,

$$Hf(x) = (w_0 * f)(x) = \lim_{\varepsilon \rightarrow 0} H^{(\varepsilon)} f(x).$$

6.3 The Fourier Multiplier

Proposition 6.3. *With the convention*

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx,$$

one has

$$\widehat{w_0}(\xi) = -i \operatorname{sgn}(\xi).$$

Proof. For $\varphi \in \mathcal{S}(\mathbb{R})$,

$$\begin{aligned} \langle \widehat{w_0}, \varphi \rangle &= \langle w_0, \widehat{\varphi} \rangle \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon \leq |\xi| \leq 1/\varepsilon} \widehat{\varphi}(\xi) \frac{d\xi}{\xi} \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \varphi(x) \left[\frac{1}{\pi} \int_{\varepsilon \leq |\xi| \leq 1/\varepsilon} e^{-2\pi i x \xi} \frac{d\xi}{\xi} \right] dx. \end{aligned}$$

The cosine term vanishes by oddness. Therefore the expression in brackets is

$$-\frac{i}{\pi} \int_{\varepsilon \leq |\xi| \leq 1/\varepsilon} \frac{\sin(2\pi x \xi)}{\xi} d\xi = -\frac{2i}{\pi} \operatorname{sgn}(x) \int_{2\pi \varepsilon |x|}^{2\pi |x|/\varepsilon} \frac{\sin u}{u} du.$$

The Dirichlet integral

$$\int_0^\infty \frac{\sin u}{u} du = \frac{\pi}{2}$$

shows that this converges pointwise for $x \neq 0$ to $-i \operatorname{sgn}(x)$. The truncated integrals are uniformly

bounded, so dominated convergence yields

$$\langle \widehat{w_0}, \varphi \rangle = \int_{\mathbb{R}} \varphi(x) (-i \operatorname{sgn}(x)) dx.$$

□

Thus

$$Hf = \left[-i \operatorname{sgn}(\xi) \widehat{f}(\xi) \right]^\vee.$$

Plancherel's theorem immediately gives

$$\|Hf\|_2 = \|f\|_2, \quad H^2 = -I, \quad H^* = -H.$$

6.4 L^p Boundedness

Theorem 6.4. *For every $1 < p < \infty$, the Hilbert transform is bounded on $L^p(\mathbb{R})$.*

Proof. By duality and $H^* = -H$, it suffices to treat $1 < p \leq 2$. Marcinkiewicz interpolation and the L^2 isometry reduce the problem to the weak-type estimate

$$|\{x : |Hf(x)| > \lambda\}| \leq \frac{C}{\lambda} \|f\|_1.$$

Perform a one-dimensional Calderón–Zygmund decomposition of f at height λ . There are pairwise disjoint dyadic intervals $\{I_j\}$ such that

$$\lambda < \frac{1}{|I_j|} \int_{I_j} |f| \leq 2\lambda.$$

Let $\Omega = \bigcup_j I_j$, and write $f = g + b$, where

$$g(x) = \begin{cases} f(x), & x \notin \Omega, \\ \frac{1}{|I_j|} \int_{I_j} f, & x \in I_j, \end{cases}$$

and

$$b = \sum_j b_j, \quad b_j = \left(f - \frac{1}{|I_j|} \int_{I_j} f \right) \mathbf{1}_{I_j}.$$

Then

$$\int_{I_j} b_j = 0, \quad |\Omega| \leq \frac{1}{\lambda} \|f\|_1.$$

For the good part, Chebyshev's inequality and the L^2 isometry give

$$|\{|Hg| > \lambda/2\}| \leq \frac{4}{\lambda^2} \|Hg\|_2^2 = \frac{4}{\lambda^2} \|g\|_2^2.$$

Outside Ω , one has $g = f$ and $|f| \leq \lambda$ almost everywhere; on each I_j , one has $|g| \leq 2\lambda$. Hence

$$\begin{aligned} \int |g|^2 &\leq \lambda \int_{\mathbb{R} \setminus \Omega} |f| + 4\lambda^2 |\Omega| \\ &\leq C\lambda \|f\|_1. \end{aligned}$$

It follows that

$$|\{|Hg| > \lambda/2\}| \leq \frac{C}{\lambda} \|f\|_1.$$

For the bad part, let $I_j^* = 2I_j$ and

$$E = \bigcup_j I_j^*.$$

Then

$$|E| \leq 2|\Omega| \leq \frac{2}{\lambda} \|f\|_1.$$

Let c_j be the center of I_j . If $x \notin I_j^*$, the cancellation $\int b_j = 0$ gives

$$\begin{aligned} |Hb_j(x)| &= \left| \frac{1}{\pi} \int_{I_j} \left(\frac{1}{x-y} - \frac{1}{x-c_j} \right) b_j(y) dy \right| \\ &\leq C \int_{I_j} \frac{|y-c_j|}{|x-c_j|^2} |b_j(y)| dy \\ &\leq C \frac{|I_j|}{|x-c_j|^2} \|b_j\|_1. \end{aligned}$$

Therefore, for $x \in E^c$,

$$|Hb(x)| \leq C \sum_j \frac{|I_j|}{|x-c_j|^2} \|b_j\|_1.$$

Chebyshev's inequality now gives

$$\begin{aligned} |\{x \notin E : |Hb(x)| > \lambda/2\}| &\leq \frac{2}{\lambda} \int_{\mathbb{R} \setminus E} |Hb(x)| dx \\ &\leq \frac{C}{\lambda} \sum_j \|b_j\|_1 \int_{|x-c_j| \geq |I_j|} \frac{|I_j|}{|x-c_j|^2} dx. \end{aligned}$$

Since

$$\int_{|x-c_j| \geq |I_j|} \frac{|I_j|}{|x-c_j|^2} dx = 2$$

and

$$\sum_j \|b_j\|_1 \leq 2 \|f\|_1,$$

we obtain

$$|\{x \notin E : |Hb(x)| > \lambda/2\}| \leq \frac{C}{\lambda} \|f\|_1.$$

Combining the good and bad estimates proves weak type $(1, 1)$, and interpolation and duality complete the proof. $\square$

7 The Fourier Transform on Euclidean Space

Throughout this section,

$$\widehat{f}(\xi) = \int_{\mathbb{R}^d} e^{-2\pi i x \cdot \xi} f(x) dx.$$

7.1 Convolution and the Multiplication Formula

Proposition 7.1. (1) If $f, g \in L^1(\mathbb{R}^d)$, then

$$\widehat{f * g}(\xi) = \widehat{f}(\xi)\widehat{g}(\xi).$$

(2) If $f, g \in \mathcal{S}(\mathbb{R}^d)$, then

$$\int_{\mathbb{R}^d} f(x)\widehat{g}(x) dx = \int_{\mathbb{R}^d} \widehat{f}(y)g(y) dy.$$

Proof. Absolute integrability permits Fubini's theorem:

$$\begin{aligned} \widehat{f * g}(\xi) &= \int e^{-2\pi i x \cdot \xi} \int f(x - y)g(y) dy dx \\ &= \int e^{-2\pi i y \cdot \xi} g(y) \int f(x - y)e^{-2\pi i (x - y) \cdot \xi} dx dy \\ &= \widehat{f}(\xi)\widehat{g}(\xi). \end{aligned}$$

Similarly,

$$\begin{aligned} \int f(x)\widehat{g}(x) dx &= \iint f(x)g(y)e^{-2\pi i y \cdot x} dy dx \\ &= \int \widehat{f}(y)g(y) dy. \end{aligned}$$

□

7.2 Basic Examples and Decay

In the sense of tempered distributions,

$$\widehat{\delta}_0 = 1.$$

If $f \in C_c^\infty(\mathbb{R}^d)$, then $\widehat{f} \in \mathcal{S}(\mathbb{R}^d)$, but $\widehat{f}$ is not compactly supported unless $f = 0$. Indeed, $\widehat{f}$ extends to an entire function. This is an elementary manifestation of the uncertainty principle: localization in one domain produces spreading in the other.

Proposition 7.2. If $f \in L^1(\mathbb{R}^d) \cap C_c^k(\mathbb{R}^d)$, then

$$|\widehat{f}(\xi)| \leq \frac{C_f}{\langle \xi \rangle^k}, \quad \langle \xi \rangle = (1 + |\xi|^2)^{1/2}.$$

Proof. Integration by parts in the first coordinate gives

$$\begin{aligned} \widehat{f}(\xi) &= \int f(x) \frac{\partial}{\partial x_1} \left(e^{-2\pi i x \cdot \xi} \right) \frac{1}{-2\pi i \xi_1} dx \\ &= \frac{1}{2\pi i \xi_1} \int \frac{\partial f}{\partial x_1}(x) e^{-2\pi i x \cdot \xi} dx. \end{aligned}$$

The boundary term vanishes because f is compactly supported. Iterate k times in a coordinate direction for which $|\xi_j|$ is maximal. Together with the trivial L^1 estimate for small ξ , this proves the result. □

7.3 The Gaussian

Proposition 7.3. *The Gaussian is fixed by the Fourier transform:*

$$\widehat{e^{-\pi|x|^2}}(\xi) = e^{-\pi|\xi|^2}.$$

Proof. The product structure and Fubini's theorem reduce the proof to one dimension. Completing the square,

$$-2\pi ix\xi - \pi x^2 = -\pi(x + i\xi)^2 - \pi\xi^2.$$

Thus

$$\widehat{e^{-\pi x^2}}(\xi) = e^{-\pi\xi^2} \int_{\mathbb{R}} e^{-\pi(x+i\xi)^2} dx.$$

The function $e^{-\pi z^2}$ is entire. Integrate it around the rectangle with vertices

$$-R, \quad R, \quad R + i\xi, \quad -R + i\xi.$$

The two vertical integrals have modulus at most

$$|\xi|e^{-\pi R^2 + \pi\xi^2},$$

which tends to zero as $R \rightarrow \infty$. Hence

$$\int_{\mathbb{R}} e^{-\pi(x+i\xi)^2} dx = \int_{\mathbb{R}} e^{-\pi x^2} dx = 1,$$

and the claim follows. □

7.4 Fourier Inversion on the Schwartz Space

Theorem 7.4. *The Fourier transform*

$$\mathcal{F} : \mathcal{S}(\mathbb{R}^d) \longrightarrow \mathcal{S}(\mathbb{R}^d)$$

is a topological isomorphism. Its inverse is

$$\mathcal{F}^{-1}f(x) = \int_{\mathbb{R}^d} e^{2\pi i x \cdot \xi} f(\xi) d\xi.$$

Proof. The formal computation

$$\mathcal{F}^{-1}\mathcal{F}f(x) = \int f(y) \left(\int e^{2\pi i(x-y) \cdot \xi} d\xi \right) dy$$

is invalid because the inner integral is not absolutely convergent. We regularize it with a Gaussian:

$$\mathcal{F}^{-1}\mathcal{F}f(x) = \lim_{\varepsilon \rightarrow 0} \int e^{2\pi i x \cdot \xi} \widehat{f}(\xi) e^{-\pi\varepsilon^2|\xi|^2} d\xi.$$

Since $\widehat{f} \in L^1$, dominated convergence justifies this limit. Fubini's theorem is now applicable:

$$\mathcal{F}^{-1}\mathcal{F}f(x) = \lim_{\varepsilon \rightarrow 0} \int f(y) \int e^{2\pi i(x-y) \cdot \xi} e^{-\pi\varepsilon^2|\xi|^2} d\xi dy.$$

The scaling law for the Gaussian gives

$$\int e^{2\pi i(x-y)\cdot\xi} e^{-\pi\varepsilon^2|\xi|^2} d\xi = \frac{1}{\varepsilon^d} e^{-\pi|(y-x)/\varepsilon|^2} =: G_\varepsilon(x-y).$$

The family $\{G_\varepsilon\}$ is an approximate identity:

$$G_\varepsilon \geq 0, \quad \int G_\varepsilon = 1,$$

and its mass concentrates at the origin. Therefore

$$\mathcal{F}^{-1}\mathcal{F}f(x) = \lim_{\varepsilon \rightarrow 0} \int f(y) G_\varepsilon(x-y) dy = f(x).$$

The standard seminorm estimates show continuity of $\mathcal{F}$ and its inverse on the Schwartz space. $\square$

7.5 Plancherel's Theorem

Theorem 7.5 (Plancherel). *The Fourier transform extends uniquely to a unitary operator on $L^2(\mathbb{R}^d)$. In particular,*

$$\|\widehat{f}\|_2 = \|f\|_2, \quad \langle \widehat{f}, \widehat{g} \rangle_{L^2} = \langle f, g \rangle_{L^2}.$$

Proof. First let $f \in \mathcal{S}$, and set

$$g = \mathcal{F}^{-1}(\overline{f}).$$

Then

$$g(x) = \int e^{2\pi i x \cdot \xi} \overline{f(\xi)} d\xi = \overline{\widehat{f}(x)},$$

and $\widehat{g} = \overline{f}$. Applying the multiplication formula,

$$\|f\|_2^2 = \int f \overline{f} = \int f \widehat{g} = \int \widehat{f} g = \int |\widehat{f}|^2 = \|\widehat{f}\|_2^2.$$

The polarization identity gives preservation of inner products. Since $\mathcal{S}$ is dense in L^2 , $\mathcal{F}$ extends uniquely to an isometry on L^2 . Fourier inversion on $\mathcal{S}$ implies that its range is dense and closed, hence all of L^2 . $\square$

7.6 The Hausdorff–Young Theorem

Theorem 7.6 (Hausdorff–Young). *If $1 \leq p \leq 2$, then the Fourier transform extends to a bounded operator*

$$\mathcal{F} : L^p(\mathbb{R}^d) \longrightarrow L^{p'}(\mathbb{R}^d), \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

Proof. At the endpoints,

$$\mathcal{F} : L^1 \rightarrow L^\infty, \quad \|\widehat{f}\|_\infty \leq \|f\|_1,$$

and Plancherel gives

$$\mathcal{F} : L^2 \rightarrow L^2$$

isometrically. Riesz–Thorin interpolation gives

$$\frac{1}{p} = \frac{1-\theta}{1} + \frac{\theta}{2}, \quad \frac{1}{q} = \frac{1-\theta}{\infty} + \frac{\theta}{2} = \frac{\theta}{2}.$$

Thus $1/p + 1/q = 1$, so $q = p'$. $\square$

Finally, the Fourier transform extends to the space of tempered distributions $\mathcal{S}'$ by duality:

$$\langle \widehat{T}, \varphi \rangle = \langle T, \widehat{\varphi} \rangle.$$

Part III

Randomization and Pointwise Fourier Convergence

8 Probability and Randomization

8.1 Basic probability theory

Probability space $(\Omega, \Sigma, \mathbb{P})$, with $A \in \Sigma$ the *events*.

INDEPENDENCE.

- A, B are independent if $\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$.
- Let $\Sigma_1, \Sigma_2 \subseteq \Sigma$ be two σ -subalgebras. Σ_1, Σ_2 are independent if for all $A \in \Sigma_1, B \in \Sigma_2$, A and B are independent.

RANDOM VARIABLES.

- A random variable is a measurable map $X : \Omega \rightarrow \mathbb{R}$.
- $\sigma(X) \subseteq \Sigma$ is the smallest σ -algebra making X measurable.
- X_1, X_2 are independent random variables if $\sigma(X_1), \sigma(X_2)$ are independent.

EXAMPLE (RADEMACHER FUNCTIONS). $\Omega = [0, 1]$, Σ = Borel algebra, $\mathbb{P}$ = Lebesgue measure. For $j \geq 0$, define $r_j : \Omega \rightarrow \mathbb{R}$ by

$$r_j(t) = \operatorname{sgn}(\sin(2\pi 2^j t)).$$

These are independent, and

$$\mathbb{P}((r_{j_i} = \varepsilon_i)_{i=1}^n) = 2^{-n}, \quad \varepsilon_i \in \{1, -1\}.$$

8.2 The Borel–Cantelli lemmas

Let $(A_j)_{j=1}^\infty$ be an infinite sequence in a probability space $(\Omega, \Sigma, \mathbb{P})$, and consider $\sum_j \mathbf{1}_{A_j}$.

If

$$\mathbb{E}\left(\sum_j \mathbf{1}_{A_j}\right) = \sum_j \mathbb{E}(\mathbf{1}_{A_j}) = \sum_j \mathbb{P}(A_j) < +\infty,$$

then $\sum_j \mathbf{1}_{A_j} < +\infty$ a.e. Note that

$$\{x \mid x \in A_j \text{ for infinitely many } j\} = \bigcap_k \bigcup_{j>k} A_j.$$

LEMMA.

1. $\sum_j \mathbb{P}(A_j) < +\infty \implies \mathbb{P}(A_j \text{ occurs infinitely often}) = 0.$
2. If $\{A_j\}$ are independent and $\sum_j \mathbb{P}(A_j) = +\infty$, then $\mathbb{P}(\bigcap_k \bigcup_{j>k} A_j) = 1.$

PROOF. Use $1 - x \leq e^{-x}$ for all $x \geq 0$. Then

$$\mathbb{P}\left(\bigcap_{j>k}^N A_j^c\right) = \prod_{j>k}^N (1 - \mathbb{P}(A_j)) \leq e^{-\sum \mathbb{P}(A_j)} \xrightarrow{N \rightarrow +\infty} 0.$$

Hence $\mathbb{P}(\bigcup_{j>k}^{+\infty} A_j) = 1$, which gives the claim. $\square$

8.3 Kolmogorov's zero-one law

THEOREM. Let X_n be independent. Then for all $A \in \mathcal{F}_\infty$ we have $\mathbb{P}(A) \in \{0, 1\}$, where

$$\mathcal{F}_\infty = \bigcap_{n=1}^{\infty} \sigma(X_n, X_{n+1}, \dots) \quad (\text{the tail } \sigma\text{-algebra / tail events}).$$

PROOF. Put $G_n = \sigma(X_1, \dots, X_n)$, which is independent of $\sigma(X_{n+1}, \dots)$. Hence any $A \in \mathcal{F}_\infty$ is independent of $\bigcup_n G_n$, i.e. $\mathcal{F}_\infty$ is independent of every G_n . Since $\mathcal{F}_\infty \subseteq \sigma(\bigcup_n G_n)$, we get

$$\mathbb{P}(A) = \mathbb{P}(A \cap A) = \mathbb{P}(A)^2. \quad \square$$

8.4 Sub-Gaussian bounds

Setting: X_i i.i.d. with $\mathbb{P}\{X_i = 1\} = \frac{1}{2}$.

LEMMA. Let $(r_j)_{j=1}^{+\infty}$ be a sequence of Rademacher functions. For all $N \geq 1$ and $(a_j)_{j=1}^N \subseteq \mathbb{C}^N$, set

$$S_N = \sum_{j=1}^N a_j r_j.$$

For all $\lambda > 0$: since $\mathbb{E}[r_j] = 0$, we have

$$\text{Var}(S_N) = \mathbb{E}[|S_N|^2] = \|a\|_{\ell^2(\mathbb{R}^N)}^2 = \sigma^2,$$

and

$$\mathbb{P}(|S_N| > \lambda \sigma) \leq 4e^{-\lambda^2/4}.$$

This measures how far S_N can be from its expectation.

PROOF. Suppose first $a_j \in \mathbb{R}$, $j = 1, \dots, N$. Let $t > 0$. By independence,

$$\mathbb{E} e^{tS_N} = \prod_{j=1}^N \mathbb{E}(e^{tr_j a_j}) = \prod_j \frac{e^{ta_j} + e^{-ta_j}}{2} = \prod_j \cosh(ta_j).$$

Using the elementary inequality

$\cosh(x) \leq e^{x^2/2}$

we obtain

$$\mathbb{E} e^{tS_N} \leq \prod_j e^{\frac{1}{2}(ta_j)^2} = e^{t^2(\sum_j a_j^2)/2} = e^{\frac{t^2}{2}\sigma^2}.$$

Therefore, by Chebyshev's inequality applied to the exponential,

$$\mathbb{P}(S_N > \lambda\sigma) = \mathbb{P}(e^{tS_N} > e^{t\lambda\sigma}) \leq e^{-t\lambda\sigma} \mathbb{E} e^{tS_N} = e^{-t\lambda\sigma + \frac{1}{2}t^2\sigma^2}.$$

Choosing $t = \frac{\lambda}{\sigma}$ gives $e^{-\lambda^2/2}$. The same argument applies to $S_N < -\lambda\sigma$, hence

$$\mathbb{P}(|S_N| > \lambda\sigma) \leq 2e^{-\lambda^2/2}.$$

COMPLEX CASE. $|S_N|^2 = |\operatorname{Re}(S_N)|^2 + |\operatorname{Im}(S_N)|^2 \geq \lambda^2\sigma^2$ forces

$$|\operatorname{Re}(S_N)|^2 \geq \frac{\lambda^2\sigma^2}{2} \quad \text{or} \quad |\operatorname{Im}(S_N)|^2 \geq \frac{\lambda^2\sigma^2}{2},$$

so the bound becomes $\leq 4e^{-\lambda^2/4}$. $\square$

8.5 Khinchine's inequality

THEOREM (KHINCHINE). For $1 \leq p < +\infty$ there exists $c = c(p) > 0$ such that

$$\frac{1}{c} \|a\|_2 \leq \|S_N\|_p \leq c \|a\|_2.$$

PROOF.

Case $p = 2$: $\|S_N\|_{L^2} = \|a\|_{\ell^2}$. WLOG $\|a\|_{\ell^2} = 1 = \sigma$.

Upper bound:

$$\mathbb{E}(|S_N|^p) = \int_0^\infty \mathbb{P}(|S_N| > \lambda) p \lambda^{p-1} d\lambda \leq \int_0^{+\infty} 4e^{-\lambda^2/4} p \lambda^{p-1} d\lambda \leq c(p) < +\infty.$$

Lower bound: Since $p \mapsto \|f\|_p$ is increasing for $p \geq 1$, it suffices to prove the case $p = 1$. Write

$$\begin{aligned} \int |S_N|^2 &= \int |S_N|^{2/3} |S_N|^{4/3} \\ &\leq \left(\int |S_N| \right)^{2/3} \left(\int |S_N|^4 \right)^{1/3} \\ &\leq C \left(\int |S_N| \right)^{2/3} \left(\int |S_N|^2 \right)^{2/3}, \end{aligned}$$

where the last step used the upper bound for $p = 4$. Hence

$$\frac{1}{c} \int |S_N|^2 \leq \left(\int |S_N| \right)^2. \quad \square$$

9 Fourier Partial Sums and Divergence

9.1 Partial sums

For $f \in L^1(\mathbb{T})$ and $N \geq 1$, let $D_N = \sum_{|n| \leq N} e(nx)$ be the Dirichlet kernel. Then

$$S_N f(x) = \sum_{n=-N}^N \widehat{f}(n) e^{-2\pi i n x} = D_N * f(x).$$

The central question is whether $S_N f$ converges to f almost everywhere.

- **1968: Carleson** — $f \in L^2$, via time-frequency analysis.
- **Hunt** — $f \in L^p$.
- Open-ended problem: $\lim_{r \rightarrow +\infty} \int_{B_r} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi \xrightarrow{\text{a.e.}} f(x)$ — higher-order Fourier analysis.

9.2 Norm convergence and uniform boundedness

PROPOSITION. Let $1 \leq p \leq \infty$. The following statements are equivalent:

1. For all $f \in L^p(\mathbb{T})$, $\|S_N f - f\|_p \rightarrow 0$;
2. $\sup_N \|S_N\|_{p \rightarrow p} < +\infty$.

PROOF.

(ii) $\Rightarrow$ (i): Take $g \in C_c^\infty$ and split

$$\|S_N f - f\|_p \leq \underbrace{\|S_N f - S_N g\|_p}_{\text{bounded operator}} + \underbrace{\|S_N g - g\|_p}_{\rightarrow 0} + \underbrace{\|g - f\|_p}_{< \varepsilon/3}.$$

(i) $\Rightarrow$ (ii): Uniform boundedness principle. $\square$

THEOREM. For $1 < p < +\infty$, $\sup_N \|S_N\|_{p \rightarrow p} < +\infty$.

PROOF. The Hilbert transform on $\mathbb{T}$ is the multiplier

$$\widehat{Hf}(n) = -i \operatorname{sgn}(n) \widehat{f}(n),$$

so that

$$Hf = \sum_{n>0} -i \widehat{f}(n) + \sum_{n<0} i \widehat{f}(n) = \sum_{n \geq 0} -i \widehat{f}(n) + \sum_{n \leq 0} i \widehat{f}(n)$$

(the added terms $-i \widehat{f}(0) + i \widehat{f}(0)$ cancel). Write $e_n(x) = e(nx)$. Then

$$Tf := \frac{1}{2}(1 + iH)(f) = \frac{1}{2}f + \frac{i}{2}Hf = \sum_{n \geq 0} \widehat{f}(n) e_n - \frac{1}{2}\widehat{f}(0),$$

i.e. T is the Riesz projection onto nonnegative frequencies. Using

$$\mathbf{1}_{[-N, N]} = \mathbf{1}_{[-N, \infty)} - \mathbf{1}_{(N, \infty)},$$

one gets

$$S_N f = \bar{e}_N T(e_N f) - e_N T(\bar{e}_N f) - \frac{1}{2}(\widehat{f}(-N)e_{-N} + \widehat{f}(N)e_N),$$

where the last (rank-two) term has L^p norm $\leq \|f\|_p$. Hence

$$\|S_N\|_{p \rightarrow p} \leq 2\|T\|_{p \rightarrow p} \leq 1 + \|H\|_{p \rightarrow p}. \quad \square$$

MAXIMAL OPERATOR. Define

$$Cf(x) = \sup_N |S_N f(x)|.$$

9.3 Kolmogorov's divergence theorem

THEOREM (KOLMOGOROV). There exists $f \in L^1(\mathbb{T})$ with $S_N f \not\rightarrow f$ a.e.; in fact there exists $f \in L^1(\mathbb{T})$ such that

$$|\{x \in \mathbb{T} \mid Cf(x) < \infty\}| = 0.$$

COROLLARY. If $S_N f \rightarrow f$ a.e. for all $f \in L^1$, then there exists $A > 0$ with

$$|\{x \in \mathbb{T} \mid C\mu(x) > \lambda\}| \leq \frac{A}{\lambda} \|\mu\| = \frac{A}{\lambda} |\mu|(\mathbb{T}),$$

for μ a complex finite Borel measure.

LEMMA. There exist probability measures μ_N on $\mathbb{T}$ with

$$\limsup_n |S_n \mu_N(x)| \gtrsim \log(N) \quad \text{a.e.}$$

Combining: $1 = |\{x \in \mathbb{T} \mid C\mu_N(x) > \log N\}| \leq \frac{A}{\log N} \text{ — contradiction!}$

9.4 The Banach principle

THEOREM. Let (T_n) be a sequence of bounded linear operators on $L^1(\mathbb{T})$, translation-invariant. Define

$$Mf(x) = \sup_{n \geq 1} |T_n f(x)|, \quad \|Mf\|_\infty < +\infty \quad \forall f \text{ trigonometric polynomial.}$$

If for every f ,

$$|\{x \mid Mf(x) < \infty\}| > 0, \quad (*)$$

then there exists A with

$$|\{x \mid Mf(x) > \lambda\}| \leq \frac{A}{\lambda} \|f\|_1.$$

SCHEME. Banach principle (together with $(*)$) $\Rightarrow \{f \mid \text{pointwise convergence holds}\}$ is a closed subset of $L^1 \Rightarrow \text{everything converges} \Rightarrow (\text{via the Corollary + Lemma above}) \text{ contradiction.}$

LEMMA. Let E_n be a sequence of measurable subsets of $\mathbb{T}$ with $\sum_{n \geq 1} |E_n| = +\infty$. Then there exists a sequence of points $(x_n)_{n \geq 1}$ such that

$$\sum_{n \geq 1} \mathbf{1}_{E_n}(x + x_n) = \infty \quad \text{a.e.}$$

9.4.1 Proof

By contradiction, assume there is a sequence $(f_j)_{j \geq 1} \subseteq L^1(\mathbb{T})$ with $\|f_j\|_{L^1} = 1$ such that for each j there is λ_j with

$$|\{x \mid Mf_j > \lambda_j\}| = |E_j| > \frac{2^j}{\lambda_j}.$$

By monotone convergence and the definition of M ,

$$\lim_{m \rightarrow +\infty} \left| \left\{ x \mid \sup_{1 \leq n \leq m} |T_n f_j(x)| > \lambda_j \right\} \right| > \frac{2^j}{\lambda_j}.$$

Hence for each j there is M_j with

$$\|T\| \geq \left| \left\{ x \mid \sup_{1 \leq n \leq M_j} |T_n f_j(x)| > \lambda_j \right\} \right| > \frac{2^j}{\lambda_j},$$

and WLOG f_j is a trigonometric polynomial, by continuity of the T_k .

(Side note: $\sigma_N f = K_N * f$, with the Fejér kernel $K_N = \frac{1}{N} \sum_{n=0}^{N-1} D_n \cdot$)

Choose integers $m_j \geq 1$ with

$$m_j \leq \frac{\lambda_j}{2^j} \leq m_j + 1 \implies \sum_{j=1}^{\infty} m_j |E_j| = \infty.$$

By the Lemma there exist points $x_{j,\ell}$, $1 \leq \ell \leq m_j$, with

$$\sum_{j=1}^{\infty} \sum_{\ell=1}^{m_j} \mathbf{1}_{E_j}(x - x_{j,\ell}) = \infty \quad \text{for a.e. } x \in \mathbb{T}.$$

Set $\delta_j = \frac{1}{j^2 m_j}$ and define the random function

$$f(x) = \sum_{j=1}^{\infty} \sum_{\ell=1}^{m_j} \pm \delta_j f_j(x - x_{j,\ell}).$$

The translations move all the bad sets together, so the construction is stable under translation; the random signs ensure that the terms do not cancel one another.

Then

$$\|f\|_{L^1} \leq \sum_{j=1}^{\infty} m_j \delta_j = \sum_j \frac{1}{j^2} < +\infty.$$

Let $x \in \mathbb{T}$ be such that $x \in E_j + x_{j,\ell}$ for infinitely many j , with $\ell = \ell(j)$; let J be the set of all such j .

Since T_n is translation invariant,

$$M[f_j(x - x_{j,\ell})] > \lambda_j \quad \text{for } x \in E_j + x_{j,\ell},$$

so there exists $n = n(j, x)$ with

$$|T_{n(j,x)} f_j(x - x_{j,\ell})| > \lambda_j \quad \text{on } E_j + x_{j,\ell}.$$

But

$$T_n f(x) = \sum_{j=1}^{\infty} \sum_{\ell=1}^{m_j} \pm \delta_j T_n f_j(x - x_{j,\ell}),$$

so

$$\begin{aligned} T_{n(j,x)} f(x) &= \sum_{k=1}^{\infty} \sum_{\ell=1}^{m_k} \pm \delta_k T_{n(j,x)} f_k(x - x_{k,\ell}) \\ &= \underbrace{\pm \delta_j T_{n(j,x)} f_j(x - x_{j,\ell})}_{> \delta_j \lambda_j \text{ in modulus}} + \underbrace{\text{other terms}}_{\text{expectation 0 with respect to the signs}}. \end{aligned}$$

Therefore

$$\mathbb{P}(|T_{n(j,x)} f(x)| > \delta_j \lambda_j) \geq \frac{1}{2} \implies \mathbb{P}(Mf = +\infty) \geq \frac{1}{2}.$$

Geometrically, half of the mass lies beyond $-\delta_j \lambda_j$, while the other half lies beyond $+\delta_j \lambda_j$.

9.4.2 The divergence event has full probability

PROOF. We show $\{Mf = +\infty\} \in \mathcal{F}_{\infty}$ (a tail event). Write

$$\begin{aligned} Mf(x) &= M \left(\sum_{k=1}^{\infty} \sum_{\ell=1}^{m_k} \pm \delta_k f_k(x - x_{k,\ell}) \right) \\ &= M \left(\underbrace{\sum_{k \geq N} \sum_{\ell=1}^{m_k} \pm \delta_k f_k(x - x_{k,\ell})}_{\text{tail}} \right) + M(\text{remainder}). \end{aligned}$$

where the remainder is a **finite** sum. Then:

- $Mf(x) = \infty \implies M(\text{tail}) = \infty$;
- $M(\text{tail}) = \infty \implies M(\text{tail}) \leq Mf(x) + M(\text{finite terms})$, so $Mf(x) = \infty$.

Hence $\{Mf(x) = \infty\} = \{M(\text{tail}) = \infty\}$, which belongs to $\mathcal{F}_{\infty}$. By the zero-one law and the bound $\geq \frac{1}{2}$ above,

$$\mathbb{P}(Mf(x) = \infty) = 1 \quad \text{for every } x.$$

Consequently

$$\begin{aligned} \int_0^1 \mathbb{P}(Mf(x) = +\infty) dx &= 1 \\ &= \int_0^1 \int_{\Omega} \mathbf{1}_{\{Mf(x)=\infty\}} d\omega dx \\ &= \int_{\Omega} \int_0^1 \mathbf{1}_{\{Mf(x)=\infty\}} dx d\omega \\ &= \mathbb{E}(|\{Mf(x) = \infty\}|). \end{aligned}$$

so $|\{Mf(x) = \infty\}| = 1$ for a.e. choice of signs ± 1 . $\square$

9.5 A measure-theoretic corollary and the construction

COROLLARY. If $\{S_N f\}$ converges for all $f \in L^1$, then there exists A' such that

$$|\{x \in \mathbb{T} \mid C\mu(x) > \lambda\}| \leq \frac{A'}{\lambda} \|\mu\|.$$

PROOF. We start from

$$|\{x \mid Cf(x) > \lambda\}| \leq \frac{A}{\lambda} \|f\|_{L^1(\mathbb{T})}.$$

Consider $f = V_N * \mu$, where

$$V_N(x) = K_N(x) + K_N(x)e(Nx) + K_N(x)e(-Nx)$$

(the de la Vallée Poussin type kernel). Since $\|K_N * \mu\|_{L^1} \leq \|\mu\|$, we get

$$\|V_N * \mu\|_{L^1} \leq 3 \|\mu\|_{L^1} \implies |\{x : |C(V_N * \mu)| > \lambda\}| \leq \frac{3A}{\lambda} \|\mu\|_M.$$

Hence

$$\lim_{M \rightarrow \infty} \left| \left\{ x \mid \sup_{1 \leq n \leq M} |S_n(V_N * \mu)(x)| > \lambda \right\} \right| \leq \frac{3A}{\lambda} \|\mu\|_M,$$

and therefore

$$\left| \left\{ x \mid \sup_{1 \leq n \leq M} \dots \right\} \right| \leq \frac{3A}{\lambda} \|\mu\|_M,$$

letting $n = M$, $n \rightarrow +\infty$. $\square$

9.5.1 A first construction

Set

$$\mu_N = \frac{1}{N} \sum_{j=1}^N \delta_{x_{j,N}}, \quad \left| x_{j,N} - \frac{j}{N} \right| < \frac{1}{N^2},$$

i.e. the points $x_{j,N}$ are tiny perturbations of the equispaced points $0, \frac{1}{N}, \frac{2}{N}, \dots, 1$.

Then

$$\begin{aligned} S_n \mu_N(x) &= D_n * \mu_N(x) = \int D_n(x-y) d\mu_N(y) \\ &= \frac{1}{N} \sum_{j=1}^N D_n(x - x_{j,N}) \\ &= \frac{1}{N} \sum_{j=1}^N \frac{\sin((2n+1)\pi(x - x_{j,N}))}{\sin(\pi(x - x_{j,N}))}. \end{aligned}$$

Choosing x_j so that $|\sin(\cdot)| \geq \frac{1}{2}$ in the numerator gives

$$\begin{aligned} |S_n \mu_N(x)| &\geq \frac{1}{2N} \sum_{j=1}^N \frac{1}{|\sin(\pi(x - x_{j,N}))|} \\ &\geq \frac{c}{N} \sum_{j=1}^N \frac{1}{j/N} = c \sum_{j=1}^N \frac{1}{j} \gtrsim c \log N. \end{aligned}$$

Remark 9.1. The estimate involving $\sin(\pi(x - x_{j,N}))$ requires the points $x_{j,N}$ to be ordered relative to x ; this is the geometric input behind the bound by j/N .

9.6 A refined almost-everywhere divergence construction

Write

$$Cf(x) = \sup_{n \geq 0} |S_n f(x)|$$

for the maximal partial-sum operator.

COROLLARY. If $\{S_N f\}$ converges a.e. (in fact: *on a set of positive measure*) for all f , then

$$|\{x \in \mathbb{T} \mid Cf(x) > \lambda\}| \leq \frac{A}{\lambda} \|f\|_{L^1} \quad (****)$$

and the same with f replaced by μ and $\|f\|_{L^1}$ by $\|\mu\|_{\mathbb{T}}$, for $\mu \in M(\mathbb{T})$.

To show that there exists $f \in L^1$ whose Fourier series diverges a.e. (on a set of positive measure), it suffices to show that (****) **fails**.

EQUIDISTRIBUTION LEMMA. Assume $(\theta_1, \dots, \theta_d) \in \mathbb{T}^d$ is incommensurate; that is, for every $(n_1, \dots, n_d) \in \mathbb{Z}^d \setminus \{0\}$,

$$n_1 \theta_1 + \dots + n_d \theta_d \notin \mathbb{Z}.$$

Then the orbit $\{(n\theta_1, \dots, n\theta_d) \bmod \mathbb{Z}^d\}$ is dense in $\mathbb{T}^d$.

CONSTRUCTION. There exists $c > 0$ such that for every N there is a probability measure μ_N on $\mathbb{T}$ with

$$\limsup_n \frac{|S_n \mu_N(x)|}{\log N} > c \quad \text{a.e. } x.$$

If so, and if (****) held, then with $\lambda_N \sim \log N$,

$$1 = |\{x \mid C\mu_N(x) \geq \lambda_N\}| \leq \frac{A}{\lambda_N} \|\mu_N\| = \frac{A}{\lambda_N} \longrightarrow 0,$$

a contradiction. $\square$

PROOF OF THE CONSTRUCTION. Set

$$\mu_N = \frac{1}{N} \sum_{j=1}^N \delta_{x_{j,N}}, \quad \left| x_{j,N} - \frac{j}{N} \right| \leq \frac{1}{N^2}, \quad j = 1, \dots, N.$$

Choose $(x_{j,N})_{j=1}^N$ to be **incommensurate** (this is possible since the set of commensurate vectors — those satisfying $n_1 x_1 + \dots + n_d x_d \in \mathbb{Z}$ — has measure zero).

Then choose n so that $(n(x - x_{j,N}))_{j=1}^N$ is also incommensurate (for a.e. x). Hence

$$((2n+1)(x - x_{j,N}))_{j=1}^N \bmod \mathbb{Z}^d \text{ is dense in } \mathbb{T}^N,$$

which implies

$$\sin((2n+1)\pi(x - x_{j,N})) \geq \frac{1}{2} \quad \text{for infinitely many } n, \text{ a.e. } x.$$

Therefore

$$\begin{aligned}
|S_n \mu_N(x)| &= |D_n * \mu_N(x)| \\
&= \left| \int_{\mathbb{T}} \frac{\sin((2n+1)\pi(x-y))}{\sin(\pi(x-y))} d\mu_N(y) \right| \\
&= \left| \frac{1}{N} \sum_j \frac{\sin((2n+1)\pi(x-x_{j,N}))}{\sin(\pi(x-x_j))} \right|. \\
&\geq \frac{1}{2N} \sum_{j=1}^N \frac{1}{|\sin(\pi(x-x_{j,N}))|} \geq \frac{c}{N} \sum_j \frac{1}{|x-x_{j,N}|} \geq \frac{c'}{N} \sum_j \frac{1}{j/N} \geq c \log N. \quad \square
\end{aligned}$$

10 Poisson Integrals and Boundary Values

10.1 The Hilbert transform and the Poisson kernel

On $\mathbb{R}$, the Hilbert transform is

$$Hf(x) = \text{p. v.} \frac{1}{\pi} \int_{\mathbb{R}} \frac{f(y)}{x-y} dy.$$

On $\mathbb{T}$, its periodic analogue is

$$Hf(x) = \text{p. v.} \frac{1}{\pi} \int_{-1/2}^{1/2} \cot\left(\frac{x-y}{2}\right) f(y) dy.$$

On the Fourier side, both are described by the same sign-type multiplier, continuous on $\mathbb{R}$ and sampled at the integer frequencies on $\mathbb{T}$.

Its connection with analytic functions can be formulated as follows. Given a real-valued $f \in L^p(\mathbb{T})$, one seeks an analytic function $F : \mathbb{D} \rightarrow \mathbb{C}$ satisfying

$$F|_{\partial\mathbb{D}} = f + ig \quad (f = \text{real part})?$$

Such a function is obtained from the Schwarz integral

$$F(z) = \int_{-1/2}^{1/2} \frac{e^{2\pi i x} + z}{e^{2\pi i x} - z} f(x) dx$$

1. By dominated convergence, F is analytic.
2. $F(0) = \int_{-1/2}^{1/2} f(x) dx = \widehat{f}(0) \in \mathbb{R}$.

Splitting into real and imaginary parts,

$$F(z) = \int_{-1/2}^{1/2} \text{Re}\left(\frac{e^{2\pi i x} + z}{e^{2\pi i x} - z}\right) f(x) dx + i \int_{-1/2}^{1/2} \text{Im}\left(\frac{e^{2\pi i x} + z}{e^{2\pi i x} - z}\right) f(x) dx.$$

With $z = re^{2\pi iy}$:

$$\begin{aligned} \frac{e^{2\pi ix} + z}{e^{2\pi ix} - z} &= \frac{e^{2\pi i(x-y)} + r}{e^{2\pi i(x-y)} - r} \\ &= \frac{(e^{2\pi i(x-y)} + r)(e^{-2\pi i(x-y)} - r)}{|e^{2\pi i(x-y)} - r|^2} \\ &= \underbrace{\frac{1 - r^2}{|e^{2\pi i(x-y)} - r|^2}}_{P_r(x-y)} - \underbrace{\frac{2ir \sin(2\pi(x-y))}{|e^{2\pi i(x-y)} - r|^2}}_{iQ_r(x-y)}. \end{aligned}$$

Hence

$$\operatorname{Re} F(re^{2\pi iy}) = P_r * f(y),$$

where $P_r(t) = P_r(-t)$, peaks at $t = 0$, and becomes singular as $r \rightarrow 1$.

PROPOSITION. P_r is an approximate identity, $0 \leq r < 1$.

PROOF. $P_r(t) \geq 0$.

$$\text{Claim: } P_r(t) = \sum_{k \in \mathbb{Z}} r^{|k|} e^{2\pi ikt}.$$

Proof of claim:

$$\begin{aligned} \sum_k r^{|k|} e^{2\pi ikt} &= 1 + \sum_{k \geq 1} r^k e^{2\pi ikt} + \sum_{k=1}^{\infty} r^k e^{-2\pi ikt} \\ &= 1 + \frac{re^{2\pi it}}{1 - re^{2\pi it}} + \frac{re^{-2\pi it}}{1 - re^{-2\pi it}} = P_r(t). \end{aligned}$$

Moreover,

$$\int_{\mathbb{T}} P_r(t) dt = \widehat{P_r}(0) = 1 = \|P_r\|_{L^1},$$

and for all $\delta > 0$,

$$\sup_{|\theta| > \delta} |P_r(t)| \rightarrow 0 \quad \text{as } r \rightarrow 1.$$

Indeed $P_r(t)$ is decreasing as a function of $|t|$, so

$$\sup_{|\theta| > \delta} |P_r(t)| = P_r(\delta) = \frac{1 - r^2}{|r - e^{2\pi i\delta}|^2} \xrightarrow{r \rightarrow 1} 0. \quad \square$$

Consequently

$$\operatorname{Re} F(re^{2\pi iy}) = P_r * f \longrightarrow f \quad \text{in } L^p.$$

10.2 Almost-everywhere convergence of Poisson integrals

CLAIM. $P_r * f \xrightarrow{r \rightarrow 1} f$ for a.e. $y \in \mathbb{T}$.

SKETCH. It suffices to show the weak-type bound

$$\left| \left\{ \sup_{r < 1} |P_r * f| > \lambda \right\} \right| \leq \frac{A}{\lambda} \|f\|_{L^1}.$$

The key pointwise domination is

$$|K * f(x)| \leq \|K\|_{L^1} Mf(x) \quad (M = \text{Hardy-Littlewood maximal function})$$

valid whenever K is even and decreasing in t (a "bump" peaked at the origin).

Baby case: $K = \sum_i a_i \mathbf{1}_{I_i}$ with all intervals I_i centered at 0. Then

$$|K * f| \leq \sum_i a_i (\mathbf{1}_{I_i} * f) \leq \sum_i a_i |I_i| Mf \leq \|K\|_{L^1} Mf.$$

General case: approximate by the baby case. $\square$

IMAGINARY PART. Similarly,

$$\operatorname{Im} F(re^{2\pi i y}) = Q_r * f \longrightarrow Hf, \quad Q_r(y) = \frac{2r \sin(2\pi y)}{|1 - re^{2\pi i y}|^2} \xrightarrow{r \rightarrow 1} \cot\left(\frac{y}{2}\right).$$

Part IV

Calderón–Zygmund Theory and BMO

11 Homogeneous Singular Integrals

11.1 Dini continuity and the Hörmander condition

Setting: $K(x) = \frac{\Omega(x/|x|)}{|x|^d}$ with Ω defined on S^{d-1} , $\int_{S^{d-1}} \Omega d\sigma = 0$, and Ω having modulus of continuity ω .

GOAL (III)' $\Rightarrow$ (III): the Dini condition

$$\int_0^1 \frac{\omega(\delta)}{\delta} d\delta < \infty$$

implies the Hörmander condition

$$\int_{|x| \geq 2|y|} |K(x-y) - K(x)| dx \leq B < \infty.$$

11.1.1 Splitting the difference

$$K(x-y) - K(x) = \frac{\Omega\left(\frac{x-y}{|x-y|}\right) - \Omega\left(\frac{x}{|x|}\right)}{|x-y|^d} + \Omega\left(\frac{x}{|x|}\right) \left[\frac{1}{|x-y|^d} - \frac{1}{|x|^d} \right].$$

The first term measures the **angular** variation of Ω , the second the **radial** variation of $|x|^{-d}$. Handle them separately.

11.1.2 Radial part

Claim: $\int_{|x| \geq 2|y|} \left| \frac{1}{|x-y|^d} - \frac{1}{|x|^d} \right| dx \leq C.$

Put $F(t) = \frac{1}{|x-ty|^d}$, $t \in [0, 1]$. By the fundamental theorem of calculus,

$$\left| \frac{1}{|x-y|^d} - \frac{1}{|x|^d} \right| = |F(1) - F(0)| \leq \int_0^1 |F'(t)| dt \leq d|y| \sup_{t \in [0, 1]} \frac{1}{|x-ty|^{d+1}}.$$

On the region $|x| \geq 2|y|$ we have $|x-ty| \geq |x| - |y| \geq \frac{|x|}{2}$, hence

$$\int_{|x| \geq 2|y|} \left| \frac{1}{|x-y|^d} - \frac{1}{|x|^d} \right| dx \leq d 2^{d+1} |y| \int_{|x| \geq 2|y|} |x|^{-d-1} dx \leq C,$$

since $\int_{|x| \geq R} |x|^{-d-1} dx \simeq R^{-1}$ and $R = 2|y|$. Note this bound is **independent of y** — this is the scale invariance of the estimate.

11.1.3 Angular part

First, the unit vectors are close when $|x| \geq 2|y|$:

$$\begin{aligned} \left| \frac{x-y}{|x-y|} - \frac{x}{|x|} \right| &= \frac{||x|(x-y) - |x-y|x||}{|x||x-y|} \\ &\leq \frac{|x||y| + |x||x| - |x-y||}{|x||x-y|} \\ &\leq \frac{2|x||y|}{|x||x-y|} \leq C \frac{|y|}{|x|}. \end{aligned}$$

where we used $||x| - |x-y|| \leq |y|$ and $|x-y| \geq |x|/2$.

Therefore $|\Omega(\frac{x-y}{|x-y|}) - \Omega(\frac{x}{|x|})| \leq \omega\left(C \frac{|y|}{|x|}\right)$, and

$$\int_{|x| \geq 2|y|} \frac{|\Omega(\frac{x-y}{|x-y|}) - \Omega(\frac{x}{|x|})|}{|x-y|^d} dx \leq C' \int_{|x| \geq 2|y|} \frac{\omega\left(C \frac{|y|}{|x|}\right)}{|x|^d} dx \leq C'' \int_0^{C/2} \frac{\omega(\delta)}{\delta} d\delta < +\infty.$$

(The last step: polar coordinates $dx = r^{d-1} dr d\sigma$ turns the integral into $c \int_{2|y|}^{\infty} \omega(C|y|/r) \frac{dr}{r}$; substituting $\delta = C|y|/r$ gives $\frac{dr}{r} = -\frac{d\delta}{\delta}$ and the range $0 < \delta \leq C/2$.)

□

11.2 Translation invariance and dilation structure

PROP. Let $T : L^2 \rightarrow L^2$ be bounded and linear, commuting with all translations,

$$T \circ \tau_x = \tau_x \circ T, \quad (\tau_x f)(y) = f(x+y).$$

Then there exists $m \in L^\infty$ such that $\widehat{Tf} = m \widehat{f}$.

11.2.1 Homogeneity of the multiplier

$K(x) = \frac{\Omega(x)}{|x|^d}$ is homogeneous of degree $-d$: $K(\varepsilon x) = \varepsilon^{-d} K(x)$.

For $\varepsilon > 0$ define the **unitary** dilation on $L^2(\mathbb{R}^d)$:

$$(D_\varepsilon f)(x) = \varepsilon^{-d/2} f\left(\frac{x}{\varepsilon}\right).$$

Let $Tf = \text{p.v. } K * f$. Then $T \circ D_\varepsilon = D_\varepsilon \circ T$:

$$\begin{aligned}
D_\varepsilon(Tf)(x) &= \varepsilon^{-d/2} T f\left(\frac{x}{\varepsilon}\right) = \varepsilon^{-d/2} \text{p.v.} \int K\left(\frac{x}{\varepsilon} - y\right) f(y) dy \\
&= \varepsilon^{d/2} \text{p.v.} \int K(x - \varepsilon y) f(y) dy & \left[K\left(\frac{x - \varepsilon y}{\varepsilon}\right) = \varepsilon^d K(x - \varepsilon y) \right] \\
&= \text{p.v.} \int K(x - z) \varepsilon^{-d/2} f\left(\frac{z}{\varepsilon}\right) dz & [z = \varepsilon y] \\
&= T D_\varepsilon f(x).
\end{aligned}$$

On the Fourier side this commutation says the multiplier m satisfies $m(\lambda\xi) = m(\xi)$, i.e. m is homogeneous of degree 0.

11.2.2 Truncated kernels

For $0 < \varepsilon < \eta < \infty$ set

$$K_{\varepsilon,\eta}(x) = \begin{cases} \Omega(x)/|x|^d, & \varepsilon \leq |x| \leq \eta, \\ 0, & \text{else.} \end{cases}$$

These are integrable, so $\widehat{K_{\varepsilon,\eta} * f} = \widehat{K_{\varepsilon,\eta}} \widehat{f}$ makes sense. The two things to prove are:

1. (i) $|\widehat{K_{\varepsilon,\eta}}| \leq A$ uniformly in ε, η ;
2. (ii) for $x \neq 0$, $\lim_{\varepsilon \rightarrow 0, \eta \rightarrow \infty} \widehat{K_{\varepsilon,\eta}}(x) = m(x)$.

11.3 Fourier multipliers of truncated kernels

Write $x = Rx'$ with $x' = x/|x|$, $R = |x|$, and $y = ry'$ with $y' = y/|y|$.

Define

$$I_{\varepsilon,\eta}(x, y') = \int_\varepsilon^\eta \frac{e^{2\pi i R r x' \cdot y'} - \cos(2\pi R r)}{r} dr.$$

The subtraction of $\cos(2\pi R r)$ is legitimate because $\int_{S^{d-1}} \Omega(y') d\sigma(y') = 0$ — it costs nothing and it makes the integral converge at $r \rightarrow 0$.

IMAGINARY PART.

$$\int_\varepsilon^\eta \frac{\sin(2\pi R r x' \cdot y')}{r} dr$$

is bounded **uniformly** in ε, η, R (Dirichlet integral), and converges to

$$\int_0^\infty \frac{\sin t}{t} dt \cdot \text{sgn}(x' \cdot y') = \frac{\pi}{2} \text{sgn}(x' \cdot y').$$

REAL PART.

$$\int_\varepsilon^\eta \frac{\cos(2\pi R r x' \cdot y') - \cos(2\pi R r)}{r} dr$$

is bounded in absolute value by $C \log \frac{1}{|x' \cdot y'|} + C$ for some constant C , and converges to $\log \frac{1}{|x' \cdot y'|}$.

ASSEMBLING. Since $\int_{S^{d-1}} \Omega d\sigma = 0$,

$$\widehat{K_{\varepsilon,\eta}}(x) = \int_{S^{d-1}} \int_\varepsilon^\eta \frac{e^{2\pi i R r x' \cdot y'}}{r} \Omega(y') dr d\sigma(y') = \int_{S^{d-1}} I_{\varepsilon,\eta}(x, y') \Omega(y') d\sigma(y').$$

Hence the uniform bound

$$|\widehat{K_{\varepsilon,\eta}}(x)| \leq \int_{S^{d-1}} |I_{\varepsilon,\eta}(x, y')| |\Omega(y')| d\sigma(y') \leq A \int_{S^{d-1}} \left[1 + \log \frac{1}{|x' \cdot y'|} \right] |\Omega(y')| d\sigma(y') \leq A',$$

the last integral being finite because $\log \frac{1}{|x' \cdot y'|}$ is integrable on the sphere.

Consequently $\widehat{K_{\varepsilon,\eta}}(x) \rightarrow m(x)$ pointwise for $x \neq 0$, and for $f \in L^2(\mathbb{R}^d)$:

$$K_{\varepsilon,\eta} * f \text{ converges in } L^2, \text{ with Fourier transform } \rightarrow m\widehat{f}.$$

(Order of limits: first $\eta \rightarrow +\infty$ with ε fixed, then $\varepsilon \rightarrow 0$.)

12 Maximal Singular Integrals

DEF. $T^*f(x) = \sup_{\varepsilon > 0} |T_\varepsilon f(x)|$, where $T_\varepsilon f(x) = \int_{|x-y| > \varepsilon} K(x-y)f(y) dy$.

THEOREM. Let Ω be as before.

- (a) If $1 \leq p < \infty$, then $\lim_{\varepsilon \rightarrow 0} T_\varepsilon f(x)$ exists for a.e. x , for every $f \in L^p$.
- (b) $T^* : L^p(\mathbb{R}^d) \rightarrow L^p(\mathbb{R}^d)$ is bounded for $1 < p < \infty$.
- (c) $|\{x \in \mathbb{R}^d : |T^*f(x)| > \lambda\}| \leq \frac{C}{\lambda} \|f\|_{L^1(\mathbb{R}^d)}$ for $f \in L^1$, $\lambda > 0$.

(a) follows from (c) by the standard density + maximal-inequality argument.

12.1 Calderón–Zygmund decomposition and the good part

Let $f \in L^1$, $\lambda > 0$. Perform the **Calderón–Zygmund decomposition at height λ** :

$$f = g + b,$$

- $|g| \leq \lambda$ (the good part),
- b supported in the bad cubes $\{Q_j\}$, with $\int_{Q_j} b = 0$ for all j ,
- $\sum_j |Q_j| \leq C \frac{\|f\|_{L^1}}{\lambda}$.

DEF. $Q_j^* = 2Q_j$ (concentric dilate), and $F^* = [\bigcup_j Q_j^*]^c$.

Split:

$$|\{x : T^*f > \lambda\}| \leq |\{x : T^*b > \frac{\lambda}{2}\}| + |\{x : T^*g > \frac{\lambda}{2}\}|.$$

The **good part** is handled by the L^2 theory (Chebyshev + L^2 boundedness of T^*):

$$|\{x : T^*g > \frac{\lambda}{2}\}| \leq C \frac{\|g\|_{L^2}^2}{\lambda^2} \leq C \frac{1}{\lambda} \|g\|_{L^1} \leq C \frac{1}{\lambda} \|f\|_{L^1},$$

using $\|g\|_{L^2}^2 \leq \|g\|_\infty \|g\|_1 \leq \lambda \|g\|_1$ and $\|g\|_1 \leq C \|f\|_1$.

It therefore remains to prove

$$|\{x : T^*b > \lambda\}| \leq C \frac{\|f\|_{L^1}}{\lambda}.$$

12.2 Cancellation estimate for the bad part

CLAIM. For all $x \in F^*$,

$$|T^*b(x)| \leq \underbrace{\sum_j \int_{Q_j} |K(x-y) - K(x-y_j)| |b(y)| dy}_{=: \Sigma(x)} + C Mf(x),$$

where y_j is the **center** of Q_j .

12.3 Consequences of the cancellation estimate

$$|\{T^*b \geq \frac{\lambda}{2}\}| \leq \underbrace{|\{x \in F^* : \Sigma(x) > \frac{\lambda}{4}\}|}_{\textcircled{1}} + \underbrace{|\{x \in F^* : Mf(x) > \frac{\lambda}{4C}\}|}_{\leq C\|f\|_1/\lambda \text{ (weak (1,1) of } M)}} + \underbrace{|(F^*)^c|}_{\leq C\|f\|_1/\lambda}$$

The third term is controlled because $|(F^*)^c| \leq \sum_j |Q_j^*| = 2^d \sum_j |Q_j| \leq C\|f\|_1/\lambda$.

For $\textcircled{1}$, use Chebyshev and Fubini:

$$\begin{aligned} \textcircled{1} &\leq \frac{C}{\lambda} \int_{F^*} |\Sigma(x)| dx = \frac{C}{\lambda} \sum_j \int_{Q_j} \underbrace{\int_{F^*} |K(x-y) - K(x-y_j)| dx}_{\leq B \text{ (Hörmander)}} |b(y)| dy \\ &\leq \frac{CB}{\lambda} \sum_j \int_{Q_j} |b(y)| dy \leq \frac{CB}{\lambda} \|f\|_{L^1}. \end{aligned}$$

The inner integral is bounded by the Hörmander constant B precisely because $x \in F^*$ means $x \notin 2Q_j$, so $|x - y_j| \geq 2|y - y_j|$ for $y \in Q_j$. $\square$

12.4 Geometric proof of the cancellation estimate

Fix $x \in F^*$ and look at

$$T_\varepsilon b(x) = \sum_j \int_{Q_j} K_\varepsilon(x-y) b(y) dy, \quad K_\varepsilon(z) = K(z) \mathbf{1}_{\{|z| > \varepsilon\}}.$$

Classify each cube Q_j by its position relative to the ball $B(x, \varepsilon)$:

CASE 1 — $Q_j \subseteq B(x, \varepsilon)$. For every $y \in Q_j$ we have $|x - y| < \varepsilon$, so $K_\varepsilon(x - y) = 0$ and

$$\int_{Q_j} K_\varepsilon(x - y) b(y) dy = 0.$$

Such cubes contribute nothing.

CASE 2 — $Q_j \cap B(x, \varepsilon) = \emptyset$. Here $K_\varepsilon(x - y) = K(x - y)$ on all of Q_j , so we may use the cancellation $\int_{Q_j} b = 0$:

$$\begin{aligned} \left| \int_{Q_j} K_\varepsilon(x - y) b(y) dy \right| &= \left| \int_{Q_j} K(x - y) b(y) dy \right| \\ &= \left| \int_{Q_j} [K(x - y) - K(x - y_j)] b(y) dy \right| \\ &\leq \int_{Q_j} |K(x - y) - K(x - y_j)| |b(y)| dy. \end{aligned}$$

These are exactly the terms of $\Sigma(x)$.

CASE 3 — THE REMAINING CASE (Q_j MEETS BOTH $B(x, \varepsilon)$ AND ITS COMPLEMENT). Then the cube straddles the sphere $\partial B(x, \varepsilon)$, and since $x \in F^*$ (so x is outside $2Q_j$, hence Q_j is not too big compared to its distance from x), we get for all $y \in Q_j$

$$\frac{\varepsilon}{10} \leq \text{dist}(x, y) \leq 10\varepsilon.$$

Therefore

$$\int_{Q_j} |K_\varepsilon(x - y)b(y)| dy \leq C \int_{Q_j} \frac{|b(y)|}{|x - y|^d} dy \leq \frac{C}{\varepsilon^d} \int_{Q_j} |b(y)| dy \leq C \frac{1}{(10\varepsilon)^d} \int_{B(x, 10\varepsilon) \cap Q_j} |b(y)| dy,$$

which is $\leq C M b(x) \leq C M f(x)$ — an average over a ball centered at x , i.e. controlled by the **maximal function**. (Only boundedly many cubes fall into Case 3.)

Summing over all j and taking $\sup_{\varepsilon > 0}$ gives the claim. $\square$

13 BMO and Endpoint Estimates

DEF. For $f \in L^1_{loc}$, $f \in \text{BMO}$ if

$$\|f\|_{\text{BMO}} := \sup_Q \frac{1}{|Q|} \int_Q |f - f_Q| dx < \infty.$$

NOTE. $f \in L^\infty \Rightarrow f \in \text{BMO}$, with $\|f\|_{\text{BMO}} \leq 2\|f\|_\infty$.

BMO is a Banach space with norm $\|\cdot\|_{\text{BMO}}$ **modulo constant functions** (constants have BMO norm 0).

13.1 Averages over nested cubes grow only linearly

OBSERVATION. Let $Q_2 \subset Q_1$ be cubes with $\ell(Q_2) = \frac{1}{2}\ell(Q_1)$, $f \in \text{BMO}$. Then

$$|f_{Q_2} - f_{Q_1}| \leq \frac{1}{|Q_2|} \int_{Q_2} |f - f_{Q_1}| \leq \frac{|Q_1|}{|Q_2|} \cdot \frac{1}{|Q_1|} \int_{Q_1} |f - f_{Q_1}| \leq 2^d \|f\|_{\text{BMO}}.$$

Iterating: if $Q_n \subset Q_{n-1} \subset \dots \subset Q_1 \subset Q_0$ with $\ell(Q_j) = \frac{1}{2}\ell(Q_{j-1})$, then

$$|f_{Q_n} - f_{Q_0}| \leq n 2^d \|f\|_{\text{BMO}}.$$

Thus geometric changes in scale affect the averages only arithmetically. This linear growth is precisely the source of the logarithmic blow-up allowed by BMO and, ultimately, of the John–Nirenberg inequality.

13.2 Equivalent norm with arbitrary constants

$$\|f\|_{\text{BMO}} = \sup_Q \inf_{c_Q} \frac{1}{|Q|} \int_Q |f - c_Q| dx.$$

NOTE. $\|f\|_{\text{BMO}} \leq \|f\|_{\text{BMO}} \leq 2\|f\|_{\text{BMO}}$.

Proof of the right inequality. For any constant c_Q ,

$$\frac{1}{|Q|} \int_Q |f - f_Q| \leq \frac{1}{|Q|} \int_Q |f - c_Q| + \frac{1}{|Q|} \int_Q |f_Q - c_Q| \leq \|f\|_{\text{BMO}} + |f_Q - c_Q| + \varepsilon,$$

and

$$|f_Q - c_Q| = \left| \frac{1}{|Q|} \int_Q (f - c_Q) \right| \leq \frac{1}{|Q|} \int_Q |f - c_Q|,$$

so taking the infimum over c_Q gives $\leq 2\|f\|_{\text{BMO}} + \varepsilon$. $\square$

The left inequality is immediate ($c_Q = f_Q$ is one admissible choice).

REMARK. Cubes $\rightsquigarrow$ balls: replacing cubes by balls gives an equivalent norm.

13.3 Where BMO sits

$$L^\infty \subsetneq \text{BMO} \subseteq \bigcap_{p \geq 1} L^p_{\text{loc}}$$

Related duality (Hardy spaces H^p):

- $(H^p)^* \simeq H^{p'}$,
- $(H^1)^* \simeq \text{BMO}$ (Fefferman duality),
- $\|f\|_{H^p} = \|f\|_{L^p} + \|Hf\|_{L^p}$.

13.4 Calderón–Zygmund operators map L^∞ to BMO

PROP. Let T be a Calderón–Zygmund operator. Then for all $f \in L^2 \cap L^\infty$,

$$\|Tf\|_{\text{BMO}} \leq C\|f\|_\infty, \quad \text{i.e.} \quad T : L^\infty \longrightarrow \text{BMO}.$$

Heuristically, a Calderón–Zygmund operator may send a bounded bump to a function with logarithmic growth. This explains why L^∞ is not preserved and why BMO is the natural endpoint space.

13.5 Local and far-field decomposition

Since $f \in L^2$, Tf is well-defined. Fix a ball $B_0 = B(x_0, R)$ and choose the constant

$$c_{B_0} = \int_{|x_0 - y| \geq 2R} K(x_0 - y) f(y) dy$$

(the "far field" evaluated at the center — this is the natural normalization, and by §7.2 we are free to use any constant).

Split $f = f \mathbf{1}_{\{|x_0 - y| \geq 2R\}} + f \mathbf{1}_{\{|x_0 - y| < 2R\}}$ (far part + near part). Write

$$\frac{1}{|B_0|} \int_{B_0} |Tf(x) - c_{B_0}| dx \leq I_{\text{far}} + I_{\text{near}},$$

where

$$I_{\text{far}} = \frac{1}{|B_0|} \int_{B_0} \left| \int_{|x_0-y| \geq 2R} (K(x-y) - K(x_0-y)) f(y) dy \right| dx,$$

$$I_{\text{near}} = \frac{1}{|B_0|} \int_{B_0} \left| T(f \mathbf{1}_{\{|x_0-y| < 2R\}})(x) \right| dx.$$

FIRST TERM (FAR FIELD — SMOOTHNESS/HÖRMANDER). For $x \in B_0$ and $|x_0 - y| \geq 2R$ we have $|x_0 - y| \geq 2|x - x_0|$, so the Hörmander condition applies:

$$I_{\text{far}} \leq \frac{\|f\|_{\infty}}{|B_0|} \int_{B_0} \int_{|x_0-y| \geq 2R} |K(x-y) - K(x_0-y)| dy dx$$

$$\leq CB \|f\|_{\infty}.$$

SECOND TERM (NEAR FIELD — HÖLDER + L^2 BOUNDEDNESS). By Cauchy–Schwarz and then $L^2 \rightarrow L^2$ boundedness of T :

$$I_{\text{near}} \leq \frac{1}{|B_0|^{1/2}} \left(\int_{B_0} |T(f \mathbf{1}_{B(x_0, 2R)})|^2 \right)^{1/2}$$

$$\leq \frac{C}{|B_0|^{1/2}} \left(\int |f \mathbf{1}_{B(x_0, 2R)}|^2 \right)^{1/2}$$


$$\leq C \|f\|_{\infty},$$

the last step because $|B(x_0, 2R)| \leq 2^d |B_0|$, so $\left(\int_{B(x_0, 2R)} |f|^2 \right)^{1/2} \leq \|f\|_{\infty} |B(x_0, 2R)|^{1/2} \leq C |B_0|^{1/2} \|f\|_{\infty}$.

Taking the supremum over all balls B_0 gives $\|Tf\|_{\text{BMO}} \leq C \|f\|_{\infty}$. $\square$

14 Calderón–Zygmund Proof Architecture

Every Calderón–Zygmund estimate decomposes along the same two-by-two grid: **which piece of f** (good/bad) $\times$ **which regime of the kernel** (x near y / x far from y).



- $g, x \approx y$ **and** $x \gg y$: the good part is bounded, so raw $L^2 \rightarrow L^2$ boundedness suffices; no cancellation needed.
- $b, x \approx y$: the singular part lives inside the bad cubes; one simply throws away $\bigcup 2Q_j$, whose measure is $\lesssim \|f\|_1 / \lambda$.

- $b, x \gg y$: away from the bad cubes, the mean-zero property of b plus the smoothness of K (Hörmander) gives the decay that makes the sum converge.

Part V

Multipliers and Littlewood–Paley Theory

15 Fourier Multipliers

15.1 The Mihlin multiplier theorem

A basic multiplier arises from the equation $\Delta u = f$. Consider the operator sending f to a second derivative of u :

$$T : f \mapsto \partial_i \partial_j u.$$

Taking Fourier transforms,

$$\widehat{Tf} = -(2\pi)^2 \xi_i \xi_j \widehat{u} = \frac{\xi_i \xi_j}{|\xi|^2} \widehat{f}(\xi).$$

So the associated multiplier is $m(\xi) = \frac{\xi_i \xi_j}{|\xi|^2}$, which is 0-homogeneous.

DEF. For $m : \mathbb{R}^d \rightarrow \mathbb{C}$ bounded, and $f \in \mathcal{S}(\mathbb{R}^d)$,

$$T_m f = (\widehat{f}(\xi) m(\xi))^\vee.$$

REM. T is an operator on $L^2(\mathbb{R}^d)$ with operator norm given by $\|m\|_\infty$.

DEF. We say $m : \mathbb{R}^d \rightarrow \mathbb{C}$ is a **Mihlin multiplier** if

$$\exists B > 0 \text{ s.t. } \forall \gamma \text{ multi-index with } |\gamma| \leq d+2, \quad \|\partial^\gamma m(\xi)\| \leq B |\xi|^{-|\gamma|}.$$

REM. If $m \in C^{d+2}(\mathbb{R}^d \setminus \{0\})$ is 0-homogeneous (i.e. $m(\lambda\xi) = m(\xi)$), then m is a Mihlin multiplier.

THEOREM 1. If m is a Mihlin multiplier, then $\exists C \sim_{d,p}$ such that for all $f \in \mathcal{S}(\mathbb{R}^d)$,

$$\|T_m f\|_p \leq CB \|f\|_p, \quad 1 < p < \infty.$$

REM. If m is a Mihlin multiplier, $\lambda > 0$, then the dilate $D_\lambda m(\xi) = m(\lambda\xi)$ is a Mihlin multiplier with the same constant B .

15.2 Reduction to Calderón–Zygmund kernels

DEF. A **Calderón–Zygmund (C–Z) kernel** is a function $K : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{C}$ such that

1. $|K(x)| \leq B |x|^{-d}$;
2. $\int_{|x| \geq 2|y|} |K(x) - K(x-y)| dy \leq B$;
3. $\int_{r < |x| < R} K(x) dx = 0$.

THEOREM (C-Z).

$$Tf = \lim_{\varepsilon \rightarrow 0} \int_{|x-y| > \varepsilon} K(x-y) f(y) dy$$

is bounded on L^p with norm $\lesssim CB$.

LEMMA 3. If $K : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{C}$ satisfies $|\nabla K| \leq B|x|^{-d-1}$, then condition (2) holds.

Proof.

$$\begin{aligned} \int_{|x| \geq 2|y|} |K(x) - K(x-y)| dy &\leq \int_{|x| \geq 2|y|} \underbrace{\int_0^1 |\nabla K(x+ty)| dt}_{|x+ty| \geq |x|/2 \geq |y|} |y| \\ &\leq \int_{|x| \geq 2|y|} B \left(\frac{|x|}{2}\right)^{-d-1} |y| \leq CB. \end{aligned}$$

Also note

$$T_m f = (m\hat{f})^\vee = m^\vee * f.$$

LEMMA 4. There exists $\psi \in C^\infty(\mathbb{R}^d)$ such that

1. $\psi \geq 0$, radial, with $\text{supp } \psi \subseteq \{\frac{1}{2} \leq |\xi| \leq 2\}$;
2. $\sum_{j \in \mathbb{Z}} \psi(2^{-j}\xi) = 1$ for $\xi \neq 0$.

Proof. Pick $\tilde{\psi}$ smooth, radial, with $\text{supp } \tilde{\psi} \subset \{\frac{1}{2} \leq |\xi| \leq 2\}$ and $\tilde{\psi} \geq 1$ on $\{\frac{2}{3} \leq |\xi| \leq \frac{4}{3}\}$. Let

$$\psi = \frac{\tilde{\psi}}{\sum_j \tilde{\psi}(2^{-j}\xi)}. \quad \square$$

LEMMA 5. Set

$$m_j(\xi) = m(\xi) \psi(2^{-j}\xi), \quad \text{so that } \sum_j m_j = m.$$

Define $K_j(x) = m_j^\vee(x)$ and $K(x) = \sum_{-N}^N K_j(x)$. Then (with constants **independent of K and N**):

1. $|K(x)| \leq CB|x|^{-d}$;
2. $|\nabla K(x)| \leq CB|x|^{-d-1}$.

15.3 Dyadic kernel estimates

CLAIM 1.

1. $|K_0(x)| \leq CB \min\{1, |x|^{-d-2}\}$;
2. $|\nabla K_0(x)| \leq CB \min\{1, |x|^{-d-2}\}$.

PROOF OF (I). It suffices to show

$$\|x_j^{d+2} K_0(x)\|_\infty \leq CB \quad \forall j, \quad \text{and} \quad \|K_0(x)\|_\infty \leq CB.$$

Now

$$\|x_j^{d+2} K_0(x)\|_\infty \leq \|(x_j^{d+2} K_0(x))^\wedge\|_{L^1} = C \|\partial_j^{d+2} m_0(\xi)\|_{L^1}.$$

Expanding the derivative of $m_0 = m \psi$:

$$\partial_j^{d+2}(m(\xi)\psi(\xi)) = \sum_{\ell=0}^{d+2} \binom{d+2}{\ell} \partial_j^\ell \psi \cdot \partial_j^{d+2-\ell} m,$$

and on $\text{supp } \psi$ (where $\frac{1}{2} \leq |\xi| \leq 2$),

$$|\partial_j^{d+2-\ell} m| \leq B |\xi|^{-(d+2-\ell)} \leq B \left(\frac{1}{2}\right)^{-(d+2-\ell)} \leq B 2^{d+2}.$$

Therefore

$$\|\partial_j^{d+2} m_0\|_{L^1} \leq C(d) B \|\mathbf{1}_{B(0,2)}\|_{L^1} \implies \text{(i)}.$$

PROOF OF (II). It suffices to show

$$\|x_j^{d+2} \partial_k K_0(x)\|_{L^\infty} \leq CB, \quad \|\partial_k K_0(x)\|_\infty \leq CB.$$

Then

$$\|x_j^{d+2} \partial_k K_0(x)\|_{L^\infty} \leq \|(x_j^{d+2} \partial_k K_0)^\wedge\|_{L^1} = C \|\partial_j^{d+2} [\xi_k m_0]\|_{L^1} \stackrel{\text{same argument}}{\leq} CB.$$

CLAIM 2.

1. $|K_j(x)| \leq CB 2^{jd} \min\{1, |2^j x|^{-d-2}\};$
2. $|\nabla K_j(x)| \leq CB 2^{j(d+1)} \min\{1, |2^j x|^{-d-2}\}.$

(The factor $2^{jd} / 2^{j(d+1)}$ comes from the change of variables $\xi \mapsto 2^{-j}\xi$; the exponent $-d-2$ comes from the $(d+2)$ -fold differentiation, as in Claim 1.)

PROOF OF THE LEMMA.

$$|K(x)| \leq \sum_{-N}^N |K_j(x)| \leq CB \sum_{j=-N}^N 2^{jd} \min\{1, |2^j x|^{-d-2}\}.$$

Split according to whether 2^j is below or above $|x|^{-1}$:

$$\begin{aligned} &\leq CB \sum_{2^j < |x|^{-1}} 2^{jd} + CB \sum_{2^j \geq |x|^{-1}} |2^j x|^{-d-2} 2^{jd} \\ &\leq CB |x|^{-d} + CB |x|^{-d-2} \sum_{2^j \geq |x|^{-1}} 2^{-2j} \leq CB |x|^{-d}. \quad \square \end{aligned}$$

(The ∇K estimate follows by the same splitting, using Claim 2(ii).)

15.4 Proof of the Mikhlin theorem

For $f \in \mathcal{S}(\mathbb{R}^d)$,

$$T_m f = \mathcal{F}^{-1}(m \hat{f}) = \mathcal{F}^{-1}\left(\sum_j m_j \hat{f}\right).$$

Writing $K^N = \sum_{-N}^N K_j$ (which is uniformly a C-Z kernel by Lemma 5),

$$\|T_m f\|_p = \|\lim f * K^N\|_p \leq \liminf \|f * K^N\|_p \leq CB \|f\|_p. \quad \blacksquare$$

16 Littlewood–Paley Theory

16.1 Dimension $d = 1$

For $f \in \mathcal{S}(\mathbb{R})$, define the **(rough) Littlewood–Paley projection**

$$\pi_j f(x) = \left(\mathbf{1}_{[2^j, 2^{j+1}] \cup (-2^{j+1}, -2^j]} \cdot \widehat{f} \right)^\vee, \quad j \in \mathbb{Z}.$$

Then (orthogonality on L^2):

$$\left\| \left(\sum_{j \in \mathbb{Z}} |\pi_j f|^2 \right)^{1/2} \right\|_{L^2(\mathbb{R})}^2 = \sum_{j \in \mathbb{Z}} \|\pi_j f\|_{L^2}^2 = \|f\|^2,$$

so heuristically $f \approx (\pi_j f)_{j \in \mathbb{Z}}$.

DEF. The rough Littlewood–Paley square function is

$$S_\pi f := \left(\sum_j |\pi_j f(x)|^2 \right)^{1/2}.$$

THEOREM. For $1 < p < +\infty$, there exists C_p such that

$$\frac{1}{C_p} \|f\|_p \leq \|S_\pi f\|_p \leq C_p \|f\|_p.$$

The mechanism behind the estimate is Khinchine's inequality:

$$\left\| \sum_{j=1}^N a_j r_j \right\|_p \approx \left(\sum a_j^2 \right)^{1/2}, \quad r_j = \text{Rademacher ("wave").}$$

Here $\pi_j f$ is a "wave" of frequency 2^j . Khinchine says: a sum of waves whose frequencies double at each step $\approx$ the square-sum of the coefficients:

$$|\pi_j f(x)| \longleftrightarrow a_j, \quad \pi_j f(x) \approx a_j e^{2\pi i 2^j x}.$$

Proof: not proven here. (One recovers π_j from **Hilbert transforms**: the sharp cutoff $\mathbf{1}_{[2^j, 2^{j+1}]}$ is a difference of shifted Hilbert-transform-type step multipliers — by shifting the Hilbert transform and subtracting, one recovers π_j .)

16.2 Higher dimensions

In higher dimensions, however, the sharp cutoff is no longer bounded.

FACT. $f \mapsto (\mathbf{1}_{B(0,1)} \widehat{f})^\vee$ is **unbounded** on L^p if $d \geq 2$ (the ball multiplier problem).

Instead, for each $j \in \mathbb{Z}$, define **smooth Littlewood–Paley projections**:

$$P_j f = (\psi_j \widehat{f})^\vee, \quad \psi \in C_c^\infty(\mathbb{R}^n \setminus \{0\}), \quad \sum_j \psi_j = \sum_j \psi(2^{-j} \xi) \equiv 1,$$

with $P_j f$ localized at frequency $\sim 2^j$. Define the smooth square function

$$S_P f = \left(\sum_j |P_j f|^2 \right)^{1/2}.$$

THEOREM. Let $1 < p < \infty$. There exists $C_{p,d}$ such that

$$\|S_P f\|_{L^p} \approx_{C_{p,d}} \|f\|_{L^p}.$$

Proof (uses Khinchine + Mikhlin).

Set $m(\xi) = \sum_{-N}^N r_j \psi_j(\xi)$.

CLAIM: $m(\xi)$ satisfies the Mikhlin multiplier condition. Indeed,

$$|D^r m(\xi)| \leq \sum_{-N}^N |D^r \psi_j(\xi)| = \sum_{-N}^N 2^{-j|r|} |D^r \psi(\xi/2^j)| \leq \sum_{-N}^N |\xi|^{-r} |D^r \psi(\xi/2^j)| \leq |\xi|^{-r} C_r,$$

using that for each ξ the supports of $\psi(\cdot/2^j)$ **overlap only ~ 3 times**.

UPPER BOUND. With $a_j = |P_j f(x)|$ and Khinchine,

$$\begin{aligned} \|S_P f\|_{L^p}^p &= \lim_{N \rightarrow \infty} \int \left(\sum_{-N}^N |P_j f|^2 \right)^{p/2} \leq C_p \int \mathbb{E} \left| \sum_j r_j P_j f(x) \right|^p dx \\ &= C_p \mathbb{E} \int \underbrace{\left| \sum_j r_j P_j f(x) \right|^p}_{=(m\widehat{f})^\vee} dx \leq C_{p,d} \|f\|_{L^p}, \end{aligned}$$

the last step by the Mikhlin theorem applied to m . This gives $\|S_P f\|_p \leq C_{p,d} \|f\|_p$.

LOWER BOUND (DUALITY). Find $\tilde{\psi}$ with $\tilde{\psi} \equiv 1$ on $\text{supp } \psi$; define $\tilde{\psi}_j$ analogously and

$$\tilde{P}_j g(x) = \mathcal{F}^{-1}(\tilde{\psi}_j \widehat{g})(x).$$

Then

$$\begin{aligned} \langle f, g \rangle &= \langle \mathcal{F} f, \mathcal{F} g \rangle = \int \widehat{f} \widehat{g} d\xi \\ &= \int \sum_j \psi_j \widehat{f} \widehat{g} d\xi \\ &= \sum_j \int \psi_j \widehat{f} \widetilde{\psi_j \widehat{g}} d\xi \\ &= \sum_j \langle \psi_j \widehat{f}, \widetilde{\psi_j \widehat{g}} \rangle = \sum_j \langle P_j f, \tilde{P}_j g \rangle. \end{aligned}$$

Hence

$$\begin{aligned} |\langle f, g \rangle| &= \left| \sum_j \langle P_j f, \tilde{P}_j g \rangle \right| \leq \int \sum_j |P_j f \cdot \tilde{P}_j g| \leq \int \left(\sum_j |P_j f|^2 \right)^{1/2} \left(\sum_j |\tilde{P}_j g|^2 \right)^{1/2} \\ &= \int S_P f \cdot \tilde{S}_P g \leq \|S_P f\|_{L^p} \|\tilde{S}_P g\|_{L^{p'}} \leq C_{d,p} \|S_P f\|_p \|g\|_{p'}. \end{aligned}$$

Taking $\|g\|_{p'} = 1$ and the supremum over such g ,

$$\|f\|_p \leq C_{p,d} \|S_P f\|_p. \quad \blacksquare$$

17 Applications of Littlewood–Paley Theory

17.1 Homogeneous Sobolev spaces and Sobolev embedding

DEF. For $s \geq 0$, the **homogeneous Sobolev space** $\dot{H}^s(\mathbb{R}^d)$ is the completion of $\mathcal{S}(\mathbb{R}^d)$ under

$$\|f\|_{\dot{H}^s(\mathbb{R}^d)} = \left(\int |\xi|^{2s} |\widehat{f}(\xi)|^2 d\xi \right)^{1/2}.$$

PROPOSITION. Let $0 \leq s < \frac{d}{2}$ and $\frac{1}{2} - \frac{1}{p} = \frac{s}{d}$. Then

$$\|f\|_{L^p(\mathbb{R}^d)} \leq C_{p,d} \|f\|_{\dot{H}^s} \quad \forall f \in \mathcal{S}(\mathbb{R}^d).$$

Proof.

SPECIAL CASE: $f \in \mathcal{S}(\mathbb{R}^d)$ with $\text{supp } \widehat{f} \subseteq \{\frac{R}{4} \leq |\xi| \leq 4R\}$. For such f ,

$$\|f\|_{\dot{H}^s}^2 = \int |\xi|^{2s} |\widehat{f}(\xi)|^2 \geq C_s R^{2s} \int |\widehat{f}|^2 = C_s R^{2s} \int |f|^2.$$

So it suffices to show

$$\|f\|_{L^p(\mathbb{R}^d)} \leq C_{p,d} R^s \|f\|_{L^2(\mathbb{R}^d)}, \quad s = d\left(\frac{1}{2} - \frac{1}{p}\right). \quad (**)$$

Define $g(x) = f(Rx) R^d \Rightarrow \widehat{g}(\xi) = \widehat{f}(\xi/R)$. Then $(**)$ is equivalent to

$$\|g\|_{L^p} \leq C_{p,d} \|g\|_{L^2}.$$

Find $\chi \equiv 1$ on $\frac{1}{4} \leq |x| \leq 4$; then $\widehat{g}(\xi) \cdot \chi = \widehat{g}$, so $g = \check{\chi} * g$. By **Young's inequality** (with $1 + \frac{1}{p} = \frac{1}{2} + \frac{1}{q}$),

$$\|g\|_{L^p} = \|\check{\chi} * g\|_p \leq \|\check{\chi}\|_q \|g\|_2.$$

GENERAL CASE:

$$\begin{aligned} \|f\|_p &\approx \|S_P f\|_p = \left\| \left(\sum_j |P_j f|^2 \right)^{1/2} \right\|_p \leq \left(\sum_j \|P_j f\|_p^2 \right)^{1/2} \\ &\lesssim \left(\sum_j \underbrace{2^{2js} \|P_j f\|_{L^2}^2} \right)^{1/2} \leq \left(\sum_j \int |\widehat{P_j f}|^2 |\xi|^{2s} \right)^{1/2} = \|f\|_{\dot{H}^s}. \quad \blacksquare \end{aligned}$$

(Middle step: apply the special case $(**)$ to $P_j f$, which has frequency $\sim 2^j$, giving $\|P_j f\|_p \lesssim 2^{js} \|P_j f\|_2$.)

17.2 Hölder spaces and their Littlewood–Paley characterization

HÖLDER SPACE.

$$C^\alpha(\mathbb{R}^n) = \left\{ f \mid [f]_\alpha = \sup_{x,y \in \mathbb{R}^d} \frac{|f(x+y) - f(x)|}{|y|^\alpha} < +\infty \right\}.$$

THEOREM. Suppose $|f(x)| \leq 1$. Then

$$f \in C^\alpha \iff \exists A \text{ s.t. } \|P_j f\|_{L^\infty(\mathbb{R}^d)} \leq A \cdot 2^{-j\alpha},$$

and moreover

$$C_{d,\alpha}^{-1} A \leq [f]_\alpha \leq C_{d,\alpha} A.$$

17.2.1 Proof

($\Rightarrow$ DIRECTION / KERNEL SIDE.) Since $P_j f = \check{\psi}_j * f$ and ψ_j is supported away from 0:

$$D^r \psi_j \in C_c^\infty(\mathbb{R}^d \setminus \{0\}) \Rightarrow D^r \psi_j(0) \equiv 0 \quad \forall r \Rightarrow \int x^r \check{\psi}_j(x) dx \equiv 0 \quad \forall r.$$

In particular $\int \check{\psi}_j = 0$, so

$$P_j f(x) = \check{\psi}_j * f = \int \check{\psi}_j(y) f(x-y) dy = \int \check{\psi}_j(y) (f(x-y) - f(x)) dy.$$

Hence, using $\check{\psi}_j(y) = 2^{jd} \check{\psi}(2^j y)$ and the change of variables $u = 2^j y$,

$$\begin{aligned} |P_j f(x)| &\leq \int |\check{\psi}_j(y)| |f(x-y) - f(x)| dy \\ &\leq [f]_\alpha \int |\check{\psi}_j(y)| |y|^\alpha dy \\ &= [f]_\alpha \int 2^{jd} |\check{\psi}(2^j y)| |y|^\alpha dy \\ &= C 2^{-j\alpha} [f]_\alpha. \end{aligned}$$

($\Leftarrow$ DIRECTION / CONVERSE.)

LEMMA (BERNSTEIN-TYPE). Define $Pf(x) = g * f$, where $f \in L^\infty$ and g satisfies $\text{supp } \hat{g} \subset \{|\xi| < R\}$. Then

$$\|\nabla Pf\|_{L^p} \leq R \cdot C_d \|Pf\|_{L^p}, \quad 1 \leq p \leq \infty.$$

Proof. Since $\text{supp}(\hat{g}) \subseteq \{|\xi| \leq R\}$, we have $\hat{g}(\xi) = \hat{g}(\xi) \cdot \chi(\xi/R)$, i.e. $g = g * \check{\chi}_R$, so $\nabla g = g * \nabla \check{\chi}_R$. With

$$\check{\chi}_R(x) = R^d \check{\chi}(Rx), \quad \nabla \check{\chi}_R(x) = R^{d+1} (\nabla \check{\chi})(Rx),$$

we get $\nabla Pf = \nabla(g * f) = \nabla \check{\chi}_R * g * f$, hence

$$\|\nabla \check{\chi}_R * g * f\|_{L^p} \leq \|\nabla \check{\chi}_R\|_{L^1} \|Pf\|_{L^p} = R \|\nabla \check{\chi}\|_{L^1} \|Pf\|_{L^p}. \quad \square$$

Now let f satisfy $\|P_j f\|_{L^\infty} \leq A 2^{-j\alpha}$, and set $g_\ell = \sum_{- \ell}^\ell P_j f$. Decompose the increment into a low-frequency gradient term and a high-frequency tail:

$$|g_\ell(x+y) - g_\ell(x)| \leq \underbrace{\sum_{|j| \leq \ell} |P_j f(x+y) - P_j f(x)|}_{L_\ell(x,y)} + \underbrace{\sum_{j > \ell} 2 \|P_j f\|_\infty}_{H_\ell}.$$

HIGH-FREQUENCY ESTIMATE.

$$H_\ell \leq 2A \sum_{j \geq \ell} 2^{-j\alpha} \lesssim A 2^{-\ell\alpha}.$$

LOW-FREQUENCY ESTIMATE.

$$\begin{aligned} |P_j f(x+y) - P_j f(x)| &= \left| \int_0^1 (\nabla P_j f) \cdot y \, dt \right| \\ &\leq \|\nabla P_j f\|_\infty |y| \leq 2^j \|P_j f\|_\infty |y| \\ &\leq A 2^{j(1-\alpha)} |y|. \end{aligned}$$

Summing,

$$L_\ell(x, y) \leq \sum_{|j| \leq \ell} A 2^{j(1-\alpha)} |y| \lesssim A 2^{\ell(1-\alpha)} |y|.$$

OPTIMIZE. Take $2^\ell = \frac{1}{|y|}$. Then

$$H_\ell \lesssim A |y|^\alpha, \quad L_\ell(x, y) \lesssim A |y|^\alpha.$$

So $|g_\ell(x+y) - g_\ell(x)| \leq CA |y|^\alpha$, i.e. $[g_\ell]_\alpha \leq CA$.

Since $g_\ell \rightarrow f$, we conclude $[f]_\alpha \leq CA$. $\blacksquare$

Part VI

Oscillatory Integrals and Fourier Restriction

18 Oscillatory Integrals and Stationary Phase

18.1 Fourier transforms of surface measures

Consider the **Schrödinger equation**

$$\begin{cases} i\partial_t u + \Delta u = 0, \\ u(x, 0) = u_0(x), \end{cases} \quad u_0 \in \mathcal{S}(\mathbb{R}^d).$$

On the Fourier side:

$$\begin{cases} i\partial_t \hat{u} - 4\pi^2 |\xi|^2 \hat{u} = 0, \\ \hat{u}(\xi, 0) = \hat{u}_0(\xi), \end{cases} \quad \implies \quad \hat{u}(\xi, t) = e^{-4\pi^2 i |\xi|^2 t} \hat{u}_0(\xi).$$

Hence

$$\begin{aligned} u &= \mathcal{F}_\xi^{-1}(e^{-4\pi^2 i |\xi|^2 t} \hat{u}_0(\xi))(x) = \int e^{2\pi i x \cdot \xi} e^{-4\pi^2 i |\xi|^2 t} \hat{u}_0(\xi) \, d\xi \\ &= \int_{\mathbb{R}^d} e^{2\pi i (x, t) \cdot (\xi, -2\pi |\xi|^2)} \hat{u}_0(\xi) \, d\xi = \int_{\mathbb{R}^{d+1}} e^{2\pi i (x, t) \cdot (\xi, \tau)} \hat{u}_0(\xi) \, d\mu(\xi, \tau), \end{aligned}$$

where μ is the measure on the paraboloid defined (as a distribution) by

$$\int F(\xi, \tau) \, d\mu(\xi, \tau) = \int_{\mathbb{R}^d} F(\xi, -2\pi |\xi|^2) \, d\xi.$$

Thus μ is supported on the downward paraboloid $\tau = -2\pi|\xi|^2$.

This representation raises the question of how rapidly the Fourier transform of the surface measure decays. At positive times the initially concentrated profile spreads and oscillates.

One can show $u_0 \in \mathcal{S}(\mathbb{R}^d) \Rightarrow u(\cdot, t) \in \mathcal{S}(\mathbb{R}^d)$ for $t > 0$:

$$u(x, t) = C_d \frac{1}{t^{d/2}} \int e^{i|x-y|^2/4t} u_0(y) dy \implies \|u(\cdot, t)\|_\infty \leq C_d t^{-d/2} \|u_0\|_{L^1}.$$

$u(\cdot, t)$ decays faster than $|x|^{-N}$ for all N , because it is Schwartz in x . Therefore

$$|\hat{\mu}(x, t)| \leq C_d \|(x, t)\|^{-d/2}, \quad d = \dim(\text{hypersurface}).$$

CONJECTURE: $|\hat{\mu}(x)| \lesssim |x|^{-d/2}$.

The conjectured decay fails for flat hypersurfaces. For example, in $\mathbb{R}^{d+1}$ take $\mu(\xi, \tau) = d\xi \times \delta_0(\tau)$. Then

$$\mathcal{F}^{-1}(u d\mu) = \int e^{2\pi i(x, t) \cdot (\xi, \tau)} u(\xi) d\xi \delta_0 d\tau = \int e^{2\pi i x \cdot \xi} u(\xi) d\xi$$

has no decay in the t direction, precisely because there is no curvature in that direction. Non-vanishing Gaussian curvature restores decay:

Theorem 18.1 (Decay of curved surface measure). *Let $M \subseteq \mathbb{R}^d$ be a hypersurface with non-vanishing Gauss curvature, and let $d\mu = \psi d\sigma$, where $\psi \in C_c^\infty$ and $d\sigma$ is surface measure. Then*

$$|\hat{\mu}(\xi)| \leq C(M, \psi) |\xi|^{-\frac{d-1}{2}}.$$

18.2 Stationary phase

The proof rests on integrals of the form

$$I(\lambda) = \int e^{i\lambda\phi(\xi)} a(\xi) d\xi, \quad a \in C_c^\infty.$$

There are two distinct regimes:

1. If $\nabla\phi \neq 0$ on $\text{supp}(a)$, repeated integration by parts gives rapid decay.
2. If $\nabla\phi(\xi_0) = 0$ for some ξ_0 , the main contribution comes from a neighbourhood of the critical point.

18.2.1 Non-stationary phase

If $\nabla\phi \neq 0$ for all $\xi \in \text{supp}(a)$, then

$$|I(\lambda)| = \left| \int e^{i\lambda\phi} a(\xi) d\xi \right| \leq C_{\phi, a, N} \lambda^{-N} \quad \forall N.$$

PROOF (INTEGRATION BY PARTS).

Case $d = 1$. Since $[e^{i\lambda\phi}]' = i\lambda\phi' e^{i\lambda\phi}$,

$$\int e^{i\lambda\phi} a = \int \frac{1}{i\lambda\phi'} [e^{i\lambda\phi}]' a = \frac{1}{i\lambda} \int [e^{i\lambda\phi}]' \frac{a}{\phi'} = -\frac{1}{i\lambda} \int e^{i\lambda\phi} \left[\frac{a}{\phi'} \right]' = \frac{1}{i\lambda} \int e^{i\lambda\phi} a_1,$$

and repeat several times (each pass gains a factor λ^{-1}).

Case $d \geq 2$.

$$\nabla e^{i\lambda\phi} = e^{i\lambda\phi} \nabla\phi \cdot i\lambda \implies \nabla\phi \cdot \nabla e^{i\lambda\phi} = e^{i\lambda\phi} |\nabla\phi|^2 i\lambda \implies e^{i\lambda\phi} = \frac{\nabla\phi \cdot \nabla e^{i\lambda\phi}}{|\nabla\phi|^2} \cdot \frac{1}{i\lambda},$$

$$\implies \int e^{i\lambda\phi} a \, d\xi = \int \frac{\nabla\phi \cdot \nabla e^{i\lambda\phi}}{|\nabla\phi|^2} \frac{a}{i\lambda} = \frac{1}{i\lambda} \left(- \int \operatorname{div} \left(a \frac{\nabla\phi}{|\nabla\phi|^2} \right) e^{i\lambda\phi} \, d\xi \right).$$

18.2.2 The stationary regime

For a stationary point we need **degeneracy** of ϕ at ξ_0 .

Example: $\phi \equiv 0 \Rightarrow I(\lambda) = \int a(\xi) \, d\xi$ — no decay.

We need $\det(D^2\phi(\xi_0)) \neq 0$.

Theorem 18.2 (Stationary phase). *Suppose $\nabla\phi(\xi_0) = 0$, $\nabla\phi \neq 0$ on $\operatorname{supp}(a) \setminus \{\xi_0\}$, and $\det D^2\phi(\xi_0) \neq 0$. Then*

$$(1) \quad |I(\lambda)| = \left| \int_{\mathbb{R}^d} e^{i\lambda\phi} a(\xi) \, d\xi \right| \leq C \lambda^{-d/2},$$

$$(2) \quad |\partial_\lambda^k (e^{-i\lambda\phi(\xi_0)} I(\lambda))| \leq C \lambda^{-\frac{d}{2}-k}.$$

PROOF. Assume $\xi_0 = 0$. Write (with $\chi = 1$ on $\{|\xi| \leq 1\}$, $\chi = 0$ on $\{|\xi| \geq 2\}$)

$$\int_{\mathbb{R}^d} e^{i\lambda\phi(\xi)} a(\xi) \, d\xi = \underbrace{\int e^{i\lambda\phi} \chi(\xi/\varepsilon) a \, d\xi}_A + \underbrace{\int (1 - \chi(\xi/\varepsilon)) e^{i\lambda\phi} a \, d\xi}_B.$$

18.2.3 Estimating the main and error terms

ONE-DIMENSIONAL MODEL.

$$A(\lambda) \leq \int |\chi(\xi/\varepsilon)| a(\xi) \, d\xi \leq \varepsilon \cdot C,$$

$$B(\lambda) \stackrel{\text{IBP}}{=} \left(\frac{1}{i\lambda} \right)^N \int e^{i\lambda\phi(\xi)} L^N((1 - \chi(\xi/\varepsilon))a(\xi)) \, d\xi, \quad Lf(\xi) = \left(\frac{f(\xi)}{\phi'(\xi)} \right)'.$$

Applying L^N produces three types of terms:

1. derivatives landing on a ;
2. derivatives landing on $(1 - \chi) \rightsquigarrow$ factors of ε^{-N} ;
3. factors from $1/\phi' \rightsquigarrow \frac{1}{|\xi|^{2N}}$ (since $\phi'(\xi) \sim |\xi|$ near 0).

Hence

$$|B(\lambda)| \leq C \int_{|\xi| \geq \varepsilon} \left(\frac{1}{|\xi|^{2N}} + \frac{\varepsilon^{-N}}{|\xi|^N} \right) d\xi \approx \varepsilon^{-(2N-1)},$$

so $|B| \lesssim \lambda^{-N} \varepsilon^{-(2N-1)}$. Balancing with $|A| \lesssim \varepsilon$ via $\varepsilon^{2N} = \lambda^{-N}$ gives

$\varepsilon = \lambda^{-1/2}.$

HIGHER DIMENSION, $d \geq 2$, $\xi_0 = 0$, $\det(D^2\phi) \neq 0 \Rightarrow |\nabla\phi(\xi)| \sim |\xi|$ (use Taylor expansion). Let χ be a smooth bump, $\chi(\xi) = 1$ if $|\xi| \leq 1$, $\chi(\xi) = 0$ if $|\xi| \geq 2$:

$$I(\lambda) = \underbrace{\int e^{i\lambda\phi} \chi(\xi/\varepsilon) a \, d\xi}_{A(\lambda)} + \underbrace{\int e^{i\lambda\phi} (1 - \chi(\xi/\varepsilon)) a \, d\xi}_{B(\lambda)}.$$

$$|A(\lambda)| \leq \int |\chi(\xi/\varepsilon) a(\xi)| \, d\xi \leq C \int \chi(\xi/\varepsilon) \, d\xi \leq C\varepsilon^d.$$

For $B(\lambda)$, integrate by parts:

$$B(\lambda) = \int e^{i\lambda\phi(\xi)} (1 - \chi(\xi/\varepsilon)) a = \left(\frac{-1}{i\lambda}\right)^N \int e^{i\lambda\phi} L^N((1 - \chi(\xi/\varepsilon)) a) \, d\xi, \quad Lf = \operatorname{div}\left(\frac{\nabla\phi f(\xi)}{|\nabla\phi|^2}\right).$$

Lemma 18.3. *On $\{|\xi| \geq \varepsilon\}$ one has*

$$\left|L^N((1 - \chi(\xi/\varepsilon)) a(\xi))\right| \leq C \frac{\varepsilon^{-N}}{|\xi|^N}.$$

Given the claim,

$$|B(\lambda)| \leq \frac{1}{\lambda^N} \int_{|\xi| \geq \varepsilon} \varepsilon^{-N} \frac{1}{|\xi|^N} \, d\xi \leq \frac{1}{\lambda^N} \varepsilon^{-N} \int_{\geq \varepsilon} r^{d-1} r^{-N} \, dr = \frac{\varepsilon^{-N}}{\lambda^N} \varepsilon^{-N+d} = \frac{\varepsilon^{-2N+d}}{\lambda^N},$$

$$\Rightarrow |I| \leq \varepsilon^d + \frac{\varepsilon^{-2N+d}}{\lambda^N}; \quad \text{take } \varepsilon \text{ with } \varepsilon^{2N} = \lambda^{-N} \Rightarrow \varepsilon = \lambda^{-1/2}.$$

PROOF OF CLAIM.

$$L^N((1 - \chi(\xi/\varepsilon)) a) = \cdots \left(\operatorname{div} \frac{\nabla\phi}{|\nabla\phi|^2} \operatorname{div} \left(\frac{\nabla\phi(1-\chi)}{|\nabla\phi|^2} a \right) \right).$$

- *Case 1:* derivative on a .
- *Case 2:* derivative on $\chi(\xi/\varepsilon) \rightsquigarrow \varepsilon^{-N}$.
- *Case 3:* on $\frac{\nabla\phi}{|\nabla\phi|^2}$:

$$\partial_\ell \left(\frac{\nabla\phi}{|\nabla\phi|^2} \right) = \frac{\nabla\partial_\ell\phi}{|\nabla\phi|^2} - \nabla\phi \cdot \frac{2\langle \nabla\partial_\ell\phi, \nabla\phi \rangle}{|\nabla\phi|^4} \sim \frac{1}{|\nabla\phi|^2} \rightsquigarrow \frac{1}{|\nabla\phi|^{2N}}.$$

18.2.4 Completion of the estimate

(Note: each step adds one more factor; the two parts together account for the N derivatives.)

$$|I| \leq C \sum_{k=0}^N \varepsilon^{-k} \left(\sum_{\ell=0}^{N-k} \frac{1}{|\nabla\phi|^{N+\ell}} \right).$$

If $|\xi| \geq \varepsilon$ (so $k + \ell \leq N$),

$$\leq C \sum_{k=0}^N \frac{1}{|\xi|^N} \sum_{\ell=0}^{N-k} \varepsilon^{-(k+\ell)} \leq \frac{\varepsilon^{-N}}{|\xi|^N} \cdot C, \quad \varepsilon^{-(k+\ell)} = \varepsilon^{-N} \cdot \underbrace{\varepsilon^{N-(k+\ell)}}_{\leq 1}.$$

PART 2 (DERIVATIVE BOUND).

$$I(\lambda) = \int e^{i\lambda\phi} a(\xi) d\xi = e^{i\lambda\phi(\xi_0)} \int_{\mathbb{R}^d} e^{i\lambda(\phi-\phi(\xi_0))} a(\xi) d\xi$$

$$\Rightarrow e^{-i\lambda\phi(\xi_0)} I(\lambda) = \int e^{i\lambda(\phi-\phi(\xi_0))} a(\xi) d\xi$$

$$\Rightarrow \frac{d}{d\lambda}(\dots) = \int e^{i\lambda(\phi-\phi(\xi_0))} (\phi-\phi(\xi_0)) a(\xi) \cdot i d\xi = \frac{-1}{i\lambda} \int e^{i\lambda(\phi-\phi(\xi_0))} \operatorname{div} \left(\frac{\nabla \phi}{|\nabla \phi|^2} i(\phi-\phi(\xi_0)) a \right) d\xi,$$

where $\phi - \phi(\xi_0) \approx |\xi|^2$ and the integrand is C_c^∞ . Each λ -derivative gains λ^{-1} .

18.3 Fourier decay of surface measures

In $\mathbb{R}^{d+1}$ let M be a smooth hypersurface, $d\mu = \psi d\sigma$ with $\psi \in C_c^\infty(\mathbb{R}^{d+1})$, $d\sigma$ the surface measure, and $N(x)$ = normal vector at x . For $\xi \in \mathbb{R}^{d+1}$:

1. If ξ is **not parallel** to $N(x)$ for all $x \in \operatorname{supp} \psi \cap M$, then $|\hat{\mu}(\xi)| \leq C_{\psi, M} |\xi|^{-k} \forall k \geq 0$.
2. If M has **non-vanishing curvature**, then $|\hat{\mu}(\xi)| \leq C_{\psi, M} |\xi|^{-d/2}$.

PROOF (SETUP). Cover $M \cap \operatorname{supp} \psi$ by balls (B_i) such that on $\frac{1}{2}B_i$, M is a graph $f_1, \dots, f_n$. Take a finite subcover and a partition of unity $\{\psi_i\}$ subordinate to (B_i) with $\psi_i = 1$ on $\frac{1}{2}B_i$, $\operatorname{supp} \psi_i \subset B_i$. Then

$$\hat{\mu}(\xi) = \int e^{-2\pi i x \cdot \xi} d\mu(x) = \sum_i \int e^{-2\pi i x \cdot \xi} \underbrace{\psi_i \psi(x) d\sigma(x)}_{d\mu_i(x)}.$$

It suffices to show $|\hat{\mu}_i(\xi)| \leq C_{\psi, M} |\xi|^{-d/2}$. Writing M as a graph $x = (y, f(y))$,

$$\hat{\mu}_i(\xi) = \int e^{-2\pi i x \cdot \xi} \psi_i \psi d\sigma = \int_{\mathbb{R}^d} e^{-2\pi i (y, f(y)) \cdot \xi} \underbrace{\psi_i(y, f(y)) \psi(y, f(y)) \sqrt{1 + |\nabla f|^2}}_{a(y)} dy,$$

which "looks like the integral before."

19 Fourier Restriction Theory

19.1 Critical points and the restriction problem

Write $\xi = \lambda\nu$:

$$\hat{\mu}_i(\xi) = \int_{\mathbb{R}^d} e^{-2\pi i \lambda (y, f(y)) \cdot \nu} \psi_i(y, f(y)) \psi(y, f(y)) \sqrt{1 + |\nabla f|^2} dy.$$

1. WHEN DOES THE PHASE HAVE A CRITICAL POINT? Consider $x \mapsto -\langle x, \xi \rangle$ restricted to M . It has a critical point at $x_0 \in M$

$$\iff \forall \text{ curves } r : (-1, 1) \rightarrow M, r(0) = x_0, t \mapsto -\langle r(t), \xi \rangle \text{ is critical at } 0$$

$$\Longleftrightarrow -\langle r'(0), \xi \rangle = 0 \Longleftrightarrow \xi \parallel N(x_0).$$

So if there is **no** critical point $\Rightarrow$ non-stationary phase (case 1 above).

2. THERE IS A CRITICAL POINT AT $x_0 \Longleftrightarrow \xi \parallel N(x_0)$.

- Only finitely many such points can occur.
- Refine the balls so that at most one ball meets each critical point:

$$\hat{\mu}_i = \int_{\mathbb{R}^n} e^{-2\pi i \lambda (y, f(y)) \cdot \nu} \underbrace{\psi_i \psi}_a dy, \quad \xi \parallel N(x_0).$$

With $\nu = N(x_0) = (-\partial_1 f, \dots, -\partial_d f, 1) / \sqrt{1 + |\nabla f(x_0)|^2}$, the phase is

$$\phi(y) = (y, f(y)) \cdot (-\partial_1 f(y_0), \dots, -\partial_d f(y_0), 1),$$

$$\partial_i \partial_j \phi = (0, \partial_{ij} f) \cdot (\dots, 1) / \sqrt{1 + |\nabla f|^2} \implies \det(\phi_{ij}) \neq 0 \Longleftrightarrow \det(\partial_{ij} f) \neq 0.$$

So non-degenerate Hessian of f (non-vanishing Gaussian curvature) $\Longleftrightarrow$ non-degenerate stationary point, giving the $\lambda^{-d/2}$ decay.

19.1.1 Formulation of the restriction problem

For $f \in L^1(\mathbb{R}^d)$, the Fourier transform $\hat{f}(\xi) = \int_{\mathbb{R}^d} f(x) e^{-2\pi i x \cdot \xi} dx$ is continuous and decays at infinity.

- $L^1(\mathbb{R}^d) \xrightarrow{\mathcal{F}} C_0(\mathbb{R}^d)$, with $\|\hat{f}\|_{L^\infty} \leq \|f\|_{L^1}$.
- If $f \in L^1 \cap L^2$, then $\|\hat{f}\|_{L^2} = \|f\|_{L^2}$ (Plancherel).
- By Riesz–Thorin: $\|\hat{f}\|_{p'} \leq \|f\|_p$ for $1 \leq p \leq 2$.

THE RESTRICTION PROBLEM asks, for given $1 \leq p \leq 2$ and $S \subseteq \mathbb{R}^d$, whether $\hat{f}|_S$ is meaningful.

- If S has positive measure, $\hat{f} \in L^{p'} \Rightarrow$ meaningful.
- If S has measure 0: for $p = 1$, $\hat{f} \in C_0(\mathbb{R}^d)$; but for $p = 2$, $\hat{f} \in L^2(\mathbb{R}^d)$ only, so restriction is **impossible**.

19.2 The L^2 restriction theorem

STEIN'S OBSERVATION. If $S = S^{d-1} \subseteq \mathbb{R}^d$, then restriction holds for $f \in L^p$, $1 < p < 2$: i.e. $\hat{f}$ has more structure in $L^{p'}$ than the trivial bound suggests.

Theorem 19.1 (L^2 restriction). *Let $d \geq 2$ and $1 \leq p < \frac{4d}{3d+1}$. Then*

$$\|\hat{f}\|_{L^2(S^{d-1}, d\sigma)} \lesssim_{d,p} \|f\|_{L^p(\mathbb{R}^d)} \quad \forall f \in \mathcal{S}(\mathbb{R}^d),$$

where $d\sigma$ is surface measure.

PROOF. Fix d, p, f :

$$\begin{aligned}\|\widehat{f}\|_{L^2(S^{d-1}, d\sigma)}^2 &= \int_{S^{d-1}} |\widehat{f}|^2 d\sigma = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{S^{d-1}} f(x) \overline{f(y)} e^{-2\pi i(x-y) \cdot \xi} dx dy d\sigma(\xi) \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f(x) \overline{f(y)} (d\sigma)^\vee(y-x) dx dy = \langle f, f * (d\sigma)^\vee \rangle.\end{aligned}$$

Now $(d\sigma)^\vee(x) \lesssim 1$ trivially. By rotation invariance, with $x = (0, \dots, |x|)$,

$$I(x) = (d\sigma)^\vee(x) = \int_{\mathbb{R}^d} e^{2\pi i \xi \cdot x} d\sigma(\xi) = \int_{\mathbb{R}^d} e^{2\pi i \xi_d \cdot |x|} d\sigma(\xi)$$

(the height function is ξ_d). By the (non-)stationary phase method, splitting the sphere into three parts — *good parts* ($\lesssim |x|^{-N}$) and *bad parts* (the two poles) — one gets

$$(d\sigma)^\vee(x) \lesssim_d |x|^{-\frac{d-1}{2}}, \quad |x| \geq 1, \quad \text{i.e.} \quad (d\sigma)^\vee(x) \lesssim_d \langle x \rangle^{-\frac{d-1}{2}}.$$

Since $\langle x \rangle^{-a} \in L^p \iff p > \frac{d}{a}$, we get $(d\sigma)^\vee \in L^q$ for $q > \frac{2d}{d-1}$.

By Young's inequality ($\|f * g\|_{L^r} \leq \|f\|_{L^p} \|g\|_{L^q}$, $1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}$),

$$\|\widehat{f}\|_{L^2(S^{d-1}, d\sigma)}^2 = \langle f, f * (d\sigma)^\vee \rangle, \quad \|f * (d\sigma)^\vee\|_{L^{p'}} \lesssim \|f\|_{L^p},$$

$$|\langle f, f * (d\sigma)^\vee \rangle| \leq \|f\|_{L^p(\mathbb{R}^d)} \|f * (d\sigma)^\vee\|_{L^{p'}(\mathbb{R}^d)} \lesssim \|f\|_{L^p}^2. \quad \blacksquare$$

Conjecture 19.2 (Stein). *For $d \geq 2$,*

$$\|\widehat{f}\|_{L^q(S^{d-1}, d\sigma)} \lesssim_{d,p,f} \|f\|_{L^p}, \quad p < \frac{2d}{d+1}, \quad q \leq \frac{d-1}{d+1} p'.$$

19.3 Restriction–extension duality

$d = 2$: SOLVED.

19.3.1 The dual formulation

Let μ be a Radon measure on $\mathbb{R}^d$, $1 \leq p, q \leq \infty$, $A > 0$. The following are equivalent.

- **(1) Restriction estimate** ($f \in C_c^\infty(\mathbb{R}^d)$):

$$\|\widehat{f}\|_{L^q(\mathbb{R}^d, \mu)} \leq A \|f\|_{L^p}.$$

- **(2) Extension estimate** ($g \in C_c^\infty$):

$$\|(g d\mu)^\vee\|_{L^{p'}} \leq A \|g\|_{L^{q'}(\mathbb{R}^d, \mu)}.$$

PROOF. (1) $\Rightarrow$ (2).

$$f \mapsto \int f \cdot \overline{(g d\mu)^\vee} dx = \int \widehat{f} g d\mu \leq \|\widehat{f}\|_{L^q(\mathbb{R}^d, \mu)} \|g\|_{L^{q'}(d\mu)} \leq A \|f\|_{L^p} \|g\|_{L^{q'}(d\mu)}$$

$$\Rightarrow (g d\mu)^\vee \in L^{p'} \text{ bounded by } A \|g\|_{L^{q'}(d\mu)}.$$

(2) $\Rightarrow$ (1).

$$g \mapsto \int f \overline{(g d\mu)^\vee} = \int \hat{f} g d\mu,$$

and

$$\begin{aligned} \|(g d\mu)^\vee\|_{L^{p'}} \|f\|_{L^p} &\leq A \|g\|_{L^{q'}(\mathbb{R}^d, \mu)} \|f\|_{L^p} \\ &\Rightarrow \|\hat{f}\|_{L^q} \leq A \|f\|_{L^p}. \end{aligned}$$

Write $R_{S^{d-1}}(p \rightarrow q)$ = "Stein's conjecture holds for p, q ". By duality,

$$\|(g d\sigma)^\vee\|_{L^{p'}} \leq A \|g\|_{L^{q'}(S^{d-1}, d\sigma)}, \quad \text{in particular } (g d\sigma)^\vee \in L^{p'}.$$

NECESSITY OF THE RANGE (adapting the bound for $(d\sigma)^\vee$ via the stationary-phase/SWM asymptotic):

$$\begin{aligned} (g d\sigma)^\vee(x) &= \frac{1}{|x|^{\frac{d-1}{2}}} \left(C_+ g\left(\frac{x}{|x|}\right) e^{2\pi i|x|} + C_- g\left(\frac{x}{|x|}\right) e^{-2\pi i|x|} \right) + O_{g,d}(|x|^{-\frac{d}{2}}), \\ &\Rightarrow \|(g d\sigma)^\vee\|_{L^{p'}}^{p'} \gtrsim \int_1^\infty r^{-\frac{d-1}{2}p'} r^{d-1} dr. \end{aligned}$$

For convergence we need $(-\frac{d-1}{2})p' + (d-1) < -1$, i.e.

$$p' > \frac{2d}{d-1} \Rightarrow \frac{1}{1-\frac{1}{p}} > \frac{2d}{d-1} \Rightarrow 1 - \frac{1}{p} < \frac{d-1}{2d} \Rightarrow \frac{1}{p} > \frac{d+1}{2d} \Rightarrow \boxed{p < \frac{2d}{d+1}}.$$

19.4 Necessary exponents, Stein's conjecture, and Kakeya

Proposition 19.3 (Preliminary L^2 restriction estimate). *Let $d \geq 2$ and $1 \leq p < \frac{4d}{3d+1}$. Then*

$$\|\hat{f}\|_{L^2(S^{d-1}, d\sigma)} \leq C_{p,d} \|f\|_{L^p(\mathbb{R}^d)} \quad \forall f \in \mathcal{S}(\mathbb{R}^d).$$

Conjecture 19.4 (Restriction to the sphere). *For $d \geq 2$,*

$$\|\hat{f}\|_{L^q(S^{d-1}, d\sigma)} \leq C_{d,p,q} \|f\|_{L^p(\mathbb{R}^d)} \quad \forall f \in C_c^\infty(\mathbb{R}^d),$$

$$\boxed{\forall p < \frac{2d}{d+1}, \quad q \leq \frac{d-1}{d+1} p'}.$$

Status: $d = 2$ is solved; for $d \geq 3$, only a partial range is known.

KAKEYA SET ($d \geq 2$): a set containing a unit line segment in every direction.

- **Conjecture:** Hausdorff dimension = d . (The notes here abbreviate; the standard Kakeya conjecture asserts full dimension d .)
- **Restriction $\Rightarrow$ Kakeya conjecture.**

19.4.1 Asymptotics of the extension operator

Assume $\|(g d\sigma)^\vee\|_{L^{p'}} \leq C\|g\|_{L^{q'}(S^{d-1}, d\sigma)}$. Then for all $g \in C_c^\infty(\mathbb{R}^d)$,

$$F(x) = (g d\sigma)^\vee(x) = \frac{1}{|x|^{\frac{d-1}{2}}} \left(C_+ g\left(\frac{x}{|x|}\right) e^{2\pi i|x|} + C_- g\left(\frac{x}{|x|}\right) e^{-2\pi i|x|} \right) + O(|x|^{-\frac{d}{2}}), \quad |x| \geq 1.$$

PROOF.

$$F(x) = \int_{S^{d-1}} g(\omega) e^{2\pi i x \cdot \omega} d\sigma(\omega).$$

Assume $x = (0, \dots, r)$ (using rotation invariance of $d\sigma$), so

$$F(x) = \int_{S^{d-1}} g(\omega) e^{2\pi i r \omega_d} d\omega.$$

Using a partition of unity localizing to the two poles and the rest:

$$F(x) = I_+(r) + I_-(r) + I_{\text{rest}}(r),$$

- (a) **Non-stationary (equator/rest):** $O(r^{-N})$.
- (b) **North pole:** parametrize $\omega(u) = (u, \sqrt{1 - |u|^2})$, $u \in \mathbb{R}^{d-1}$, $|u| \leq \delta$.

Then

$$d\sigma(u) = \sqrt{\det(D_u \omega(u)^\top D_u \omega(u))} du, \quad D_u \omega(u)^\top D_u \omega(u) = \text{Id} + O(|u|^2),$$

$$\omega_d = 1 - \frac{1}{2}|u|^2 + O(|u|^4) = \sqrt{1 - |u|^2},$$

$$a(u) = \chi_+(u) g(\omega(u)) (1 + O(|u|^2)), \quad \phi(u) = 2\pi \omega_d(u),$$

so $\phi(0) = 2\pi$, $\nabla \phi(0) = 0$, $D^2 \phi(0) = -2\pi \text{Id}$ (a non-degenerate maximum). Refined stationary phase (Lemma 1 below) then yields the stated asymptotic. ■

19.5 A refined stationary-phase lemma

Lemma 19.5 (Refined stationary phase).

$$\int e^{i\lambda \phi(u)} a(u) du = e^{i\lambda \phi(0)} e^{i\frac{\pi}{4} \text{sgn}(D^2 \phi(0))} \left| \det\left(\frac{1}{2\pi} D^2 \phi(0)\right) \right|^{-\frac{1}{2}} (a(0) + O(\lambda^{-1})).$$

Applying it at the north pole (with large parameter r , $\phi(0) = 1$, $D^2 \phi(0) = -2\pi \text{Id}$, $\text{sgn} = -(d-1)$):

$$I_+(r) = e^{2\pi i r} C_d r^{-\frac{d-1}{2}} (g(0, \dots, 1) + O(r^{-1})), \quad C_d = e^{-i\frac{\pi}{4}(d-1)} (2\pi)^{-\frac{d-1}{2}}.$$