

a
Half-lives of the steps, against the dosing intervals
8 h
12 h
24 h daily
48 h
168 h weekly
half-lives per 24 h
miR-7
miR-7, upper bound
miR-29b
miR-29c
median miRNA
1.7 h
5.0 h
7.0 h
10.6 h
34.0 h
miRNA
14.1
4.8
3.4
2.3
0.7
SNCA mRNA
BACE1 mRNA
6.5 h
15.2 h
mRNA
3.7
1.6
α-synuclein
BACE1
brain proteome, median
24.0 h
18.6 h
195 h
Protein
1.0
1.3
0.1
Aβ42 clearance, controls
Aβ42 clearance, AD
9.1 h
13.1 h
Aβ42
2.6
1.8
1 h
10 h
100 h
1000 h
b
Washout
Level, multiple of baseline
within 10% of baseline
Time after a 10-fold bolus (h)
c
Cost of instability
daily
miR-7 (1.7 h)
miR-7, upper bound (5 h)
miR-29b (7 h)
miR-29c (10.6 h)
median miRNA (34 h)
Peak / average
Dosing interval (h)
d
Stabilisation for daily dosing
miR-7
miR-7 (5 h)
miR-29b
miR-29c
20.0×
6.8×
4.9×
3.2×
Fold increase in half-life