
a
Alzheimer's disease
Parkinson's disease
Primary: one estimate per study
Marker-type-neutral selection
Confidence intervals as published
Only studies with a reported interval
Leave one study out (each dot)
2000 random accuracy-blind selections
Every eligible estimate (ignores correlation)
Plus search-update study, RT-qPCR
Plus search-update study, ORCA-Cas
0.82 [0.77–0.85]
0.81 [0.76–0.84]
0.82 [0.77–0.85]
0.83 [0.73–0.89]
0.81–0.83
0.80 [0.79–0.82]
0.79 [0.77–0.81]
0.82 [0.78–0.85]
0.82 [0.78–0.86]
0.80 [0.74–0.85]
0.78 [0.73–0.83]
0.81 [0.75–0.85]
0.77 [0.66–0.85]
0.79–0.81
0.79 [0.77–0.81]
0.78 [0.75–0.80]
0.6 0.7 0.8 0.9 1.0
Pooled AUC
b
Parkinson's disease
Primary selection
Neutral selection
Alzheimer's disease
Primary selection
Neutral selection
single (n)
panel (n)
panel vs single
0.76 (12)
0.85 (7)
p < 0.0001
0.77 (16)
0.84 (3)
p = 0.009
0.81 (12)
0.83 (6)
p = 0.65
0.81 (14)
0.81 (4)
p = 0.43
0.5 0.6 0.7 0.8 0.9 1.0
Pooled AUC (95% CI); square = single microRNA, diamond = panel; n studies in brackets