
DEPOSITED STRUCTURES


renders of the deposited coordinates, as illustration: no result is read from them



Argonaute 2 with a miRNA guide

6N4O · X-ray, 2.9 Å
guide, seed (red) and target


AD PD



BACE1 with an inhibitor

4D8C · X-ray, 2.07 Å
cut open; catalytic Asp pair in red


AD



Aβ42 fibril

5OQV · cryo-EM, 4.0 Å
seen down the fibril axis

AD



α-synuclein fibril

6CU7 · cryo-EM, 3.5 Å (FSC 0.5)
seen down the fibril axis

PD

repression strength not measured
BACE1 about 50% in cells (K047)

ALZHEIMER

Mimic
bolus D
every T h

miR-29a, b, c
b 7 h, c 10.6 h
a: no decay in 12 h
K017–K019

BACE1 mRNA
half-life 15.2 h
5 isoforms pooled
K067, K068

BACE1 protein
half-life ≈19 h
BACE1:APP = 1:54
K025, K055, K056

Aβ42 monomers
production 6.6 %/h
clearance 5.3 %/h
in AD; K001–K004

Aβ42 nucleation
orders 2 and 2
M* 10 nM
rate constants: gap
K012–K014, K037

repression strength not measured
SNCA 30% and 43% in cells (K058, K063)

PARKINSON

Mimic
bolus D
every T h

miR-7
half-life 1.7 h
upper bound 5 h
Cyrano degrades it
K020–K023

SNCA mRNA
half-life 6.5 h
rat neurons
K066

α-synuclein
21.6 µM in a bouton
half-life ≥ 24 h
K026, K054

Elongation
saturating
m½ 46–50 µM
K009, K010

Nucleation
gated by pH:
none at neutral pH
K011, K042

dashed outline: no measured rate constants dashed bar: repression