



Review

The role of signalling and the cytoskeleton during Vaccinia Virus egress



Flavia Leite, Michael Way*

Cell Motility Laboratory, London Research Institute, Cancer Research UK, 44 Lincoln's Inn Fields, London, WC2A 3LY, UK

ARTICLE INFO

Article history:

Received 20 November 2014
Received in revised form 26 January 2015
Accepted 26 January 2015
Available online 11 February 2015

Keywords:

Vaccinia virus
Signalling
Actin
Microtubules
Transport
Viral spread

ABSTRACT

Viruses are obligate intracellular parasites that are critically dependent on their hosts to replicate and generate new progeny. To achieve this goal, viruses have evolved numerous elegant strategies to subvert and utilise the different cellular machineries and processes of their unwilling hosts. Moreover, they often accomplish this feat with a surprisingly limited number of proteins. Among the different systems of the cell, the cytoskeleton is often one of the first to be hijacked as it provides a convenient transport system for viruses to reach their site of replication with relative ease. At the latter stages of their replication cycle, the cytoskeleton also provides an efficient means for newly assembled viral progeny to reach the plasma membrane and leave the infected cell. In this review we discuss how Vaccinia virus takes advantage of the microtubule and actin cytoskeletons of its host to promote the spread of infection into neighboring cells. In particular, we highlight how analysis of actin-based motility of Vaccinia has provided unprecedented insights into how a phosphotyrosine-based signalling network is assembled and functions to stimulate Arp2/3 complex-dependent actin polymerization. We also suggest that the formin FHOD1 promotes actin-based motility of the virus by capping the fast growing ends of actin filaments rather than directly promoting filament assembly. We have come a long way since 1976, when electron micrographs of vaccinia-infected cells implicated the actin cytoskeleton in promoting viral spread. Nevertheless, there are still many unanswered questions concerning the role of signalling and the host cytoskeleton in promoting viral spread and pathogenesis.

© 2015 Elsevier B.V. All rights reserved.

Contents

1. Introduction	88
2. Microtubule-based transport of Vaccinia	88
2.1. IMV and IEV move on microtubules	88
2.2. The basis of kinesin-1 recruitment to IEV	88
3. Getting through the actin cortex	91
3.1. F11 enhances viral spread by inhibiting RhoA signalling	91
3.2. F11 is a PDZ scaffolding protein that binds Myosin-9A to inhibit RhoA	91
4. Vaccinia induced actin polymerization enhances the spread of infection	92
4.1. Src and Abl phosphorylation of A36 induces actin polymerization	92
4.2. Phosphorylated A36 recruits a signalling network	92
4.3. Molecular dissection of the A36 signalling network	94
4.4. Cdc42 enhances Vaccinia actin tail formation	94
4.5. The role of FHOD1 in actin tail assembly	94
4.6. Actin polymerization powers viral spread	96
4.7. Conclusions	96
Acknowledgements	96
References	96

* Corresponding author. Tel.: +44 0 207 269 3733; fax: +44 0 207 269 3581.
E-mail address: michael.way@cancer.org.uk (M. Way).

1. Introduction

The hijacking of the cytoskeleton is a common strategy employed by viruses infecting virtually all organisms including bacteria, plants and animals (Dodding and Way, 2011; Erb and Pogliano, 2013; Niehl et al., 2013; Taylor et al., 2011). However, one of the most striking examples of viral subversion of the host actin and microtubule cytoskeletons occurs during Vaccinia virus infection (Dodding and Way, 2011; Welch and Way, 2013). Vaccinia is a large double stranded DNA virus, which replicates exclusively in the cytoplasm of infected cells and is the most studied member of the Orthopoxviridae (Moss, 2007). Vaccinia is perhaps best known for its use as the vaccine to protect against smallpox, a deadly human disease caused by its close relative Variola virus (Jacobs et al., 2009; Walsh and Dolin, 2011). Smallpox was eradicated more than 30 years ago. Nevertheless, Vaccinia is increasingly being used as a vaccine vector for a wide range of different diseases as well as for oncolytic therapies (Jacobs et al., 2009; Kirn and Thorne, 2009; Thorne, 2011; Volz and Sutter, 2013; Walsh and Dolin, 2011). The vaccinia virus genome consists of ~200 kbp encoding for some 260 proteins, only about 80 of which, end up in infectious intracellular mature virus (IMV) particles (Chung et al., 2006; Resch et al., 2007; Yoder et al., 2006). This large coding capacity, which allows Vaccinia to infect and replicate in many different cell types, is in part due to its complex replication cycle, which involves the assembly of two morphologically distinct types of cytoplasmic virus particles (Fig. 1). The large genome also reflects the prodigious number of viral proteins Vaccinia uses to inhibit or suppress the antiviral activity of its host at all stages of its replication cycle (Haller et al., 2014). This includes inhibiting apoptosis of infected cells before new viral progeny are assembled and minimizing detection by the host immune system (Bahar et al., 2011; Mohamed and McFadden, 2009; Postigo and Ferrer, 2009).

After binding to the cell membrane, virus entry occurs either by direct fusion with the plasma membrane (Carter et al., 2005; Law et al., 2006) or by low-pH endosomal entry pathway (Huang et al., 2008; Townsley et al., 2006). Moreover, Vaccinia actually promotes its uptake by stimulating actin-dependent macropinocytosis (Mercer and Helenius, 2008; Mercer et al., 2010; Schmidt et al., 2011). Having gained access to the cell cytoplasm, expression of early proteins is initiated allowing viral cores to uncoat and release their DNA (Kilcher et al., 2014; Mercer et al., 2012; Schmidt et al., 2013). This early protein expression is required for viral DNA replication, which occurs in viral factories located in a perinuclear region near the microtubule-organizing centre of the infected cell (Ploubidou et al., 2000; Roberts and Smith, 2008). Only after DNA replication does intermediate and late gene expression start, resulting in the assembly of intracellular mature virus (IMV) particles. IMV which represent the majority of viral progeny are infectious but are only released when the infected cell undergoes lysis (Roberts and Smith, 2008). Alternatively, some IMV can become intracellular enveloped virus (IEV) by being 'wrapped' by membrane cisternae derived from trans-Golgi or endosomal compartments containing a subset of viral proteins (Roberts and Smith, 2008; Smith et al., 2002). The molecular basis for this envelopment remains to be established, but it involves multiple integral viral membrane proteins as well as the Vaccinia E2 and F12 proteins (Dodding et al., 2009; Domi et al., 2008; Roper et al., 1998; Röttger et al., 1999; Sanderson et al., 1998; Smith et al., 2002; Wolffe et al., 1997; Wolffe et al., 1998). Once formed, IEV are transported to the cell periphery on microtubules by kinesin-1 before fusing with the plasma membrane (Fig. 1). Fusion of IEV with the plasma membrane results in two outcomes that have a different impact on the subsequent spread of infection. The extracellular enveloped virus (EEV), which are infectious and released from the cell, promote the long-range spread of vaccinia. Alternatively, after their

fusion with the plasma membrane, some virions remain attached to the outside of the cell and are known as the cell-associated enveloped virus (CEV). It is the CEV that are responsible for the local actin-dependent cell-to-cell spread of vaccinia (Fig. 1). In this review, we will discuss our current understanding of how vaccinia uses and manipulates the cytoskeleton of the cell to enhance its spread.

2. Microtubule-based transport of Vaccinia

2.1. IMV and IEV move on microtubules

Microtubule-based transport is the primary way in which cargoes are moved over micron distances in a directed fashion throughout the cell (Franker and Hoogenraad, 2013; Fu and Holzbaur, 2014; Stephens, 2012). It is perhaps not surprising then that viruses have developed numerous strategies to take advantage of this cellular transport system at all stages of their infection cycles (Dodding and Way, 2011; Greber and Way, 2006; Radtke et al., 2006). Moreover, the ability of large viruses, such as Vaccinia, to hijack this rapid and efficient transport system is essential, as their size precludes their movement by diffusion (Greber and Way, 2006; Sodeik, 2000). For the virologist, examining how viruses use the microtubule cytoskeleton and its associated motors is necessary to understand how the infection cycle is established, as well as the mechanisms underlying viral replication, assembly and spread. For the cell biologist, the same analysis promises to uncover fundamental insights into the molecular basis of microtubule motor recruitment and regulation.

Immunofluorescence analysis of vaccinia-infected cells reveals that intracellular mature virions (IMV) are capable of dispersing from their peri-nuclear site of assembly throughout the cell at the latter stages of the viral replication cycle (Fig. 2). This movement to the cell periphery is more apparent in viral backgrounds that result in an absence of IEV formation (Fig. 2). The extent of viral dispersal in fixed samples provided the first suggestion that IMV are capable of moving on microtubules, as it has been calculated that they would only diffuse 10 μm in ~5 h (Sodeik, 2000). Subsequent live-cell imaging of infected cells demonstrated that YFP-tagged IMV are capable of undergoing rapid, linear movements at speeds approaching 3 $\mu\text{m}/\text{s}$ (Ward, 2005). IMV movement is saltatory in nature and was abolished when cells were treated with the microtubule depolymerizing agent nocodazole. Direct imaging of IMV moving on microtubules still remains to be performed and the identity of the motor responsible for transporting IMV has yet to be established. In contrast, GFP-tagged IEV have been imaged rapidly moving in a microtubule dependent fashion as well as along microtubules towards the cell periphery prior to their fusion with the plasma membrane (Dodding et al., 2009; Geada et al., 2001; Herrero-Martinez et al., 2005; Hollinshead et al., 2001; Rietdorf et al., 2001; Ward and Moss, 2001a; Ward and Moss, 2001b). IEV movement is dependent on kinesin-1 (also known as Kif5B or conventional kinesin), which uses the power of ATP hydrolysis to transport cargoes towards the plus end of microtubules usually located at the cell periphery (Dodding and Way, 2011; Rietdorf et al., 2001).

2.2. The basis of kinesin-1 recruitment to IEV

Kinesin-1 is recruited to IEV through an interaction between the tetratricopeptide repeats (TPR) of its light chain (KLC) and A36, an integral IEV membrane protein that has a cytoplasmic domain of ~195 residues exposed on the surface of the virus (Röttger et al., 1999; van Eijl et al., 2000; Ward and Moss, 2004). Fluorescence resonance energy transfer (FRET) experiments have confirmed that




Fig. 1. Schematic representation of viral egress. Once assembled the intracellular mature virus (IMV) can be moved to the cell periphery on microtubules nucleated from the microtubule organizing center (MTOC). The motor responsible for this movement remains to be identified. Alternatively, a proportion of IMV can become enveloped by membrane cisternae derived from trans-Golgi containing a subset of viral proteins. This results in the formation of the intracellular enveloped virus (IEV), which are then transported to the cell periphery on microtubules by kinesin-1. On reaching the cell boundary, IEV traverse the cortical actin network before fusing with the plasma membrane. This fusion results in the formation of the cell associated enveloped virus (CEV). CEV, which remain attached to the cell transiently recruit AP-2 and clathrin prior to inducing actin polymerization and the assembly of an actin tail. Alternatively, CEV can be directly released from the cell to form the infectious extracellular enveloped virus (EEV).

A36 and KLC can interact with each other in infected cells (Jeshtadi et al., 2010). Deletion of the gene encoding A36 from the vaccinia genome has no impact on IEV formation (Röttger et al., 1999; Sanderson et al., 1998; van Eijl et al., 2000), but it does impair viral spread from their perinuclear site of assembly to the plasma membrane (Rietdorf et al., 2001; Ward and Moss, 2001a; Ward et al.,

2003). Consequently, an absence of A36 during infection reduces the cell-to-cell spread of vaccinia, as the virus is less likely to reach and fuse with the plasma membrane (Ward and Moss, 2001a). Deletion of the gene encoding ECTV-Mos-42, an A36 orthologue in Ectromelia virus, the causative agent of mousepox, also results in reduced cell-to-cell spread (Lynn et al., 2012).




Fig. 2. Immunofluorescence images of HeLa cells infected for 8 h with WR or the recombinant Δ B5R virus, which does not form IEV as it lacks the viral protein B5. All forms of the virus are detected with an antibody against A27 (left). CEV are detected with an antibody against B5 (middle). The actin cytoskeleton is labelled with phalloidin, which only detects actin filaments (right). In WR infected cells, CEV induce the formation of comet shaped actin tails. In contrast, in the Δ B5R-infected cells there are only prominent actin stress fibres. Nevertheless, dispersal of IMV towards the cell periphery still occurs.

Using a combination of yeast two hybrid and *in vitro* GST pull-down assays, it was demonstrated that residues 81–111 of A36 were sufficient to bind the KLC tetratricopeptide repeats (Ward and Moss, 2004). Examination of this region of A36 reveals it contains a short WD motif present in the KLC binding site of Calsyntenin-1 (Araki et al., 2007; Dodding and Way, 2009; Konecna et al., 2006; Morgan et al., 2010). An additional KLC binding WE motif first identified in Caytaxin and γ -BAR (also known as Gadkin) (Aoyama et al., 2009; Schmidt et al., 2009) also occurs at residues 64–65 of A36 (Dodding et al., 2011; Morgan et al., 2010). Mutation of the WE motif in A36 deficient in its ability to stimulate actin polymerization (see Section 4.1), results in a small but significant decrease in IEV spread to the cell periphery at 8 h post infection (Dodding et al., 2011). In contrast, loss of the WD motif leads to a dramatic albeit not total reduction in viral spread at 8 h post infection. Live cell imaging reveals that this reduction is due to a decrease in the IEV run length. This represents the average distance travelled along the microtubule between pauses, from 6.44 ± 0.37 to 2.58 ± 0.14 μ m rather than a change in the velocity of the virus (Dodding et al., 2011). Immunofluorescence analysis suggests this change is due to a decrease in the ability of the virus to recruit and/or retain kinesin-1 (Dodding et al., 2011). Nevertheless, by 11 h post infection, significant numbers of IEV are able to reach the cell periphery (See Fig. 3B Dodding et al., 2011). A similar result was observed by Morgan et al. (2010), who found that over expression of an A36 WD motif alanine mutant in cells infected with the Δ A36R virus, which does not encode A36 allowed IEV to reach the plasma membrane and induce actin tails at 12 h post infection (Morgan et al., 2010). In contrast, mutation of both the WD and WE motifs results in an inability of IEV to recruit kinesin-1 and an absence of viral spread, as seen with the Δ A36R virus (Dodding et al., 2011).

In addition to A36, it has been proposed that the viral protein F12 is also essential for microtubule-mediated transport of IEV (Morgan et al., 2010; van Eijl et al., 2002). F12 associates with IEV when they are moving on microtubules and its loss results in peri-nuclear accumulation of the virus (Dodding and Way, 2009; Morgan et al., 2010; van Eijl et al., 2002; Zhang et al., 2000). Our ultrastructural analyses of Δ F12L infected cells, however, suggests that the loss of IEV spread in the absence of F12 is because the protein is required for their formation (Dodding and Way, 2009). Moreover, live cell imaging reveals that the small numbers of IEV that do assemble can still undergo linear movements at ~ 1 μ m/s (Dodding and Way, 2009). While we cannot formally rule out the participation of F12 in microtubule-based transport of IEV, live cell imaging clearly demonstrates F12 is not essential for this process. Furthermore, recent analysis has revealed that F12 has a structural fold of an inactive DNA polymerase (Yutin et al., 2014), rather than structural similarity to the kinesin light chain as predicted by Morgan et al. (Morgan et al., 2010). Nevertheless, F12 does contain a potential kinesin-1 WD binding motif (Morgan et al., 2010). In contrast to A36, it still remains to be demonstrated if this motif directly binds to the TPR of the kinesin-1 light chain. However, over expression of an F12 WD motif mutant in cells infected with the Δ F12L virus does not rescue kinesin-1 recruitment and IEV egress (Morgan et al., 2010). It is still not clear if this phenotype is due to a defect in IEV assembly rather than microtubule transport *per se*. It is also possible that the mutation impacts on the structure of F12 and/or its interaction with E2 and/or A36, although the function of these interactions remains to be elucidated (Dodding and Way, 2009; Johnston and Ward, 2008). Notwithstanding the above, the current overexpression study clearly indicates that the WD motif in F12 is important for the function of the protein during the virus




Fig. 3. Schematic representation of the mechanism by which F11 down regulates RhoA signalling. An intramolecular interaction between the central PDZ-like domain and the C-terminal PDZ binding motif (PBM) prevents F11 from interacting with RhoA. In its “open” conformation the PDZ domain interacts with the RhoGAP Myosin-9A, while GTP bound RhoA binds towards the C-terminus of the protein. The GAP activity of Myosin-9A enhances the hydrolysis of GTP to GDP, resulting in the release of RhoA and inhibition of its signalling.

replication cycle. The generation and detailed analysis of a recombinant virus expressing the F12 WD mutant will hopefully help clarify the exact role played by this potential kinesin-1 binding site during IEV assembly and transport.

3. Getting through the actin cortex

3.1. F11 enhances viral spread by inhibiting RhoA signalling

When IEV reach the cell periphery, they cannot fuse with the plasma membrane until they traverse the cortical actin cytoskeleton, a dense mesh of actin filaments associated with the cytoplasmic face of the plasma membrane (Biro et al., 2013; Charras et al., 2006; Clark et al., 2013; Fritzsche et al., 2013). The actin cortex provides a cell with mechanical resilience (Clark et al., 2013; Fritzsche et al., 2013; Gauthier et al., 2012; Salbreux et al., 2012; Tinevez et al., 2009) and plays an important role in regulating the morphology of the cell during a variety of cellular processes such as migration, mitotic rounding and cytokinesis (Bovellan et al., 2014; Matthews et al., 2012; Sedzinski et al., 2011). However, from a viral perspective, the cortical actin cytoskeleton represents a significant physical barrier, not only to IEV egress, but also to many other viruses during both their entry and exit from the cell (Radtke et al., 2006; Van den Broeke et al., 2014). Fortunately for Vaccinia, it encodes F11, which inhibits the activity of the small GTPase RhoA (Arakawa et al., 2007b; Cordeiro et al., 2009; Valderrama et al., 2006). F11 was originally identified as a viral protein required for vaccinia-induced cell migration (Morales et al., 2008; Valderrama

et al., 2006). However, its ability to inhibit RhoA signalling also plays an important role in promoting viral release and spread in cell culture as well as *in vivo* in an intranasal mouse model of infection (Arakawa et al., 2007a; Cordeiro et al., 2009). F11 mediated inhibition of RhoA signalling increases the dynamics and targetting of microtubules to the cell periphery, which enhances the ability of IEV to reach the plasma membrane (Arakawa et al., 2007b). F11 also makes it easier for the virus to get through the cortical actin network by preventing RhoA from stimulating mDia induced actin polymerization (Arakawa et al., 2007a). Lastly, by stimulating infected cell migration and the formation of long neurite-like projections, F11 enhances the spread of infection by increasing viral contact with more non-infected cells (Cordeiro et al., 2009; Valderrama et al., 2006). The ability of F11 to promote viral spread has also been elegantly illustrated using Myxoma virus, which lacks a related homologue (Irwin and Evans, 2012). Irwin et al. found that a recombinant Myxoma virus expressing F11 is more effective in its cell-to-cell spread, as plaques expand ~6 fold faster and are four times larger than controls after 4 days (Irwin and Evans, 2012). The expression of F11 also improves the oncolytic efficacy of Myxoma by increasing viral spread in tumours and prolonging the survival of mice (Irwin et al., 2013). The conservation of F11 homologues in chordopoxviruses (www.poxvirus.org) suggests that inhibiting RhoA signalling represents a common strategy used by poxviruses to enhance their cell-to-cell spread (Handa et al., 2013).

3.2. F11 is a PDZ scaffolding protein that binds Myosin-9A to inhibit RhoA

F11 binds directly to RhoA using a short sequence motif that is also present in its downstream effector ROCK (Valderrama et al., 2006). Consistent with this, F11 only interacts with active GTP-bound RhoA (Cordeiro et al., 2009). This allows F11 to sequester active RhoA to inhibit its signalling, but does not explain how the protein reduces the steady state level of GTP-bound RhoA during vaccinia infection (Arakawa et al., 2007a; Cordeiro et al., 2009). More recently, the mechanism by which F11 down regulates RhoA signalling has been uncovered (Figure 3). We found F11 contains a central PDZ-like domain that is required to down regulate RhoA signalling and enhance viral spread (Handa et al., 2013). PDZ domains are 80–90 residues in length with ~ 30% sequence identity and are conserved from bacteria to man, with the human genome encoding ~ 270 different PDZ domains (Harris and Lim, 2001; Ivarsson, 2012; Luck et al., 2012; Nourry et al., 2003; Ponting, 1997; Subbaiah et al., 2011). PDZ domains most frequently bind short peptide motifs, known as PDZ binding motifs (PBMs) located at the very C-terminus of proteins. However, they can also interact with internal peptide motifs and lipids as well as hetero- and/or homo dimerize (Harris and Lim, 2001; Ivarsson, 2012; Luck et al., 2012; Nourry et al., 2003; Subbaiah et al., 2011). PDZ containing proteins participate in a wide range of fundamental cellular processes including adhesion, polarity, proliferation, apoptosis and signalling. It is not surprising, given their role in so many cellular processes, that viruses frequently target their function using viral proteins containing PDZ binding motifs (Davey et al., 2011; Javier and Rice, 2011). Curiously though, there is no evidence that viruses manipulate their hosts using virally encoded PDZ domain containing proteins (Ivarsson, 2012; Subbaiah et al., 2011). F11 therefore represents the first example of a viral protein containing a functional PDZ-like domain.

In addition to its PDZ domain, F11 also has a class II PDZ binding motif (PBM) at its extreme C-terminus (Handa et al., 2013). The interaction of this PBM with the central PDZ domain inhibits the ability of F11 to bind RhoA, presumably by inducing a conformation that occludes the RhoA binding site (Figure 3). The C-terminal PBM also has additional roles that go beyond regulating the interaction

of F11 with RhoA, as its deletion reduces virus release to a level that is equivalent to the loss of the complete protein (Handa et al., 2013). This phenotype may be related to a disruption of the localization of F11 within the infected cell, as presence of the C-terminal PBM enhances its association with membranes.

The PDZ domain of F11 is required to reduce the level of active RhoA during infection (Handa et al., 2013). Given this, we wondered if the F11 PDZ domain down regulates RhoA by interacting with a class II PBM containing GTPase activating protein (GAP). GAPs, which have an over representation of PDZ binding motifs compared to other protein families, inactivate Rho GTPases by stimulating the hydrolysis of GTP to GDP (Garcia-Mata and Burridge, 2007; Giallourakis et al., 2006). Pulldown assays demonstrate that the PDZ domain of F11 interacts with the PBM of the GAPs β -Chimaerin and Myosin-9A, which regulate Rac and RhoA respectively (Chieragatti et al., 1998; Handa et al., 2013; Yang and Kazanietz, 2007). Consistent with a possible role in promoting viral spread, Myosin-9A regulates cortical actin in a RhoA-dependent fashion during cell-cell adhesion (Omelchenko and Hall, 2012). Subsequent analysis demonstrated that RNAi mediated loss of Myosin-9A during infection leads to reduced viral release and spread, as RhoA signalling is no longer inhibited at 8 h post infection. The ability of Myosin-9A to enhance viral spread is dependent on its GAP activity as well as its ability to bind the F11 PDZ domain using its C-terminal PBM. Interestingly, it also depends on the ability of F11 to bind RhoA. Furthermore, pulldown assays demonstrate the presence of a tri-molecular complex between F11, RhoA and Myosin-9A during infection. This suggests that F11 is acting as a PDZ scaffolding protein to bring Myosin-9A and RhoA together to inactivate the GTPase (Fig. 3).

4. Vaccinia induced actin polymerization enhances the spread of infection

4.1. Src and Abl phosphorylation of A36 induces actin polymerization

Nearly 40 years ago, analysis of vaccinia infected cells in the electron microscope revealed the presence of virus particles on the tip of large microvilli projecting from the plasma membrane (Stokes, 1976). These projections appeared late in infection and contained actin, α -actinin, fimbrin and filamin but not tropomyosin or myosin (Hiller et al., 1981; Hiller et al., 1979; Krempien et al., 1981). These studies were essentially forgotten until 1995, when it was demonstrated that vaccinia is moved by the power of actin polymerization on the tips of actin tails before extending out into neighbouring non-infected cells (Cudmore et al., 1995; Cudmore et al., 1996). Subsequent studies demonstrated that actin polymerization and tail formation is induced by CEV, which are formed when IEV fuse with the plasma membrane after undergoing kinesin-1 driven microtubule transport (Cudmore et al., 1996; Hollinshead et al., 2001; Humphries et al., 2012; Rietdorf et al., 2001; Ward and Moss, 2001a; Ward and Moss, 2001b) (Fig. 1).

Vaccinia induced actin polymerization is dependent on Src and Abl family kinase mediated phosphorylation of the IEV protein A36 (Frischknecht et al., 1999; Newsome et al., 2004; Newsome et al., 2006; Reeves et al., 2005) (Fig. 4). Deletion of the A36R gene and loss of A36 expression does not affect IEV assembly but does lead to an absence of actin tail formation and a small plaque phenotype, indicative of a defect in viral spread (Röttger et al., 1999; Sanderson et al., 1998; Wolffe et al., 1998). A36 is highly conserved in orthopoxvirus genomes, suggesting that virus-induced actin polymerization at the plasma membrane is widely used by mammalian poxviruses to enhance their cell-to-cell spread. Indeed, monkeypox and variola viruses induce Src and Abl dependent actin

tails, while the A36 homologue of ectromelia virus enhances viral spread by inducing actin tail formation (Lynn et al., 2012; Reeves et al., 2011). Moreover, the ability to stimulate actin polymerization is not just restricted to orthopoxviruses, as other divergent vertebrate poxviruses (*Chordopoxviridae*) can also induce actin tails (Dodding and Way, 2009; Duteyrat et al., 2006; Law et al., 2004). In these cases, a divergent family of viral proteins related to YL126 a transmembrane protein of Yaba-Like Disease virus (yatapoxvirus) is used to stimulate actin polymerization (Dodding and Way, 2009). These YL126 orthologues bear little or no sequence homology to A36, but they stimulate actin polymerization in a tyrosine phosphorylation dependent manner via the same Grb2, Nck, WIP and N-WASP signalling network as Vaccinia virus (Dodding and Way, 2009) (see Section 4.2).

Vaccinia A36 is a type Ib transmembrane protein with a cytoplasmic domain of ~ 195 residues on the IEV surface (Röttger et al., 1999; van Eijl et al., 2000). When IEV fuse with the plasma membrane, A36 becomes localized beneath CEV but retains the same topology with respect to the cytoplasm of the cell (Smith et al., 2002; van Eijl et al., 2000) (Fig. 4). Src recruitment and phosphorylation of A36 only occurs after viral fusion with the plasma membrane, even though its cytoplasmic domain is fully exposed on the IEV surface (Newsome et al., 2004). This implied that one or more of the four integral viral membrane proteins (A33, A34, A56 and B5) exposed on the surface of CEV (Smith et al., 2002), are stimulating an outside-in signal to locally activate Src kinase. We still do not have a molecular understanding of how CEV signal back into the cell, but the SCR4 domain of B5 is essential to activate Src, phosphorylate A36 and induce actin tails (Newsome et al., 2004).

4.2. Phosphorylated A36 recruits a signalling network

The ubiquitous Src and Abl family kinase members (Src, Fyn, Yes, Abl and Arg) are recruited by CEV and capable of stimulating actin tail formation by virtue of their ability to phosphorylate tyrosine 112 of A36 (Frischknecht et al., 1999; Newsome et al., 2004; Newsome et al., 2006; Reeves et al., 2005) (Fig. 4). Additional tyrosine kinases might also be capable of phosphorylating A36, however, inhibition of Src and Abl family kinases or mutation of Y112 of A36 is sufficient to prevent vaccinia actin tail formation (Frischknecht et al., 1999; Newsome et al., 2004; Reeves et al., 2005; Scaplehorn et al., 2002; Ward and Moss, 2001a). Consistent with this, Abl, Arg, Fyn, Src and Yes can all phosphorylate tyrosine 112 of A36 *in vitro* (Newsome et al., 2006). When phosphorylated, tyrosine 112 of A36 recruits the adapter Nck to the virus by interacting directly with its SH2 domain (Frischknecht et al., 1999; Scaplehorn et al., 2002). Nck is essential for Vaccinia induced-actin polymerization as no actin tails are formed in mouse embryo fibroblasts (MEFs) lacking Nck (Weisswange et al., 2009). The presence of Nck beneath the CEV results in the recruitment of WIP and N-WASP, the latter of which activates the Arp2/3 complex to stimulate actin polymerization (Donnelly et al., 2013; Frischknecht et al., 1999; Moreau et al., 2000; Snapper et al., 2001; Weisswange et al., 2009; Zettl and Way, 2002) (Fig. 5). Nck is not the only SH2/SH3 adapter that is recruited by the virus (Scaplehorn et al., 2002). When phosphorylated by Src, Fyn, Abl and Arg but not Yes, tyrosine 132 of A36 is able to recruit Grb2 to the virus by binding its SH2 domain (Newsome et al., 2006; Scaplehorn et al., 2002) (Fig. 4 and 5). The recruitment of Grb2 is dependent on both the phosphorylation of tyrosine 132 of A36 and the presence of the proline rich region of N-WASP, which presumably interacts with the SH3 domains of Grb2 (Scaplehorn et al., 2002; Weisswange et al., 2009). In contrast to tyrosine 112, phosphorylation of tyrosine 132 is not required for vaccinia-induced actin polymerization, although its presence does enhance actin tail formation (Scaplehorn et al., 2002; Ward and Moss, 2001a; Weisswange et al., 2009). Analysis of the dynamics




Fig. 4. Schematic of the steps involved in assembling the signalling network responsible for stimulating Arp2/3 complex dependent actin polymerization beneath CEV.

of GFP-tagged Nck, WIP and N-WASP during vaccinia actin-based motility reveals that Grb2 helps stabilize the signalling complex, which may in part account for the enhanced actin tail formation (Weisswange et al., 2009).

4.3. Molecular dissection of the A36 signalling network

The robust recruitment of Nck, Grb2, WIP and N-WASP to the virus, coupled with the ability to quantify both the number and speed of actin tails as well as the dynamics of proteins in the system make vaccinia an excellent model to understand how a phosphotyrosine-based signalling network activates Arp2/3-dependent actin polymerization (Donnelly et al., 2013; Weisswange et al., 2009; Welch and Way, 2013) (Fig. 5). Another strength of the system is the ability of vaccinia to infect mouse embryo fibroblasts (MEFs) derived from knockout mice lacking individual proteins such as Nck, WIP or N-WASP (Donnelly et al., 2013; Snapper et al., 2001; Weisswange et al., 2009). An initial over expression study in HeLa cells suggested that WIP and N-WASP are recruited to the virus as a complex (Moreau et al., 2000). Subsequent analysis using MEFs derived from mice lacking N-WASP confirmed that its presence is required for the recruitment of WIP to the virus (Donnelly et al., 2013; Snapper et al., 2001; Weisswange et al., 2009). The presence of WIP and/or its homologue WIRE is also required for the recruitment of N-WASP (Donnelly et al., 2013). Collectively, these data demonstrate that vaccinia recruits WIP and N-WASP as a complex (Fig. 4). In contrast, Nck is still localized on the virus in the absence of WIP or N-WASP, indicating it is upstream of the WIP:N-WASP complex (Donnelly et al., 2013; Snapper et al., 2001; Weisswange et al., 2009). We envisage that Nck engages with tyrosine 112 of A36 when it is phosphorylated and then recruits a WIP:N-WASP complex to the virus, which subsequently associates with Grb2 interacting with phosphorylated tyrosine 132 of A36. This explains why Grb2 is not recruited to the virus in the absence of N-WASP and why Nck is essential for vaccinia actin tail formation (Weisswange et al., 2009).

A combination of biochemical approaches and expression of mutants in cells lacking endogenous Nck, WIP or N-WASP, demonstrate that an interaction of the second SH3 domain of Nck with WIP is essential for actin tail formation, as it is required to recruit the WIP:N-WASP complex (Donnelly et al., 2013). Consistent with this, the recruitment of N-WASP to the virus depends on its interaction with WIP rather than Nck. The first and third SH3 domains of Nck are essential to stimulate actin tails even though they are not required to recruit the WIP:N-WASP complex. Curiously, however, an interaction between N-WASP and Nck is not required for virus induced actin polymerization even in the absence of Grb2 (Donnelly et al., 2013). Taken together, these data suggested that an additional factor is also able to stimulate the ability of N-WASP to activate the Arp2/3 complex and induce actin polymerization beneath CEV. This additional component may also explain why the turnover dynamics of N-WASP is ~3.4 times slower than that of Nck and WIP (half life of turnover is 2.7 rather than 0.8 s), even though they are responsible for its recruitment to the virus (Weisswange et al., 2009).

4.4. Cdc42 enhances Vaccinia actin tail formation

Our previous observations demonstrated that Cdc42, a well-known activator of N-WASP, was recruited to vaccinia inducing actin tails (Moreau et al., 2000). However, expression of dominant negative Cdc42 appeared to have no impact on actin tail formation, based on assessing for the presence of a single actin tail or not in infected cells (Moreau et al., 2000). Given this stringent quantification criterion and in light of the results of Donnelly et al. (2013), we re-examined the potential role of Cdc42 in vaccinia actin tail

formation. We found there was a significant decrease in the number of actin tails in cells either lacking Cdc42 or expressing dominant negative Cdc42 or when N-WASP was unable to interact with Cdc42 (Humphries et al., 2014). Moreover, the inability of N-WASP to bind Cdc42 has a greater impact on actin tail formation than the loss of Nck binding, suggesting Cdc42 is more important than Nck in stabilizing and/or activating N-WASP. Consistent with this, a loss of Cdc42 binding results in a significantly faster turnover of N-WASP than that observed when its interaction with Nck is abolished (half life of turnover is 1.18 rather than 2.41 s) (Donnelly et al., 2013; Humphries et al., 2014). However, Nck still contributes to vaccinia induced actin polymerization, as far fewer actin tails are formed when N-WASP cannot bind both proteins compared to Cdc42 alone. FRAP analysis also reveals that the turnover of N-WASP is essentially the same as Nck and WIP when it cannot interact with Nck and Cdc42 (Humphries et al., 2014). Moreover, actin tail formation is largely abolished in the absence of Grb2 recruitment, indicating that Arp2/3-dependent actin based motility of vaccinia is principally driven by Cdc42 and Nck-dependent activation of N-WASP (Humphries et al., 2014).

In contrast to Nck, Cdc42 is not recruited to the virus in the absence of N-WASP (Humphries et al., 2014). This suggested that activation and recruitment of Cdc42 is downstream of the presence of the WIP:N-WASP complex (Figure 4). Consistent with this, live cell imaging reveals that the accumulation of Cdc42 on the virus is delayed by ~5 s compared to N-WASP, which begins to be recruited ~10 s before actin polymerization is initiated (Humphries et al., 2014). This delay presumably reflects the activation of Cdc42 by a RhoGEF beneath the virus, as N-WASP can only interact with GTP-bound Cdc42 (Miki et al., 1998). The identification of the RhoGEF responsible for activating Cdc42 was greatly facilitated by the observation that immediately following fusion with the plasma membrane, vaccinia transiently recruits AP-2 and clathrin prior to actin tail formation (Humphries et al., 2012). Clathrin appears to have an organizational role, as its recruitment enhances polarization of A36 and N-WASP beneath the CEV making it easier for the virus to induce and sustain actin polymerization (Humphries et al., 2012; Humphries and Way, 2013). We found that the virus activates Cdc42 by recruiting intersectin-1, a RhoGEF that interacts with both AP-2 and N-WASP (Humphries et al., 2014; Hunter et al., 2013; Hussain et al., 2001). Furthermore, the recruitment of intersectin-1 to CEV is upstream of actin tail formation, as it occurs in the absence of N-WASP. Loss of intersectin-1 or its GEF activity results in a similar reduction in actin tail number as expressing dominant negative Cdc42 (Humphries et al., 2014). Moreover, this decrease is not additive, suggesting that intersectin-1 and Cdc42 are acting in the same pathway to enhance vaccinia actin tail formation. In contrast to Nck, however, the recruitment of Cdc42 is not essential for vaccinia actin tail formation. Nevertheless, Cdc42 does collaborate with Nck to enhance the ability of the virus to induce actin polymerization by stabilizing and/or activating N-WASP (Humphries et al., 2014). We now have to understand how CEV recruit intersectin-1 and whether it is essential for AP-2 and clathrin recruitment prior to actin tail formation.

4.5. The role of FHOD1 in actin tail assembly

Cdc42 is not the only Rho GTPase that has been shown to enhance actin tail formation and viral spread. By performing a siRNA screen against actin cytoskeletal regulators Alvarez and Agaisse (2013) demonstrated that the formin, FHOD1 promotes the cell-to-cell spread of vaccinia. Loss of FHOD1 results in ~2.5 fold reduction in the number of tails induced by CEV as well as a 30% decrease in their velocity, suggesting the protein enhances formation and elongation of actin tails (Alvarez and Agaisse, 2013). The ability of FHOD1 to enhance actin tail formation is dependent on




Fig. 5. Immunofluorescence image showing the recruitment of GFP-tagged Nck, Grb2, WIP and N-WASP (green) to the tip of actin tails (red) in a HeLa cell infected for 8 h with WR. The scale bar is 5 μ m.

its FH1, FH2 and GBD domains, which interact with profilin:actin complexes, the fast growing end of actin filaments and the RhoGTPase Rac respectively (Schonichen et al., 2006; Schulte et al., 2008). Consistent with a role for the FH1 and GBD domains, RNAi mediated knockdown of profilin or Rac phenocopied the impact of FHOD1 depletion on actin tail number and speed (Alvarez and Agaisse, 2013). The authors also found that active Rac was enriched in the membrane surrounding projecting actin tails in an A36-independent fashion. Furthermore, FHOD1 recruitment to vaccinia actin tails was dependent on Rac as well as N-WASP (Alvarez and Agaisse, 2013). In contrast to Nck and N-WASP, but in line with intersectin-1 and Cdc42, Rac and FHOD1 enhance but are not essential for vaccinia actin tail formation.

Based on their observations, Alvarez and Agaisse suggest that FHOD1 enhances actin tail formation and speed by recruiting profilin:actin complexes and enhancing the nucleation and elongation of the barbed end of actin filaments. They propose that FHOD1 might enhance the growth of actin filaments nucleated by the Arp2/3 complex or provide the initial actin filaments from which Arp2/3 induces new filament assembly. We favour a different mechanism. Ground breaking *in vitro* reconstitution experiments with pure proteins from the Carlier laboratory have demonstrated that Arp2/3 driven actin based motility is critically dependent on the presence of an actin filament severing protein such as ADF/cofilin as well as a barbed end capping protein (Loisel et al., 1999). In contrast, the presence of profilin in the *in vitro* motility

assays is not essential, although it does enhance the rate of movement. More recently, *in vitro* assays have shown that FHOD1 caps the fast growing or barbed end of actin filaments in an FH2 dependent fashion but does not nucleate or promote filament elongation (Schonichen et al., 2013). Results from *in vitro* biochemical assays may not always reflect the situation in more complex *in vivo* setting. However, we think that the observations of Alvarez and Agaisse are consistent with the hypothesis that FHOD1 is enhancing vaccinia actin tail formation by capping actin filaments and providing profilin:actin complexes for filament growth induced by the Arp2/3 complex. First, FHOD1 is found throughout the actin tail and not immediately beneath the CEV as seen for N-WASP. Such a localization is consistent with FHOD1 binding the barbed ends of actin filaments within the tails. Second, the deletion of the FH2 domain, which is responsible for its actin filament capping activity, results in an absence of FHOD1 recruitment to actin tails. Third, fluorescent actin incorporation experiments have demonstrated that actin polymerization only occurs immediately beneath the virus and that the filaments in the tail do not grow as they are capped (Cudmore et al., 1996). Nevertheless, our analysis using both antibodies and GFP-tagged clones has failed to detect capping protein (also known as CapZ) or Eps8 in vaccinia induced actin tails (Way lab unpublished). Lastly, the presence of FHOD1 in actin tails is dependent on N-WASP. However, this does not demonstrate that the interaction is direct or rule out that the recruitment of FHOD1 is actually dependent on the Arp2/3 complex or the actin filaments

it nucleates. Future experiments are clearly required to determine the exact function of FHOD1 in vaccinia actin tail formation and uncover how the virus activates Rac.

4.6. Actin polymerization powers viral spread

We now know the main players required for CEV-induced actin tail formation. However, we are still a long way from having a complete molecular understanding of exactly how the actin cytoskeleton promotes the release and spread of Vaccinia. More than 30 years ago it was demonstrated that inhibition of actin polymerization with cytochalasin D inhibits release of vaccinia from infected cells (Payne and Kristensson, 1982). More recently, it was found that A36 phosphorylation and actin polymerization promotes virus release by driving CEV out of invaginations in the plasma membrane (Horsington et al., 2013). The activity of Abl and Arg but not Src family kinases also promotes the release of CEV from infected cells (Reeves et al., 2005). Consistent with this, treatment of mice with the Abl-family kinase inhibitor Gleevec/STI-571/Imatinib reduces viral spread and promotes their survival from an otherwise lethal infection (Reeves et al., 2005; Reeves et al., 2011). The activity of Abl and Arg clearly enhances A36-dependent release of Vaccinia from infected cells (Horsington et al., 2013). In addition, both kinases also promote viral release independently of their ability to phosphorylate A36 (Horsington et al., 2013). The identity of this additional Abl and Arg substrate that presumably promotes actin-dependent viral release remains to be established. However, an attractive candidate is N-WASP, as its ability to activate Arp2/3-dependent actin polymerization is enhanced by Arg and Abl mediated phosphorylation (Burton et al., 2005; Miller et al., 2010). Another reason to think N-WASP might be the target is because the release of Vaccinia from infected cells has strong parallels with exocytosis, which also involves N-WASP (Gasman et al., 2004). Furthermore, the recruitment of the phosphoinositide 5-phosphatase SHIP2, a negative regulator of viral release that is not involved in actin tail formation depends on its SH2 domain and the presence of N-WASP (McNulty et al., 2011).

A36 induced actin tails help propel Vaccinia into neighbouring cells (see Fig. 4 (Cudmore et al., 1995) to enhance the spread of infection (Röttger et al., 1999; Sanderson et al., 1998; Ward and Moss, 2001a; Wolffe et al., 1998). However, it was unclear how Vaccinia-induced actin polymerization promoted viral spread in a cell monolayer faster than the virus replication cycle. This mystery has now been solved (Doceul et al., 2010). Live cell imaging of viral spread during plaque formation reveals that if a virus (CEV or EEV) lands on a recently infected neighbouring cell, it is not internalized but rather induces an actin tail. The power of actin polymerization then propels the virus across the outer surface of the infected cell and onto adjacent non-infected cells, where internalization can occur. This mechanism allows the virus to ignore already infected cells, thus enhancing the rate of viral spread through the cell monolayer. The ability of the virus to surf across the surface of an infected cell is dependent on the presence of A33 and A36 in the plasma membrane (Doceul et al., 2010). The SCR4 domain of B5, which is involved in activating Src (Newsome et al., 2004), is also required for viral surfing (Doceul et al., 2012). These observations suggest that the surfing virus is effectively mimicking the normal process of actin tail formation by inducing Src and Abl kinase activation and tyrosine phosphorylation of A36 when it interacts with the extracellular domain of A33 on plasma membrane of the infected cell.

4.7. Conclusions

Studies from many different labs have collectively demonstrated that the ability of Vaccinia to stimulate actin polymerization

at the plasma membrane plays an important role in promoting the spread of infection. Moreover, unravelling exactly how Vaccinia stimulates actin polymerization has provided unprecedented insights into how a co-operative phosphotyrosine-based Nck and N-WASP signalling network activated by Src and Abl family kinases is assembled, regulated and functions at the molecular level to stimulate Arp2/3 dependent actin polymerization. Nevertheless, there are still many outstanding questions, including how the ability of Vaccinia to stimulate actin polymerization contributes to viral spread and pathogenesis in animal models. Furthermore, mechanistic basis of Src and Abl family kinase activation at the plasma membrane is yet to be understood. Actin is not the only component of the cytoskeleton that is usurped by Vaccinia. In the absence of cell lysis, microtubule-based transport plays an essential role in viral egress, as it facilitates the virus reaching the plasma membrane. The importance of microtubule transport during Vaccinia infection was realised more than a decade ago, and yet our molecular understanding of this process lags far behind actin-based motility of the virus. This is probably in part because of the speed and transient nature of microtubule-based movement as compared to the robust actin-based motility of Vaccinia, which is some 5–10 times slower, making live imaging less problematic. Nevertheless, in recent years, studies have begun to uncover the molecular basis of kinesin-1 recruitment to IEV. Further analysis of the role of microtubule transport in infected cells promises to provide important fundamental insights into mechanisms regulating the recruitment and activity of kinesin-1. Moreover, such studies have the potential to uncover how additional motors participate in other aspects of virus replication including IMV transport and IEV assembly. Such analyses will also undoubtedly uncover general principals underlying the recruitment and regulation of motors by cellular cargoes outside the context of infection. The next phase of research into the role of signalling and the host cytoskeleton in promoting the spread of Vaccinia clearly promises to provide interesting insights for the virologist and cell biologist alike.

Acknowledgements

We thank Drs. David Barry and Jasmine Abella in the Way lab, London Research Institute, Cancer Research UK for comments on the MS. F.L. and M.W. are supported by funding from Cancer Research UK.

References

- Alvarez, D.E., Agaisse, H., 2013. The formin FHOD1 and the small GTPase Rac1 promote vaccinia virus actin-based motility. *J. Cell. Biol.* 202 (7), 1075–1090.
- Aoyama, T., Hata, S., Nakao, T., Tanigawa, Y., Oka, C., Kawauchi, M., 2009. Cayman ataxia protein caytaxin is transported by kinesin along neurites through binding to kinesin light chains. *J. Cell. Sci.* 122 (Pt 22), 4177–4185.
- Arakawa, Y., Cordeiro, J.V., Schleich, S., Newsome, T., Way, M., 2007a. The release of Vaccinia virus from infected cells requires RhoA-mDia modulation of cortical actin. *Cell Host Microbe* 1, 227–240.
- Arakawa, Y., Cordeiro, J.V., Way, M., 2007b. F11L-mediated inhibition of RhoA-mDia signaling stimulates microtubule dynamics during vaccinia virus infection. *Cell Host Microbe* 1, 213–226.
- Araki, Y., Kawano, T., Taru, H., Saito, Y., Wada, S., Miyamoto, K., Kobayashi, H., Ishikawa, H.O., Ohsugi, Y., Yamamoto, T., Matsuno, K., Kinjo, M., Suzuki, T., 2007. The novel cargo Alcadin induces vesicle association of kinesin-1 motor components and activates axonal transport. *Embo J.* 26 (6), 1475–1486.
- Bahar, M.W., Graham, S.C., Chen, R.A., Cooray, S., Smith, G.L., Stuart, D.I., Grimes, J.M., 2011. How vaccinia virus has evolved to subvert the host immune response. *J. Struct. Biol.* 175, 127–134.
- Biro, M., Romeo, Y., Kroschwald, S., Bovellan, M., Boden, A., Tcherkezian, J., Roux, P.P., Charras, G., Paluch, E.K., 2013. Cell cortex composition and homeostasis resolved by integrating proteomics and quantitative imaging. *Cytoskeleton (Hoboken)* 70, 741–754.
- Bovellan, M., Romeo, Y., Biro, M., Boden, A., Chugh, P., Yonis, A., Vaghela, M., Fritzsche, M., Moulding, D., Thorogate, R., Jegou, A., Thrasher, A.J., Romet-Lemonne, G., Roux, P.P., Paluch, E.K., Charras, G., 2014. Cellular control of cortical actin nucleation. *Curr. Biol.* 24 (14), 1628–1635.

- Burton, E.A., Oliver, T.N., Pendergast, A.M., 2005. *Abl* kinases regulate actin comet tail elongation via an N-WASP-dependent pathway. *Mol. Cell. Biol.* 25 (20), 8834–8843.
- Carter, G.C., Law, M., Hollinshead, M., Smith, G.L., 2005. Entry of the vaccinia virus intracellular mature virion and its interactions with glycosaminoglycans. *J. Gen. Virol.* 86 (Pt 5), 1279–1290.
- Charras, G.T., Hu, C.K., Coughlin, M., Mitchison, T.J., 2006. Reassembly of contractile actin cortex in cell blebs. *J. Cell. Biol.* 175 (3), 477–490.
- Chieriegatti, E., Gartner, A., Stoffer, H.E., Bahler, M., 1998. Myr 7 is a novel myosin IX-RhoGAP expressed in rat brain. *J. Cell. Sci.* 111 (Pt 24), 3597–3608.
- Chung, C.S., Chen, C.H., Ho, M.Y., Huang, C.Y., Liao, C.L., Chang, W., 2006. Vaccinia virus proteome: identification of proteins in vaccinia virus intracellular mature virion particles. *J. Virol.* 80 (5), 2127–2140.
- Clark, A.G., Dierkes, K., Paluch, E.K., 2013. Monitoring actin cortex thickness in live cells. *Biophys. J.* 105 (3), 570–580.
- Cordeiro, J.V., Guerra, S., Arakawa, Y., Dodding, M.P., Esteban, M., Way, M., 2009. F11-mediated inhibition of RhoA signalling enhances the spread of vaccinia virus in vitro and in vivo in an intranasal mouse model of infection. *PLoS One* 4 (12), e8506.
- Cudmore, S., Cossart, P., Griffiths, G., Way, M., 1995. Actin-based motility of vaccinia virus. *Nature* 378 (6557), 636–638.
- Cudmore, S., Reckmann, I., Griffiths, G., Way, M., 1996. Vaccinia virus: a model system for actin-membrane interactions. *J. Cell Sci.* 109, 1739–1747.
- Davey, N.E., Trave, G., Gibson, T.J., 2011. How viruses hijack cell regulation. *Trends Biochem. Sci.* 36 (3), 159–169.
- Doceul, V., Hollinshead, M., Breiman, A., Laval, K., Smith, G.L., 2012. Protein B5 is required on extracellular enveloped vaccinia virus for repulsion of superinfecting virions. *J. Gen. Virol.* 93 (Pt 9), 1876–1886.
- Doceul, V., Hollinshead, M., van der Linden, L., Smith, G.L., 2010. Repulsion of superinfecting virions: a mechanism for rapid virus spread. *Science* 327 (5967), 873–876.
- Dodding, M.P., Mitter, R., Humphries, A.C., Way, M., 2011. A kinesin-1 binding motif in vaccinia virus that is widespread throughout the human genome. *Embo J.* 30 (22), 4523–4538.
- Dodding, M.P., Newsome, T.P., Collinson, L.M., Edwards, C., Way, M., 2009. An E2-F12 complex is required for IEV morphogenesis during vaccinia infection. *Cell. Microbiol.* 11, 808–824.
- Dodding, M.P., Way, M., 2009. Nck- and N-WASP-dependent actin-based motility is conserved in divergent vertebrate poxviruses. *Cell Host Microbe* 6 (6), 536–550.
- Dodding, M.P., Way, M., 2011. Coupling viruses to dynein and kinesin-1. *Embo J.* 30 (17), 3527–3539.
- Domí, A., Weisberg, A.S., Moss, B., 2008. Vaccinia virus E2L null mutants exhibit a major reduction in extracellular virion formation and virus spread. *J. Virol.* 82 (9), 4215–4226.
- Donnelly, S.K., Weisswange, I., Zettl, M., Way, M., 2013. WIP provides an essential link between Nck and N-WASP during Arp2/3-dependent actin polymerization. *Curr. Biol.* 23 (11), 999–1006.
- Duteyrat, J.L., Gelfi, J., Bertagnoli, S., 2006. Ultrastructural study of myxoma virus morphogenesis. *Arch. Virol.* 151 (11), 2161–2180.
- Erb, M.L., Pogliano, J., 2013. Cytoskeletal proteins participate in conserved viral strategies across kingdoms of life. *Curr. Opin. Microbiol.* 16 (6), 786–789.
- Franker, M.A., Hoogenraad, C.C., 2013. Microtubule-based transport – basic mechanisms, traffic rules and role in neurological pathogenesis. *J. Cell Sci.* 126 (Pt 11), 2319–2329.
- Frischknecht, F., Moreau, V., Röttger, S., Gonfloni, S., Reckmann, I., Superti-Furga, G., Way, M., 1999. Actin based motility of vaccinia mimics receptor tyrosine kinase signalling. *Nature* 401, 926–929.
- Fritzsche, M., Lewalle, A., Duke, T., Kruse, K., Charras, G., 2013. Analysis of turnover dynamics of the submembranous actin cortex. *Mol. Biol. Cell.* 24, 757–767.
- Fu, M.M., Holzbaur, E.L., 2014. Integrated regulation of motor-driven organelle transport by scaffolding proteins. *Trends Cell Biol.*
- García-Mata, R., Burridge, K., 2007. Catching a GEF by its tail. *Trends Cell Biol.* 17 (1), 36–43.
- Gasman, S., Chasserot-Golaz, S., Malacombe, M., Way, M., Bader, M.F., 2004. Regulated exocytosis in neuroendocrine cells: a role for subplasmalemmal Cdc42/N-WASP-induced actin filaments. *Mol. Biol. Cell.* 15 (2), 520–531.
- Gauthier, N.C., Masters, T.A., Sheetz, M.P., 2012. Mechanical feedback between membrane tension and dynamics. *Trends Cell Biol.* 22 (10), 527–535.
- Geada, M.M., Galindo, I., Lorenzo, M.M., Perdiguer, B., Blasco, R., 2001. Movements of vaccinia virus intracellular enveloped virions with GFP tagged to the F13L envelope protein. *J. Gen. Virol.* 82 (Pt 11), 2747–2760.
- Giallourakis, C., Cao, Z., Green, T., Wachtel, H., Xie, X., Lopez-Illasaca, M., Daly, M., Rioux, J., Xavier, R., 2006. A molecular-properties-based approach to understanding PDZ domain proteins and PDZ ligands. *Genome Res.* 16 (8), 1056–1072.
- Greber, U.F., Way, M., 2006. A superhighway to virus infection. *Cell* 124 (4), 741–754.
- Haller, S.L., Peng, C., McFadden, G., Rothenburg, S., 2014. Poxviruses and the evolution of host range and virulence. *Infect. Genet. Evol.* 21, 15–40.
- Handa, Y., Durkin, C.H., Dodding, M.P., Way, M., 2013. Vaccinia virus F11 promotes viral spread by acting as a PDZ-containing scaffolding protein to bind myosin-9A and inhibit RhoA signaling. *Cell Host Microbe* 14 (1), 51–62.
- Harris, B.Z., Lim, W.A., 2001. Mechanism and role of PDZ domains in signaling complex assembly. *J. Cell Sci.* 114 (Pt 18), 3219–3231.
- Herrero-Martínez, E., Roberts, K.L., Hollinshead, M., Smith, G.L., 2005. Vaccinia virus intracellular enveloped virions move to the cell periphery on microtubules in the absence of the A36R protein. *J. Gen. Virol.* 86 (Pt 11), 2961–2968.
- Hiller, G., Jungwirth, C., Weber, K., 1981. Fluorescence microscopical analysis of the life cycle of vaccinia virus in the chick embryo fibroblasts. *Exp. Cell Res.* 132, 81–87.
- Hiller, G., Weber, K., Schneider, L., Parajsz, C., Jungwirth, C., 1979. Interaction of assembled progeny pox viruses with the cellular cytoskeleton. *Virology* 98, 142–153.
- Hollinshead, M., Rodger, G., Van Eijl, H., Law, M., Hollinshead, R., Vaux, D.J., Smith, G.L., 2001. Vaccinia virus utilizes microtubules for movement to the cell surface. *J. Cell Biol.* 154 (2), 389–402.
- Horsington, J., Lynn, H., Turnbull, L., Cheng, D., Braet, F., Diefenbach, R.J., Whitchurch, C.B., Karupiah, G., Newsome, T.P., 2013. A36-dependent actin filament nucleation promotes release of vaccinia virus. *PLoS Pathog.* 9 (3), e1003239.
- Huang, C.Y., Lu, T.Y., Bair, C.H., Chang, Y.S., Jwo, J.K., Chang, W., 2008. A novel cellular protein, VPEF, facilitates vaccinia virus penetration into HeLa cells through fluid phase endocytosis. *J. Virol.* 82 (16), 7988–7999.
- Humphries, A.C., Dodding, M.P., Barry, D.J., Collinson, L.M., Durkin, C.H., Way, M., 2012. Clathrin potentiates vaccinia-induced actin polymerization to facilitate viral spread. *Cell Host Microbe* 12 (3), 346–359.
- Humphries, A.C., Donnelly, S.K., Way, M., 2014. Cdc42 and the Rho GEF intersectin-1 collaborate with Nck to promote N-WASP-dependent actin polymerisation. *J. Cell Sci.* 127, 673–685.
- Humphries, A.C., Way, M., 2013. The non-canonical roles of clathrin and actin in pathogen internalization, egress and spread. *Nat. Rev. Microbiol.* 11 (8), 551–560.
- Hunter, M.P., Russo, A., O'Bryan, J.P., 2013. Emerging Roles for Intersectin (ITSN) in Regulating Signaling and Disease Pathways. *Int. J. Mol. Sci.* 14 (4), 7829–7852.
- Hussain, N.K., Jenna, S., Glogauer, M., Quinn, C.C., Wasiak, S., Guipponi, M., Antonarakis, S.E., Kay, B.K., Stossel, T.P., Lamarque-Vane, N., McPherson, P.S., 2001. Endocytic protein intersectin-1 regulates actin assembly via Cdc42 and N-WASP. *Nat. Cell Biol.* 3 (10), 927–932.
- Irwin, C.R., Evans, D.H., 2012. Modulation of the myxoma virus plaque phenotype by vaccinia virus protein F11. *J. Virol.* 86 (13), 7167–7179.
- Irwin, C.R., Favis, N.A., Agopowicz, K.C., Hitt, M.M., Evans, D.H., 2013. Myxoma virus oncolytic efficiency can be enhanced through chemical or genetic disruption of the actin cytoskeleton. *PLoS One* 8 (12), e84134.
- Ivarsson, Y., 2012. Plasticity of PDZ domains in ligand recognition and signaling. *FEBS Lett.* 586 (17), 2638–2647.
- Jacobs, B.L., Langland, J.O., Kibler, K.V., Denzler, K.L., White, S.D., Holecek, S.A., Wong, S., Huynh, T., Baskin, C.R., 2009. Vaccinia virus vaccines: past, present and future. *Antiviral Res.* 84 (1), 1–13.
- Javier, R.T., Rice, A.P., 2011. Emerging theme: cellular PDZ proteins as common targets of pathogenic viruses. *J. Virol.* 85 (22), 11544–11556.
- Jeshtadi, A., Burgos, P., Stubbs, C.D., Parker, A.W., King, L.A., Skinner, M.A., Botchway, S.W., 2010. Interaction of poxvirus intracellular mature virion proteins with the TPR domain of kinesin light chain in live infected cells revealed by two-photon-induced fluorescence resonance energy transfer fluorescence lifetime imaging microscopy. *J. Virol.* 84 (24), 12886–12894.
- Johnston, S.C., Ward, B.M., 2008. The vaccinia virus protein F12 associates with IEV through an interaction with A36. *J. Virol.* 83, 1708–1717.
- Kilcher, S., Schmidt, F.I., Schneider, C., Kopf, M., Helenius, A., Mercer, J., 2014. siRNA screen of early poxvirus genes identifies the AAA+ ATPase D5 as the virus genome-uncoating factor. *Cell Host Microbe* 15 (1), 103–112.
- Kirn, D.H., Thorne, S.H., 2009. Targeted and armed oncolytic poxviruses: a novel multi-mechanistic therapeutic class for cancer. *Nat. Rev. Cancer* 9 (1), 64–71.
- Konecna, A., Frischknecht, R., Kinter, J., Ludwig, A., Steuble, M., Meskenaitė, V., Indermuhle, M., Engel, M., Cen, C., Mateos, J.M., Streit, P., Sonderegger, P., 2006. Calsynenin-1 docks vesicular cargo to kinesin-1. *Mol. Biol. Cell* 17 (8), 3651–3663.
- Krempien, U., Schneider, L., Hiller, G., Weber, K., Katz, E., Jungwirth, C., 1981. Conditions for pox virus-specific microvilli formation studied during synchronized virus assembly. *Virology* 113 (2), 556–564.
- Law, M., Carter, G.C., Roberts, K.L., Hollinshead, M., Smith, G.L., 2006. Ligand-induced and nonfusogenic dissolution of a viral membrane. *Proc. Natl. Acad. Sci. USA* 103 (15), 5989–5994.
- Law, M., Hollinshead, M., Lee, H.J., Smith, G.L., 2004. Yaba-like disease virus protein Y144R, a member of the complement control protein family, is present on enveloped virions that are associated with virus-induced actin tails. *J. Gen. Virol.* 85 (Pt 5), 1279–1290.
- Loisel, T.P., Boujemaa, R., Pantaloni, D., Carlier, M.-F., 1999. Reconstitution of actin based motility of *Listeria* and *Shigella* using pure proteins. *Nature* 401, 613–616.
- Luck, K., Charbonnier, S., Trave, G., 2012. The emerging contribution of sequence context to the specificity of protein interactions mediated by PDZ domains. *FEBS Lett.* 586 (17), 2648–2661.
- Lynn, H., Horsington, J., Ter, L.K., Han, S., Chew, Y.L., Diefenbach, R.J., Way, M., Chaudhri, G., Karupiah, G., Newsome, T.P., 2012. Loss of cytoskeletal transport during egress critically attenuates ectromelia virus infection in vivo. *J. Virol.* 86 (13), 7427–7443.
- Matthews, H.K., Delabre, U., Rohn, J.L., Guck, J., Kunda, P., Baum, B., 2012. Changes in Ect2 localization couple actomyosin-dependent cell shape changes to mitotic progression. *Dev. Cell* 23 (2), 371–383.
- McNulty, S., Powell, K., Erneux, C., Kalman, D., 2011. The host phosphoinositide 5-phosphatase SHIP2 regulates dissemination of vaccinia virus. *J. Virol.* 85 (14), 7402–7410.
- Mercer, J., Helenius, A., 2008. Vaccinia virus uses macropinocytosis and apoptotic mimicry to enter host cells. *Science* 320 (5875), 531–535.

- Mercer, J., Knebel, S., Schmidt, F.I., Crouse, J., Burkard, C., Helenius, A., 2010. Vaccinia virus strains use distinct forms of macropinocytosis for host-cell entry. *Proc. Natl. Acad. Sci. USA* 107 (20), 9346–9351.
- Mercer, J., Snijder, B., Sacher, R., Burkard, C., Bleck, C.K., Stahlberg, H., Pelkmans, L., Helenius, A., 2012. RNAi screening reveals proteasome- and Cullin3-dependent stages in vaccinia virus infection. *Cell Rep.* 2 (4), 1036–1047.
- Miki, H., Sasaki, T., Takai, Y., Takenawa, T., 1998. Induction of filopodium formation by a WASP-related actin-depolymerizing protein N-WASP. *Nature* 391, 93–96.
- Miller, M.M., Lapetina, S., MacGrath, S.M., Sfakianos, M.K., Pollard, T.D., Koleske, A.J., 2010. Regulation of actin polymerization and adhesion-dependent cell edge protrusion by the Abl-related gene (Arg) tyrosine kinase and N-WASP. *Biochemistry* 49 (10), 2227–2234.
- Mohamed, M.R., McFadden, G., 2009. NFkB inhibitors: strategies from poxviruses. *Cell Cycle* 8 (19), 3125–3132.
- Morales, I., Carbajal, M.A., Bohn, S., Holzer, D., Kato, S.E., Greco, F.A., Moussatche, N., Locker, J.K., 2008. The vaccinia virus F11 gene product facilitates cell detachment and promotes migration. *Traffic* 9, 1283–1298.
- Moreau, V., Frischknecht, F., Reckmann, I., Vincentelli, R., Rabut, G., Stewart, D., Way, M., 2000. A complex of N-WASP and WIP integrates signalling cascades that lead to actin polymerization. *Nat. Cell Biol.* 2, 441–448.
- Morgan, G.W., Hollinshead, M., Ferguson, B.J., Murphy, B.J., Carpentier, D.C., Smith, G.L., 2010. Vaccinia protein F12 has structural similarity to kinesin light chain and contains a motor binding motif required for virion export. *PLoS Pathog.* 6 (2), e1000785.
- Moss, B., 2007. Poxviridae: the viruses and their replication. In: Knipe, D.M., Howley, P.M. (Eds.), *Fields Virology*, 5th ed. Lippincott Williams and Wilkins, New York, pp. 2905–2945.
- Newsome, T.P., Scaplehorn, N., Way, M., 2004. SRC mediates a switch from microtubule- to actin-based motility of vaccinia virus. *Science* 306 (5693), 124–129.
- Newsome, T.P., Weisswange, I., Frischknecht, F., Way, M., 2006. Abl collaborates with Src family kinases to stimulate actin-based motility of vaccinia virus. *Cell Microbiol.* 8 (2), 233–241.
- Niehl, A., Pena, E.J., Amari, K., Heinlein, M., 2013. Microtubules in viral replication and transport. *Plant J.* 75 (2), 290–308.
- Nourry, C., Grant, S.G., Borg, J.P., 2003. PDZ domain proteins: plug and play! *Sci. STKE* 2003 (179), RE7.
- Omelchenko, T., Hall, A., 2012. Myosin-IXA regulates collective epithelial cell migration by targeting RhoGAP activity to cell-cell junctions. *Curr. Biol.* 22 (4), 278–288.
- Payne, L.G., Kristensson, K., 1982. The effect of cytochalasin D and monensin on enveloped vaccinia virus release. *Arch. Virol.* 74, 11–20.
- Ploubidou, A., Moreau, V., Ashman, K., Reckmann, I., Gonzalez, C., Way, M., 2000. Vaccinia virus infection disrupts microtubule organization and centrosome function. *EMBO J.* 19, 3932–3944.
- Ponting, C.P., 1997. Evidence for PDZ domains in bacteria, yeast, and plants. *Protein Sci.* 6 (2), 464–468.
- Postigo, A., Ferrer, P.E., 2009. Viral inhibitors reveal overlapping themes in regulation of cell death and innate immunity. *Microbes Infect.* 11 (13), 1071–1078.
- Radtko, K., Dohner, K., Sodeik, B., 2006. Viral interactions with the cytoskeleton: a hitchhiker's guide to the cell. *Cell Microbiol.* 8 (3), 387–400.
- Reeves, P.M., Bommaribus, B., Lebeis, S., Christensen, J., Swimm, A., Chahroudi, A., Chavan, R., Feinberg, M.B., Veach, D., Bornmann, W., Sherman, M., Kalman, D., 2005. Disabling poxvirus pathogenesis by inhibition of Abl-family tyrosine kinases. *Nat. Med.* 11 (7), 731–739.
- Reeves, P.M., Smith, S.K., Olson, V.A., Thorne, S.H., Bornmann, W., Damon, I.K., Kalman, D., 2011. Variola and monkeypox viruses utilize conserved mechanisms of virion motility and release that depend on abl and SRC family tyrosine kinases. *J. Virol.* 85 (1), 21–31.
- Resch, W., Hixson, K.K., Moore, R.J., Lipton, M.S., Moss, B., 2007. Protein composition of the vaccinia virus mature virion. *Virology* 358 (1), 233–247.
- Rietdorf, J., Ploubidou, A., Reckmann, I., Holmström, A., Frischknecht, F., Zettl, M., Zimmermann, T., Way, M., 2001. Kinesin dependent movement on microtubules precedes actin based motility of vaccinia virus. *Nat. Cell Biol.* 3, 992–1000.
- Roberts, K.L., Smith, G.L., 2008. Vaccinia virus morphogenesis and dissemination. *Trends Microbiol.* 10, 472–479.
- Roper, R.L., Wolffe, E.J., Weisberg, A., Moss, B., 1998. The envelope protein encoded by the A33R gene is required for formation of actin-containing microvilli and efficient cell-to-cell spread of vaccinia virus. *J. Virol.* 72, 4192–4204.
- Röttger, S., Frischknecht, F., Reckmann, I., Smith, G.L., Way, M., 1999. Interactions between vaccinia virus IEV membrane proteins and their roles in IEV assembly and actin tail formation. *J. Virol.* 73, 2863–2875.
- Salbreux, G., Charras, G., Paluch, E., 2012. Actin cortex mechanics and cellular morphogenesis. *Trends Cell Biol.* 22 (10), 536–545.
- Sanderson, C.M., Frischknecht, F., Way, M., Hollinshead, M., Smith, G.L., 1998. Roles of vaccinia virus EEV-specific proteins in intracellular actin tail formation and low pH-induced cell-cell fusion. *J. Gen. Virol.* 79, 1415–1425.
- Scaplehorn, N., Holmstrom, A., Moreau, V., Frischknecht, F., Reckmann, I., Way, M., 2002. Grb2 and nck act cooperatively to promote actin-based motility of vaccinia virus. *Curr. Biol.* 12 (9), 740–745.
- Schmidt, F.I., Bleck, C.K., Helenius, A., Mercer, J., 2011. Vaccinia extracellular virions enter cells by macropinocytosis and acid-activated membrane rupture. *EMBO J.* 30 (17), 3647–3661.
- Schmidt, F.I., Bleck, C.K., Reh, L., Novy, K., Wollscheid, B., Helenius, A., Stahlberg, H., Mercer, J., 2013. Vaccinia virus entry is followed by core activation and proteasome-mediated release of the immunomodulatory effector VH1 from lateral bodies. *Cell Rep.* 4 (3), 464–476.
- Schmidt, M.R., Maritzen, T., Kukhtina, V., Higman, V.A., Doglio, L., Barak, N.N., Strauss, H., Oschkinat, H., Dotti, C.G., Hauke, V., 2009. Regulation of endosomal membrane traffic by a Gadin/AP-1/kinesin KIF5 complex. *Proc. Natl. Acad. Sci. USA* 106, 15344–15349.
- Schonichen, A., Alexander, M., Gasteier, J.E., Cuesta, F.E., Fackler, O.T., Geyer, M., 2006. Biochemical characterization of the diaphanous autoregulatory interaction in the formin homology protein FHOD1. *J. Biol. Chem.* 281 (8), 5084–5093.
- Schonichen, A., Mannherz, H.G., Behrmann, E., Mazur, A.J., Kuhn, S., Silvan, U., Schoenenberger, C.A., Fackler, O.T., Raunser, S., Dehmelt, L., Geyer, M., 2013. FHOD1 is a combined actin filament capping and bundling factor that selectively associates with actin arcs and stress fibers. *J. Cell Sci.* 126 (Pt 8), 1891–1901.
- Schulte, A., Stolp, B., Schonichen, A., Pylypenko, O., Rak, A., Fackler, O.T., Geyer, M., 2008. The human formin FHOD1 contains a bipartite structure of FH3 and GTPase-binding domains required for activation. *Structure* 16 (9), 1313–1323.
- Sedzinski, J., Biro, M., Oswald, A., Tinevez, J.Y., Salbreux, G., Paluch, E., 2011. Polar actomyosin contractility destabilizes the position of the cytokinetic furrow. *Nature* 476 (7361), 462–466.
- Smith, G.L., Vanderplassen, A., Law, M., 2002. The formation and function of extracellular enveloped vaccinia virus. *J. Gen. Virol.* 83, 2915–2931.
- Snapper, S.B., Takeshima, F., Anton, I., Liu, C.H., Thomas, S.M., Nguyen, D., Dudley, D., Fraser, H., Purich, D., Lopez-Illasaca, M., Klein, C., Davidson, L., Bronson, R., Mulligan, R.C., Southwick, F., Geha, R., Goldberg, M.B., Rosen, F.S., Hartwig, J.H., Alt, F.W., 2001. N-WASP deficiency reveals distinct pathways for cell surface projections and microbial actin-based motility. *Nat. Cell Biol.* 3 (10), 897–904.
- Sodeik, B., 2000. Mechanisms of viral transport in the cytoplasm. *Trends Microbiol.* 8, 465–472.
- Stephens, D.J., 2012. Functional coupling of microtubules to membranes - implications for membrane structure and dynamics. *J. Cell Sci.* 125 (Pt 12), 2795–2804.
- Stokes, G.V., 1976. High-voltage electron microscope study of the release of vaccinia virus from whole cells. *J. Virol.* 18 (2), 636–643.
- Subbiah, V.K., Kranjec, C., Thomas, M., Banks, L., 2011. PDZ domains: the building blocks regulating tumorigenesis. *Biochem. J.* 439 (2), 195–205.
- Taylor, M.P., Koyuncu, O.O., Enquist, L.W., 2011. Subversion of the actin cytoskeleton during viral infection. *Nat. Rev. Microbiol.* 9 (6), 427–439.
- Thorne, S.H., 2011. Immunotherapeutic potential of oncolytic vaccinia virus. *Immunol. Res.* 50 (2–3), 286–293.
- Tinevez, J.Y., Schulze, U., Salbreux, G., Roensch, J., Joanny, J.F., Paluch, E., 2009. Role of cortical tension in bleb growth. *Proc. Natl. Acad. Sci. USA* 106 (44), 18581–18586.
- Townsend, A.C., Weisberg, A.S., Wagenaar, T.R., Moss, B., 2006. Vaccinia virus entry into cells via a low-pH-dependent endosomal pathway. *J. Virol.* 80 (18), 8899–8908.
- Valderrama, F., Cordeiro, J.V., Schleich, S., Frischknecht, F., Way, M., 2006. Vaccinia virus-induced cell motility requires F11L-mediated inhibition of RhoA signaling. *Science* 311 (5759), 377–381.
- Van den Broeke, C., Jacob, T., Favoreel, H.W., 2014. Rho'ing in and out of cells: viral interactions with Rho GTPase signaling. *Small GTPases* 5, e28318.
- van Eijl, H., Hollinshead, M., Rodger, G., Zhang, W.H., Smith, G.L., 2002. The vaccinia virus F12L protein is associated with intracellular enveloped virus particles and is required for their egress to the cell surface. *J. Gen. Virol.* 83 (Pt 1), 195–207.
- van Eijl, H., Hollinshead, M., Smith, G.L., 2000. The vaccinia virus A36R protein is a type Ib membrane protein present on intracellular but not extracellular enveloped virus particles. *Virology* 271 (1), 26–36.
- Volz, A., Sutter, G., 2013. Protective efficacy of Modified Vaccinia virus Ankara in preclinical studies. *Vaccine* 31 (39), 4235–4240.
- Walsh, S.R., Dolin, R., 2011. Vaccinia viruses: vaccines against smallpox and vectors against infectious diseases and tumors. *Exp. Rev. Vaccines* 10 (8), 1221–1240.
- Ward, B.M., 2005. Visualization and characterization of the intracellular movement of vaccinia virus intracellular mature virions. *J. Virol.* 79 (8), 4755–4763.
- Ward, B.M., Moss, B., 2001a. Vaccinia virus intracellular movement is associated with microtubules and independent of actin tails. *J. Virol.* 75 (23), 11651–11663.
- Ward, B.M., Moss, B., 2001b. Visualization of intracellular movement of vaccinia virus virions containing a green fluorescent protein-B5R membrane protein chimera. *J. Virol.* 75 (10), 4802–4813.
- Ward, B.M., Moss, B., 2004. Vaccinia virus A36R membrane protein provides a direct link between intracellular enveloped virions and the microtubule motor kinesin. *J. Virol.* 78 (5), 2486–2493.
- Ward, B.M., Weisberg, A.S., Moss, B., 2003. Mapping and functional analysis of interaction sites within the cytoplasmic domains of the vaccinia virus A33R and A36R envelope proteins. *J. Virol.* 77 (7), 4113–4126.
- Weisswange, I., Newsome, T.P., Schleich, S., Way, M., 2009. The rate of N-WASP exchange limits the extent of ARP2/3-complex-dependent actin-based motility. *Nature* 458 (7234), 87–91.
- Welch, M.D., Way, M., 2013. Arp2/3-mediated actin-based motility: a tail of pathogen abuse. *Cell Host Microbe* 14 (3), 242–255.
- Wolffe, E.J., Katz, E., Weisberg, A., Moss, B., 1997. The A34R glycoprotein gene is required for induction of specialized actin-containing microvilli and efficient cell-to-cell transmission of vaccinia virus. *J. Virol.* 71, 3904–3915.
- Wolffe, E.J., Weisberg, A.S., Moss, B., 1998. Role for the vaccinia virus A36R outer envelope protein in the formation of virus-tipped actin-containing microvilli and cell-to-cell virus spread. *Virology* 25, 20–26.
- Yang, C., Kazanietz, M.G., 2007. Chimaerins: GAPs that bridge diacylglycerol signalling and the small G-protein Rac. *Biochem. J.* 403 (1), 1–12.

- Yoder, J.D., Chen, T.S., Gagnier, C.R., Vemulapalli, S., Maier, C.S., Hruby, D.E., 2006. [Pox proteomics: mass spectrometry analysis and identification of Vaccinia virion proteins](#). *Viol. J.* 3, 10.
- Yutin, N., Faure, G., Koonin, E.V., Mushegian, A.R., 2014. [Chordopoxvirus protein F12 implicated in enveloped virion morphogenesis is an inactivated DNA polymerase](#). *Biol. Direct* 9 (1), 22.
- Zettl, M., Way, M., 2002. [The WH1 and EVH1 domains of WASP and Ena/VASP family members bind distinct sequence motifs](#). *Curr. Biol.* 12 (18), 1617–1622.
- Zhang, W.H., Wilcock, D., Smith, G.L., 2000. [Vaccinia virus F12L protein is required for actin tail formation, normal plaque size, and virulence](#). *J. Virol.* 74 (24), 11654–11662.