
ELECTROLYSIS #4

HOW DOES ELECTROLYSIS WORK?

The process of electrolysis involve three things:

- The external circuit as in the power supply
- · Reactions at the electrodes
- · Reactions in the electrolyte



NOTE:

When the cations and the anions gain or lose electron(s) at the electrodes, they form **atoms** or **molecules**. So, we say the cations and the anions are **discharged**.

Electrolysis Of Molten Ionic Compounds -

lonic compounds are sometimes called Binary compounds. A binary ionic compound which contains **only two elements**, mainly a <u>metal cation</u> and a <u>non-metal anion</u>.

When a binary ionic compound is heated, melted and undergoes electrolysis, <u>a metal and a non-metal elements are</u> formed as products.

2. Electrolysis Of Molten Potassium Iodide -

IONS PRESENT	K ⁺ , I ⁻
CATIONS	K+
ANIONS	Ĭ.
EQUATION AT CATHODE	$K^+ + e^- \longrightarrow K$ (REDUCTION)
EQUATION AT ANODE	$2I^{-} - 2e^{-} \longrightarrow I_{2}$ (OXIDATION)

$$\Rightarrow$$
 25 \longrightarrow I₂ + 2e⁻

3. Electrolysis Of Molten Lithium Bromide -

IONS PRESENT	Li+	, Br-	****		13
CATIONS	L_i^+				
ANIONS	Br-				
EQUATION AT CATHODE	Li+	+ e-	→ Li	(REDUCTION)	
EQUATION AT ANODE	288-	- 2e-	$\rightarrow \beta \gamma_{2}$	(OXIDATION)	
> 20r - Or2 + 2e-					

IONS PRESENT	Pb2+	B1-	
CATIONS	fb ²⁺		
ANIONS	Br-		
EQUATION AT CATHODE	Pb2+	+ 2e⁻ → Pb	(REDUCTION)
EQUATION AT ANODE	2Br -	→ Br ₂ + Ze ⁻	(OX IDATION)

5. Electrolysis Of Magnesium Nitride liquid -

IONS PRESENT	Mg2+,	N ³⁻	
CATIONS	Mg2+		
ANIONS	N ³⁻		
EQUATION AT CATHODE	Mg2+	+ 2e⁻ → Ng	(REDUCTION)
EQUATION AT ANODE	2 N ³⁻	- 6 e → N2	(OX IDATION)

$$\stackrel{>}{>}$$
 2N³⁻ \longrightarrow N₂ + 6e⁻

6. Electrolysis Of Aluminium Oxide liquid Lo Al₂ O₃ (L)

IONS PRESENT	A13+ O2-
CATIONS	A13+
ANIONS	O ²⁻
EQUATION AT CATHODE	$Al^{3+} + 3e^{-} \longrightarrow Al$ (REDUCTION)
EQUATION AT ANODE	20^{2-} - $4e^{-}$ \rightarrow O_2 (0XIDATION)

$$\Rightarrow$$
 202 - \rightarrow 02 + 4e

Electrolysis Of Aqueous Solutions Of Compounds -

Aqueous solutions of compounds means it is a <u>mixture</u> of the compound in water. For instance, an aqueous solution of Sodium Chloride is a mixture of Sodium Chloride and Water.

In an aqueous solution, more than one type of cations and one type of anions are present. During electrolysis, only one type of cation and one type of anion is preferentially discharged over the other. This is known as Selective Discharge Of Ions.

During the electrolysis of an aqueous solution, <u>if the electrodes are **inert / unreactive**</u>, the ions discharged at the electrode depends on three factors:

- Selective Discharge Of Cations
- 2. Selective Discharge Of Anions
- 3. Effect Of Concentration Of The Selective Discharge Of Anions

1. SELECTIVE DISCHARGE OF CATIONS -

The more reactive a metal, the greater its tendency to form its positive ion. Thus, during electrolysis,

- Ions of the more reactive metals, such as Potassium and Sodium, will remain as positive ions and will not get attracted and discharged at the Cathode.
- Ions of the **less** reactive metals, such as Copper and Silver and also the ion of Hydrogen <u>will not remain as positive ions</u> and will get attracted and discharged at the Cathode.

The ease of discharge of cations thus depends on the <u>position the metal (producing the cation) in the reactivity series</u>:

The ease of discharge of cations thus depends on the <u>position the metal (producing the cation) in the reactivity series</u>:

```
DIFFICULT
Potassium ion, K+
Sodium ion, Na+
Lithium ion, Li+
Calcium ion, Ca2+
                          EASE
                                         DISCHARGE
Magnesium ion, Mg2+
Aluminium ion, Al3+
                                 INCREASES -
Zinc ion, Zn2+
Iron ion, Fe2+
Lead ion, Pb2+
Hydrogen ion, H+
Copper ion, Cu2+
Silver ion, Ag+
```

2. SELECTIVE DISCHARGE OF ANIONS -

The ease of discharge of anions will depend on the position of the anion in the following list:

