

INDIAN INSTITUTE OF TECHNOLOGY, GUWAHATI

PRE-FINAL YEAR : 6<sup>th</sup> SEMESTER

# Measure Theory

*Prof. Pratyooosh KUMAR*

*Written by*  
Animesh Renanse

March 3, 2021

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Few introductory definitions . . . . .	2
1.2	Homework - I . . . . .	3
1.3	Measure of an Interval . . . . .	5
1.3.1	Outer Lebesgue Measure . . . . .	5
<b>2</b>	<b>Measures</b>	<b>6</b>
2.1	Algebras & Sigma-Algebras . . . . .	6
2.2	Measures . . . . .	8
2.2.1	Elementary Properties of Measures . . . . .	8
2.3	Outer Measures . . . . .	11
2.4	Exercises - II . . . . .	13
2.5	Lebesgue Measurability. . . . .	17
2.6	Does $\lambda^*(E) = 0$ implies $E$ is countable? . . . . .	21
2.6.1	Properties of Cantor Set . . . . .	21
2.7	A Non-measurable Set . . . . .	22
2.8	Regularity . . . . .	23
<b>3</b>	<b>Measurable Functions</b>	<b>25</b>
3.1	Almost Everywhere Property. . . . .	29
3.1.1	A striking result . . . . .	29
3.2	Cantor Set, again. . . . .	30
3.3	Sequence of Functions approximating a Measurable Function. . . . .	32
3.3.1	Replacing <i>simple</i> functions by <i>step</i> functions <sup>1</sup> . . . . .	35
3.4	Egorov's Theorem . . . . .	37
3.5	Lusin's Theorem . . . . .	38
<b>4</b>	<b>Integration of Measurable Functions</b>	<b>40</b>
4.1	Integration of Non-Negative Measurable Functions . . . . .	41
4.2	Monotone Convergence Theorem . . . . .	42
4.3	Fatou's Lemma . . . . .	45
4.4	Integration of General Real-Valued Measurable Functions . . . . .	46
4.4.1	Basic Properties of General Lebesgue Integral . . . . .	46
4.5	Dominated Convergence Theorem . . . . .	48

---

<sup>1</sup>Unfortunately, this section onwards, we would tentatively only work with the Lebesgue Measure space  $(\mathbb{R}, \mathcal{M}_{\lambda^*}, \lambda)$  for pedagogical reasons of the instructor, so the abstract versions of the following results would need to wait.

# 1 Introduction

Measure Theory is a field of mathematics which deals with formalizing the notion of a *measure* of a given quantity. Traditional notion of a *measure* fails in lots of complicated scenario. Hence, we need a rigorous definition of it. The work in this direction was forwarded by Lebesgue and others in the dawn of 20<sup>th</sup> century.

## 1.1 Few introductory definitions

These are few of the basic definitions that one might remember from real analysis.

- **Limit Points** :  $x \in X$  is called a *limit point* of a subset  $S \subseteq X$  if  $\forall r > 0, \exists a \neq x$  such that  $a \in S \cap B_r(x)$ . That is, ball of any size  $r$  around  $x$  contains atleast one point of  $S$ .
- **Isolated Points** :  $y \in S$  is called an *isolated point* of a subset  $S \subseteq X$  if  $\exists r > 0$  such that  $(B_r(y) \setminus \{y\}) \cap S = \Phi$ . That is,  $B_r(y)$  contains no other point of  $S$  apart from  $y$ .
  - Also note that every point of closure  $\bar{S}$  is either a limit point or an isolated point of  $S$ .
  - More specifically, any subset of  $\mathbb{R}^d$  is closed if and only if it contains all of it's limit points.
- **Perfect Set** :  $A$  is called a perfect set if  $A = A'$  where  $A'$  is the set of all *limit points* of  $A$ . More conveniently, if  $A$  does not contain any isolated points then it is a perfect set.  $\mathbb{R}$  is a perfect set.
- **Symmetric Difference** :  $A$  and  $B$  are two sets then symmetric difference is  $A \Delta B = (A \setminus B) \cup (B \setminus A)$ .
- **Power Set** : Collection of all subsets of a set  $S$ , written as  $P(S)$ .
- **Lower Bound** : A *lower bound* of a subset  $S$  of a poset  $(P, \leq)$  is an element  $a \in P$  such that  $a \leq x$  for all  $x \in S$ .
- **Infimum** : A lower bound  $p \in P$  is called an *infimum* of  $S$  if for all lower bounds  $y$  of  $S$  in  $P$ ,  $y \leq p$ .
- **Limit Infimum** : For a sequence  $\{x_n\}$ , limit inferior is defined by:

$$\begin{aligned} \liminf_{n \rightarrow \infty} x_n &= \lim_{n \rightarrow \infty} \left( \inf_{m \geq n} x_m \right) \\ &= \sup_{n \geq 0} \inf_{m \geq n} x_m \\ &= \sup \{ \inf \{ x_m \mid m \geq n \} \mid n \geq 0 \}. \end{aligned} \tag{1}$$

- **Upper Bound** : An *upper bound* of a subset  $S$  of a poset  $P$  is an element  $b \in P$  such that  $b \geq x$  for all  $x \in S$ .
- **Supremum** : An upper bound  $u \in P$  is called a *supremum* of  $S$  if for all upper bounds  $z$  of  $S$  in  $P$ ,  $z \geq u$ .
- **Limit Supremum** : For a sequence  $\{x_n\}$ , limit supremum is defined by:

$$\begin{aligned} \limsup_{n \rightarrow \infty} x_n &= \lim_{n \rightarrow \infty} \left( \sup_{m \geq n} x_m \right) \\ &= \inf_{n \geq 0} \sup_{m \geq n} x_m \\ &= \inf \{ \sup \{ x_m \mid m \geq n \} \mid n \geq 0 \} \end{aligned} \tag{2}$$

- **Limit** : Consider the sequence  $\{x_n\}$  in  $[-\infty, +\infty]$ , then  $\lim_n x_n$  is defined as

$$\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} x_n.$$

- **Lower Sum** :  $l(f, \mathcal{P})$  is the sum of the minimum functional values at the partition. That is,

$$l(f, \mathcal{P}) = \sum_{i=0}^{n-1} m_i (a_{i+1} - a_i)$$

where  $m_i = \inf \{ f(x) \mid x \in [a_{i-1}, a_i] \}$ .

- **Upper Sum** : Similarly,

$$u(f, \mathcal{P}) = \sum_{i=0}^{n-1} M_i(a_{i+1} - a_i)$$

where  $M_i = \sup\{f(x) \mid x \in [a_{i-1}, a_i]\}$ .

Remember that the function is *Riemann Integrable* if  $l(f, \mathcal{P}) = u(f, \mathcal{P})$ .

- **Countable Sets** : Note the following,

1. *Cardinality* : Sets  $X$  and  $Y$  have the same cardinality if there exists a bijection from  $X$  to  $Y$ .
2. *Finite Set* : A set is finite if it is empty or it has the same cardinality as  $\{1, 2, \dots, n\}$  for some  $n \in \mathbb{N}$ .
3. *Countably Infinite* : If the set has the same cardinality as  $\mathbb{N}$ .
4. *Enumeration* : An enumeration of a countably infinite set  $X$  is a bijection of  $\mathbb{N}$  onto  $X$ . That is, an enumeration is an infinite sequence  $\{x_n\}$  such that each of the  $x_i$ 's are in  $X$  and each element of  $X$  is  $x_i$  for some  $i$ .
5. *Countable* : A set is countable if it is finite or countably infinite. For example,  $\mathbb{N}$  is countable,  $\mathbb{Q}$  is also countable (!),  $\mathbb{R} \setminus \mathbb{Q}$  (irrationals) is not countable,  $\mathbb{R}$  is not countable.

- **Totally Bounded** : A subset  $B \subseteq X$  is *totally bounded* when it can be covered by a finite number of  $r$ -balls for all  $r > 0$ . That is,

$$\forall r > 0, \exists N \in \mathbb{N}, \exists a_1, \dots, a_N \in X \text{ such that } B \subseteq \bigcup_{n=1}^N B_r(a_n)$$

- **Compact Set** : A set  $K$  is said to be *compact* when given any cover of balls of possibly unequal radii, there is a finite sub-collection of them that still covers the set  $K$ . That is,

$$K \subseteq \bigcup_i B_{r_i}(a_i) \implies \exists i_1, \dots, i_N, K \subseteq \bigcup_{n=1}^N B_{r_{i_n}}(a_{i_n})$$

Note that *compact metric spaces are totally bounded* (!). Also, *compact sets are closed*.

- **Essential Supremum** : For a sequence  $\{x_n\}$ , essential supremum is defined as,

The problem begins with Riemann Integrable functions when we see that functions like Dirichlet function (1 on irrational and 0 on rational points) can become *measurable* even when the function is not continuous! This motivates the need of a formal notion of a *measure*.

## 1.2 Homework - I

1. Every open set in  $\mathbb{R}$  can be written as disjoint union of open intervals.

*Proof.* Let  $G \subseteq \mathbb{R}$  be a open subset. Now by definition of an open subset, we have that for any  $x \in G$ , there exists atleast one open subset  $U$  such that  $x \in U \subseteq G$ . Now consider the following union of all such open subsets of  $x$ ,

$$U_x = \bigcup_{x \in U \subseteq G} U$$

It's now easy to see that  $U_x$  is the largest such subset of  $G$ , as any other  $V \subseteq G$  such that  $x \in V$  is by definition contained in  $U_x$ . Moreover,  $U_x$  is an interval as it is an arbitrary union of open intervals. Now, define the following relation on  $G$ :

$$y \sim x \iff y \in U_x$$

Now we clearly have that  $x \in U_x$  (reflexive); for  $y \sim U_x$  we have  $U \subseteq U_x$  such that  $x, y \in U$ , hence  $x \in U_y$  (symmetric); for  $x \in U_y$  and  $y \in U_z$ , we have that  $x, y, z \in U_y$ , since  $z \in U_y \subseteq G$  so  $U_y \subseteq U_z$ , so  $x \in U_z$  (transitive). Hence  $\sim$  is an equivalence relation, hence  $\sim$  partitions the set  $G$ . Denote the set of all equivalence classes as  $\mathcal{I}$  so we get

$$G = \bigcup_{I \in \mathcal{I}} I$$

such that  $I_1 \cap I_2 = \emptyset$  for any  $I_1, I_2 \in \mathcal{I}$ . Now note that for any  $I \in \mathcal{I}$  is open because each  $I$  is generated by the relation  $\sim$  such that  $y \sim x$  iff  $y \in U_x$ . Hence for any  $z \in I$ , we have  $z \in U_x \subseteq G$  where  $U_x$  is open. Therefore, we have  $G = \bigcup_{I \in \mathcal{I}} I$  for disjoint open intervals in  $\mathcal{I}$ . ■

2. Prove that every non-empty perfect subset of  $\mathbb{R}$  (or  $\mathbb{R}^n$ ) is uncountable. That is, if  $A = A'$  then  $A$  is uncountable.

*Proof.* Take  $A \subseteq \mathbb{R}$  to be a perfect subset. Since  $A$  is perfect, therefore, it must contain all of its limit points or, equivalently, contains no isolated points. Clearly, then,  $A$  cannot be finite, but can only be countably infinite or uncountable. If it is uncountable, then the proof is over. If  $A$  is countably infinite, then we can write  $A$  as the following :

$$A = \{a_1, a_2, \dots\}.$$

Construct a ball around  $a_{i_1}$  of any radius  $r_1 > 0$ . Since  $A$  is perfect, therefore  $\exists a_{i_2} \in B_{r_1}(a_{i_1}) \cap A = C_1$ . Similarly, for some  $r_2 > 0$ , we have  $a_{i_3} \in B_{r_2}(a_{i_2}) \cap B_{r_1}(a_{i_1}) \cap A = C_2$  such that  $a_{i_1} \notin C_2$  and so on. In general, we would have the following,

$$a_{i_{n+1}} \in \left( \bigcap_{j=1}^n B_{r_j}(a_{i_j}) \right) \cap A = C_n.$$

Now, consider  $C = \bigcap_n C_n$ . Since  $C_{n+1} \subseteq C_n$ , therefore  $C \neq \Phi$ . But,  $a_i \notin C$  for any  $i \in \mathbb{N}$  as  $a_i \notin C_{i+1}$ . Therefore we have a contradiction. Hence  $A$  cannot be countably infinite, it must only be uncountable. ■

3. In the definition of Lebesgue Outer Measure on  $\mathbb{R}$ , one can instead take  $\mathcal{C}_A$  to be collection of infinite sequences of the any form from  $\{[a_n, b_n]\}, \{(a_n, b_n)\}$  or  $\{(a_n, b_n]\}$ .

*Proof.* Refer Proof of Proposition 9. ■

4. Show the following:

$$\bigcup_{n=1}^N E_n = \bigcup_{n=1}^N \left( E_n \cap \left( \bigcup_{k < n} E_k \right)^c \right)$$

*Proof.* Take  $x \in \bigcup_{n=1}^N E_n$ . Then  $\exists E_k$  for some  $a$  such that  $x \in E_a$ . Now, clearly,  $x \in E_a \subseteq (\bigcup_{k < a} E_k)^c$ , hence  $x \in (E_a \cap (\bigcup_{k < a} E_k)^c)$ . Hence, we have  $\bigcup_{n=1}^N E_n \subseteq \bigcup_{n=1}^N (E_n \cap (\bigcup_{k < n} E_k)^c)$ . The converse is easy to see too. ■

## 1.3 Measure of an Interval

In the definition of Riemann Integral, we see that a function is Riemann Integrable if  $u(f, \mathcal{P}) = l(f, \mathcal{P})$ . But it depends upon the definition of the *length of an interval*.

Note that in the traditional notion of *measure*, the number of points has no contribution in length. Hence the question becomes:

*How to measure a set  $A$  in  $\mathbb{R}$ ?*

For this, we define *Outer Measure*.

### 1.3.1 Outer Lebesgue Measure

The concept of this topic began from the following question : If we consider the cauchy sequence of Riemann Integrable functions on  $\mathbb{R}$ , will it converge to a Riemann Integrable function?

We follow the following notion of measure (atleast for the first part of the course). We do approximation by approximation.

Suppose  $A \subseteq \mathbb{R}$ . We define  $m^*(A)$  as the *Outer Lebesgue Measure* defined by

$$m^*(A) = \inf \left\{ \sum_{i=1}^{\infty} l(I_i) \mid A \subseteq \bigcup_i I_i \right\}$$

where  $I_i = [a_i, b_i)$ . Since we are taking infimum of the the collection, hence Note that it is approximating measure from outside. Also note that this is different from the definition of measurable sets.

Outer measure is a function  $m^* : P(\mathbb{R}) \rightarrow [0, \infty]$ .

## 2 Measures

### 2.1 Algebras & Sigma-Algebras

★ **Definition 1. (Algebra/Field)** Let  $X$  be an arbitrary set. A collection  $\mathcal{A} \subseteq P(X)$  of subsets of  $X$  is an algebra on  $X$  if:

- $X \in \mathcal{A}$ .
- $A \in \mathcal{A} \implies A^c \in \mathcal{A}$ .
- For each finite sequence  $A_1, A_2, \dots, A_n \in \mathcal{A}$  implies that

$$\bigcup_{i=1}^n A_i \in \mathcal{A}$$

- For each finite sequence  $A_1, A_2, \dots, A_n \in \mathcal{A}$  implies that

$$\bigcap_{i=1}^n A_i \in \mathcal{A}$$

★ **Definition 2. ( $\sigma$ -Algebra/ $\sigma$ -Field)** Let  $X$  be an arbitrary set. A collection  $\mathcal{A} \subseteq P(X)$  of subsets of  $X$  is a  $\sigma$ -algebra on  $X$  if:

- $X \in \mathcal{A}$ .
- $A \in \mathcal{A} \implies A^c \in \mathcal{A}$ .
- For each infinite sequence  $\{A_i\}$  such that  $A_i \in \mathcal{A}$ , it implies that

$$\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$$

- For each infinite sequence  $\{A_i\}$  such that  $A_i \in \mathcal{A}$ , it implies that

$$\bigcap_{i=1}^{\infty} A_i \in \mathcal{A}$$

✓ **Proposition 1.** Let  $X$  be a set. Then the intersection of an arbitrary non-empty collection of  $\sigma$ -algebras on  $X$  is a  $\sigma$ -algebra on  $X$ .

*Proof.* Consider a collection  $\mathcal{C}$  of  $\sigma$ -algebras on  $X$ . Denote  $\mathcal{A} = \bigcap \mathcal{C}$  as intersection of all  $\sigma$ -algebras in  $\mathcal{C}$ . We can now easily see that any subset in  $\mathcal{A}$  would be present in every  $\sigma$ -algebra present in collection  $\mathcal{C}$ , hence, it would obey all properties of a  $\sigma$ -algebra. Therefore,  $\mathcal{A}$  is a  $\sigma$ -algebra. ■

→ **Corollary 1.** Let  $X$  be a set and let  $\mathcal{F} \subseteq P(X)$  be a family of subsets of  $X$ . Then there exists a smallest  $\sigma$ -algebra on  $X$  that includes  $\mathcal{F}$ .

*Proof.* Consider any given family  $\mathcal{F} \subseteq P(X)$  and just take intersection of the family  $\mathcal{C}$  of all  $\sigma$ -algebras which contains  $\mathcal{F}$  to construct this smallest  $\sigma$ -algebra. ■

★ **Definition 3. (Generated  $\sigma$ -algebra)** The smallest  $\sigma$ -algebra on  $X$  containing a given family  $\mathcal{F} \subseteq P(X)$  of subsets is called the  $\sigma$ -algebra generated by  $\mathcal{F}$ , denoted as  $\sigma(\mathcal{F})$ .

★ **Definition 4. (Borel  $\sigma$ -algebra on  $\mathbb{R}^d$ )** It is the  $\sigma$ -algebra on  $\mathbb{R}^d$  generated by the collection of all open subsets of  $\mathbb{R}^d$ , denoted as  $\mathcal{B}(\mathbb{R}^d)$ .

★ **Definition 5. (Borel Subsets of  $\mathbb{R}^d$ )** Any  $A \subseteq \mathbb{R}^d$  is called a Borel subset of  $\mathbb{R}^d$  if  $A \in \mathcal{B}(\mathbb{R}^d)$ .

✓ **Proposition 2.** The Borel  $\sigma$ -algebra on  $\mathbb{R}$ ,  $\mathcal{B}(\mathbb{R})$ , of Borel subsets of  $\mathbb{R}$  is generated by each of the following collection of sets:

1. The collection of all closed subsets of  $\mathbb{R}$ .
2. The collection of all subintervals of  $\mathbb{R}$  of the form  $(-\infty, b]$ .
3. The collection of all subintervals of  $\mathbb{R}$  of the form  $(a, b]$ .

*Proof.* To show all of these, consider the three  $\sigma$ -algebras  $\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3$  corresponding to conditions 1, 2 & 3 respectively and try to prove  $\mathcal{A}_3 \subseteq \mathcal{A}_2 \subseteq \mathcal{A}_1 \subseteq \mathcal{B}(\mathbb{R})$  together with  $\mathcal{B}(\mathbb{R}) \subseteq \mathcal{A}_3$ . The first three inclusions are trivial to see. For the case that  $\mathcal{B}(\mathbb{R}) \subseteq \mathcal{A}_3$ , simply note that any open subset can be made by unions of the sets of form  $(a, b]$  and by Homework-I,1, each open set is union of open subsets. ■

✓ **Proposition 3.** The  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R}^d)$  of Borel subsets of  $\mathbb{R}^d$  is generated by each of the following collections:

1. The collection of all closed subsets of  $\mathbb{R}^d$ .
2. The collection of all closed half-spaces in  $\mathbb{R}^d$  that have the form  $\{(x_1, \dots, x_d) \mid x_i \leq b\}$  for some index  $i$  and some  $b \in \mathbb{R}$ .
3. The collection of all rectangles in  $\mathbb{R}^d$  that have the form

$$\{(x_1, \dots, x_d) \mid a_i < x_i \leq b_i \text{ for } i = 1, \dots, d\}$$

*Proof.* Almost the same as in Proposition 2.  $\mathcal{A}_1 \subseteq \mathcal{B}(\mathbb{R}^d)$  trivially by definition.  $\mathcal{A}_2 \subseteq \mathcal{A}_1$  as  $\{(x_1, \dots, x_d) \mid x_i \leq b\}$  is closed itself.  $\mathcal{A}_3 \subseteq \mathcal{A}_2$  by the observation that  $\{(x_1, \dots, x_d) \mid a_i < x_i \leq b_i\}$  is made by the difference of two subsets of the form  $\{(x_1, \dots, x_d) \mid x_i \leq b_i\}$  and  $\{(x_1, \dots, x_d) \mid x_i > a_i\}$ , the latter is the complement of a certain subset in  $\mathcal{A}_2$ , moreover,  $\{(x_1, \dots, x_d) \mid a_i < x_i \leq b_i \text{ for } i = 1, \dots, d\}$  is then constructed by intersection of  $d$  such subsets. Finally,  $\mathcal{B}(\mathbb{R}^d) \subseteq \mathcal{A}_3$  can be seen via the fact that open subsets in  $\mathbb{R}^d$  are made by union of rectangles of type 3 and as such, they are called open subsets. ■

*Conditions for an algebra to become a  $\sigma$ -algebra.*

✓ **Proposition 4.** Let  $X$  be a set and let  $\mathcal{A}$  be an algebra on  $X$ . Then,  $\mathcal{A}$  is a  $\sigma$ -algebra on  $X$  if **either**

- $\mathcal{A}$  is closed under the formation of **unions** of *increasing* sequence of sets, or,
- $\mathcal{A}$  is closed under the formation of **intersections** of *decreasing* sequence of sets.

*Proof.* Take any countably infinite collection of subsets  $A_1, A_2, \dots \in \mathcal{A}$  where  $\mathcal{A}$  is an algebra. Due to the definition of an algebra, we have that  $C_n = \bigcup_{i=1}^n A_i \in \mathcal{A}$  for any  $n \geq 1 \in \mathbb{Z}_+$ . Now note that  $C_1 \subseteq C_2 \subseteq \dots$ , that is, the sequence  $\{C_n\}$  forms an increasing sequence of sets. Hence, by the requirement of the question, we have that  $\bigcup_{i=1}^{\infty} C_i \in \mathcal{A}$ . But then we also have that  $\bigcup_{i=1}^{\infty} A_i \subseteq \bigcup_{i=1}^{\infty} C_i \in \mathcal{A}$ . Hence we have the required condition for part 1. For part 2, we can see that  $C_1^c \supseteq C_2^c \supseteq \dots$  is a decreasing sequence of sets. Then we must have, by the requirement of the question, that  $\bigcap_{i=1}^{\infty} C_i^c = (\bigcup_{i=1}^{\infty} C_i)^c \in \mathcal{A}$ . But then by definition of algebra, we must have  $\bigcup_{i=1}^{\infty} C_i \in \mathcal{A}$ , which already contains the countably infinite union  $\bigcup_{i=1}^{\infty} A_i$ . ■



## 2.2 Measures

★ **Definition 6. (Countably Additive Function)** Let  $X$  be a set and  $\mathcal{A}$  be a  $\sigma$ -algebra on  $X$ . Function  $\mu : \mathcal{A} \rightarrow [0, +\infty]$  is said to be *countably additive* if it satisfies:

$$\mu \left( \bigcup_{i=1}^{\infty} A_i \right) = \sum_{i=1}^{\infty} \mu(A_i)$$

for **each** infinite sequence  $\{A_i\}$  of **disjoint** sets in  $\mathcal{A}$ .

★ **Definition 7. (Measure)** A *measure* on  $\mathcal{A}$  is a function  $\mu : \mathcal{A} \rightarrow [0, +\infty]$  that is **countably additive** and satisfies:

$$\mu(\Phi) = 0.$$

**Remark.** This is sometimes also referred as *countably additive measure* on  $\mathcal{A}$ .

★ **Definition 8.** We have following definitions to compactly represent above definitions:

1. (**Measure Space**) If  $X$  is a set,  $\mathcal{A}$  is a  $\sigma$ -algebra on  $X$  and if  $\mu$  is a measure on  $\mathcal{A}$ , then the triple  $(X, \mathcal{A}, \mu)$  is called a *measure space*.
2. (**Measurable Space**) If  $X$  is a set and  $\mathcal{A}$  is a  $\sigma$ -algebra on  $X$ , then the pair  $(X, \mathcal{A})$  is called a *measurable space*.

✓ **Proposition 5.** Let  $(X, \mathcal{A}, \mu)$  be a measure space and let  $A, B \in \mathcal{A}$  such that  $A \subseteq B$ . Then,

- We have  $\mu(A) \leq \mu(B)$ .
- Additionally, if  $A$  satisfies that  $\mu(A) < +\infty$ , then:

$$\mu(B - A) = \mu(B) - \mu(A).$$

*Proof.* Note that  $A$  and  $B \cap A^c$  are disjoint sets in the sigma algebra  $\mathcal{A}$ . Hence we can write, by countably additive property of  $\mu$ , that:

$$\begin{aligned} \mu(A \cup (B \cap A^c)) &= \mu(B) \\ &= \mu(A) + \mu(B \cap A^c) \end{aligned}$$

Since  $\mu(B \cap A^c) \geq 0$ , hence  $\mu(A) \leq \mu(B)$ . Moreover, if  $\mu(A) < \infty$ , then we can additionally write  $\mu(B \cap A^c) = \mu(B) - \mu(A)$ . ■

★ **Definition 9.** Let  $\mu$  be a measure on a measurable space  $(X, \mathcal{A})$ . Then,

- (**Finite Measure**) If  $\mu(X) < +\infty$ .
- ( **$\sigma$ -Finite Measure**) If  $X = \bigcup_i A_i$  where  $A_i \in \mathcal{A}$  such that  $\mu(A_i) < +\infty$  for all  $i \in \mathbb{N}$ .

**Remark.** In other words, a subset  $A \in \mathcal{A}$  is  *$\sigma$ -finite* if it is a union of a countable sequence of sets that are in  $\mathcal{A}$  and are of finite measure under  $\mu$ .

### 2.2.1 Elementary Properties of Measures

✓ **Proposition 6.** Let  $(X, \mathcal{A}, \mu)$  be a measure space. If  $\{A_k\}$  is an **arbitrary** sequence of sets that belong to  $\mathcal{A}$ , then,

$$\mu \left( \bigcup_{k=1}^{\infty} A_k \right) \leq \sum_{k=1}^{\infty} \mu(A_k).$$

*Proof.* Denote  $B_1 = A_1$  and  $B_i = A_i \cap \left( \bigcup_{k=1}^{i-1} A_k \right)^c$ . Note that  $B_i$  and  $B_j$  are disjoint for distinct  $i$  and  $j$ . Since  $\{A_k\} \in \mathcal{A}$ , therefore  $\{B_i\} \in \mathcal{A}$ . Moreover,  $\bigcup_{i=1}^{\infty} B_i = \bigcup_{k=1}^{\infty} A_k$  by construction. We then get,

$$\begin{aligned} \mu \left( \bigcup_{k=1}^{\infty} A_k \right) &= \mu \left( \bigcup_{i=1}^{\infty} B_i \right) = \sum_{i=1}^{\infty} \mu(B_i) \\ &\leq \sum_{i=1}^{\infty} \mu(A_i) \quad (\because B_i \subseteq A_i \text{ by construction.}) \end{aligned}$$

Hence proved. ■

✓ **Proposition 7.** Let  $(X, \mathcal{A}, \mu)$  be a measure space.

- If  $\{A_k\}$  is an **increasing** sequence of sets in  $\mathcal{A}$ , then

$$\mu \left( \bigcup_k A_k \right) = \lim_k \mu(A_k).$$

- If  $\{A_k\}$  is a **decreasing** sequence of sets in  $\mathcal{A}$  and if  $\mu(A_n) < +\infty$  holds for some  $n$ , then

$$\mu \left( \bigcap_k A_k \right) = \lim_k \mu(A_k).$$

*Proof.* Consider  $\{A_i\} \in \mathcal{A}$  be an increasing sequence of subsets of  $X$ . Define  $B_1 = A_1$  and  $B_i = A_i - A_{i-1} = A_i \cap A_{i-1}^c$ . Clearly,  $B_i$  are disjoint for distinct  $i$ . Hence, we can now write,

$$\mu \left( \bigcup_i B_i \right) = \sum_{i=1}^{\infty} \mu(B_i).$$

Moreover,  $\bigcup_i B_i = \bigcup_k A_k$  and  $B_i = \bigcup_{j=1}^i A_j$ , therefore we can modify the above as

$$\mu \left( \bigcup_k A_k \right) = \lim_{k \rightarrow \infty} \sum_{i=1}^k \mu \left( \bigcup_{j=1}^i A_j \right) = \lim_{k \rightarrow \infty} \mu \left( \bigcup_{i=1}^k \bigcup_{j=1}^i A_j \right) = \lim_{k \rightarrow \infty} \mu(A_k)$$

Now, consider  $\{A_k\}$  to be a decreasing sequence of subsets in  $\mathcal{A}$ . Take  $n \in \mathbb{N}$  such that  $\mu(A_n) < +\infty$ . Denote  $B_i = A_n - A_{n+i}$ . We have each  $B_i$  to be disjoint by construction for any two  $i$ 's and  $\{B_i\}$  is increasing. Moreover,  $\bigcup_k B_k = A_n - \bigcap_i A_{n+i}$ . Hence by first part, we have:

$$\begin{aligned} \mu \left( \bigcup_k B_k \right) &= \mu \left( A_n - \bigcap_i A_{n+i} \right) \\ &= \mu(A_n) - \mu \left( \bigcap_i A_{n+i} \right) \quad (\text{Since } \{A_k\} \text{ is decreasing}) \\ &= \lim_{k \rightarrow \infty} \mu(B_k) = \lim_{k \rightarrow \infty} \mu(A_n - A_{n+k}) \quad (\text{Part 1.}) \\ &= \mu(A_n) - \lim_{k \rightarrow \infty} \mu(A_{n+k}) \end{aligned}$$

Hence, we get the second result (from 2<sup>nd</sup> and 4<sup>th</sup> equality). ■

---

The following partial converse of Proposition 7 helps in checking whether finitely additive measure is countably additive.

✓ **Proposition 8.** Let  $(X, \mathcal{A})$  be a measurable space and let  $\mu$  be a **finitely additive** measure on  $(X, \mathcal{A})$ . Then  $\mu$  is a countably additive measure if **either**

- The following holds for each **increasing sequence**  $\{A_k\}$  of sets that belong to  $\mathcal{A}$ :

$$\lim_k \mu(A_k) = \mu \left( \bigcup_k A_k \right)$$

or,

- The following holds for each **decreasing sequence**  $\{A_k\}$  that belongs to  $\mathcal{A}$  which **satisfies**  $\bigcap_k A_k = \Phi$ :

$$\lim_k \mu(A_k) = 0.$$

*Proof.* Consider any disjoint sequence of sets  $\{A_i\}$  in  $\mathcal{A}$ . Let the first part be true. We can construct an increasing sequence from  $\{A_i\}$  by considering a new sequence  $\{B_i\}$  where  $B_k = \bigcup_{j=1}^k A_j$ . Now, finite additivity of  $\mu$  means that  $\mu(B_k) = \mu \left( \bigcup_{j=1}^k A_j \right) = \sum_{j=1}^k \mu(A_j)$ .

Also, by the condition of part 1, we have  $\lim_{k \rightarrow \infty} \mu(B_k) = \mu(\bigcup_k B_k) = \mu(\bigcup_k A_k)$ . Hence proved part 1.

For part 2, we simply have for any disjoint sequence  $\{A_i\}$  in  $\mathcal{A}$ , a decreasing sequence  $\{B_k\}$  given as  $B_k = \bigcup_{i=k}^{\infty} A_i$ . Again, we can write,

$$\begin{aligned} \mu\left(\bigcup_{i=1}^{\infty} A_i\right) &= \mu\left(\left(\bigcup_{i=1}^k A_i\right) \cup \left(\bigcup_{j=k+1}^{\infty} A_j\right)\right) \\ &= \mu\left(\left(\bigcup_{i=1}^k A_i\right) \cup B_{k+1}\right) \\ &= \mu\left(\bigcup_{i=1}^k A_i\right) + \mu(B_{k+1}) \quad (\text{Since these two are disjoint.}) \\ &= \sum_{i=1}^k \mu(A_i) + \mu(B_{k+1}) \end{aligned}$$

Taking  $\lim_{k \rightarrow \infty}$  both sides then yields us

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i) + 0$$

as  $\lim_{k \rightarrow \infty} \mu(B_k) = 0$  is given in the assumption of part 2. ■

---

## 2.3 Outer Measures

★ **Definition 10. (Outer Measure)** Let  $X$  be a set and let  $\mathcal{P}(X)$  be the collection of all subsets of  $X$ . An *outer measure* on  $X$  is a function  $\mu^* : \mathcal{P}(X) \rightarrow [0, +\infty]$  such that:

- For the empty set  $\Phi$ ,

$$\mu^*(\Phi) = 0$$

- If  $A \subseteq B \subseteq X$ , then

$$\mu^*(A) \leq \mu^*(B).$$

- If  $\{A_n\}$  is an infinite sequence of subsets of  $X$ , then

$$\mu^*\left(\bigcup_n A_n\right) \leq \sum_n \mu^*(A_n)$$

★ **Definition 11. (Lebesgue Outer Measure on  $\mathbb{R}$ )** For each subset  $A \subseteq \mathbb{R}$ , let  $\mathcal{C}_A$  be the set of all infinite sequences  $\{(a_i, b_i)\}$  of bounded open intervals such that  $A \subseteq \bigcup_i (a_i, b_i)$ . That is,

$$\mathcal{C}_A = \{ \{(a_i, b_i)\} \mid A \subseteq \bigcup_i (a_i, b_i) \text{ and } a_i, b_i \in \mathbb{R} \}$$

Then,  $\lambda^* : \mathcal{P}(\mathbb{R}) \rightarrow [0, +\infty]$  is the Lebesgue outer measure, defined by:

$$\lambda^*(A) = \inf \left\{ \sum_i (b_i - a_i) \mid \{(a_i, b_i)\} \in \mathcal{C}_A \right\} \quad (3)$$

To verify that  $\lambda^*$  is indeed an outer measure.

✓ **Proposition 9.** Lebesgue outer measure on  $\mathbb{R}$  is an outer measure and it assigns to each subinterval of  $\mathbb{R}$  its length.

*Proof.* Denote  $\mathcal{C}_A = \{ \{(a_i, b_i)\} \mid A \subseteq \bigcup_i (a_i, b_i) \}$ . To show that  $\lambda^*$  is an outer measure, we first need to show that  $\lambda^*(\Phi) = 0$ . For that, consider the set of all infinite sequences  $\{(a_i, b_i)\} \in \mathcal{C}_\Phi$ , that is (trivially)  $\Phi \subseteq \bigcup_i (a_i, b_i)$ , such that  $\sum_i (b_i - a_i) < \epsilon$  for all  $\epsilon > 0$ . Then, if we denote  $\mathcal{L}_A = \{ \sum_i (b_i - a_i) \mid \{(a_i, b_i)\} \in \mathcal{C}_A \}$ , then  $\inf \mathcal{L}_\Phi = 0$  as for any lower bound  $l$  of  $\mathcal{L}_A$ , if  $l > 0$  then  $\exists \{(a_i, b_i)\} \in \mathcal{C}_\Phi$  such that  $\sum_i (b_i - a_i) < l$ , hence  $l \leq 0$ , or  $\inf \mathcal{L}_\Phi = 0$ .

Second, we need to show that if  $A \subseteq B \subseteq X$ , then  $\lambda^*(A) \leq \lambda^*(B)$ . For this, consider  $A \subseteq B$ . Clearly, we have that  $\mathcal{C}_B \subseteq \mathcal{C}_A$ , therefore  $\mathcal{L}_B \subseteq \mathcal{L}_A$  and hence  $\inf \mathcal{L}_B \geq \inf \mathcal{L}_A$ .

Third, we need to show that for any infinite sequence  $\{A_n\}$  of subsets of  $X$ ,

$$\lambda^*\left(\bigcup_n A_n\right) \leq \sum_n \lambda^*(A_n)$$

For this, consider the Lebesgue outer measure of  $A_n$ , that is,  $\lambda^*(A_n)$ . We must have, that for any infinite sequence  $\{(a_{n,i}, b_{n,i})\} \in \mathcal{C}_{A_n}$ , that

$$\sum_{i=1}^{\infty} (b_{n,i} - a_{n,i}) \geq \lambda^*(A_n).$$

Hence, consider that the difference is upper bounded according to  $n$ , that is the sequence  $\{(a_{n,i}, b_{n,i})\} \in \mathcal{C}_{A_n}$  is such that,

$$\sum_{i=1}^{\infty} (b_{n,i} - a_{n,i}) - \lambda^*(A_n) \leq \epsilon/2^n.$$

Now, we can cover the entire  $\bigcup_i A_i$  by the union of the above intervals, that is,

$$\bigcup_i A_i \subseteq \bigcup_n \bigcup_i (a_{n,i}, b_{n,i}).$$

Now, we know that

$$\lambda^*\left(\bigcup_i A_i\right) = \inf \mathcal{L}_{\bigcup_i A_i}.$$

But since

$$\sum_n \sum_i (b_{n,i} - a_{n,i}) \in \mathcal{L}_{\bigcup_i A_i},$$

and

$$\sum_n \left( \sum_i (b_{n,i} - a_{n,i}) - \lambda^*(A_n) \right) \leq \sum_n \epsilon / 2^n$$

which is equal to

$$\sum_n \sum_i (b_{n,i} - a_{n,i}) - \sum_n \lambda^*(A_n) \leq \epsilon \times 1$$

or,

$$\sum_n \sum_i (b_{n,i} - a_{n,i}) \leq \sum_n \lambda^*(A_n) + \epsilon$$

and since  $\lambda^*(\bigcup_i A_i) = \inf \mathcal{L}_{\bigcup_i A_i}$ , therefore,

$$\lambda^*\left(\bigcup_i A_i\right) \leq \sum_n \sum_i (b_{n,i} - a_{n,i}) \leq \sum_n \lambda^*(A_n)$$

Hence proved.

Now, we need to show that  $\lambda^*$  assigns each subinterval it's length.

For this first show that  $\lambda^*([a, b]) \leq b - a$ . This is easy to show if we take,

$$[a, b] = \bigcup_i (a_i, b_i)$$

where  $(a_1, b_1) = (a, b)$ ,  $(a_i, b_i) = (a - \epsilon/2^i, a)$  for all even  $i$  and  $(a_j, b_j) = (b, b + \epsilon/2^j)$  for all odd  $j$ . Now,

$$\begin{aligned} \sum_i (b_i - a_i) &= (b - a) + \sum_{i=2,4,\dots} \epsilon/2^i + \sum_{i=3,5,\dots} \epsilon/2^i \\ &= b - a + \sum_{i=1,2,\dots} \epsilon/2^i \\ &= b - a + \epsilon \end{aligned}$$

therefore  $\lambda^*([a, b]) = \inf \mathcal{L}_{[a,b]} \leq b - a + \epsilon$  for all  $\epsilon > 0$ , hence  $\lambda^*([a, b]) \leq b - a$ .

Now, to show the converse that  $b - a \leq \lambda^*([a, b])$ , we first note that  $[a, b]$  is compact, so for any infinite cover  $\{(a_i, b_i)\} \in \mathcal{C}_{[a,b]}$ , there exists a finite subcover  $\{(a_i, b_i)\}_{i=1}^n$  of  $[a, b]$ . Now, since  $\lambda^*$  is an outer measure, therefore,

$$b - a \leq \sum_{i=1}^n \lambda^*((a_i, b_i)) \leq \sum_{i=1}^{\infty} \lambda^*((a_i, b_i)) \in \mathcal{L}_{[a,b]}$$

Therefore,  $b - a$  is a lower bound of  $\mathcal{L}_{[a,b]}$  and hence  $b - a \leq \inf \mathcal{L}_{[a,b]} = \lambda^*([a, b])$ .

Hence  $\lambda^*([a, b]) = b - a$ .

Now since, one can construct subintervals of the form  $(a, b]$  or  $[a, b)$  from the following manner:

$$(a, b] \subseteq (a, b) \cup \left( \bigcup_n [b, b + \epsilon/2^n] \right)$$

from which we get that  $\lambda^*((a, b]) \leq b - a$  and also,

$$[a, b] \subseteq (a, b] \cup \left( \bigcup_n [a - \epsilon/2^n, a] \right)$$

which yields  $b - a \leq \lambda^*((a, b])$ . Similarly for  $(-\infty, b]$  to show that  $\lambda^*((-\infty, b]) = +\infty$ . ■

## 2.4 Exercises - II

1. *Proof.* Note that  $\mathbb{Q}$  is dense, that is  $\overline{\mathbb{Q}} = \mathbb{R}$ . Therefore,  $\overline{\mathbb{Q} \cap [0, 1]} = [0, 1]$ . Now, since *closure of union is union of closures*<sup>2</sup>, then we get,

$$\mathbb{Q} \cap [0, 1] \subseteq \overline{\mathbb{Q} \cap [0, 1]} = [0, 1] \subseteq \bigcup_{n=1}^m \overline{(a_n, b_n]} = \bigcup_{n=1}^m [a_n, b_n]$$

Now since  $\lambda^*$  is an outer measure, therefore,

$$\lambda^*([0, 1]) = 1 \leq \lambda^*\left(\bigcup_{n=1}^m [a_n, b_n]\right) \leq \sum_{n=1}^m l(I_n)$$

because  $[0, 1]$  is a subinterval, closure of unions is union of closures and infinite subadditivity implies finite subadditivity. Hence proved. ■

2. *Proof.* Since  $\lambda^*(A) = \inf \mathcal{L}_A$  where

$$\mathcal{L}_A = \left\{ \sum_i (b_i - a_i) \mid \{(a_i, b_i)\} \in \mathcal{C}_A \right\}$$

and

$$\mathcal{C}_A = \{ \{(a_i, b_i)\} \mid A \subseteq \bigcup_n (a_n, b_n) \}.$$

Therefore, for any  $\epsilon > 0$ , and if we write  $\lambda^*(A) + \epsilon$ , then  $\exists \{(a_i, b_i)\} \in \mathcal{C}_A$  such that  $\lambda^*(A) + \epsilon \geq \sum_i (b_i - a_i) \geq \lambda^*(\bigcup_i (a_i, b_i))$ . We can clearly see that  $\bigcup_i (a_i, b_i)$  is union of open intervals and since arbitrary union of open intervals is open, therefore, we have the union of the sequence  $\{(a_i, b_i)\}$  as the needed open interval. ■

3. *Proof.* From the previous question 2, we have that for  $A \subseteq \mathbb{R}$  and for any  $\epsilon > 0$ , there exists an open subset  $O \supseteq A$  such that  $\lambda^*(O) \leq \lambda^*(A) + \epsilon$ . Now, consider the sequence  $\epsilon_n = 1/n$  and the corresponding open subsets  $O_n$  containing  $A$ . Clearly,  $A \subseteq \bigcap_n O_n$ . Since  $G = \bigcap_n O_n$  is countable intersection of open subsets, therefore  $G$  is a  $G_\delta$  set. Now, note that  $G = \bigcap_n O_n \subseteq O_m$  for any  $m \in \mathbb{N}$ . Therefore,  $G \setminus E \subseteq O_m \setminus E$  for any  $m \in \mathbb{N}$ . Hence,

$$\begin{aligned} \lambda^*(G \setminus E) &\leq \lambda^*(O_m \setminus E) \\ &= \lambda^*(O_m) - \lambda^*(E) \quad (\text{Since } E \subseteq O_m) \\ &\leq \frac{1}{n} \end{aligned}$$

for all  $n \in \mathbb{N}$ . Therefore  $\lambda^*(G \setminus E) = 0$  which implies that

$$\lambda^*(G) = \lambda^*(E)$$

as  $E \subseteq G$ . ■

4. *Proof.* Take any  $B \subseteq \mathbb{R}$ . We are given a subset  $A \subseteq \mathbb{R}$  such that  $\lambda^*(A) = 0$ . Note that

$$\lambda^*(B) \leq \lambda^*(A \cup B) \leq \lambda^*(A) + \lambda^*(B) = \lambda^*(B)$$

Hence we already have that  $\lambda^*(B) = \lambda^*(A \cup B)$ . Now to show that  $\lambda^*(A \cup B) = \lambda^*(B - A)$ , first note that  $B - A = B \cap A^c$ . Clearly,  $B \cap A^c \subseteq B$ , therefore,

$$\lambda^*(B \cap A^c) \leq \lambda^*(B) = \lambda^*(A \cup B).$$

Now, to show the converse to complete the proof, we note that  $A \cup B = A \cup (B \cap A^c)$ , hence, by countable sub-additivity, we finally have,

$$\lambda^*(A \cup (B \cap A^c)) = \lambda^*(A \cup B) = \lambda^*(B) \leq \lambda^*(A) + \lambda^*(B \cap A^c) = \lambda^*(B \cap A^c)$$

Hence the proof is complete. ■

5. *Proof.* Take  $A \subseteq \mathbb{R}$  such that it is countable. Now,  $A$  can either be finite or countably infinite. Let's first take  $A$  to be finite, so that  $\exists n \in \mathbb{N}$  such that

$$A = \{a_1, a_2, \dots, a_n\}.$$

Now, consider  $\{(a_i, b_i)\} \in \mathcal{C}_A$  such that it covers each point with an exponentially decreasing interval, as follows:

$$\{I_j\} = \left\{ (a_1 - \frac{\epsilon}{2}, a_1 + \frac{\epsilon}{2}), \dots, (a_n - \frac{\epsilon}{2}, a_n + \frac{\epsilon}{2}), (a_1 - \frac{\epsilon}{2^2}, a_1 + \frac{\epsilon}{2^2}), \dots \right\}$$

<sup>2</sup>Proof done in diary : 3<sup>rd</sup> March, 2018 - The date of reference.

In other words,

$$\{(a_i, b_i)\} = \left\{ \left( a_{i^{9\%n}+1} - \frac{\epsilon}{2^{\lceil i^{9\%n} \rceil}}, a_{i^{9\%n}+1} + \frac{\epsilon}{2^{\lceil i^{9\%n} \rceil}} \right) \right\}$$

We clearly have that this sequence  $\{(a_i, b_i)\} \in \mathcal{C}_A$  and since  $\lambda^*(A) = \inf \mathcal{L}_A$ , therefore we must have that

$$\lambda^*(A) \leq \sum_i (b_i - a_i) = \sum_i \frac{2\epsilon}{2^{\lceil i^{9\%n} \rceil}} = 4\epsilon = \epsilon'$$

Now since  $\lambda^* : \mathcal{P}(\mathbb{R}) \rightarrow [0, +\infty]$ , therefore,  $\lambda^*(A) = 0$  as  $\epsilon > 0$  is arbitrary. So far, we have proved that finite set has outer measure zero. For countably infinite set, we would have  $A = \{a_1, a_2, \dots\}$ . On this, consider the cover  $\{(a_i, b_i)\} = \left\{ \left( a_i - \frac{\epsilon}{2^{i+1}}, a_i + \frac{\epsilon}{2^{i+1}} \right) \right\}$ . Clearly, this sequence  $\{(a_i, b_i)\}$  is in  $\mathcal{C}_A$ . Moreover,

$$\lambda^*(A) \leq \sum_i (b_i - a_i) = \sum_i \frac{\epsilon}{2^i} = \epsilon$$

Now since  $\epsilon > 0$  is arbitrary, therefore,  $\lambda^*(A) = 0$  even for countably infinite set  $A$ .

Now since  $\lambda^*(\mathbb{R}) \neq 0$ , therefore  $\mathbb{R}$  is not countable. ■

6. *Proof.* Take any open interval  $(a, b)$ . We can see that

$$(a, b) \subseteq (a, b]$$

and

$$(a, b] \subseteq (a, b) \cup \bigcup_n (b, b + \epsilon/2^n]$$

Therefore,

$$\begin{aligned} \lambda^*((a, b)) &\leq \lambda^*((a, b]) \\ &= b - a. \end{aligned}$$

and

$$\begin{aligned} \lambda^*((a, b]) &= b - a \leq \lambda^* \left( (a, b) \cup \bigcup_n (b, b + \epsilon/2^n] \right) \\ &\leq \lambda^*((a, b)) + \sum_n (b - (b + \epsilon/2^n)) \\ &= \lambda^*((a, b)) + \sum_n (b - b - \epsilon/2^n) \\ &= \lambda^*((a, b)) + \sum_n (-\epsilon/2^n) \\ &= \lambda^*((a, b)). \end{aligned}$$

Hence,  $\lambda^*((a, b)) = \lambda^*((a, b]) = b - a$ . Hence, one can take interval  $\{(a_n, b_n)\}$  as the sequence of intervals in the definition of Lebesgue Outer Measure as well, instead of  $\{(a_n, b_n]\}$ . ■

7. *Proof.* Consider  $\mathcal{C}_A = \{\{(a_i, b_i)\} \mid A \subseteq \bigcup_i (a_i, b_i)\}$  and  $\mathcal{C}_{\alpha A} = \{\{(c_i, d_i)\} \mid \alpha A \subseteq \bigcup_i (c_i, d_i)\}$ . We will show that

$$\text{To Show : } \mathcal{C}_{\alpha A} = \alpha \mathcal{C}_A = \left\{ \{\alpha(a_i, b_i)\} \mid A \subseteq \bigcup_i (a_i, b_i) \right\}.$$

First, take any infinite sequence  $\{(a_i, b_i)\} \in \mathcal{C}_A$ . Then  $\{\alpha(a_i, b_i)\} \in \mathcal{C}_{\alpha A}$  because if  $A \subseteq \bigcup_i (a_i, b_i)$ , then  $\alpha A \subseteq \bigcup_i \alpha(a_i, b_i)$  where  $\alpha(a_i, b_i) = (\alpha a_i, \alpha b_i)$  if  $\alpha > 0$  or  $(\alpha b_i, \alpha a_i)$  if  $\alpha < 0$ . Hence,

$$\alpha \mathcal{C}_A \subseteq \mathcal{C}_{\alpha A}.$$

To see the converse, first note that the case of  $\alpha = 0$  is trivial, so we would take  $\alpha \neq 0$ . Now, take  $\{(c_i, d_i)\} \in \mathcal{C}_{\alpha A}$ . Hence  $\alpha A \subseteq \bigcup_i (c_i, d_i)$ . Now,  $\frac{1}{\alpha} \cdot \alpha A = A \subseteq \bigcup_i \frac{1}{\alpha} (c_i, d_i)$ . Therefore,  $\{\frac{1}{\alpha} (c_i, d_i)\} \in \mathcal{C}_A \implies \{(c_i, d_i)\} \in \alpha \mathcal{C}_A$ . Hence  $\mathcal{C}_{\alpha A} \subseteq \alpha \mathcal{C}_A$ . Hence  $\mathcal{C}_{\alpha A} = \alpha \mathcal{C}_A$ . Hence, we get that  $\mathcal{L}_{\alpha A} = |\alpha| \mathcal{L}_A$  because  $\sum_i l(\alpha I_n) = \sum_i \alpha(b_i - a_i)$  if  $\alpha > 0$  or  $\sum_i \alpha(a_i - b_i)$  if  $\alpha < 0$ . Hence we can write  $\alpha \mathcal{L}_A = \{\sum_i |\alpha| (b_i - a_i) \mid \{(a_i, b_i)\} \in \mathcal{C}_A\}$ . Now,  $\inf \mathcal{L}_{\alpha A} = \inf |\alpha| \mathcal{L}_A \implies \lambda^*(\alpha A) = |\alpha| \lambda^*(A)$ . ■

8. *Proof.*<sup>3</sup> We have that  $B$  is the set of all points of  $\mathbb{R}$  which are present in infinite number of subsets  $A_n$ . Now consider  $x \in B$ . Then, there exists  $\{n\}$  such that the first element of the sequence  $n_1$  is such that  $x \in A_{n_1}$ , or  $x \in \bigcup_{n \geq n_1} A_n$ . Then, for any  $x \in B$ , we have  $x \in \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n$ . Hence,

$$B \subseteq \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n.$$

Now, take  $x \in \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n$ . Then  $x \in \bigcup_{n \geq m} A_n$  for all  $m \in \mathbb{N}$ . This implies that  $x \in A_{k_m}$  for some  $k_m \geq m$  for all  $m$ . This implies that  $x \in A_n$  for infinitely many  $n$ . Hence  $x \in B$ , so that,

$$B = \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n.$$

We now show the following:

$$\text{To Show : } \left( \bigcup_{n \geq m_0} A_n \right) \setminus \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) = \bigcup_{n \geq m_0} A_n. \quad (4)$$

for any  $m_0 \in \mathbb{N}$ . To show this, we first note that the following is easily deduced:

$$\left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \subseteq \bigcup_n A_n$$

because for any  $x \in \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right)$ ,  $\exists k \in \mathbb{N}$  such that  $x \in \bigcup_{n \geq k} A_n \subseteq \bigcup_n A_n$ . Now, note the following:

$$\begin{aligned} \bigcup_{n=1}^{m-1} A_n &\subseteq \left( \bigcup_{n \geq m} A_n \right)^c \\ \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{m-1} A_n &\subseteq \bigcup_{m=1}^{\infty} \left( \bigcup_{n \geq m} A_n \right)^c \\ \bigcup_n A_n &\subseteq \bigcup_{m=1}^{\infty} \left( \bigcup_{n \geq m} A_n \right)^c \end{aligned} \quad (5)$$

Therefore, if we now expand Eq. 4, we get the following:

$$\begin{aligned} \left( \bigcup_{n \geq m_0} A_n \right) \setminus \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) &= \bigcup_{n \geq m_0} A_n \cap \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right)^c \\ &= \bigcup_{n \geq m_0} A_n \cap \left( \bigcup_{m=1}^{\infty} \bigcap_{n \geq m} A_n^c \right)^c \\ &= \bigcup_{n \geq m_0} A_n \cap \left( \bigcup_{m=1}^{\infty} \left( \bigcup_{n \geq m} A_n \right)^c \right) \\ &= \bigcup_{n \geq m_0} A_n \quad (\text{Equation 5}) \end{aligned} \quad (6)$$

Therefore,

$$\begin{aligned} \lambda^* \left( \left( \bigcup_{n \geq m_0} A_n \right) \setminus \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \right) &= \lambda^* \left( \bigcup_{n \geq m_0} A_n \right) - \lambda^* \left( \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \right) \\ \lambda^* \left( \left( \bigcup_{n \geq m_0} A_n \right) \setminus \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \right) - \lambda^* \left( \bigcup_{n \geq m_0} A_n \right) &= \lambda^* \left( \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \right) \\ 0 &= \lambda^* \left( \left( \bigcap_{m=1}^{\infty} \bigcup_{n \geq m} A_n \right) \right) = \lambda^*(B) \quad (\text{From Eq. 6}) \end{aligned}$$

Hence, proved. ■

---

<sup>3</sup>This is the Borel-Cantelli Lemma(!)



9. *Proof.* **This proof is wrong, write the correct solution.** We divide the proof as follows:

**Act 1.**  $A$  and  $B$  are disjoint.

Take two subsets  $A, B \subseteq \mathbb{R}$  such that  $d(A, B) > 0$ . Now, take any arbitrary pair  $a, b$  such that  $a \in A$  and  $b \in B$ . Due to the fact that  $d(A, B) = \inf\{|x - y| \mid x \in A, y \in B\}$ , therefore we have

$$\begin{aligned} |a - b| &\geq \inf\{|x - y| \mid x \in X, y \in Y\} \\ &> 0 \quad (\text{As per question}). \end{aligned}$$

Therefore, for any arbitrary  $a \in A$  and  $b \in B$ , we have that  $|a - b| > 0$ . This implies that either  $a < b$  or  $a > b$ , but NOT  $a = b$  for any pair  $a, b$ . This further implies that for any  $a \in A$  there does not exist  $b \in B$  such that  $a = b$ . Hence  $A \cap B = \Phi$ .

**Act 2.**  $\lambda^*$  for disjoint sets is countably additive.

Now we show that for any subsets  $A, B \subseteq \mathbb{R}$  satisfying  $A \cap B = \Phi$ , we have

$$\lambda^*(A \cup B) = \lambda^*(A) + \lambda^*(B).$$

For this, we first trivially have due to countable sub-additivity of  $\lambda^*$  that  $\lambda^*(A \cup B) \leq \lambda^*(A) + \lambda^*(B)$ . All that remains to be shown is

$$\lambda^*(A \cup B) \geq \lambda^*(A) + \lambda^*(B).$$

For this, first consider the set relation which is trivial to show

$$A \cup B \supseteq A \setminus (A \cap B) \cup B \setminus (A \cap B) \cup (A \cap B).$$

as for any  $x \in A \setminus (A \cap B) \cup B \setminus (A \cap B) \cup (A \cap B)$ , either  $x \in A$  or  $x \in B$ . Now, due to the fact that  $A \cap B = \Phi$ , we get the following conclusion:

$$\begin{aligned} B \setminus (A \cap B) &= B \cap (A \cap B)^c \\ &= B \cap (A^c \cup B^c) \\ &= (B \cap A^c) \cup (B \cap B^c) \\ &= B \cup \Phi \quad (B \subseteq A^c) \\ &= B \end{aligned}$$

Similarly,  $A \setminus (A \cap B) = A$ . Combining both of these results, we get,

$$\begin{aligned} \lambda^*(A \cup B) &= \lambda^*(A \setminus (A \cap B) \cup B \setminus (A \cap B) \cup (A \cap B)) \\ &\leq \lambda^*(A) + \lambda^*(B) + \lambda^*(\Phi) \\ &= \lambda^*(A) + \lambda^*(B) \quad (\lambda^*(\Phi) = 0) \end{aligned}$$

Hence we have that  $\lambda^*(A \cup B) = \lambda^*(A) + \lambda^*(B)$  for disjoint sets  $A$  and  $B$  of  $\mathbb{R}$ . Therefore, for the sets  $A$  and  $B$  satisfying the question's criterion, we have the required result. ■

## 2.5 Lebesgue Measurability.

★ **Definition 12.** ( $\mu^*$ -measurable subset) Let  $X$  be a set and let  $\mu^*$  be an outer measure on  $X$ . A subset  $B \subseteq X$  is  $\mu^*$ -measurable if:

$$\mu^*(A) = \mu^*(A \cap B) + \mu^*(A \cap B^c)$$

holds for all subsets  $A \subseteq X$ .

★ **Definition 13.** (Lebesgue Measurable subset of  $\mathbb{R}$ ) A subset  $B \subseteq \mathbb{R}$  is called a Lebesgue measurable subset of  $\mathbb{R}$  if  $B$  is  $\lambda^*$ -measurable. That is, for any  $A \subseteq \mathbb{R}$ , we must have:

$$\lambda^*(A) = \lambda^*(A \cap B) + \lambda^*(A \cap B^c)$$

**Remark.** Important to note are the following:

- Due to sub-additivity of  $\mu^*$  and  $A \subseteq (A \cap B) \cup (A \cap B^c)$ , we already have that

$$\mu^*(A) \leq \mu^*(A \cap B) + \mu^*(A \cap B^c)$$

for any subsets  $A, B \subseteq X$ .

- ★ Due to the above fact, all that remains to be shown to ascertain that  $B \subseteq \mathbb{R}$  is  $\mu^*$ -measurable is to show the following converse:

$$\mu^*(A) \geq \mu^*(A \cap B) + \mu^*(A \cap B^c).$$

for all  $A \subseteq X$ .

✓ **Proposition 10.** Let  $X$  be a set and let  $\mu^*$  be an outer measure on  $X$ . Then each subset  $B \subseteq X$  that satisfies  $\mu^*(B) = 0$  or that satisfies  $\mu^*(B^c) = 0$  is  $\mu^*$ -measurable.

*Proof.* This result actually proves that for subset  $B \subseteq X$  which has zero outer measure under  $\mu^*$ , any other subset  $A \subseteq X$  would be such that  $\mu^*(A \cap B) = 0$  (!) After proving this, and from the remark above, we would just be left to show that if  $\mu^*(B) = 0$ , then  $\mu^*(A) \geq \mu^*(A \cap B) + \mu^*(A \cap B^c)$ . We show the former here, from which the latter follows naturally.

Consider  $B \subseteq X$  such that  $\mu^*(B) = 0$ . It's true that  $A \cap B \subseteq B$ . Now since  $\mu^*$  is an outer measure on  $X$ , therefore, we must have  $\mu^*(A \cap B) \leq \mu^*(B) = 0$ . This implies that  $\mu^*(A \cap B) = 0$ . Now, we would see that the required condition follows naturally from the previous. First, note the following:

$$A \cap B \subseteq A \text{ and } A \cap B^c \subseteq A.$$

Hence, we can write:

$$\mu^*(A \cap B) \leq \mu^*(A) \text{ and } \mu^*(A \cap B^c) \leq \mu^*(A).$$

Now if  $\mu^*(B) = 0$ , then  $\mu^*(A \cap B) = 0$  and then in the second inequality, we would have:

$$\mu^*(A \cap B^c) + \mu^*(A \cap B) \leq \mu^*(A) + 0$$

Or, if  $\mu^*(B^c) = 0$ , then  $\mu^*(A \cap B^c) = 0$  and then in the first inequality, we would have:

$$\mu^*(A \cap B) + \mu^*(A \cap B^c) \leq \mu^*(A) + 0.$$

Hence,  $B$  is  $\mu^*$ -measurable for any  $B \subseteq X$  which satisfies that either  $\mu^*(B) = 0$  or  $\mu^*(B^c) = 0$ . ■

*The following theorem is a fundamental fact about outer measures.*

**Theorem 1.** Let  $X$  be a set, let  $\mu^*$  be an outer measure on  $X$  and let  $\mathcal{M}_{\mu^*}$  be the collection of all  $\mu^*$ -measurable subsets of  $X$ . Then,

- $\mathcal{M}_{\mu^*}$  is a  $\sigma$ -algebra.
- The restriction of  $\mu^*$  to  $\mathcal{M}_{\mu^*}$  is a measure on  $\mathcal{M}_{\mu^*}$ .

*Proof.* **Act 1.**  $\mathcal{M}_{\mu^*}$  is an algebra.

First, it is clear that  $X, \Phi \in \mathcal{M}_{\mu^*}$  from Proposition 10, because  $\mu^*(\Phi) = \mu^*(X^c) = 0$ . Now, if  $B \in \mathcal{M}_{\mu^*}$ , then  $\mu^*(A) = \mu^*(A \cap B) + \mu^*(A \cap B^c) \forall A \subseteq X$ . But if we replace  $B$  by  $B^c$  in the above, we would get the same equation, hence  $B^c \in \mathcal{M}_{\mu^*}$ . So  $\mathcal{M}_{\mu^*}$  is closed under complements. Now, to show closed nature under finite unions, we take any two subsets  $B_1, B_2 \in \mathcal{M}_{\mu^*}$  and show that  $A \cup B \in \mathcal{M}_{\mu^*}$ . First we have

$$\begin{aligned}\mu^*(A) &= \mu^*(A \cap B_1) + \mu^*(A \cap B_1^c) \\ &= \mu^*(A \cap B_2) + \mu^*(A \cap B_2^c)\end{aligned}$$

for any  $A \subseteq X$ . Now, we see that from the fact that  $B_1 \in \mathcal{M}_{\mu^*}$ ,

$$\begin{aligned}\mu^*(A \cap (B_1 \cup B_2)) &= \mu^*(A \cap (B_1 \cup B_2) \cap B_1) + \mu^*(A \cap (B_1 \cup B_2) \cap B_1^c) \\ &= \mu^*(A \cap B_1) + \mu^*(A \cap B_2 \cap B_1^c)\end{aligned}$$

Similarly, we have from the fact  $B_2 \in \mathcal{M}_{\mu^*}$ ,

$$\begin{aligned}\mu^*(A \cap (B_1 \cup B_2)^c) &= \mu^*(A \cap (B_1 \cup B_2)^c \cap B_2) + \mu^*(A \cap (B_1 \cup B_2)^c \cap B_2^c) \\ &= \mu^*(A \cap B_1^c \cap B_2^c \cap B_2) + \mu^*(A \cap B_1^c \cap B_2^c \cap B_2^c) \\ &= \mu^*(\Phi) + \mu^*(A \cap B_1^c \cap B_2^c) \\ &= \mu^*(A \cap (B_1 \cup B_2)^c)\end{aligned}$$

Now, adding the above results yield,

$$\begin{aligned}\mu^*(A \cap (B_1 \cup B_2)^c) + \mu^*(A \cap (B_1 \cup B_2)) &= \mu^*(A \cap (B_1 \cup B_2)^c) + \mu^*(A \cap B_1) + \mu^*(A \cap B_2 \cap B_1^c) \\ &= \mu^*(A \cap B_1^c \cap B_2^c) + \mu^*(A \cap B_1^c \cap B_2) + \mu^*(A \cap B_1) \\ &= \mu^*(A \cap B_1^c) + \mu^*(A \cap B_1) \\ &= \mu^*(A).\end{aligned}$$

Hence,  $B_1 \cup B_2$  is  $\mu^*$ -measurable, so  $B_1 \cup B_2 \in \mathcal{M}_{\mu^*}$ . Now, we can, for a finite collection of subsets in  $\mathcal{M}_{\mu^*}$ , we can proceed like above, to show that  $\mathcal{M}_{\mu^*}$  is closed under finite union, hence showing that  $\mathcal{M}_{\mu^*}$  is an algebra.

**Act 2.**  $\mathcal{M}_{\mu^*}$  is a  $\sigma$ -algebra.

All that is left to show that  $\mathcal{M}_{\mu^*}$  is a  $\sigma$ -algebra is to show that it is closed under countable union. We have already proved closed nature under finite union. We extend it via induction principle. Suppose  $\{B_i\}$  is a sequence of disjoint subsets in  $\mathcal{M}_{\mu^*}$ . For this, we first prove<sup>4</sup> using induction that, for all  $A \subseteq X$  and  $n \in \mathbb{N}$ ,

$$\text{To Prove : } \mu^*(A) = \sum_{i=1}^n \mu^*(A \cap B_i) + \mu^*\left(A \cap \left(\bigcap_{i=1}^n B_i^c\right)\right) \quad (7)$$

For the case when  $n = 1$ , we see that it Eq. 7 reduces to  $\mu^*(A) = \mu^*(A \cap B_1) + \mu^*(A \cap B_1^c)$ . But since  $B_i \in \mathcal{M}_{\mu^*} \forall i \in \mathbb{N}$ , therefore this is trivially true. Now, by the induction principle, we assume that Eq. 7 is true uptill  $n$  and then we try to prove it for  $n + 1$  step. For this, since  $B_{n+1} \in \mathcal{M}_{\mu^*}$  is disjoint to all other  $B_i$ 's, we have,

$$\begin{aligned}\mu^*\left(A \cap \bigcap_{i=1}^n B_i^c\right) &= \mu^*\left(\left(A \cap \bigcap_{i=1}^n B_i^c\right) \cap B_{n+1}\right) + \mu^*\left(\left(A \cap \bigcap_{i=1}^n B_i^c\right) \cap B_{n+1}^c\right) \\ &= \mu^*(A \cap B_{n+1}) + \mu^*\left(A \cap \bigcap_{i=1}^{n+1} B_i^c\right)\end{aligned}$$

where the last line follows from the fact that each  $B_i$  is disjoint to other  $B_j$ 's, hence each  $B_j^c$  would contain  $B_i$  and therefore  $B_{n+1} \subseteq \bigcap_{i=1}^n B_i^c$ . Now, substituting the above equation in Eq. 7 gives,

<sup>4</sup>But why to prove Eq. 7? The motivation for Eq. 7 comes from Part 1. More specifically, notice in the equation where we added  $\mu^*(A \cap (B_1 \cup B_2)^c)$  and  $\mu^*(A \cap (B_1 \cup B_2))$ . Note it's 2<sup>nd</sup> line, this is the case when  $n = 2$  in Eq. 7 combined with the fact that  $B_i$ 's are disjoint. Now why to take  $B_i$ 's to be disjoint? The reason for this comes from the fact that for any infinite sequence of subsets  $\{A_i\}$ , one can construct infinite sequence of disjoint subsets, that is :  $A_1, A_2 \cap A_1^c, A_3 \cap (A_1 \cup A_2)^c, \dots$  and it's union is again  $\bigcup_n A_n$ . Hence if we prove that a disjoint infinite sequence is closed under union, then we could prove that any infinite sequence of subsets is closed under union too!

$$\begin{aligned}
\mu^*(A) &= \sum_{i=1}^n \mu^*(A \cap B_i) + \mu^*(A \cap B_{n+1}) + \mu^*\left(A \cap \bigcap_{i=1}^{n+1} B_i^c\right) \\
&= \sum_{i=1}^{n+1} \mu^*(A \cap B_i) + \mu^*\left(A \cap \bigcap_{i=1}^{n+1} B_i^c\right)
\end{aligned}$$

Hence, by induction principle, Eq. 7 is true for all  $n \in \mathbb{N}$ . Hence, now we can write,

$$\begin{aligned}
\mu^*(A) &\geq \sum_{i=1}^{\infty} \mu^*(A \cap B_i) + \mu^*\left(A \cap \bigcap_{i=1}^{\infty} B_i^c\right) \\
&= \sum_{i=1}^{\infty} \mu^*(A \cap B_i) + \mu^*\left(A \cap \left(\bigcup_{i=1}^{\infty} B_i\right)^c\right)
\end{aligned}$$

Now, to prove that  $\bigcup_i B_i \in \mathcal{M}_{\mu^*}$ , we need to show

$$\text{To Show : } \mu^*(A) \geq \mu^*\left(A \cap \bigcup_i B_i\right) + \mu^*\left(A \cap \left(\bigcup_i B_i\right)^c\right)$$

This comes from previous result as follows:

$$\begin{aligned}
\mu^*(A) &\geq \sum_{i=1}^{\infty} \mu^*(A \cap B_i) + \mu^*\left(A \cap \left(\bigcup_{i=1}^{\infty} B_i\right)^c\right) \\
&\geq \mu^*\left(\bigcup_{i=1}^{\infty} (A \cap B_i)\right) + \mu^*\left(A \cap \left(\bigcup_{i=1}^{\infty} B_i\right)^c\right) \\
&= \mu^*\left(A \cap \bigcup_{i=1}^{\infty} B_i\right) + \mu^*\left(A \cap \left(\bigcup_{i=1}^{\infty} B_i\right)^c\right)
\end{aligned} \tag{8}$$

Therefore,  $\bigcup_i B_i \in \mathcal{M}_{\mu^*}$ . Now, as the previous footnote mentions, for every infinite sequence  $\{C_i\}$  in  $\mathcal{M}_{\mu^*}$ , we have a disjoint sequence of subsets as  $C_1, C_2 \cap C_1^c, C_3 \cap C_2^c \cap C_1^c, \dots$ . Now, this disjoint sequence is closed under union as we just showed and since union of this disjoint sequence is equal to the union of  $\{C_i\}$ , hence  $\bigcup_i C_i \in \mathcal{M}_{\mu^*}$  for any sequence  $\{C_i\}$  in  $\mathcal{M}_{\mu^*}$ . Thus,  $\mathcal{M}_{\mu^*}$  is a  $\sigma$ -algebra.

**Act 3.**  $\mu^*$  restricted to  $\mathcal{M}_{\mu^*}$  is a measure.

Consider  $\{B_n\}$  be an infinite sequence of subsets in  $\mathcal{M}_{\mu^*}$ . Now, by finite subadditivity, we trivially have

$$\mu^*\left(\bigcup_i B_i\right) \leq \sum_i \mu^*(B_i)$$

Moreover, from Part 2 and setting  $A = \bigcup_i B_i$ , we get:

$$\begin{aligned}
\mu^*\left(\bigcup_i B_i\right) &\geq \sum_j \mu^*\left(\bigcup_i B_i \cap B_j\right) + \mu^*\left(\bigcup_i B_i \cap \left(\bigcup_i B_i\right)^c\right) \\
&= \sum_j \mu^*(B_j) + \mu^*(\Phi) \\
&= \sum_j \mu^*(B_j).
\end{aligned}$$

We hence have the complete proof. ■

✓ **Proposition 11.** Every interval of form  $(-\infty, b]$  is Lebesgue measurable.

*Proof.* All we need to show that for  $B = (-\infty, b]$  and for all  $A \subseteq \mathbb{R}$ ,

$$\text{To Show: } \lambda^*(A) \geq \lambda^*(A \cap B) + \lambda^*(A \cap B^c).$$

Consider  $\{(a_i, b_i)\} \in \mathcal{C}_A$ . Therefore  $A \subseteq \bigcup_i (a_i, b_i)$ . Clearly, this implies that

$$\lambda^*(A) \leq \sum_i (b_i - a_i).$$

Equivalently,  $\exists \epsilon > 0$  such that,

$$\lambda^*(A) + \epsilon \geq \sum_i (b_i - a_i).$$

Now, it's also clear that  $(a_k, b_k) \cap B$  and  $(a_k, b_k) \cap B^c$  are disjoint for any  $k \in \mathbb{N}$ . But we know that **for disjoint  $\lambda^*$ -measurable sets  $C$  and  $D$ , it's true that  $\lambda^*(C \cup D) = \lambda^*(C) + \lambda^*(D)$ , which can be extended for countable union**<sup>5</sup>. Hence, we can write:

$$(b_k - a_k) = \lambda^*((a_k, b_k)) = \lambda^*((a_k, b_k) \cap B \cup (a_k, b_k) \cap B^c) = \lambda^*((a_k, b_k) \cap B) + \lambda^*((a_k, b_k) \cap B^c).$$

Now since  $A \subseteq \bigcup_i (a_i, b_i)$ , therefore  $A \cap B \subseteq \bigcup_i (a_i, b_i) \cap B$ . Similarly,  $A \cap B^c \subseteq \bigcup_i (a_i, b_i) \cap B^c$ . This implies that,

$$\begin{aligned} \lambda^*(A \cap B) &\leq \lambda^*\left(\bigcup_i (a_i, b_i) \cap B\right) \\ \lambda^*(A \cap B^c) &\leq \lambda^*\left(\bigcup_i (a_i, b_i) \cap B^c\right) \end{aligned}$$

Adding these equations, we get the desired result:

$$\begin{aligned} \lambda^*(A \cap B) + \lambda^*(A \cap B^c) &\leq \lambda^*\left(\bigcup_i (a_i, b_i) \cap B\right) + \lambda^*\left(\bigcup_i (a_i, b_i) \cap B^c\right) \\ &\leq \sum_i \lambda^*((a_i, b_i) \cap B) + \sum_i \lambda^*((a_i, b_i) \cap B^c) \\ &= \sum_i \lambda^*((a_i, b_i)) \quad (\text{Shown above.}) \\ &= \sum_i (b_i - a_i) \\ &\leq \lambda^*(A) + \epsilon \end{aligned}$$

which is true for all  $\epsilon > 0$ , therefore

$$\lambda^*(A \cap B) + \lambda^*(A \cap B^c) \leq \lambda^*(A).$$

■

✓ **Proposition 12.** Every Borel subset of  $\mathbb{R}$  is Lebesgue Measurable.

*Proof.* From Proposition 11, we have that the subsets of the form  $(-\infty, b]$  are Lebesgue measurable. Remember that a subset  $B \subseteq \mathbb{R}$  is called borel if  $B \in \mathcal{B}(\mathbb{R})$ ; that is, it is present in the smallest  $\sigma$ -algebra generated by open subsets of  $\mathbb{R}$ .

Now, since  $(-\infty, b]$  is Lebesgue measurable, therefore  $(-\infty, b] \in \mathcal{M}_{\lambda^*}$ . But since  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R})$  is closed under complements, complement of open subset is closed and  $(-\infty, b]$  is closed, therefore all intervals of form  $(-\infty, b]$  are also in  $\mathcal{B}(\mathbb{R})$ . Since  $\mathcal{B}(\mathbb{R})$  is the smallest  $\sigma$ -algebra containing intervals of form  $(-\infty, b]$ , therefore  $\mathcal{B}(\mathbb{R}) \subseteq \mathcal{M}_{\lambda^*}$ . ■

<sup>5</sup>We need it's proof too.

★ **Definition 14. (Lebesgue Measure)** The restriction of Lebesgue outer measure on  $\mathbb{R}$  to the collection  $\mathcal{M}_{\lambda}^*$  of Lebesgue measurable subsets of  $\mathbb{R}$  is called *Lebesgue Measure*. It would be denoted by  $\lambda$ . Hence, we would work with the measure space  $(\mathbb{R}, \mathcal{M}_{\lambda}^*, \lambda)$ <sup>6</sup>.

## 2.6 Does $\lambda^*(E) = 0$ implies $E$ is countable?

We would construct today a set which has measure 0, but not countable(!).

1. Take  $E_0 = [0, 1]$ .
2. Remove  $(1/3, 2/3)$  from  $E_0$  to form  $E_1 = [0, 1/3] \cup [2/3, 1]$ .
3. Proceed in the same way to form  $E_2 = [0, 1/9] \cup [2/9, 1/3] \cup [2/3, 7/9] \cup [8/9, 1]$ .
4. At  $n^{th}$  step,  $E_n$  contains  $2^n$  subintervals and each of which is of length  $\frac{1}{3^n}$ .
5. We clearly have  $E_0 \supset E_1 \supset E_2 \supset \dots$ .
6. Here, note that each  $E_n$  is a closed and compact subset of  $\mathbb{R}$ .
7. The set

$$P = \bigcap_{n=0}^{\infty} E_n \text{ is known as } \mathbf{Cantor Set}.$$

### 2.6.1 Properties of Cantor Set

✓ **Proposition 13.** Lebesgue Measure of Cantor Set is 0.

*Proof.* Note that Cantor Set is Lebesgue measurable as it is countable intersection of closed sets, hence it is present in the Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R})$  and hence is also in  $\mathcal{M}_{\lambda}^*$  (Proposition 12). Hence, instead of  $\lambda^*$ , we can now write  $\lambda$  as  $P \in \mathcal{M}_{\lambda}^*$ . Now, measure of Cantor set  $P$  can be written as:

$$\begin{aligned} \lambda(P) &= \lambda\left(\bigcap_n E_n\right) \\ &= \lim_{n \rightarrow \infty} \lambda(E_n) \quad (\text{Proposition 7}) \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{3^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1.5^n} \\ &= 0. \end{aligned}$$

■

✓ **Proposition 14.** Cantor Set is Uncountable(!)

*Proof.* We will show that there exists a bijection between Cantor Set and an uncountable set, specifically ternary system. For this, consider the ternary representation of every number in  $[0, 1]$ . What this means is that every number in  $[0, 1]$  can be represented only using the numbers 0, 1 and 2. Hence, one write  $\frac{1}{3}$  as 0.1 and  $\frac{2}{3}$  as 0.2. Now,  $(1/3, 2/3) = E_1^c \cap [0, 1]$  is the set that has been removed from the process of creating  $E_1$  from  $E_0$ . Clearly, every number in this  $E_1^c \cap [0, 1]$  is of the form 0.1... where ... are all combinations of 0, 1 and 2. Therefore, we are now left with the  $E_1$  that has all the numbers represented as 0.0... or 0.2...

As we saw in the generation of  $E_1$ , the generation of  $E_2$  from  $E_1$  would hence involve removing numbers of the forms 0.01... and 0.21... And hence  $E_2$  would then be the set of numbers whose first two decimal places are restricted to NOT have the digit 1; that is,  $E_2$  would be of form 0.02..., 0.00..., 0.20..., 0.22...

Continuing like this, we see that  $E_n$  would have in ternary representation, all those numbers whose first  $n$  digits are NOT 1. Hence, for any  $p \in P$ ,  $p$  would have the ternary representation constructed only from 0 and 2, but NOT 1.

Now, consider the map  $f : P \rightarrow [0, 1]$  such that  $f(p)$  replaces each occurrence of 2 by 1 in the ternary representation of  $p$ . We now show that this map is surjective(!) so that  $P$  has atleast as many elements as  $[0, 1]$ . To show this, take any  $x \in [0, 1]$  in its ternary form, and replace all 1 by 2 and denote it as  $x'$ . Clearly,  $x'$  would be in  $P$  as  $x'$  has all decimal digits generated by 0 and 2. But  $f(x')$  would be opposite action

<sup>6</sup>From this point on-wards, whenever this text mentions that a given set is measurable in space  $(X, \mathcal{A}, \mu)$ , it must be assumed that the given set is in  $\mathcal{A}$ , given that there is no ambiguity.

and would be equal to  $x$ . Therefore, we showed that for any  $x \in [0, 1]$ ,  $\exists x' \in P$  such that  $f(x') = x$ . Hence  $f$  is surjective. Therefore  $P$  has atleast as many elements as  $[0, 1]$ . But since  $P \subseteq [0, 1]$  therefore  $P$  has atmost as many elements as  $[0, 1]$ . This dichotomy suggests that

*Cantor Set has as many elements as in  $[0, 1]$  (!)*

But since  $[0, 1]$  is uncountable, therefore,  $P$  is uncountable. ■

---

With this, we conclude that **for any set  $E \subseteq \mathbb{R}$ , if  $\lambda^*(E) = 0$ , then it's NOT necessarily true that  $E$  is countable.**  
 We now see an extremely interesting example of a Non-measurable set.

## 2.7 A Non-measurable Set

⊗ **Theorem 2.** There is a subset of  $\mathbb{R}$  that is not Lebesgue Measurable<sup>7</sup>.

*Proof.* We construct the proof in the following Acts:

**Act 1.** *Equivalence Relation on  $\mathbb{R}$ .*

Construct the following relation  $\sim$  on  $\mathbb{R}$ :

$$x \sim y \equiv x - y \in \mathbb{Q}.$$

Clearly,  $\sim$  is reflexive as  $x - x = 0$  is rational; it is also symmetric as negative of a rational is also a rational number; and it is also transitive as if  $x - y$  and  $y - z$  is rational, then  $x - y + y - z = x - z$  is sum of two rationals, which is also rational. Hence  $\sim$  is an equivalence relation. Therefore  $\sim$  partitions the whole  $\mathbb{R}$  into equivalence classes. Note that each equivalence class of  $x$  would consist elements of the form  $\mathbb{Q} + x$ . But since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , therefore  $\mathbb{Q} + x$ , that is each equivalence class, is dense in  $\mathbb{R}$ .

Now, each equivalence class clearly intersects  $(0, 1)$ , therefore, inducing the Axiom of Choice on the set of all equivalence classes, we can form a subset  $E \subset (0, 1)$  which contains exactly one element from each of the equivalence classes. We will later prove that  $E$  is not Lebesgue Measurable.

**Act 2.**  *$E$  satisfies certain properties.*

Consider the set  $\mathbb{Q} \cap (-1, 1)$ . Clearly, this is countable as it's subset of  $\mathbb{Q}$ . Then, consider  $\{r_n\}$  to be the enumeration of  $\mathbb{Q} \cap (-1, 1)$ . Construct the sequence of subsets  $E_n = E + r_n$ . We now verify that  $\{E_n\}$  satisfies the following properties:

1. The sets  $E_n$  are disjoint.
2.  $\bigcup_n E_n$  is a subset of the interval  $(-1, 2)$ .
3. The interval  $(0, 1)$  is included in  $\bigcup_n E_n$ .

*Property 1:* Assume that  $E_n \cap E_m \neq \Phi$  for some  $n, m \in \mathbb{N}$  such that  $n \neq m$ . Then  $\exists e_1, e_2 \in E$  such that  $e_1 + r_n = e_2 + r_m$  which means that  $e_1 - e_2 = r_m - r_n \in \mathbb{Q}$ . But this cannot happen as  $e_1, e_2$  are elements of  $E$  and  $E$  contains exactly one element from the equivalence class of  $\sim$  intersected with  $(0, 1)$ . Therefore  $e_1 - e_2 \notin \mathbb{Q}$ . Which is a contradiction. Hence  $E_n \cap E_m = \Phi$  for all  $n, m \in \mathbb{N}$  such that  $n \neq m$ .

*Property 2:* Take  $x \in \bigcup_n E_n$ . This implies that  $x \in E_m$  for some  $m \in \mathbb{N}$ . But  $E_m = E + r_m = \{e + r_m \mid e \in E\}$ . Since  $E \subset (0, 1)$  and  $r_m \in \mathbb{Q} \cap (-1, 1) \subset (-1, 1)$ , therefore  $x \in E + r_m \subseteq (-1, 2)$ . Hence  $\bigcup_n E_n \subseteq (-1, 2)$ .

*Property 3:* Take any  $x \in (0, 1)$ . Now take the  $e \in E$  such that  $x \sim e$ , or  $x - e \in \mathbb{Q}$ . Hence  $x \in \mathbb{Q} + e$ . That is  $x = r + e$ . But since  $0 < e < 1$  and  $0 < x < 1$ , therefore  $r = x - e \in \mathbb{Q} \cap (-1, 1)$ . Hence  $x \in E + r$  and if we denote  $r = r_n$  for some  $n \in \mathbb{N}$ , we get  $x \in E + r_n = E_n$ , therefore  $x \in \bigcup_i E_i$ . Hence  $(0, 1) \subseteq \bigcup_i E_i$ .

**Act 3.**  *$E$  is Not Lebesgue Measurable.*

Assume that  $E$  is in-fact Lebesgue Measurable. Now since  $E_n$  are disjoint (Property 1), therefore we can write:

$$\lambda \left( \bigcup_n E_n \right) = \sum_n \lambda(E_n).$$

Now, since **Lebesgue Measure is translation invariant**<sup>8</sup>, therefore  $\lambda(E_n) = \lambda(E + r_n) = \lambda(E)$ . Two cases now arise for  $\lambda(\bigcup_n E_n)$ :

1. If  $\lambda(E) = 0$ : Then  $\lambda(\bigcup_n E_n) = 0$ . But

$$\lambda((-1, 2)) = 3 \leq \lambda \left( \bigcup_n E_n \right) \quad (\text{Property 3}).$$

Therefore we have a contradiction.

---

<sup>7</sup>See [1] for more information.

<sup>8</sup>Proof?

2. If  $\lambda(E) \neq 0$ : Then  $\lambda(\bigcup_n E_n) = \sum_n \lambda(E) = +\infty$ . But

$$\lambda\left(\bigcup_n E_n\right) \leq \lambda((-1, 2)) = 3 \quad (\text{Property 2}).$$

We again have a contradiction.

Hence, the set  $E$  is just not Lebesgue Measurable! ■

## 2.8 Regularity

First consider the following proposition.

✓ **Proposition 15.** Consider  $E \subseteq \mathbb{R}$ . The following statements are equivalent:

1.  $E$  is Lebesgue measurable.
2.  $\forall \epsilon > 0, \exists$  an open set  $O$  such that

$$E \subseteq O \text{ and } \lambda^*(O \setminus E) < \epsilon.$$

3.  $\exists$  a  $G_\delta$  set  $G$  such that

$$E \subseteq G \text{ and } \lambda^*(G \setminus E) = 0.$$

*Proof.* The equivalence of each statement is as follows:

**1  $\implies$  2.** Consider  $E \subseteq \mathbb{R}$  to be Lebesgue measurable. As Question 2 in Exercise 2 shows, for any  $E \subseteq \mathbb{R}$  and any  $\epsilon > 0$ , there exists open set  $U$  such that  $E \subseteq U$  which satisfies

$$\lambda^*(U) \leq \lambda^*(E) + \epsilon.$$

Now since  $E \subseteq U$ , therefore,

$$\begin{aligned} \lambda^*(U \setminus E) &= \lambda^*(U) - \lambda^*(E) \\ &\leq \epsilon \end{aligned}$$

**2  $\implies$  3.** Similarly, the Question 3 of Exercise 2 shows that there exists a  $G_\delta$  set  $G$  such that  $E \subseteq G$  which satisfies  $\lambda^*(E) = \lambda^*(G)$ . This directly means that  $\lambda^*(G \setminus E) = 0$  because  $E \subseteq G$  so  $\lambda^*(G \setminus E) = \lambda^*(G) - \lambda^*(E)$ .

**3  $\implies$  1.** Since  $G$  is  $G_\delta$  set therefore it is intersection of open sets in  $\mathbb{R}$ . Now since any open set in  $\mathbb{R}$  is an union of open intervals (Homework I, 1) which is Lebesgue Measurable and therefore  $G$  is Lebesgue Measurable. Now, we can write  $E$  as

$$E = G \setminus (G \setminus E)$$

where  $G \setminus E$  is such that (from Statement 3)  $\lambda^*(G \setminus E) = 0$ , therefore, by Proposition 10,  $G \setminus E$  is Lebesgue Measurable. Hence  $E$  is also Lebesgue Measurable. ■

Now, consider the next proposition, which is dual of the above.

✓ **Proposition 16.** Consider  $E \subseteq \mathbb{R}$ . The following statements are equivalent:

1.  $E$  is Lebesgue measurable.
2.  $\forall \epsilon > 0, \exists$  closed set  $C$  such that

$$C \subseteq E \text{ and } \lambda^*(E \setminus C) < \epsilon.$$

3.  $\exists$  a  $F_\sigma$  set  $F$  such that

$$F \subseteq E \text{ and } \lambda^*(E \setminus F) = 0.$$



*Proof.* Implications are as follows:

**1  $\implies$  2.** Suppose  $E \subseteq \mathbb{R}$  is Lebesgue Measurable. Note that if  $E$  is Lebesgue Measurable (that is  $E \in \mathcal{M}_{\lambda^*}$ ), then  $E^c$  is also Lebesgue Measurable as  $\mathcal{M}_{\lambda^*}$  is a  $\sigma$ -algebra (Theorem 1). Hence, using Proposition 15 on  $E^c$  gives us an open set  $O$  for all  $\epsilon > 0$  such that  $E^c \subseteq O$  and  $\lambda^*(O \setminus E^c) < \epsilon$ . Now let's take it's complement. Therefore,  $C = O^c \subseteq E$  where  $C$  is clearly closed. Now,  $E \setminus O^c = O \setminus E^c$ <sup>9</sup>. Now,

$$\begin{aligned}\lambda^*(E \setminus O^c) &= \lambda^*(O \setminus E^c) \\ &< \epsilon\end{aligned}$$

which proves the first implication.

**2  $\implies$  3.** From Proposition 15, we have that  $\exists$  a  $G_\delta$  set  $G$  such that  $E^c \subseteq G$  and  $\lambda^*(G \setminus E^c) = 0$ . Note that the complement of countable intersection of open sets is countable union of closed sets. Therefore,  $F = G^c$  is an  $F_\sigma$  set. Now,  $G^c \subseteq (E^c)^c = E$ . Now, we know that  $E \setminus G^c = G \setminus E^c$ . Therefore, we have the result as follows:

$$\begin{aligned}\lambda^*(E \setminus G^c) &= \lambda^*(G \setminus E^c) \\ &= 0.\end{aligned}$$

**3  $\implies$  1.** Since  $F$  is an  $F_\sigma$  set, therefore,  $F \in \mathcal{M}_{\lambda^*}$ . Moreover, as Statement 2 show,  $\lambda^*(E \setminus F) = 0$ , thus by Proposition 10,  $E \setminus F \in \mathcal{M}_{\lambda^*}$ . Since,

$$E = F \cup (E \setminus F)$$

that is  $E$  is union of two Lebesgue Measurable sets, therefore  $E \in \mathcal{M}_{\lambda^*}$ , completing the proof. ■

**★ Definition 15. (Complete Measure Space)** The measure space  $(X, \mathcal{A}, \mu)$  is complete if the for any  $A \in \mathcal{A}$  such that  $\mu(A) = 0$  implies that for any subset  $B \subseteq A$ ,

$$\mu(B) = 0.$$

**Remark.** Trivial to see are the following:

- Hence, if  $\mu^*$  is an outer measure defined on  $X$ , then the space  $(X, \mathcal{M}_{\mu^*}, \mu^*)$  is complete (follows from Proposition 10).
- This means that the Lebesgue outer measure restricted to Lebesgue measurable subsets of  $\mathbb{R}$ ,  $(\mathbb{R}, \mathcal{M}_{\lambda^*}, \lambda)$  is complete.

**★ Definition 16. (Completion of a Measure Space)** Let  $(X, \mathcal{A})$  be a measurable space and let  $\mu$  be a measure on  $\mathcal{A}$ . The completion of  $\mathcal{A}$  under  $\mu$  is the collection  $\mathcal{A}_\mu$  of subsets  $A \subseteq X$  for which there are sets  $E$  and  $F$  in  $\mathcal{A}$  such that

$$E \subseteq A \subseteq F$$

and

$$\mu(E - F) = 0$$
<sup>10</sup>.

<sup>9</sup>It's not difficult to see as for any  $x \in E \setminus O^c$ ,  $x \in E$  but  $x \notin O^c$ . Therefore,  $x \in O$  but  $x \notin E^c$ , that is  $x \in O \setminus E^c$ . Similarly for the converse.

<sup>10</sup>Note that, in Exercise III, Q. 2, we proved that for any  $A \in \mathcal{A}$ , this is trivially true. That is, all  $\mathcal{A}$ -measurable subsets are  $\mathcal{A}_\mu$ -measurable. In particular,  $E$  was a  $F_\sigma$  set and  $F$  was a  $G_\delta$  set.

### 3 Measurable Functions

We now see the definition and basic properties of Measurable Functions, which would later be used to define Lebesgue Integral.

**★ Definition 17. (Measurable Function)** Let  $(X, \mathcal{A})$  be a measurable space and let  $A \subseteq X$  which is in  $\mathcal{A}$ . The function  $f : A \rightarrow [-\infty, +\infty]$ , is called a Measurable function<sup>11</sup> if

$$\{x \mid f(x) > \alpha\} \text{ for any } \alpha \in \mathbb{R} \text{ is Measurable (belongs in } \mathcal{A} \text{).}$$

**Remark.** Please note that the function  $f$  defined above has a measurable domain.

✓ **Proposition 17.** Let  $(X, \mathcal{A})$  be a measurable space and  $A \in \mathcal{A}$ . Let  $f : A \rightarrow [-\infty, +\infty]$  be a function. Then, the following statements are equivalent:

1.  $f$  is a Measurable Function.
2. For all  $\alpha \in \mathbb{R}$ , the set  $\{x \mid f(x) \geq \alpha\} \in \mathcal{A}$ .
3. For all  $\alpha \in \mathbb{R}$ , the set  $\{x \mid f(x) < \alpha\} \in \mathcal{A}$ .
4. For all  $\alpha \in \mathbb{R}$ , the set  $\{x \mid f(x) \leq \alpha\} \in \mathcal{A}$ .

*Proof.* The equivalence is shown as follows:

**1  $\implies$  2.** Since  $f$  is a measurable, therefore for all  $\alpha \in \mathbb{R}$ , the set  $\{x \mid f(x) > \alpha\} \in \mathcal{A}$ . This means that  $C_{\alpha - \frac{1}{n}} = \{x \mid f(x) > \alpha - \frac{1}{n}\} \in \mathcal{A}$  for all  $n \in \mathbb{N}$ . Now, the following set

$$C = \bigcap_n C_{\alpha - \frac{1}{n}} = \{x \mid f(x) \geq \alpha\}.$$

is Measurable as  $C \in \mathcal{A}$  because  $C_{\alpha - \frac{1}{n}} \in \mathcal{A}$  for any  $n \in \mathbb{N}$ , hence the countable intersection would also be in  $\mathcal{A}$ , hence measurable.

**2  $\implies$  3.** Since  $\{x \mid f(x) \geq \alpha\} \in \mathcal{A}$ , therefore it's complement  $\{x \mid f(x) < \alpha\} \in \mathcal{A}$  for any  $\alpha \in \mathbb{R}$ .

**3  $\implies$  4.** Since  $\{x \mid f(x) < \alpha\} \in \mathcal{A}$  for any  $\alpha \in \mathbb{R}$ , thus,  $C_{\alpha + \frac{1}{n}} = \{x \mid f(x) < \alpha + \frac{1}{n}\} \in \mathcal{A}$  for all  $n \in \mathbb{N}$ , hence

$$C = \bigcap_n C_{\alpha + \frac{1}{n}} = \{x \mid f(x) \leq \alpha\}$$

and since each  $C_{\alpha + \frac{1}{n}} \in \mathcal{A}$ , therefore  $C \in \mathcal{A}$ .

**4  $\implies$  1.** Since  $\{x \mid f(x) \leq \alpha\} \in \mathcal{A}$  then it's complement  $\{x \mid f(x) > \alpha\}$  for all  $\alpha \in \mathbb{R}$ , making  $f$  measurable. ■

✓ **Proposition 18.** The following are basic examples of measurable functions:

- If  $f$  is a Measurable function, then the set  $\{x \mid f(x) = \alpha\}$  is measurable for all  $\alpha \in \mathbb{R}$ .
- Constant functions are Measurable.
- The characteristic function  $\chi_A$  defined by:

$$\chi_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases}$$

is Measurable if and only if  $A$  is measurable.

- Continuous functions are Measurable.
- Let  $(X, \mathcal{A})$  be a measurable space. If  $f$  and  $g$  are Measurable functions on  $X$ , then the sets

$$\begin{aligned} &\{x \in X \mid f(x) \neq g(x)\} \\ &\{x \in X \mid f(x) < g(x)\} \end{aligned}$$

are measurable (belongs to  $\mathcal{A}$ ).

- <sup>12</sup> Monotone functions are Measurable.

<sup>11</sup>One writes  $f$  as  $\mathcal{A}$ -measurable function to denote the  $\sigma$ -algebra over whose subset the function  $f$  is defined.

<sup>12</sup>Question 3 of Exercise 3.

- <sup>13</sup>Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a differentiable function. Then  $f'$  is a  $\lambda$ -Measurable function.

*Proof.* The **first** example is trivial to see in light of Proposition 17 by taking intersection of  $\{x \mid f(x) \leq \alpha\}$  and  $\{x \mid f(x) \geq \alpha\}$ , both of which are measurable.

For **second**, consider the constant function  $f(x) = b \forall x \in \mathbb{R}$ . Now, for all  $\alpha \in \mathbb{R}$ , consider the set  $f^{-1}((\alpha, \infty)) = \{x \mid f(x) > \alpha\}$ . If  $b > \alpha$ , then we are done, if  $b \leq \alpha$ , then by previous result,  $\{x \mid f(x) \leq \alpha\}$  is also measurable (equal to  $\mathbb{R}$  and  $\mathbb{R} \in \mathcal{A}$ ).

For **third** example, consider the set  $\chi_A^{-1}(\alpha, \infty) = \{x \mid \chi_A(x) > \alpha\}$  for any  $\alpha \in \mathbb{R}$ . If  $\alpha > 1$ , then  $f^{-1}(\alpha, \infty) = \Phi \in \mathcal{A}$ . If  $\alpha = 1$ , then  $f^{-1}[\alpha, \infty) = A$ , since  $\chi_A(x)$  is given measurable, hence  $A$  is measurable. Now, Assume that  $A$  is measurable. Then consider the set  $\chi_A^{-1}(\alpha, \infty)$  for any  $\alpha \in \mathbb{R}$ . As we saw previously, the case for  $\alpha > 1$  is trivial. For  $0 < \alpha \leq 1$ ,  $\chi_A^{-1}(\alpha, \infty) = A \in \mathcal{A}$ . Finally, for  $\alpha \leq 0$ ,  $\chi_A^{-1}(-\infty, \alpha] = \Phi \in \mathcal{A}$ . Thus,  $\chi_A$  is measurable.

For **fourth**, since  $f$  is continuous (so inverse of open sets is open, by definition), therefore  $f^{-1}(\alpha, \infty)$  is open in  $\mathbb{R}$ , hence it must be Borel, hence measurable for any  $\alpha \in \mathbb{R}$ .

For **fifth**, since  $f$  and  $g$  are measurable. Then due to next Proposition 19, we know that  $f - g$  is also measurable. This means that for any  $\alpha \in \mathbb{R}$ ,

$$\{x \in X \mid f(x) - g(x) < \alpha\}$$

is measurable. Now set  $\alpha = 0$  to get the result. Moreover, from this, we also get that  $\{x \in X \mid f(x) - g(x) > 0\}$  is also measurable. Hence,

$$\{x \in X \mid f(x) - g(x) \neq 0\} = \{x \in X \mid f(x) - g(x) < 0\} \cup \{x \in X \mid f(x) - g(x) > 0\}$$

is also measurable.

For **sixth**, we proceed as follows:

Consider the function  $f : A \rightarrow \mathbb{R}$  where  $A \in \mathcal{M}_{\lambda}^*$  to be monotone. Now, consider the following two sets for any  $\alpha \in \mathbb{R}$ :

$$\begin{aligned} A_1 &= \{x \in A \mid f(x) > \alpha\} \\ A_2 &= (f^{-1}(\alpha), \infty) \cap A \end{aligned}$$

Now, take any  $x \in A_1$ , then  $f(x) > \alpha \implies x > f^{-1}(\alpha)$ . Now if  $f^{-1}(\alpha) \cap A = \Phi$ , then  $\{x \in A \mid f(x) > \alpha\} = \Phi$  which is trivially measurable and we would be done. If however  $f^{-1}(\alpha) \cap A \neq \Phi$ , then  $f^{-1}(\alpha) = \{y \in A \mid f(y) = \alpha\}$  so that  $f(y) > \alpha$  implies that  $y > f^{-1}(\alpha)$  so that  $f(y) > f(f^{-1}(\alpha)) = \alpha$ . Therefore,  $x > f^{-1}(\alpha)$ , that is  $x \in A_2$ , proving that  $A_1 \subseteq A_2$ .

Similarly, take  $x \in A_2$ , therefore

$$\begin{aligned} x &> f^{-1}(\alpha) \\ f(x) &> f(f^{-1}(\alpha)) \\ f(x) &> \alpha \\ x &\in \{x \mid f(x) > \alpha\} \\ x &\in A_1. \end{aligned}$$

Therefore  $A_2 \subseteq A_1$ . Hence,  $A_1 = A_2$ . But since  $(f^{-1}(\alpha), \infty)$  is an interval, hence measurable and  $A$  is given measurable, therefore  $A_2 = (f^{-1}(\alpha), \infty) \cap A$  is measurable, which makes  $A_1 = \{x \mid f(x) > \alpha\} = A_2$  measurable for all  $\alpha \in \mathbb{R}$ .

For **seventh**, the result is simple to see since we are given that  $f$  is  $\lambda$ -Measurable due to continuity (see Statement 4). Therefore, we can define the sequence of functions  $\{f_n\}$  as follows:

$$f_n(x) = \frac{f\left(x + \frac{1}{n}\right) - f(x)}{\frac{1}{n}} \forall x \in \mathbb{R}.$$

As we can see,  $f_n$  is  $\lambda$ -Measurable due to Proposition 19. Hence, we can see that because  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$  for any  $x \in \mathbb{R}$ , and since  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ , therefore  $f_n \rightarrow f'$  is  $\lambda$ -Measurable (Proposition 21). ■

---

<sup>13</sup>Question 4 of Exercise 3.

✓ **Proposition 19.** Let  $(X, \mathcal{A})$  be a measurable space and let  $A \in \mathcal{A}$ . Consider two Measurable functions  $f, g : A \rightarrow [0, +\infty]$  and  $c \in \mathbb{R}$ . Then,

1.  $f + c$ ,
2.  $f \pm g$
3.  $cf$ ,
4.  $fg$

are also Measurable.

*Proof.* 1. Since  $f$  is measurable, therefore the set  $\{x \mid f(x) > \alpha - c\} = \{x \mid f(x) + c \geq \alpha\}$  is measurable for any  $\alpha \in \mathbb{R}$ .

2. Both  $f$  and  $g$  are given measurable. The set  $(f + g)^{-1}(\alpha, \infty)$  can be written as:

$$\begin{aligned} (f + g)^{-1}(\alpha, \infty) &= \{x \mid f(x) + g(x) > \alpha\} \\ &= \{x \mid f(x) > \alpha - g(x)\} \\ &= \{x \mid f(x) > b\} \end{aligned}$$

where  $b \in [-\infty, \alpha]$ . Note that the case where  $g(x) = +\infty$  is trivial as  $f(x) > \alpha - (+\infty) \equiv f(x) > -\infty$ , which is by definition of co-domain of  $f$ . Now since  $\{x \mid f(x) > b\}$  is measurable for any  $b \in \mathbb{R} \supset (-\infty, \alpha]$  for any  $\alpha \in \mathbb{R}$ , therefore  $(f + g)^{-1}(\alpha, \infty)$  is measurable for any  $\alpha \in \mathbb{R}$ .

3. Note that for  $c = 0$ , the function becomes constant and hence measurable (Proposition 18). Consider the set  $(cf)^{-1}(\alpha, \infty)$ . We can write this as follows,

$$\begin{aligned} (cf)^{-1}(\alpha, \infty) &= \{x \mid cf(x) > \alpha\} \\ &= \{x \mid f(x) > \alpha/c\} \end{aligned}$$

where  $c \neq 0$ . Since  $f$  is measurable, therefore  $\{x \mid f(x) > \alpha/c\}$  is also measurable for any  $\alpha \in \mathbb{R}$ . Hence  $cf$  is measurable.

4. Consider the set  $(f^2)^{-1}(-\infty, \alpha)$  for any  $\alpha \in \mathbb{R}$ .

$$\begin{aligned} (f^2)^{-1}(-\infty, \alpha) &= \{x \mid f^2(x) < \alpha\} \\ &= \{x \mid -\sqrt{\alpha} < f(x) < \sqrt{\alpha}\} \\ &= \{x \mid f(x) < \sqrt{\alpha}\} \cap \{x \mid f(x) > -\sqrt{\alpha}\} \end{aligned}$$

Therefore if  $f$  is Measurable, then  $f^2$  is Measurable. With this, we can simply write  $fg$  as:

$$fg = \frac{(f + g)^2 - (f - g)^2}{4}$$

which, by previous results (2 & 3), is measurable. ■

✓ **Proposition 20.** <sup>14</sup> Let  $(X, \mathcal{A})$  be a measurable space. Consider a function  $f : A \rightarrow \mathbb{R}$  where  $A \in \mathcal{A}$ . Then the following are equivalent:

1.  $f$  is a  $\mathcal{A}$ -Measurable function.
2.  $f^{-1}(U)$  is a measurable set  $\forall$  open sets  $U \subseteq \mathbb{R}$ .
3.  $f^{-1}(C)$  is a measurable set  $\forall$  closed sets  $C \subseteq \mathbb{R}$ .
4.  $f^{-1}(B)$  is a measurable set  $\forall$  borel sets  $B \in \mathcal{B}(\mathbb{R})$ .

*Proof.* The proof is exactly the same as of Proposition 26. ■

★ **Definition 18. (Sequence of fuctions)** If  $\{f_n\}$  is a sequence of  $[-\infty, +\infty]$  valued functions on  $A$ , then  $\sup_n f_n : A \rightarrow [-\infty, +\infty]$  is defined by

$$\left( \sup_n f_n \right) (x) = \sup \{f_n(x) \mid n \in \mathbb{N}\}.$$

<sup>14</sup>Question 1 of Exercise 3.

**Remark.** One similarly defines the following:

- The infimum:

$$\left(\inf_n f_n\right)(x) = \inf\{f_n(x) \mid n \in \mathbb{N}\}.$$

- The limit supremum:

$$\left(\limsup_n f_n\right)(x) = \limsup\{f_n(x) \mid n \in \mathbb{N}\}.$$

- The limit infimum:

$$\left(\liminf_n f_n\right)(x) = \liminf\{f_n(x) \mid n \in \mathbb{N}\}.$$

- The limit:

$$\left(\lim_n f_n\right)(x) = \lim\{f_n(x) \mid n \in \mathbb{N}\}.$$

✓ **Proposition 21.** Let  $(X, \mathcal{A})$  be a measurable space and let  $A \in \mathcal{A}$ . Consider  $\{f_n\}$  be a sequence of  $[-\infty, +\infty]$ -valued measurable functions on  $A$ . Then,

1. The functions  $\sup_n f_n$  and  $\inf_n f_n$  are measurable.
2. The functions  $\limsup_n f_n$  and  $\liminf_n f_n$  are measurable.
3. The function  $\lim_n f_n$  (whose domain is  $\{x \in A \mid \limsup_n f_n(x) = \liminf_n f_n(x)\}$ ) is measurable.

*Proof.* Note that the set  $(\sup_n f_n)^{-1}(-\infty, \alpha] = \{x \in A \mid (\sup_n f_n)(x) \leq \alpha\} = \bigcap_n \{x \in A \mid f_n(x) \leq \alpha\}$ . Therefore  $\sup_n f_n$  is measurable. Similarly,  $(\inf_n f_n)^{-1}(-\infty, \alpha) = \{x \in A \mid (\inf_n f_n)(x) < \alpha\} = \bigcup_n \{x \in A \mid f_n(x) < \alpha\}$ . Now, denote  $g_k = \sup_{n \geq k} f_n$  and  $h_k = \inf_{n \geq k} f_n$ . But since  $\limsup_n f_n = \inf_{n \geq 0} \sup_{k \geq n} f_k = \inf_{n \geq 0} g_n$  and  $\{g_n\}$  is measurable by 1<sup>st</sup> property, therefore  $\limsup_n f_n$  is also measurable, similarly for  $\liminf_n f_n$ . ■

### 3.1 Almost Everywhere Property.

★ **Definition 19.** ( $\mu$ -almost everywhere) Let  $(X, \mathcal{A}, \mu)$  be a measure space. A property  $P$  of points of  $X$  is said to hold  $\mu$ -almost everywhere if the set

$$N = \{x \in X \mid P \text{ does not hold for } x\}$$

has measure zero. That is,

$$\mu(N) = 0.$$

**Remark.** Note that it's not necessary for the set  $N$  to belong in  $\mathcal{A}$ . The only requirement is for the set  $N$  to be contained in a set  $F \in \mathcal{A}$  and then  $\mu(F) = 0$  (which automatically implies that  $\mu^*(N) = 0$ ).

But, if  $\mu$  is complete then  $N \in \mathcal{A}$ . See Definition 15.

★ **Definition 20.** (Almost everywhere convergence) If  $\{f_n\}$  is a sequence of functions on  $X$  and  $f$  is a function on  $X$ , then

$$\{f_n\} \longrightarrow f \text{ almost everywhere.}$$

if the set

$$\{x \in X \mid f(x) \neq \lim_n f_n(x)\}$$

is of measure zero.

✓ **Proposition 22.** Let  $(X, \mathcal{A}, \mu)$  be a measure space and let  $f$  and  $g$  be extended real valued functions on  $X$  that are equal almost everywhere. If  $\mu$  is **complete** and if  $f$  is Measurable, then  $g$  is also Measurable.

*Proof.* Consider the region of non-equality as

$$N = \{x \mid f(x) \neq g(x)\}.$$

Given to us is the fact that  $\mu^*(N) = 0$  and since  $\mu$  is complete, so  $N \in \mathcal{A}$ . Now, consider the following for any  $\alpha \in \mathbb{R}$ :

$$\{x \mid g(x) \geq \alpha\} = (\{x \mid g(x) \geq \alpha\} \cap N) \cup (\{x \mid g(x) \geq \alpha\} \cap N^c).$$

Denote the set  $A = \{x \mid g(x) \geq \alpha\} \cap N$  and  $B = \{x \mid g(x) \geq \alpha\} \cap N^c$ . Since for any  $x \in (\{x \mid g(x) \geq \alpha\} \cap N^c)$ ,  $f(x) = g(x)$ , therefore, we can equivalently write  $B = (\{x \mid f(x) \geq \alpha\} \cap N^c)$ . Now  $N^c \in \mathcal{A}$  and due to Measurability of  $f$ ,  $\{x \mid f(x) \geq \alpha\} \in \mathcal{A}$ . Hence  $B \in \mathcal{A}$ . Finally, due to  $\{x \mid g(x) \geq \alpha\} \cap N \subseteq N$  and  $\mu$  being complete with  $\mu(N) = 0$ , we get  $\{x \mid g(x) \geq \alpha\} \cap N \in \mathcal{A}$ , completing the proof. ■

✓ **Proposition 23.** Let  $(X, \mathcal{A}, \mu)$  be a measure space, let  $\{f_n\}$  be sequence of extended real valued functions on  $X$  and let  $f$  be an extended real valued function on  $X$  such that

$$\{f_n\} \longrightarrow f \text{ almost everywhere.}$$

If  $\mu$  is **complete** and if each  $f_n$  is Measurable, then  $f$  is Measurable.

*Proof.* As Proposition 21 shows,  $\liminf_n f_n$  and  $\limsup_n f_n$  are Measurable. As the given condition shows,  $\liminf_n f_n$  is equal to  $f$  for almost all  $X$ . Hence Proposition 22 implies that  $f$  is also Measurable. ■

#### 3.1.1 A striking result

The following is an interesting result about Lebesgue Measurable functions on  $\mathbb{R}$ .

✓ **Proposition 24.** Ci

### 3.2 Cantor Set, again.

With the new tool in hand (measurable functions), we now turn back to the ever-interesting Cantor set, this time, to prove the sheer size of the  $\sigma$ -algebra  $\mathcal{M}_{\lambda^*}$  in comparison to the Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R})$ . In particular we show that  $\mathcal{B}(\mathbb{R}) \subsetneq \mathcal{M}_{\lambda^*}$ .

But before that, we look at following results:

✓ **Proposition 25.** Show that the function  $\phi$  defined by

$$\begin{aligned} \phi : [0, 1] &\longrightarrow P(\text{Cantor Set}), \text{ such that,} \\ \phi(\alpha) &= \sum_{n=1}^{\infty} \frac{2b_n}{3^n} \text{ for } \alpha \in [0, 1] \end{aligned}$$

where  $b_n \in \{0, 1\} \forall n \in \mathbb{N}$  is Measurable in  $\mathcal{M}_{\lambda^*}$ .

*Proof.* Note that  $\phi(\alpha)$  thus maps a decimal number to it's binary representation  $\{b_n\}$ . First, we define the following function:

$$\begin{aligned} \phi_n : [0, 1] &\longrightarrow \{0, 1\} \\ \phi_n(\alpha) &= b_n. \end{aligned}$$

That is,  $\phi_n$  maps  $\alpha$  to it's  $n^{th}$  binary digit. We can see that  $\phi_n(\alpha)$  can be written as the following:

$$\phi_n(\alpha) = \chi_{E_n} = \begin{cases} 1 & \text{if } \alpha \in E_n \\ 0 & \text{otherwise.} \end{cases}$$

where  $E_n$  is the intersection of countable sequence of sub-intervals of  $[0, 1]$ . Hence  $E_n$  is a Lebesgue measurable subset of  $\mathbb{R}$ , so it is in  $\mathcal{M}_{\lambda^*}$ .

But, as Proposition 18, statement 3 shows,  $\chi_{E_n} = \phi_n$  is then a Measurable function.

Now, the following arguments:

$$\begin{aligned} \frac{2}{3^n} \phi_n(\alpha) &= \frac{2b_n}{3^n} \text{ is measurable (Proposition 19).} \\ \left\{ \frac{2\phi_n(\alpha)}{3^n} \right\} &\text{ is a sequence of measurable functions.} \\ \left\{ \sum_{k=1}^n \frac{2\phi_k(\alpha)}{3^k} \right\} &\text{ is also a sequence of measurable functions (Proposition 19).} \\ \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{2\phi_k(\alpha)}{3^k} &\text{ is a measurable function (Proposition 21).} \end{aligned}$$

Hence the function which maps each real from  $[0, 1]$  to it's binary representation is measurable. ■

✓ **Proposition 26.** Let  $(X, \mathcal{A})$  be a measurable space. If  $f$  is a  $\mathcal{A}$ -Measurable function on  $A$  and  $B \in \mathcal{B}(\mathbb{R})$ , then  $f^{-1}(B) \in \mathcal{A}$ .

*Proof.* Denote  $\mathcal{D}$  be the following set:

$$\mathcal{D} = \{B \subseteq \mathbb{R} \mid f^{-1}(B) \in \mathcal{A}\}.$$

Now, note that,

1. Since  $f^{-1}(\mathbb{R}) = A \in \mathcal{A}$ , therefore  $\mathbb{R} \in \mathcal{D}$ .
2. If  $B \in \mathcal{D}$ , then

$$B^c = \mathbb{R} \cap B^c$$

and

$$\begin{aligned} f^{-1}(B^c) &= f^{-1}(\mathbb{R} \cap B^c) \\ &= f^{-1}(\mathbb{R}) \cap f^{-1}(B^c) \\ &= A \cap (f^{-1}(B))^c \end{aligned}$$

Now since  $A \in \mathcal{A}$  and  $f^{-1}(B) \in \mathcal{A}$  because  $B \in \mathcal{D}$ , therefore  $f^{-1}(B^c) \in \mathcal{A}$  so that  $B^c \in \mathcal{D}$ .

3. We know that from the basic results of set functions that

$$f^{-1}\left(\bigcup_n B_n\right) = \bigcup_n f^{-1}(B_n)$$

Hence  $\mathcal{D}$  is a  $\sigma$ -algebra on  $\mathbb{R}$  (!) Now, due to measurability of  $f$ , we know that the set  $\{x \mid f(x) > \alpha\}$  is in  $\mathcal{A}$ , which is equivalent to saying that  $f^{-1}(\alpha, \infty) \in \mathcal{A}$ . This hence means that  $(\alpha, \infty) \in \mathcal{D}$  for any  $\alpha \in \mathbb{R}$ . Proposition 2 showed that a  $\sigma$ -algebra generated by such subsets of  $\mathbb{R}$  is  $\mathcal{B}(\mathbb{R})$ . Hence, for any  $B \in \mathcal{B}(\mathbb{R})$ , we have that  $B \in \mathcal{D}$ . Therefore for any Borel set  $B$ ,  $f^{-1}(B) \in \mathcal{A}$ <sup>15</sup>. ■

---

**WHY IS THIS SUBSECTION STILL NOT COMPLETE!?**

---

<sup>15</sup>This is a very interesting way to prove such a statement. Notice how we analyzed the set of all possible subsets of  $\mathbb{R}$  for which  $f^{-1}(B) \in \mathcal{A}$  right from the start!



### 3.3 Sequence of Functions approximating a Measurable Function.

We now show that any measurable function can be defined in terms of a simple function and a step function. For this, we first define what we mean by simple functions in Definition 21. Before that, let's see few more interesting-but-basic properties of measurable functions.

✓ **Proposition 27.** Let  $(X, \mathcal{A})$  be a measurable space and  $f$  be an extended real valued function on  $A \in \mathcal{A}$ . Define the following:

$$f^+(x) = \max(f(x), 0) \text{ and } f^-(x) = -\min(f(x), 0).$$

Then,  $f$  is measurable if and only if  $f^+$  and  $f^-$  both are measurable on  $A$ .

*Proof.* If  $f$  is measurable, then  $\{x \mid f(x) \geq \alpha\}$  is measurable. Note that  $f^+(x) \geq 0$ . Hence, for the case when  $\alpha > 0$ , the set  $\{x \mid f^+(x) \geq \alpha\} = \{x \mid f(x) \geq \alpha\}$  which is measurable due to measurability of  $f$ . Similarly, if  $\alpha = 0$ , then  $\{x \mid f^+(x) \geq 0\} = \{x \mid f(x) > 0\} \cup \{x \mid f(x) = 0\}$  in which both sets are measurable in view of Proposition 18. Finally, for  $\alpha < 0$ , we have  $\{x \mid f^+(x) > \alpha\} = \{x \mid f^+(x) \geq 0\}$  which again is measurable. Now,  $f^-(x) = -\min(f(x), 0) = \max(-f(x), 0)$  and since  $-f$  is also measurable (Proposition 19), therefore if  $f$  is a measurable function, then  $f^+$  and  $f^-$  are both measurable functions too. To show the converse, note that  $f = f^+ - f^-$  and since both are measurable, therefore  $f$  is also measurable (Proposition 19). ■

**Remark.** Due to the above result, we can hence deduce that if  $f$  is a  $\mathcal{A}$ -Measurable function then,

$$|f| = f^+ + f^- \text{ is a Measurable function on } A.$$

✓ **Proposition 28.** Let  $(X, \mathcal{A})$  be a measurable space and  $A \in \mathcal{A}$ . Let  $f : A \rightarrow [-\infty, +\infty]$ . Then,

1. If  $f$  is  $\mathcal{A}$ -Measurable and if  $B$  is a subset of  $A$ , then the restriction  $f_B$  of  $f$  to  $B$  is also  $\mathcal{A}$ -Measurable.
2. If  $\{B_n\}$  is a sequence of sets that belong to  $\mathcal{A}$  such that  $A = \bigcup_n B_n$  and  $f_{B_n}$  is  $\mathcal{A}$ -Measurable for each  $n$ , then  $f$  is also  $\mathcal{A}$ -Measurable.

*Proof.* The first result follows directly from the following observation:

$$\{x \in B \mid f_B(x) > \alpha\} = B \cap \{x \in A \mid f(x) > \alpha\}$$

and the second result follows from the following:

$$\{x \in A \mid f(x) > \alpha\} = \bigcup_n \{x \in B_n \mid f_{B_n}(x) > \alpha\}.$$

both for any  $\alpha \in \mathbb{R}$ . ■

★ **Definition 21. (Simple Function)** A function is called simple if it has only finitely many values. Equivalently, we say that  $f$  is simple if we can write it as the following:

$$f = \sum_{k=1}^N \alpha_k \chi_{E_k}, \quad \alpha_k \in \mathbb{R}$$

where each  $E_k$  is a measurable set of finite measure.

**Remark.** Note that

- If  $E_k$  are intervals, then we say  $f$  to be a **step function**.

The following Proposition asserts that **any Measurable function can be approximated by an increasing sequence of simple functions**.

✓ **Proposition 29.** Let  $(X, \mathcal{A})$  be a measurable space and let  $A \in \mathcal{A}$  with  $f : A \rightarrow [0, +\infty]$  be a Measurable function on  $A$ . Then there exists a sequence  $\{f_n\}$  of **simple**  $[0, +\infty)$ -valued Measurable functions on  $A$  that satisfy

$$f_1(x) \leq f_2(x) \leq f_3(x) \leq \dots$$

and

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

for any  $x \in A$ .

*Proof.* For the proof, construct the following sequence of sets, by dividing the whole interval  $[0, n]$  for any  $n \in \mathbb{N}$  into  $n2^n$  number of intervals each of length  $\frac{1}{2^n}$  and denote the following set:

$$A_{n,k} = \left\{ x \in A \mid \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n} \right\}$$

for any  $n \in \mathbb{N}$  and  $k = 1, 2, \dots, n2^n$ . With this construction, we can now define the following function for each  $n$ :

$$\phi_n : A \rightarrow [0, \infty), \text{ defined as}$$

$$\phi_n(x) = \begin{cases} \frac{k-1}{2^n} & \text{if } x \in A_{n,k} \text{ for any } k = 1, 2, \dots, n2^n \\ n & \text{if } x \in A - \bigcup_k A_{n,k}. \end{cases}$$

Note that we can alternatively write  $\phi_n(x)$  as the following (with more clarity):

$$\phi_n(x) = \begin{cases} \frac{k-1}{2^n} & \text{if } f(x) \leq n, \text{ where } \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n} \text{ for some } k \in \{1, 2, \dots, n2^n\} \\ n & \text{if } f(x) > n. \end{cases}$$

We now show that  $\phi_n(x) \leq \phi_{n+1}(x) \forall x \in A$ . Let's first show this for  $f(x) \leq n$ .

If  $f(x) \leq n$ , then,

$$\phi_n(x) = \frac{k_0-1}{2^n} \text{ for some } k_0 \in \{1, 2, \dots, n2^n\}$$

such that  $\frac{k_0-1}{2^n} \leq f(x) < \frac{k_0}{2^n}$ . Now, two cases arises:

- **If  $\frac{k_0-1}{2^n} \leq f(x) < \frac{2k_0-1}{2^{n+1}}$**  : This is just the case that  $f(x)$  lies in the first half of the interval  $\left[\frac{k_0-1}{2^n}, \frac{k_0}{2^n}\right]$ . Hence, in this case we get that:

$$\frac{k_0-1}{2^n} = \frac{2k_0-2}{2^{n+1}} \leq f(x) < \frac{2k_0-1}{2^{n+1}}$$

such that  $\phi_n(x) = \frac{k_0-1}{2^n} = \phi_{n+1}(x)$ .

- **If  $\frac{2k_0-1}{2^{n+1}} \leq f(x) < \frac{k_0}{2^n}$**  : This is the case when  $f(x)$  lies in the second half of the interval. In this case, we see that,

$$\frac{2k_0-1}{2^{n+1}} \leq f(x) < \frac{2k_0}{2^{n+1}} = \frac{k_0}{2^n}$$

so that  $\phi_n(x) = \frac{k_0-1}{2^n} = \frac{2k_0-2}{2^{n+1}} < \frac{2k_0-1}{2^{n+1}} = \phi_{n+1}(x)$ .

Hence from both the cases, we have  $\phi_n(x) \leq \phi_{n+1}(x)$  for all  $x \in A$  such that  $f(x) \leq n$ . One can similarly see the same result for  $n < f(x) \leq n+1$  and for  $f(x) > n+1$ ,  $\phi_n(x) \leq \phi_{n+1}(x)$  follows trivially. Hence, we have proved that  $\forall n \in \mathbb{N}$  and  $x \in A$ ,

$$\phi_n(x) \leq \phi_{n+1}(x). \quad (9)$$

Now, one can write the function  $\phi_n$  as the following combination too:

$$\phi_n(x) = \sum_{k=1}^{n2^n} \frac{k-1}{2^n} \chi_{A_{n,k}} + n \chi_{A - \bigcup_k A_{n,k}} \quad (10)$$

Due to the above representation of  $\phi_n$ , the following steps becomes easier (& interesting) to see.

Now, first note that  $A_{n,k}$  is a measurable set because it's intersection of two measurable sets. Moreover,  $A - \bigcup_k A_{n,k}$  is also a measurable set. Hence, in view of Proposition 18, Statement 3, we get that  $\phi_n(x)$  is a Measurable function for any  $n \in \mathbb{N}$ . Therefore,  $\{\phi_n\}$  is a sequence of Measurable functions adhering (9). We again find two cases:

- **If  $f$  is finite** : Now since  $f$  is finite, therefore  $\exists n_0 \in \mathbb{N}$  such that  $f(x) \leq n_0$ . Hence, one can further deduce the following for all  $n \geq n_0$  (hence  $f(x) \leq n_0 \leq n$ ),

$$\begin{aligned} f(x) - \phi_n(x) &= f(x) - \frac{k-1}{2^n} \text{ for some } k \in \{1, 2, \dots, n2^n\} \text{ such that } \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n} \\ &< \frac{1}{2^n} \end{aligned}$$

Hence, as  $n \rightarrow \infty$ ,  $|f(x) - \phi_n(x)| \rightarrow 0$ .

- If  $f$  is infinite for some  $x \in A$  : If  $f$  is infinite, then  $\forall n \in \mathbb{N}, f(x) > n$ . Hence,

$$\phi_n(x) = n \text{ for all } n \in \mathbb{N}.$$

Therefore  $\lim_{n \rightarrow \infty} \phi_n(x) = +\infty = f(x)$  for particular  $x \in A$  where  $f$  is infinity.

Hence, in both cases,  $\{\phi_n\}$  converges to  $f$ . The proof is therefore complete. ■

The following can be considered as an **important corollary of the above Proposition**.

✓ **Proposition 30.** Let  $(X, \mathcal{A})$  be a measurable space and let  $A \in \mathcal{A}$  with  $f : A \rightarrow [-\infty, +\infty]$  be a Measurable function on  $A$ . Then there exists a sequence  $\{f_n\}$  of **simple**  $(-\infty, +\infty)$ -valued Measurable functions on  $A$  that satisfy

$$|f_1(x)| \leq |f_2(x)| \leq |f_3(x)| \leq \dots$$

and

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

for any  $x \in A$ .

*Proof.* Since  $f$  is a Measurable function, therefore  $f^+$  and  $f^-$  are Measurable functions too (Proposition 27). Now, since any function  $f$  can be written as

$$f = f^+ - f^-$$

therefore, by Proposition 29, we have two sequences  $\{f_n^{(1)}\}$  and  $\{f_n^{(2)}\}$  such that

$$f_n^{(1)} \longrightarrow f^+ \text{ and } f_n^{(2)} \longrightarrow f^-$$

where  $f_1^{(1)}(x) \leq f_2^{(1)}(x) \leq \dots$  and  $f_1^{(2)}(x) \leq f_2^{(2)}(x) \leq \dots$ . Denote

$$f_n(x) = f_n^{(1)}(x) - f_n^{(2)}(x)$$

Therefore, we see that

$$|f_n(x)| = f_n^{(1)}(x) + f_n^{(2)}(x) \leq f_{n+1}^{(1)}(x) + f_{n+1}^{(2)}(x) = |f_{n+1}(x)|$$

Now, we can deduce that

$$\begin{aligned} |f(x) - f_n(x)| &= |f^+(x) - f^-(x) - f_n^{(1)}(x) + f_n^{(2)}(x)| \\ &= |f^+(x) - f_n^{(1)}(x) - (f^-(x) - f_n^{(2)}(x))| \\ &\leq |f^+(x) - f_n^{(1)}(x)| + |f^-(x) - f_n^{(2)}(x)| \\ &\rightarrow 0 + 0 \end{aligned}$$

Hence proved. ■

### 3.3.1 Replacing simple functions by step functions<sup>16</sup>.

We now prove a similar result akin to Proposition 29, where we show that *any measurable function can be approximated by a sequence of step functions, almost everywhere*. But before that, we prove a **basic fact about Lebesgue measurable sets with finite measure**.

✓ **Proposition 31.** For any  $\lambda$ -measurable set  $E$  of finite measure and a given  $\epsilon > 0$ , there exists a finite sequence of open intervals  $\{I_n\}_{n=1}^N$  such that

$$\lambda \left( E \Delta \left( \bigcup_{n=1}^N I_n \right) \right) < \epsilon.$$

*Proof.* Take any  $\epsilon > 0$ , then we have for any set  $E \subseteq \mathbb{R}$ , a sequence of open intervals  $\{I_n\}$  such that  $E \subseteq \bigcup_n I_n$  and  $\lambda(\bigcup_n I_n) \leq \lambda(E) + \epsilon$  or  $\lambda(\bigcup_n I_n \setminus E) \leq \epsilon < 2\epsilon$ . Now since  $\{I_n\}$  is a disjoint sequence, therefore,  $\lambda(\bigcup_n I_n) = \sum_n \lambda(I_n)$  and due to the fact that  $\lambda(E) < +\infty$ , we get that  $\sum_n \lambda(I_n) < +\infty$ .

Now, since  $\lambda(E) < +\infty$ , therefore the sum  $\sum_n \lambda(I_n) < +\infty$ , hence,  $\exists N \in \mathbb{N}$  such that  $\sum_{n=N+1}^{\infty} \lambda(I_n) < \epsilon$ . With this  $N$ , we now see that:

$$\begin{aligned} \lambda \left( E \Delta \bigcup_{n=1}^N I_n \right) &= \lambda \left( E \setminus \bigcup_{n=1}^N I_n \right) + \lambda \left( \bigcup_{n=1}^N I_n \setminus E \right) \quad (\text{both are disjoint.}) \\ &\leq \lambda \left( E \setminus \bigcup_{n=1}^N I_n \right) + \lambda \left( \bigcup_n I_n \setminus E \right) \\ &= \lambda \left( E \setminus \bigcup_{n=1}^N I_n \right) + \lambda \left( \bigcup_n I_n \setminus E \right) \\ &= \lambda \left( E \cap \left( \bigcup_{n=1}^N I_n \right)^c \right) + \lambda \left( \bigcup_n I_n \setminus E \right) \\ &\leq \lambda \left( \bigcup_{n=N+1}^{\infty} I_n \right) + \lambda \left( \bigcup_n I_n \setminus E \right) \quad \because E \cap \left( \bigcup_{n=1}^N I_n \right)^c \subseteq \bigcup_{n=N+1}^{\infty} I_n \\ &\leq \epsilon + \epsilon = 2\epsilon \end{aligned}$$

Hence, we get that for any finite Lebesgue measurable set  $E$ , for all  $\epsilon > 0$ ,  $\exists$  a sequence of open intervals  $\{I_n\}_{n=1}^N$  such that their symmetric difference is a set with measure  $\leq \epsilon$ . ■

✓ **Proposition 32.** Consider  $(\mathbb{R}, \mathcal{M}_{\lambda^*})$  to be the Lebesgue measurable space and  $A \in \mathcal{M}_{\lambda^*}$ . Let  $f : A \rightarrow [-\infty, +\infty]$  be a  $\lambda$ -Measurable function. Then there exists a sequence of step functions  $\{\phi_k\}$  such that

$$\phi_k \longrightarrow f \text{ almost everywhere.}$$

*Proof.* We will prove first that for any characteristic function, there exists a sequence of step functions converging to it. Let  $g = \chi_A$  be the characteristic function on  $A$ . Continuing from Proposition 31, we see that if we write the **step-function**  $\psi$  as

$$\psi = \sum_{k=1}^N \chi_{I_k}$$

where  $\{I_k\}$  is the set of open intervals such that  $\lambda \left( A \Delta \left( \bigcup_{k=1}^N I_k \right) \right) < \epsilon$  for a given  $\epsilon > 0$ , from Proposition 31, then we get that the set  $\{x \mid g(x) \neq \psi(x)\}$  has upper bound on its measure given as follows:

$$\begin{aligned} \left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) = \psi(x) \right\} &\subseteq A \cap \bigcup_{k=1}^N I_k \quad \because g(x) = \psi(x) \text{ iff } x \in \bigcup_{k=1}^N I_k \text{ and } 0 \text{ for other } x \in A \\ \left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) \neq \psi(x) \right\} &\supseteq \left( A \cap \bigcup_{k=1}^N I_k \right)^c \supseteq A \Delta \bigcup_{k=1}^N I_k. \end{aligned}$$

Similarly, it's easy to see that for any  $x$  such that  $g(x) \neq \psi(x)$ , we have  $x \in A \Delta \bigcup_{k=1}^N I_k$  so that we get,

$$\left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) \neq \psi(x) \right\} \subseteq A \Delta \bigcup_{k=1}^N I_k.$$

<sup>16</sup>Unfortunately, this section onwards, we would tentatively only work with the Lebesgue Measure space  $(\mathbb{R}, \mathcal{M}_{\lambda^*}, \lambda)$  for pedagogical reasons of the instructor, so the abstract versions of the following results would need to wait.

Hence,

$$\left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) \neq \psi(x) \right\} = A \Delta \bigcup_{k=1}^N I_k.$$

Therefore,

$$\lambda \left( \left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) \neq \psi(x) \right\} \right) < \epsilon$$

Therefore, for every  $n \geq 1$ , there exists a step-function  $\psi_n$  so that the set  $E_n = \left\{ x \in A \cup \left( \bigcup_{k=1}^N I_k \right) \mid g(x) \neq \psi_n(x) \right\}$  is such that

$$\lambda(E_n) < \frac{1}{2^n}.$$

Now, **define** the following two sets:

$$F_n = \bigcup_{j=n+1}^{\infty} E_j \quad (\text{a decreasing sequence})$$

$$F = \bigcap_{k=1}^{\infty} F_k.$$

For the set  $F_n$ , observe that

$$\begin{aligned} \lambda(F_n) &= \lambda \left( \bigcup_{j=n+1}^{\infty} E_j \right) \leq \sum_{j=n+1}^{\infty} \lambda(E_j) \\ &< \sum_{j=n+1}^{\infty} \frac{1}{2^j} \\ &= \frac{1}{2^n} \end{aligned}$$

and for set  $F$ ,

$$\begin{aligned} \lambda(F) &= \lambda \left( \bigcap_{k=1}^{\infty} F_k \right) = \lim_{k \rightarrow \infty} \lambda(F_k) \quad \because \text{Proposition 7 \& } \{F_k\} \text{ is measurable \& decreasing.} \\ &= 0. \end{aligned}$$

Note that  $\{F_k\}$  is measurable because any  $E_i$  is itself measurable because of Proposition 18, Statement 5.

Now,

$$\psi_n(x) \longrightarrow g(x) \quad \forall x \in F^c$$

because  $F^c = F_1^c$  since  $\{F_k\}$  is a decreasing sequence, therefore  $F^c$  is the set where  $g(x)$  satisfies with the limit step function.

Finally,  $\psi_n \not\rightarrow f \quad \forall x \in F$ , but since  $\lambda(F) = 0$ , hence

$$\psi_n \longrightarrow g \text{ almost everywhere}$$

Now, what we have proved so far is that for any characteristic function  $g = \chi_A$  on a measurable set, there exists a sequence of step functions converging to it point-wise almost everywhere. Since from Proposition 29, there exists a sequence of simple functions converging to  $f$ , and since a simple function  $h = \sum_{i=1}^M \alpha_i \chi_{E_i}$  is a finite combination of characteristics functions over measurable sets, therefore there exists a sequence of step functions converging to  $f$  almost everywhere.

In particular if  $\psi_n^i \longrightarrow \chi_{E_i}$  almost everywhere, then  $\sum_{i=1}^M \alpha_i \psi_n^i \longrightarrow \sum_{i=1}^M \alpha_i \chi_{E_i} = h$ . Now by Proposition 29, there exists the sequence  $\{h_n\}$  of simple functions converging to  $f$ . Since

$$K_n = \left\{ \sum_{i=1}^{M_n} \alpha_i \psi_n^i \right\} \longrightarrow h_n \text{ almost everywhere,}$$

where note that  $K_n = \sum_{i=1}^{M_n} \alpha_i \psi_n^i$  is a step function because  $\psi_n^i$  is a step function and there are finitely many ( $M_n$ ) of them, and

$$\{h_n\} \longrightarrow f$$


therefore,

$$K_n \longrightarrow f \text{ almost everywhere.}$$

Hence proved. ■

### 3.4 Egorov's Theorem

We now discuss a very important result in the theory of measurable functions named after Dmitri Fyodorovich Egorov, who published this result in 1911, thus establishing **a condition required for uniform convergence of a point-wise convergent sequence of Measurable functions**.

 **Theorem 3. (Egorov's Theorem)** Let  $(\mathbb{R}, \mathcal{M}_{\lambda^*}, \lambda)$  be the Lebesgue measure space on  $\mathbb{R}$ . Suppose  $\{f_k\}$  is a sequence of real-valued, Lebesgue Measurable functions on  $E \in \mathcal{M}_{\lambda^*}$  where  $\lambda(E) < +\infty$ . If

$$f_k \longrightarrow f \text{ pointwise on } E,$$

<sup>17</sup> Then for each  $\epsilon > 0$ , there exists a **closed** set  $A_\epsilon \subset E$  such that

1.  $\lambda(E \setminus A_\epsilon) < \epsilon$ , and
2.  $f_k \longrightarrow f$  **uniformly** on  $A_\epsilon$ .

*Proof.* We break down the proof in the following 3 parts.

**Act 1. A Basic Construction.**

For each pair of integers  $n, k$ , construct the following set:

$$E_k^n = \left\{ x \in E : |f_j(x) - f(x)| < \frac{1}{n}, \forall j > k \right\}.$$

Now, fix  $n$ , so that we have the following observations:

$$E_k^n \subseteq E_{k+1}^n \tag{11}$$

and since  $f_k \longrightarrow f$  point-wise, therefore

$$\lim_{k \rightarrow \infty} \bigcup_{i=1}^k E_i^n = E. \tag{12}$$

Hence

$$\lambda(E \setminus E_k^n) \longrightarrow 0 \text{ as } k \rightarrow \infty.$$

Note that the above result utilizes the fact that  $\lambda(E) < +\infty$ . Now by the above, we can say that  $\exists k_n$  such that

$$\lambda(E \setminus E_{k_n}^n) < \frac{1}{2^n}$$

which, by definition of  $E_k^n$  implies that

$$|f_j(x) - f(x)| < \frac{1}{n} \text{ whenever } j > k_n \text{ and } x \in E_{k_n}^n.$$

**Act 2. Constructing  $A_\epsilon$ .**

Now choose  $N \in \mathbb{N}$  such that

$$\sum_{n=N}^{\infty} \frac{1}{2^n} < \frac{\epsilon}{2}$$

and define

$$\tilde{A}_\epsilon = \bigcap_{n=N}^{\infty} E_{k_n}^n \tag{13}$$

---

<sup>17</sup>From Proposition 21, the limit of a sequence of Measurable functions is also Measurable, hence there's no point in writing extraneously the requirement for  $f$  to be also Measurable.

We now observe that

$$\begin{aligned}
\lambda(E \setminus \tilde{A}_\epsilon) &= \lambda\left(E \cap \bigcup_{n=N}^{\infty} (E_{k_n}^n)^c\right) \\
&= \lambda\left(\bigcup_{n=N}^{\infty} E \cap (E_{k_n}^n)^c\right) \\
&\leq \sum_{n=N}^{\infty} \lambda(E \cap (E_{k_n}^n)^c) \\
&= \sum_{n=N}^{\infty} \lambda(E \setminus E_{k_n}^n) \\
&< \sum_{n=N}^{\infty} \frac{1}{2^n} \\
&< \frac{\epsilon}{2}
\end{aligned}$$

**Act 3.** *Finalé.*

We now claim and prove the following:

Claim :  $f_k \longrightarrow f$  uniformly on  $\tilde{A}_\epsilon$ .

For this, let  $\delta > 0$  and choose  $n' \geq N$  such that  $\frac{1}{n'} < \delta$ . Then

$$\text{if } x \in \tilde{A}_\epsilon \implies x \in E_{k_{n'}}^{n'} \implies |f_j(x) - f(x)| < \frac{1}{n'} < \delta, \quad \forall j > k_{n'}. \quad (14)$$

Note that this is just the definition of uniform convergence.

Finally, note that  $E_k^n$  is a Lebesgue measurable set due to Proposition 18, Statement 5. Hence,  $\tilde{A}_\epsilon$  is measurable. Now, by Proposition 16, Statement 2, there exists a closed set  $A_\epsilon \subset \tilde{A}_\epsilon$  such that

$$\lambda(\tilde{A}_\epsilon \setminus A_\epsilon) < \frac{\epsilon}{2}.$$


Now,

$$\begin{aligned}
\epsilon &> \lambda(E \setminus \tilde{A}_\epsilon) + \lambda(\tilde{A}_\epsilon \setminus A_\epsilon) \\
&\geq \lambda(E \setminus \tilde{A}_\epsilon \cup \tilde{A}_\epsilon \setminus A_\epsilon) \\
&= \lambda(E \setminus A_\epsilon).
\end{aligned}$$

Now, by (14), we see that  $f_k \longrightarrow f$  uniformly for all  $x \in A_\epsilon \subset \tilde{A}_\epsilon$  such that  $\lambda(E \setminus A_\epsilon) < \epsilon$  and  $A_\epsilon$  is closed. Proof is now complete. ■

### 3.5 Luzin's Theorem

The following is the final important result on the basic theory of measurable functions, attributed to Nikolai Nikolaevich Luzin who penned this theorem around 1912.

 **Theorem 4. (Lusin's Theorem)** Consider the Lebesgue measure space  $(\mathbb{R}, \mathcal{M}_{\lambda^*}, \lambda)$ . Suppose  $f$  is a real-valued, Lebesgue Measurable function defined over a Lebesgue measurable set  $E$  with finite measure. Then for all  $\epsilon > 0$ , there exists a **closed** set  $F_\epsilon \subset E$  with

1.  $\lambda(E \setminus F_\epsilon) < \epsilon$ , and
2. The restriction  $f|_{F_\epsilon}$  of  $f$  over  $F_\epsilon$  is **continuous**.

*Proof.* From the Proposition 32, we have a sequence  $\{f_n\}$  of step functions such that

$$f_n \longrightarrow f \text{ almost everywhere.}$$

Now, consider, for example the characteristic function over an interval  $\chi_{[a,b]}$ . Then, we can define a function  $\phi(x)$  for any  $\delta > 0$  as follows:

$$\phi(x) = \begin{cases} 0 & x < a \\ \frac{x-a}{\delta/2} & a \leq x \leq a + \frac{\delta}{2} \\ 1 & a + \frac{\delta}{2} \leq x \leq b - \frac{\delta}{2} \\ \frac{b-x}{\delta/2} & b - \frac{\delta}{2} \leq x \leq b \\ 0 & x > b \end{cases}$$

which then satisfies

$$\{x \in \mathbb{R} \mid \phi(x) \neq \chi_{[a,b]}\} = \left(a, a + \frac{\delta}{2}\right) \cup \left(b - \frac{\delta}{2}, b\right)$$

which then implies that,

$$\lambda \left( \{x \in \mathbb{R} \mid \phi(x) \neq \chi_{[a,b]}\} \right) = \lambda \left( \left(a, a + \frac{\delta}{2}\right) \cup \left(b - \frac{\delta}{2}, b\right) \right) = \delta.$$

Note that  $\phi(x)$  is also continuous over all  $\mathbb{R}$ . Hence, for any step function (finite sum of  $\chi_{[a,b]}$ -type functions) and  $\delta > 0$ , one can construct a continuous function which does not satisfies with the step function on a set with measure  $< \delta$ .

Hence, for step-functions  $\{f_n\}$ , corresponding to each  $f_n$ ,  $\exists$  a continuous function  $\phi_n$  and a set  $E_n$  such that

$$E_n = \{x \mid \phi_n(x) \neq f_n(x)\} \text{ and } \lambda(E_n) < \frac{1}{2^n}.$$

Now, for all  $\epsilon > 0$ , there exists a  $N \in \mathbb{N}$  such that

$$\sum_{n \geq N} \frac{1}{2^n} < \frac{\epsilon}{3}.$$

With the above fact, construct the set  $F'$  as follows:

$$F' = \left( A_{\frac{\epsilon}{3}} \setminus \bigcup_{n \geq N} E_n \right)$$

where  $A_{\frac{\epsilon}{3}}$  is the closed subset  $A_{\frac{\epsilon}{3}} \subset E$  such that **1.**  $\lambda(E \setminus A_{\frac{\epsilon}{3}}) < \frac{\epsilon}{3}$  and **2.**  $f_n \rightarrow f$  **uniformly** on  $A_{\frac{\epsilon}{3}}$ . This is guaranteed by Theorem 3 (Egorov's Theorem).

Note that  $f_n|_{F'}$  is **continuous**  $\forall n \geq N$  because for any  $x \in F' \implies \phi_n(x) = f_n(x) \forall n \geq N$  and since  $\phi_n$  are already continuous  $\forall n \in \mathbb{N}$ .

Furthermore, since  $F' \subset E$  and  $f_n \rightarrow f$  uniformly, then the restriction  $f_n|_{F'}$  is continuous and converges uniformly to  $f|_{F'}$ , which **due to uniform convergence, is also continuous!**

Now, note that  $E_n$ 's are measurable sets due to Proposition 18, Statement 5. Similarly, since  $A_{\frac{\epsilon}{3}}$  is closed, therefore it is also measurable. Hence,  $F'$  is measurable.

Now by Proposition 16, there exists a closed set  $F_\epsilon \subset F'$  such that  $\lambda(F' \setminus F_\epsilon) < \frac{\epsilon}{3}$ . Note that because  $F_\epsilon \subset F'$  and  $f|_{F'}$  is continuous, therefore the restriction  $f|_{F_\epsilon}$  is also continuous.

Finally, combining

1.  $\sum_{n \geq N} \frac{1}{2^n} < \frac{\epsilon}{3}$ ,
2.  $\lambda(E \setminus A_{\frac{\epsilon}{3}}) < \frac{\epsilon}{3}$ ,
3.  $\lambda(F' \setminus F_\epsilon) < \frac{\epsilon}{3}$

it can be easily seen that

$$(E \setminus A_{\frac{\epsilon}{3}}) \cup (F' \setminus F_\epsilon) = E \setminus F_\epsilon$$

Hence,

$$\begin{aligned} \lambda(E \setminus F_\epsilon) &= \lambda \left( (E \setminus A_{\frac{\epsilon}{3}}) \cup (F' \setminus F_\epsilon) \right) \\ &\leq \lambda(E \setminus A_{\frac{\epsilon}{3}}) + \lambda(F' \setminus F_\epsilon) \\ &< \frac{2\epsilon}{3} \\ &< \epsilon. \end{aligned}$$

which completes the proof. ■



## 4 Integration of Measurable Functions

Let's first remind ourselves of the basic definition of a Riemann Integrable function. If we say that the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is Riemann Integrable, then the integral of  $f$  on  $[a, b]$ , written as  $\int_a^b f$ , is given by the following two constructions on a partition  $P$  of  $[a, b]$ ,

- Lower Sum :

$$L(f, P) = \sum_i m_i (a_i - a_{i-1}) \text{ where } m_i = \inf_{x \in [a_{i-1}, a_i]} f(x)$$

- Upper Sum:

$$U(f, P) = \sum_i M_i (a_i - a_{i-1}) \text{ where } M_i = \sup_{x \in [a_{i-1}, a_i]} f(x)$$

so that

$$\int_a^b f = L(f, P) = U(f, P).$$

The chain of observations begins now. One can easily write the Lower and Upper Sum as the following simple functions (remember that the partition is finitely many)

$$\begin{aligned} L(f, P) &= \sum_i m_i \lambda([a_{i-1}, a_i]) \\ U(f, P) &= \sum_i M_i \lambda([a_{i-1}, a_i]) \end{aligned}$$

Or, equivalently, we can define a lower step function as follows:

$$\phi_P = \sum_i m_i \chi_{[a_{i-1}, a_i]}$$

so that the Riemann Integral is simply

$$\int_a^b f(x) dx = \sup_P \int_a^b \phi_P(x) dx$$

where supremum is defined over all partitions. But since, by definition,  $\phi_P(x) \leq f(x) \forall x \in \mathbb{R}$ , we can alternatively define Riemann Integral as

$$\int_a^b f(x) dx = \sup_{\phi_P \leq f} \int_a^b \phi_P(x) dx \tag{15}$$

where the supremum is defined for all step functions on any partition  $P$ .

This definition presented in (15) provides the motivation for extending the definition of Integration from Riemann to Lebesgue. In particular, note the definition of  $\phi_P$ , usual measure on the intervals is applied in Riemann's definition. But, since we know that Borel  $\sigma$ -algebra is a proper subset of  $\mathcal{M}_{\lambda^*}$ , then it just makes sense to replace  $a_{i-1} - a_i$  by  $\lambda([a_{i-1}, a_i])$  in the motivation that it might generalize the notion of integration.

Therefore, the following definitions of integration for non-negative functions are motivated:

## 4.1 Integration of Non-Negative Measurable Functions

★ **Definition 22. (Lebesgue Integral of a Simple Function)** Consider  $\phi : \mathbb{R} \rightarrow [0, +\infty)$  be a simple function as

$$\phi = \sum_{i=1}^N \alpha_i \chi_{E_i} \text{ where } \alpha_i \geq 0 \text{ and } \lambda(E_i) < +\infty$$

Then, the Lebesgue integral of  $\phi$  is defined as

$$\int \phi dx = \sum_{i=1}^N \alpha_i \lambda(E_i)$$

★ **Definition 23. (Lebesgue Integral of a Measurable Function)** Suppose  $f : \mathbb{R} \rightarrow [0, +\infty)$  is a  $\lambda$ -Measurable function, then the Lebesgue Integral of  $f$  is defined as

$$\star \int f dx = \sup_{\phi \leq f} \int \phi dx \text{ where } \phi \text{ are the simple functions } \leq f. \star$$

★ **Definition 24. (Lebesgue Integral over a measurable set)** Consider  $f : \mathbb{R} \rightarrow [0, +\infty)$  to be a  $\lambda$ -Measurable function and  $E \subseteq \mathbb{R}$  is Lebesgue measurable. Then,

$$\int_E f dx = \int f \cdot \chi_E dx$$

**Remark.** Therefore, the integral of a non-negative measurable function over a measurable set is given by the integral<sup>18</sup> of restriction of  $f$  to it and zero otherwise.

✓ **Proposition 33.** Consider the two  $\lambda$ -Measurable functions  $f, g : \mathbb{R} \rightarrow [0, +\infty)$  and  $\phi : \mathbb{R} \rightarrow [0, +\infty)$  be a simple-function, then the Lebesgue Integral has the following properties:

1. Consider two Lebesgue measurable subsets  $A$  and  $B$  of  $\mathbb{R}$  such that  $A \cap B = \emptyset$ . Then,

$$\int_{A \cup B} \phi dx = \int_A \phi dx + \int_B \phi dx.$$

2. For any  $\alpha \in \mathbb{R}$ ,

$$\int \alpha f dx = \alpha \int f dx.$$

3. Integration for positive valued Measurable functions is therefore distributive:

$$\int (f + g) dx = \int f dx + \int g dx.$$

4. If  $f(x) \leq g(x)$  holds for all  $x \in \mathbb{R}$ , then

$$\int f dx \leq \int g dx.$$

5. Consider  $A$  and  $B$  be Lebesgue measurable subsets of  $\mathbb{R}$  such that  $A \subseteq B$ . Then,

$$\int_A f dx \leq \int_B f dx.$$

*Proof.* **Part 1:** Since  $\phi$  is simple, therefore we can write

$$\phi = \sum_{i=1}^N \alpha_i \chi_{E_i}.$$

<sup>18</sup>From now on, any instance of *integral* should be presupposed by Lebesgue integral, of-course, unless otherwise stated, in this text.

Now, by definition

$$\begin{aligned}
\int_{A \cup B} \phi dx &= \sum_{i=1}^N \alpha_i \lambda(E_i \cap (A \cup B)) \\
&= \sum_{i=1}^N \alpha_i \lambda((E_i \cap A) \cup (E_i \cap B)) \\
&= \sum_{i=1}^N \alpha_i \lambda(E_i \cap A) + \alpha_i \lambda(E_i \cap B) && \because E_i \cap A \text{ and } E_i \cap B \text{ are disjoint.} \\
&= \int_A \phi dx + \int_B \phi dx
\end{aligned}$$

**Part 2 & 3 :** Can be seen easily from Theorem 5.

**Part 4 :** Note that we define

$$\int f dx = \sup_{\phi \leq f} \int \phi dx \text{ where } \phi \text{ are simple functions.}$$

Therefore, for any  $\phi \leq f$ , due to given condition  $f \leq g$ , we would have  $\phi \leq g$ . Hence,

$$\int \phi dx \leq \int g dx$$

Since this is true for all simple  $\phi \leq f$ , therefore  $\sup_{\phi \leq f} \int \phi dx \leq \int g dx$ , proving the result.

**Part 5 :** Consider the following:

$$\begin{aligned}
\int_A f dx &= \int f \chi_A dx \\
&\leq \int f \chi_B dx && \because \chi_A \leq \chi_B, \text{ then apply 4.} \\
&= \int_B f dx
\end{aligned}$$

Hence proved. ■

## 4.2 Monotone Convergence Theorem

This is arguably one of the most important theorem in Integration theory,

⊗ **Theorem 5. (Monotone Convergence Theorem)** Consider a sequence  $\{f_n\}$  of  $\mathbb{R} \rightarrow [0, +\infty)$  of  $\lambda$ -Measurable functions which satisfies

$$f_n(x) \leq f_{n+1}(x) \quad \forall x \in \mathbb{R} \text{ and } n$$

and suppose  $\lim_{n \rightarrow \infty} f_n$  exists. Then,

$$\int \lim_{n \rightarrow \infty} f_n = \lim_{n \rightarrow \infty} \int f_n.$$

*Proof.* Since  $f_n \leq f_{n+1}$ , therefore,

$$\int f_n \leq \int f_{n+1} \leq \int \lim_{n \rightarrow \infty} f_n.$$

Hence,

$$\lim_{n \rightarrow \infty} \int f_n \leq \int \lim_{n \rightarrow \infty} f_n.$$

Therefore we have proved one inequality.

Now to prove the other inequality, consider any simple function  $\phi \leq \lim_{n \rightarrow \infty} f_n$ . If we can show that  $\int f_n \geq \int \phi$  for any  $n \in \mathbb{N}$ , then we are done. To this goal, consider  $\alpha \in (0, 1)$ . Construct the set<sup>19</sup>

$$E_n = \{x \mid f_n(x) \geq \alpha \phi(x)\}.$$

<sup>19</sup>After reading the proof, it should appear striking to the reader on actually how much the proof depends on this construction. Both the claims in the following page utilizes this construction  $E_n$  to full extent! Hence, it is advised (by Instructor) to pursue such effective constructions in the problem sheets and your own proofs.

Clearly,

$$E_n \subseteq E_{n+1} \quad \forall n \in \mathbb{N}.$$

Now,

$$\int f_n \geq \int_{E_n} f_n \geq \alpha \int_{E_n} \phi. \quad (16)$$

Moreover, we can see that

$$\text{Claim 1: } \bigcup_n E_n = \mathbb{R}.$$

This is easy to see as follows:

$$\text{Take } x \in \bigcup_n E_n \implies x \in E_{i_0} \text{ for some } i_0 \in \mathbb{N}.$$

$$\implies x \in \mathbb{R}$$

$\because E_n$  are subsets of  $\mathbb{R}$ .

$$\text{Take } x \in \mathbb{R} \implies \text{Either (1) } x \in \{x \mid f_n(x) - \alpha\phi(x) \geq 0\} \text{ or (2) } x \in \{x \mid f_n(x) - \alpha\phi(x) < 0\} \text{ for any } n \in \mathbb{N}.$$

$$\implies \text{If (1), then } x \in E_n, \text{ else if (2), then } \because \phi \leq \lim_{n \rightarrow \infty} f_n, \exists n' \text{ s.t. } x \in E_{n'}$$

$$\implies x \in \bigcup_n E_n.$$

Hence Claim 1.

Next, we can also see that

$$\text{Claim 2: } \int_{E_n} \phi \longrightarrow \int \phi$$

This can be seen by expanding the *simplicity* of  $\phi$  as follows:

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{E_n} \phi &= \sum_{i=1}^N a_i \lim_{n \rightarrow \infty} \lambda(A_i \cap E_n) \\ &= \sum_{i=1}^N a_i \lambda\left(\bigcup_n A_i \cap E_n\right) && \because \{A_i \cap E_n\}_n \text{ is increasing, use Proposition 7.} \\ &= \sum_{i=1}^N a_i \lambda\left(A_i \cap \bigcup_n E_n\right) \\ &= \sum_{i=1}^N a_i \lambda(A_i \cap \mathbb{R}) && \text{Claim 1.} \\ &= \sum_{i=1}^N a_i \lambda(A_i) = \int \phi && \text{Hence Claim 2.} \end{aligned}$$

Finally, take limit in (16) to get:

$$\begin{aligned} \lim_{n \rightarrow \infty} \int f_n &\geq \lim_{n \rightarrow \infty} \int_{E_n} f_n \geq \alpha \lim_{n \rightarrow \infty} \int_{E_n} \phi \\ &= \alpha \int \phi && \text{Claim 2.} \\ \lim_{n \rightarrow \infty} \int f_n &\geq \int \phi && \because 0 < \alpha < 1 \text{ is arbitrary.} \end{aligned}$$

Hence, **for any** simple function  $\phi \leq \lim_{n \rightarrow \infty} f_n$ , we have concluded that  $\int \phi \leq \lim_{n \rightarrow \infty} \int f_n$ , hence it must be true that

$$\int \lim_{n \rightarrow \infty} f_n = \sup_{\phi \leq \lim_{n \rightarrow \infty} f_n} \int \phi \leq \lim_{n \rightarrow \infty} \int f_n.$$

Combining the converse inequality at the beginning, we hence get the desired result. ■

✓ **Proposition 34.** Consider a Lebesgue Measurable function  $f : \mathbb{R} \rightarrow [0, +\infty)$ . Then,

$$\int f dx = 0 \iff f \equiv 0 \text{ almost everywhere.}$$

*Proof.* **L**  $\implies$  **R** : Consider  $f$  is a non-negative real-valued function whose integral is zero. Construct the set,

$$E_n = \left\{ x \mid f(x) \geq \frac{1}{n} \right\}.$$

In order to show that  $f \equiv 0$  almost everywhere, it is hence sufficient to show that  $\lambda(E_n) = 0 \forall n \in \mathbb{N}$  because it equivalently proves that the measure of the set where  $f$  is greater than zero is zero.

Now, consider the following function

$$g_n = \frac{1}{n} \chi_{E_n}.$$

Clearly, because  $g_n$  is a simple function and

$$\frac{1}{n} \chi_{E_n}(x) = \begin{cases} \frac{1}{n} & \text{if } f(x) \geq \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$$

which clearly means that  $\frac{1}{n} \chi_{E_n} \leq f$ , therefore,

$$\begin{aligned} \int f &= 0 = \sup_{\phi \leq f} \int \phi \\ &\geq \int \frac{1}{n} \chi_{E_n} \\ &= \frac{1}{n} \lambda(E_n) \\ \implies \lambda(E_n) &= 0 \forall n \in \mathbb{N}. \end{aligned}$$

**R**  $\implies$  **L** : If a non-negative real-valued Measurable function  $f$  is 0 almost everywhere, then for any simple function  $\phi \leq f$ ,  $\phi$  must also be 0 almost everywhere, so that

$$\begin{aligned} \int \phi &= \sum_{i=1}^N \alpha_i \lambda(E_i) \\ &= 0 \end{aligned}$$

Since this is true for any simple  $\phi \leq f$ , therefore the supremum of all such  $\int \phi$  must also be zero, to make  $\int f = 0$ . ■

---

### 4.3 Fatou's Lemma



**Theorem 6.** If  $\{f_n\}$  is a sequence of Lebesgue Measurable functions from  $\mathbb{R}$  to  $[0, +\infty)$ , then,

$$\int \liminf_n f_n \leq \liminf_n \int f_n.$$

*Proof.* We will use Monotone Convergence Theorem to prove this result. Define

$$g_k = \inf_{n \geq k} f_n$$

Therefore  $g_k \leq g_{k+1}$  with  $g_k \leq f_n \forall n \geq k$ . Then,

$$\int g_k \leq \int f_n \forall n \geq k.$$

This implies that

$$\int g_k \leq \inf_{n \geq k} \int f_n.$$

Now, by MCT,

$$\int \lim_k g_k = \lim_k \int g_k$$

Therefore

$$\begin{aligned} \lim_{k \rightarrow \infty} \inf_{n \geq k} \int f_n &= \lim_k \inf_k \int f_k \geq \lim_{k \rightarrow \infty} \int g_k \\ &= \int \lim_{k \rightarrow \infty} g_k \\ &= \int \lim_{k \rightarrow \infty} \inf_{n \geq k} f_n \\ &= \int \liminf_k f_k \end{aligned}$$

Hence Proved. ■

**Remark.** In fact,

Fatou's Lemma  $\iff$  Monotone Convergence Theorem.

---

## 4.4 Integration of General Real-Valued Measurable Functions

With the notion of integration of non-negative Measurable function in place, it's not difficult to see how can one extend the same notion to Measurable functions which takes value in the whole real line.

★ **Definition 25. (Lebesgue Integral of a Real-Valued Measurable Function)** Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$  to be a Measurable function such that

1.  $\int f^+ dx < \infty$ , and
2.  $\int f^- dx < \infty$ .

If the above two conditions are satisfied, then  $f$  is called Lebesgue Integrable. Then, the Lebesgue Integral of  $f$  is defined as

$$\star \boxed{\int f dx = \int f^+ dx - \int f^- dx} \star$$

**Remark.** It's important to note that the Integral  $\int f dx = \int f^+ dx - \int f^- dx$  is easily defined for any measurable function, but  $f$  is called Lebesgue Integral only when it's value is finite!

★ **Definition 26. (Lebesgue Integral over a Measurable Set)** Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$  is Measurable,  $f \cdot \chi_E$  is an Lebesgue Integrable function and  $E \subseteq \mathbb{R}$  is also measurable. Then,

$$\int_E f dx = \int f \cdot \chi_E dx.$$

### 4.4.1 Basic Properties of General Lebesgue Integral

The following properties are direct extensions of Proposition 33 to the bigger class of Lebesgue Integrable functions.

✓ **Proposition 35.** Consider  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  to be Lebesgue Integrable functions. Then,

1. For any  $\alpha \in \mathbb{R}$ , we have:

$$\int \alpha f dx = \alpha \int f dx.$$

2.  $f + g$  is also Lebesgue Integrable, with

$$\int (f + g) dx = \int f dx + \int g dx.$$

3. If  $f \equiv 0$  almost everywhere on  $\mathbb{R}$ , then,

$$\int f dx = 0.$$

4. If  $f \leq g$  almost everywhere on  $\mathbb{R}$ , then,

$$\int f dx \leq \int g dx.$$

5. If  $A$  and  $B$  are measurable sets such that  $A \cap B = \Phi$ , then,

$$\int_{A \cup B} f dx = \int_A f dx + \int_B f dx.$$

*Proof.* **S1:** Consider the case that  $\alpha \geq 0$ . Then,

$$\begin{aligned} (\alpha f)^+ &= \max(\alpha f, 0) = \alpha \max(f, 0) = \alpha f^+ \\ (\alpha f)^- &= -\min(\alpha f, 0) = -\alpha \min(f, 0) = \alpha f^- \end{aligned}$$

and since  $\int f^+ dx < \infty$  and  $\int f^- dx < \infty$ , therefore  $\alpha f$  is also Lebesgue Integrable, with the integral given as

$$\begin{aligned} \int \alpha f &= \int \alpha f^+ - \int \alpha f^- \\ &= \alpha \left( \int f^+ - \int f^- \right) \\ &= \alpha \int f \end{aligned}$$

Now consider that  $\alpha < 0$ , then

$$\begin{aligned}(\alpha f)^+ &= \max(\alpha f, 0) = -|\alpha| \min(f, 0) = |\alpha| f^- \\(\alpha f)^- &= -\min(\alpha f, 0) = |\alpha| \max(f, 0) = |\alpha| f^+\end{aligned}$$

Hence,  $\alpha f$  is again Lebesgue Integrable, with the integral calculated as:

$$\int \alpha f = \int (\alpha f)^+ - \int (\alpha f)^- = |\alpha| \left( \int f^- - \int f^+ \right) = -|\alpha| \int f = \alpha \int f.$$

**S2** : First,

$$\begin{aligned}(f+g)^+ &\leq f^+ + g^+ \\(f+g)^- &\leq f^- + g^-\end{aligned}$$

for all  $x \in \mathbb{R}$ , so that  $f+g$  is Lebesgue Integrable. Now,

$$\begin{aligned}f+g &= (f+g)^+ - (f+g)^- \\&= f^+ - f^- + g^+ - g^-\end{aligned}$$

therefore,

$$\begin{aligned}(f+g)^+ - (f+g)^- &= f^+ - f^- + g^+ - g^- \\(f+g)^+ + f^- + g^- &= (f+g)^- + f^+ + g^+ \\\int (f+g)^+ + f^- + g^- &= \int (f+g)^- + f^+ + g^+ \\\int (f+g)^+ + \int f^- + \int g^- &= \int (f+g)^- + \int f^+ + \int g^+ \quad (\because \text{of Proposition 33, S3.}) \\\int (f+g)^+ - \int (f+g)^- &= \int f^+ - \int f^- + \int g^+ - \int g^- \\\int (f+g)^+ - \int (f+g)^- &= \int f + \int g\end{aligned}$$

**S3** : Given to us is that  $f \equiv 0$  almost everywhere. This means that

$$\{x \in \mathbb{R} \mid f(x) \neq 0\} \text{ is of measure } 0.$$

We can write it equivalently as the union of the following two disjoint sets

$$\begin{aligned}\{x \in \mathbb{R} \mid f(x) \neq 0\} &= \{x \mid f(x) > 0\} \cup \{x \in \mathbb{R} \mid f(x) < 0\}. \\\lambda(\{x \in \mathbb{R} \mid f(x) \neq 0\}) &= \lambda(\{x \mid f^+(x) > 0\}) + \lambda(\{x \mid f^-(x) < 0\}) \\&= 0\end{aligned}$$

Since measure is positive valued by definition, therefore, these two have to be individually be zero. That is,

$$\lambda(\{x \mid f^+(x) > 0\}) = \lambda(\{x \mid f^-(x) < 0\}) = 0$$

Now, by Proposition 34, we get that

$$\int f^+ = \int f^- = 0$$

which implies that

$$\int f = \int f^+ - \int f^- = 0.$$

**S4** :

■

✓ **Proposition 36.** If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a Lebesgue Integrable Function, then,

$$\left| \int f dx \right| \leq \int |f| dx$$

*Proof.*

■



## 4.5 Dominated Convergence Theorem

**Theorem 7.** Let  $\{f_n\}$  be a sequence of Measurable functions such that there exists a Lebesgue Integrable function  $g$  which satisfies

$$|f_n| \leq g \quad \forall n.$$

Suppose that the limit  $\lim_{n \rightarrow \infty} f_n$  exists. Then  $\lim_{n \rightarrow \infty} f_n$  is Lebesgue Integrable and,

$$\lim_{n \rightarrow \infty} \int f_n dx = \int \lim_{n \rightarrow \infty} f_n dx.$$

*Proof.* Since  $\{f_n\}$  is a sequence of Measurable functions, therefore,  $\lim_{n \rightarrow \infty} f_n = f$  is also Measurable and  $|f|$  is bounded by  $g$ . But since  $g$  is Lebesgue Integrable, and  $f_n$  and  $f$  are bounded by  $g$ , then each  $f_n$  and also  $f$  are also Lebesgue Integrable (Trivial to see).

Now,  $\{g + f_n\}$  is a sequence of Measurable functions. Moreover, since  $f_n \leq g$  for all  $n$ , therefore  $f_n + g \geq 0$ , so that  $\{f_n + g\}$  is a sequence of non-negative Measurable functions.

Now using Fatou's Lemma (Theorem 6), we get,

$$\begin{aligned} \int \liminf_n (g + f_n) dx &\leq \liminf_n \int (g + f_n) dx \\ \int (g + \liminf_n f_n) dx &\leq \int g dx + \liminf_n \int f_n dx \\ \int g dx + \int \liminf_n f_n dx &\leq \int g dx + \liminf_n \int f_n dx \\ \int \liminf_n f_n dx &\leq \liminf_n \int f_n dx && \because g \text{ is L.I., so } \int g dx < \infty \\ \int f dx &\leq \liminf_n \int f_n dx && \because \limsup_n x_n = \liminf_n x_n = \lim_n x_n. \end{aligned}$$

Similarly, since  $\{g - f_n\}$  is also a sequence of non-negative Measurable functions, therefore we can use Fatou's Lemma to conclude:

$$\begin{aligned} \int \liminf_n (g - f_n) dx &\leq \liminf_n \int (g - f_n) dx \\ \int \liminf_n (-f_n) dx &\leq \liminf_n \left( - \int f_n dx \right) \\ - \int \limsup_n f_n dx &\leq - \limsup_n \int f_n dx && \because \liminf_n (-x_n) = - \limsup_n x_n. \\ \int f dx &\geq \limsup_n \int f_n dx \end{aligned}$$

We hence have that

$$\limsup_n \int f_n dx \leq \int f dx \leq \liminf_n \int f_n dx$$

But it is also true that

$$\liminf_n \int f_n dx \leq \limsup_n \int f_n dx.$$

Hence,

$$\liminf_n \int f_n dx = \limsup_n \int f_n dx = \lim_n \int f_n dx = \int f dx$$

Hence proved. ■

✓ **Proposition 37.** Consider  $\{f_n\}$  to be a sequence of Lebesgue Integrable functions such that

$$\sum_{n=1}^{\infty} \int |f_n| dx < \infty.$$

Then,

1. The series

$$\sum_{n=1}^{\infty} f_n(x) \text{ converges almost everywhere on } \mathbb{R}.$$

2. The sum

$$f = \sum_{n=1}^{\infty} f_n \text{ is Lebesgue Integrable.}$$

3. The integral is

$$\int \sum_{n=1}^{\infty} f_n dx = \sum_{n=1}^{\infty} \int f_n dx.$$

*Proof.* **S1.** Denote the following:

$$\varphi = \sum_{n=1}^{\infty} |f_n|$$

Clearly,  $\varphi$  is a non-negative Measurable function. Since we know that Lebesgue Integral for non-negative functions is countably additive, therefore,

$$\int \varphi dx = \sum_{n=1}^{\infty} \int |f_n| dx < \infty.$$

Now, if  $\int \varphi < \infty$ , then  $\varphi$  is finite almost everywhere on  $\mathbb{R}$ <sup>20</sup>. Now, since  $\sum_{n=1}^{\infty} f_n$  is absolutely convergent almost everywhere (last line), hence it is convergent almost everywhere too on  $\mathbb{R}$ .

**S2.** Since  $|\sum_{n=1}^{\infty} f_n| \leq \sum_{n=1}^{\infty} |f_n| = \varphi < \infty$  (almost everywhere) and since we can modify the set where  $\varphi$  is not defined (infinite) arbitrarily to make a new function which would be Measurable and equal to  $\sum_{n=1}^{\infty} f_n$  almost everywhere, therefore  $\sum_{n=1}^{\infty} f_n$  would be Measurable.

**S3.** Define

$$\phi_n = \sum_{i=1}^n f_i.$$

Clearly,  $\phi_n \leq |\sum_{i=1}^n f_i| \leq \sum_{i=1}^n |f_i| \leq \sum_{i=1}^{\infty} |f_i| = \varphi$ . Therefore,  $\phi_n$  is a sequence of Measurable functions and  $\phi_n \leq \varphi$  where  $\varphi$  is an Integrable function (given). Therefore, using Dominated Convergence Theorem (7), we get,

$$\begin{aligned} \lim_n \int \phi_n dx &= \int \lim_n \phi_n dx \\ \lim_n \int \sum_{i=1}^n f_i dx &= \int \lim_n \sum_{i=1}^n f_i dx \\ \lim_n \sum_{i=1}^n \int f_i dx &= \int \sum_{i=1}^{\infty} f_i dx && \because \text{Proposition 35, S2} \\ \sum_{i=1}^{\infty} \int f_i dx &= \int \sum_{i=1}^{\infty} f_i dx. \end{aligned}$$

■

Hence proved.

<sup>20</sup>For a non-negative Measurable function  $f$  with given that  $\int f dx < \infty$ , the set  $E = \{x \in \mathbb{R} \mid f(x) = \infty\}$  together with supposition that  $\lambda(E) > 0$  is such that; since  $\int f dx = \sup_{\phi \leq f} \int \phi$ , therefore, if we take  $\phi = n\chi_E$  for any  $n > 0$  then  $n\chi_E < f$ . Hence  $\int f dx > n\lambda(E)$  for all  $n$ , so that  $\int f dx = \infty$ . But it's a contradiction to  $\int f dx < \infty$ . Therefore  $\lambda(E) = 0$ .

# References

- [1] Solovay, R. (1970). A Model of Set-Theory in Which Every Set of Reals is Lebesgue Measurable. *Annals of Mathematics*, 92(1), second series, 1-56.