How was find all formulas of maths

This book belong to:

A free owl project

Index

Note: This book is in developing

1. Initial part **5**
- **1.1. Introduction** **5**
  - **1.1.1. A free owl project** **5**
  - **1.1.2. Book** **5**
- **1.2. Error Zone** **6**

2. Algebra **7**
- **2.1. Linear algebra** **7**
  - **2.1.1. Summation** **7**
- **2.2. Elementary algebra** **8**
  - **2.2.1. Notable Identities** **8**
  - **2.2.2. Property of Fractions** **9**

3. Mathematical analysis **11**
- **3.1. Calculus** **11**
  - **2.1.1. Limits** **11**
  - **2.1.2. Derivative** **12**
  - **2.1.3. Integral** **15**

4. Final Part **13**
- **4.1. Reference** **13**
- **4.2. Make by** **13**

Chapter 1

Initial part

1.1 Introduction

1.1.1 A free owl project

Goal

Collect all the knowledge of the world in only one site

Differentiation of other projects

- This project is Open Source, to protect the books of this project of malicious changes the people that want to collaborate, they have to create a pull request that is like a proposal of a modification that have to be accept you can send it by email `lucasvarelacorrea@gmail.com` or github `https://github.com/White-Mask-230/A-free-owl`
- The norms of this project are more soft than other similar projects. There will be accept all changes except the ones are out of topic or they are malicious changes
- This books can be use as primary source
- This project focus in demonstrated the information no only store it
- It is respected the work of other persons that' s why anyone can remove anything that have been create by other person but it can be say that is wrong
- It is allowed for the creators of a paper to put things of his paper in the books of this project
- The rules of the project or of the books can receive proposals to be change with a explication of why is better than the new rules writing a email to `lucasvarelacorrea@gmail.com`

1.1.2 Book

Goal

This book have the goal to collect how was find all the formulas of maths.

Problems solve

- Avoid the mathematics become a dogma, for that is proof all the formulas instead of just thinking that they are correct
- There is a lot of information relate to the mathematics in website, journals and books. This make difficult to the people that want to learn all the actual maths, to make it more easy is was make this book which will contain all the original information that have this sources.
- Verification of information different of the classical peer review

Book nomenclature

The Error Zone is where you will find all the formulas that a person find a error.

Before every formula will be his name and then his id present like this ("chapter-id"."id-chapter-formula") then it can have some comments that the author want to give to the reader:

- The formula is was find in a paper cited in the reference: ["id-reference"]
- The formula is a definition but there is any reference: [x]
- Is a popular formula but it doesn' t find how was make: (x)
- The formula was wrote less than a month: (*)
- The formula was wrote more than a month: (+)
- The formula is cited in the Error Zone: (-)
- Use a formula that is not demonstrated: (?)

The editor will write clarifications like this

- Note: "text-here"

1.2 Error Zone

Chapter 2

Algebra

2.1 Linear algebra

2.1.1 Summation

Definition summation (id 2.1) (*)

$$\sum_{i=j}^{n} f(i) = f(j) + f(j+1) + \cdots + f(n-1) + f(n)$$

Change limits

Index shift (id 2.2) (*)

$$\sum_{i=j-1}^{n-1} f(i+1) = f(j-1+1) + f(j-1+2) + \cdots + f(n-2+1) + f(n-1+1)$$

$$= f(j) + f(j+1) + \cdots + f(n-1) + f(n)$$

$$= \sum_{i=j}^{n} f(i)$$

Arithmetic Operations

Summation of the addition or subtraction of two function (id 2.3) (*)

$$\sum_{i=j}^{n} f(i)+g(i) = f(j)+g(j)+f(j+1)+g(j+1)+\cdots+f(n-1)+g(n-1)+f(n)+g(n)$$

$$= f(j) + f(j+1) + \cdots + f(n-1) + f(n) + g(j) + g(j+1) + \cdots + g(n-1) + g(n)$$

$$= \sum_{i=j}^{n} f(i) + \sum_{i=j}^{n} g(i)$$

With Constants

Summation of a constant (id 2.4) (*)

Note $C_1 = C_2 = \cdots = C_{n-1} = C_n$

$$\sum_{i=1}^{n} C = C_1 + C_2 + \cdots + C_{n-1} + C_n = C \cdot n$$

Summation of a constant and a multiplication (id 2.5) (*)

$$\sum_{i=j}^{n} C \cdot f(i) = C \cdot f(j) + C \cdot f(j+1) + \cdots + C \cdot f(n-1) + C \cdot f(n)$$

$$= C[f(j) + f(j+1) + \cdots + f(n-1) + f(n)]$$

$$= C \sum_{i=j}^{n} f(i)$$

2.2 Elementary Algebra

2.2.1 Notable Identities

Distributive property (id 2.6) (*)

$$a(b+c) = \sum_{i=1}^{a} b + c$$

$$= \sum_{i=1}^{a} b + \sum_{i=1}^{a} c$$

$$= a \cdot b + a \cdot c$$

Square of a summation or difference (id 2.7) (*)

$$(a+b)^2 = a + b \cdot a + b$$

$$= \sum_{i=1}^{a+b} a + b$$

$$= \sum_{i=1}^{a+b} a + \sum_{i=1}^{a+b} b$$

$$= a(a+b) \pm b(a+b)$$

$$= a^2 + a \cdot b + a \cdot b + b^2$$

$$= a^2 + 2 \cdot a \cdot b + b^2$$

Sum by difference (id 2.8) (*)

$$a + b \cdot a - b = \sum_{i=1}^{a+b} a - b$$

$$= \sum_{i=1}^{a+b} a - \sum_{i=1}^{a+b} b$$

$$= a(a+b) - b(a+b)$$

$$= a^2 + b \cdot a - b \cdot a - b^2$$

$$= a^2 - b^2$$

2.2.2 Property of Fractions

Reciprocal of the fraction in the denominator (id 2.9) (*)
Credits to:

- Url: `https://es.quora.com/Cul-es-la-demostracin-de-que-frac-y-frac-z-x-frac-yx-z/answer/Diego-Cabrales`
- Language: Spanish
- Title question: ¿Cuál es la demostración de que $\frac{y}{\frac{z}{x}} = \frac{yx}{z}$?
- Resolve by: Diego Cabrales

$$\frac{x}{\frac{y}{z}} = \frac{x}{\frac{y}{z}} \cdot \frac{z}{z}$$

$$= \frac{x \cdot z}{\frac{y}{z} \cdot z}$$

$$= \frac{x \cdot z}{y}$$

Chapter 3

Mathematical analysis

3.1 Calculus

3.1.1 Limits

Note: In this section I don' t find information of how was find this formulas

Properties

If only constant (id 3.1) [1] (x)

$$\lim_{x \to c} k = k$$

If only the term that approximates to (id 3.2) [1] (x)

$$\lim_{x \to c} x = c$$

If there is a constant and a function (id 3.3) [1](x)

$$\lim_{x \to c} k \cdot f(x) = k \lim_{x \to c} f(x)$$

If there are two functions that are $\pm$ (id 3.4) [1] (x)

$$\lim_{x \to c} [f(x) \pm g(x)] = \lim_{x \to c} f(x) \pm \lim_{x \to c} g(x)$$

If there are two functions that multiply (id 3.5) [1] (x)

$$\lim_{x \to c} [f(x) \cdot g(x)] = \lim_{x \to c} f(x) \cdot \lim_{x \to c} g(x)$$

If there are two functions that divide (id 3.6) [1] (x)

$$\lim_{x \to c} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \to c} f(x)}{\lim_{x \to c} g(x)}$$

If one function composite the other function (id 3.7) [1] (x)

$$\lim_{x \to c} f(g(x)) = f(\lim_{x \to c} g(x))$$

Theorem

Squeeze Theorem $f(x) \leq g(x) \leq h(x)$ (x) [1] (id 3.8)

$$\lim_{x \to c} f(x) = \lim_{x \to c} h(x) = L \to \lim_{x \to c} g(x) = L$$

Definition functions or numbers

Definition of a derivative (id 3.9) (x)

$$\frac{d}{dx} f(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

Definition of a integral (id 3.10) (x)

$$\int_a^b f(x)dx = \lim_{n \to \infty} \frac{b-a}{n} \sum_{i=1}^{n} f\left(a + i\frac{b-a}{n}\right)$$

3.1.2 Derivative

With Constants

Derivative of constant multiply and function (id 3.11) (*)

$$\frac{d}{dx}[Cf(x)] = \lim_{h \to 0} \frac{C \cdot f(x+h) - C \cdot f(x)}{h}$$

$$= C \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

$$= C \frac{d}{dx} f(x)$$

Derivative of a constant (id 3.12) (*)

$$\frac{d}{dx}[C] = \lim_{h \to 0} \frac{C - C}{h}$$

$$= 0$$

Arithmetic Operations

Derivative of the addition or subtraction of two functions (id 3.13) (*)

$$\frac{d}{dx}[f(x) \pm g(x)] = \lim_{h \to 0} \frac{f(x+h) \pm g(x+h) - f(x) - g(x)}{h}$$

$$= \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} \pm \lim_{h \to 0} \frac{g(x+h) - g(x)}{h}$$

$$= \frac{d}{dx} f(x) \pm \frac{d}{dx} g(x)$$

Derivative of a multiply of two functions (id 3.14) (*)

$$\frac{d}{dx}[f(x) \cdot g(x)] = \lim_{h \to 0} \frac{f(x+h) \cdot g(x+h) - f(x) \cdot g(x)}{h}$$

Note: addition and subtraction of $f(x) \cdot g(x+h)$

$$= \lim_{h \to 0} \frac{f(x+h) \cdot g(x+h) - f(x) \cdot g(x) + f(x) \cdot g(x+h) - f(x)g(x+h)}{h}$$

$$= \lim_{h \to 0} \frac{[f(x+h) \cdot g(x+h) - f(x) \cdot g(x+h)] + [f(x) \cdot g(x+h) - f(x) \cdot g(x)]}{h}$$

$$= \lim_{h \to 0} \frac{f(x+h) \cdot g(x+h) - f(x) \cdot g(x+h)}{h} + \lim_{h \to 0} \frac{f(x) \cdot g(x+h) - f(x) \cdot g(x)}{h}$$

$$= \lim_{h \to 0} \frac{g(x+h) \cdot [f(x+h) - f(x)]}{h} + \lim_{h \to 0} \frac{f(x) \cdot [g(x+h) - g(x)]}{h}$$

$$= \lim_{h \to 0} g(x+h) \cdot \frac{f(x+h) - f(x)}{h} + \lim_{h \to 0} f(x) \cdot \frac{g(x+h) - g(x)}{h}$$

$$= g(x) \cdot \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} + f(x) \cdot \lim_{h \to 0} \frac{g(x+h) - g(x)}{h}$$

$$= g(x) \cdot \frac{d}{dx}[f(x)] + f(x) \cdot \frac{d}{dx}[g(x)]$$

Derivative of a function elevate by a constant (id 3.15) (?)

Note: Credits to

- Url: `https://www.youtube.com/watch?v=dZnc3PtNaN4&t=304s`
- Title: Proof: `d/dx(x^n)` | Taking derivatives | Differential Calculus | Khan Academy
- Make by: Khan Academy

$$\frac{d}{dx} f^n(x) = \lim_{h \to 0} \frac{f^n(x+h) - f^n(x)}{h}$$

$$= \lim_{h \to 0} \frac{-f^n(x) + \sum_{k=0}^{n} \binom{n}{k} f^{n-k}(x+h) \cdot h^k}{h}$$

$$= \lim_{h \to 0} \sum_{k=1}^{n} \frac{\binom{n}{k} f^{n-k}(x+h) \cdot h^k}{h}$$

$$= \lim_{h \to 0} \sum_{k=1}^{n} \binom{n}{k} f^{n-k}(x+h) \cdot h^{k-1}$$

$$= \sum_{k=1}^{n} \lim_{h \to 0} \binom{n}{k} f^{n-k}(x+h) \cdot h^{k-1}$$

$$= \lim_{h\to 0}\left[\binom{n}{1}f^{n-1}(x+h)h^{1-1} + \binom{n}{2}f^{n-2}(x+h)h^{2-1} + \cdots + \binom{n}{k}f^{n-k}(x+h)h^{k-1}\right]$$

$$= \lim_{h\to 0}\binom{n}{1}f^{n-1}(x+h)h^{1-1} + \lim_{h\to 0}\binom{n}{2}f^{n-2}(x+h)h^{2-1} + \cdots + \lim_{h\to 0}\binom{n}{k}f^{n-k}(x+h)h^{k-1}$$

$$= n \cdot f^{n-1}(x)$$

Derivative of a division of two functions (id 3.16) (*)

$$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{d}{dx}[f(x)\cdot g^{-1}(x)]$$

$$= g^{-1}(x)\cdot\frac{d}{dx}f(x) + f(x)\cdot\frac{d}{dx}g^{-1}(x)$$

$$= g^{-1}(x)\frac{d}{dx}f(x) - f(x)\cdot g^{-2}(x)\cdot\frac{d}{dx}g(x)$$

$$= \frac{g(x)\frac{d}{dx}f(x) - f(x)\cdot\frac{d}{dx}g(x)}{g^2(x)}$$

Derivative of a natural logarithm (id 3.17) (*)

$$\frac{d}{dx}\ln(x) = \lim_{h\to 0}\frac{\ln(x+h)-\ln(x)}{h}$$

$$= \lim_{h\to 0}\frac{\ln\left(\frac{x+h}{x}\right)}{h}$$

$$= \lim_{h\to 0}\frac{\ln\left(1+\frac{h}{x}\right)}{h}$$

$$= \lim_{h\to 0}\frac{x}{x}\frac{\ln\left(1+\frac{h}{x}\right)}{h}$$

$$= \lim_{h\to 0}\frac{1}{x}\frac{\ln(1+\frac{h}{x})}{\frac{h}{x}}$$

$$= \lim_{u\to 0}\frac{1}{x}\frac{\ln(1+u)}{u}$$

$$= \lim_{u\to 0}\ln(1+u)\cdot\lim_{u\to 0}\frac{1}{u\cdot x}$$

$$= \ln(1)\cdot\lim_{u\to 0}\frac{1}{u\cdot x}$$

$$= \frac{1}{x}$$

Definition functions or numbers

Function digamma (id 3.17) (?)

$$\psi(x) = \frac{d}{dx}\ln[\Gamma(x)] = \frac{\frac{d}{dx}\Gamma(x)}{\Gamma(x)}$$

3.1.3 Integrals

Constants

Integral of the multiply of a function and a constant (id 3.17) (*)

$$\int_a^b C \cdot f(x)dx = \lim_{n\to\infty} \frac{b-a}{n} \sum_{i=1}^{n} C \cdot f\left(a + i\frac{b-a}{n}\right)$$

$$= C \cdot \lim_{n\to\infty} \frac{b-a}{n} \sum_{i=1}^{n} f\left(a + i\frac{b-a}{n}\right)$$

$$= C \cdot \int_a^b f(x)dx$$

Arithmetic Operations

Integral of a addition and a subtraction of two functions (id 3.18) (*)

$$\int_a^b f(x) \pm g(x)dx = \lim_{n\to\infty} \frac{b-a}{n} \left[\sum_{i=1}^{n} f\left(a + i\frac{b-a}{n}\right) \pm g\left(a + i\frac{b-a}{n}\right)\right]$$

$$= \lim_{n\to\infty} \frac{b-a}{n} \left[\sum_{i=1}^{n} f\left(a + i\frac{b-a}{n}\right) \pm \sum_{i=1}^{n} g\left(a + i\frac{b-a}{n}\right)\right]$$

$$= \lim_{n\to\infty} \frac{b-a}{n} \sum_{i=1}^{n} f\left(a + i\frac{b-a}{n}\right) \pm \lim_{n\to\infty} \frac{b-a}{n} \sum_{i=1}^{n} g\left(a + i\frac{b-a}{n}\right)$$

$$= \int_a^b f(x)dx \pm \int_a^b g(x)dx$$

Chapter 4

Final part

4.1 Reference

[1] Alan Stein. Properties of Limits: https://www2.math.uconn.edu/~stein/math115/Slides/math115-130notes.pdf

4.2 Make by

Lucas Varela Correa

Nickname: White-Mask-230
Github: https://github.com/White-Mask-230
Contact: lucasvarelacorrea@gmail.com