


Peak fused change evidence






Estimated change onset (alarmed pixels)








Fraction of AOI above threshold, per time bin





Alarmed area (ha)






South Docks: VH (cross-pol) vs AOI median (dB); grey = Nov-Mar
zone mean
brightest 5%


South Docks: NDVI vs AOI median (clear scenes)


South Docks: Landsat surface temp vs AOI median (K)
zone mean
hottest 5%



North Airfield: VH (cross-pol) vs AOI median (dB); grey = Nov-Mar
zone mean
brightest 5%


North Airfield: NDVI vs AOI median (clear scenes)


North Airfield: Landsat surface temp vs AOI median (K)
zone mean
hottest 5%


Winter Landsat surface temperature relative to the AOI median (red = warmer than surroundings)

Change clusters by evidence class (numbers = id, clusters of 1 ha or more)



Largest clusters, strongest evidence first

id	ha	evidence	onset	status	VH z	NDVI z
6	4.5	optical-only	2025-12-29	absorbed	-	-6.3
10	1.6	optical-only	2020-09-26	absorbed	-	+4.7
8	1.2	optical-only	2023-08-17	absorbed	-	+4.4
9	1.1	optical-only	2021-08-22	absorbed	-	+6.1
7	0.7	optical-only	2021-09-01	absorbed	-	+6.9
11	0.6	optical-only	2021-09-01	absorbed	-	+5.9
12	1.2	weak	2025-12-04	absorbed	-	-1.2
3	0.8	weak	2024-08-26	absorbed	-	+1.4
2	0.7	weak	2020-11-25	absorbed	-	+0.6
1	0.6	weak	2020-11-25	absorbed	-	+1.1
4	0.4	weak	2023-05-24	absorbed	-	+2.0
5	0.4	weak	2025-12-04	absorbed	-	+1.6


Scene availability (one tick per scene; grey = Nov-Mar)
thermal (189)
optical (430)
SAR asc84 (1)
2019
2020
2021
2022
2023
2024
2025
2026
Scenes per year
SAR asc84
optical
thermal
0
10
20
30
40
50
2018
2019
2020
2021
2022
2023
2024
2025
2026