


Kaplan–Meier for Appetite loss by Study

Study + 1997 (n=735) + 2000 (n=1564) + 2001 (n=226) + 2004 (n=213)



Number at risk

| Study | Time (days) | Count |
|---------|-------------|-------|
| Study 1 | 0-200 | 728 |
| | 200-400 | 323 |
| | 400-600 | 263 |
| | 600-800 | 180 |
| | 800-1000 | 153 |
| | 1000-1200 | 9 |
| | 1200-1400 | 4 |
| | 1400-1600 | 2 |
| | 1600-1800 | 2 |
| | 1800-2000 | 0 |
| Study 2 | 0-200 | 1541 |
| | 200-400 | 660 |
| | 400-600 | 529 |
| | 600-800 | 360 |
| | 800-1000 | 334 |
| | 1000-1200 | 27 |
| | 1200-1400 | 12 |
| | 1400-1600 | 0 |
| | 1600-1800 | 0 |
| | 1800-2000 | 0 |
| Study 3 | 0-200 | 215 |
| | 200-400 | 11 |
| | 400-600 | 0 |
| | 600-800 | 0 |
| | 800-1000 | 0 |
| | 1000-1200 | 0 |
| | 1200-1400 | 0 |
| | 1400-1600 | 0 |
| | 1600-1800 | 0 |
| | 1800-2000 | 0 |
| Study 4 | 0-200 | 206 |
| | 200-400 | 26 |
| | 400-600 | 4 |
| | 600-800 | 0 |
| | 800-1000 | 0 |
| | 1000-1200 | 0 |
| | 1200-1400 | 0 |
| | 1400-1600 | 0 |
| | 1600-1800 | 0 |
| | 1800-2000 | 0 |