

Probability & Statistics

APM_2F010_EP

Lecture Notes

University: Ecole Polytechnique

Lecturer: Prof. Tardy Yoan

Student: Nikolay Kharovskiy

Semester: semester 3

Date: September 21, 2026

These are personal lecture notes and may contain errors. For the official course material consult the lecturer.

Contents

1	Lecture 1	2
2	Lecture 2	3
2.1	General Probability Spaces	3
2.2	The Borel σ -algebra	4
2.3	Random Variables on General Probability Spaces	4
2.4	The Law (Distribution) of a Random Variable	5
2.5	Conditional Probability and Independence	5
3	Lecture 3	7
3.1	Important Distributions	7
3.2	Relations Between Distributions	8
3.3	Expectation	9
3.4	Variance and Covariance	11

1 Lecture 1

Theorem 1: Fubini positive

Let $f : X \times Y \rightarrow [0, \infty)$, $g : X \rightarrow [0, \infty)$ and $h : Y \rightarrow [0, \infty)$ be piecewise continuous. Then:

$$\int_Y \left(\int_X f(x, y) g(x) dx \right) h(y) dy = \int_X \left(\int_Y f(x, y) h(y) dy \right) g(x) dx$$

- Works without integrability assumptions.
- Analogue: reordering double series with nonnegative terms.

Theorem 2: Fubini - Tonelli

Let $f : X \times Y \rightarrow \mathbb{R}$ (not necessarily positive), and let $g : X \rightarrow [0, \infty)$ and $h : Y \rightarrow [0, \infty)$ be piecewise continuous. If:

$$\int_Y \left(\int_X |f(x, y)| g(x) dx \right) h(y) dy < \infty$$

Then:

$$\int_Y \left(\int_X f(x, y) g(x) dx \right) h(y) dy = \int_X \left(\int_Y f(x, y) h(y) dy \right) g(x) dx$$

- Key assumption: integrable.

What is a probability?

Example 3: Rolling a fair dice

- Possible outcomes $\Omega = \{1, \dots, 6\}$.
- All outcomes have probability $1/6$.
- An event is “The result is odd”.

Proposition 4: Discrete probability space

$(\Omega, \mathbb{P})$ is a discrete prob. space. $\mathbb{P}$ is called the discrete prob. on Ω . If $\forall A \subset \Omega$, $\mathbb{P}(A) = \frac{\text{Card}(A)}{\text{Card}(\Omega)}$, then $\mathbb{P}$ is called the uniform discrete probability.

Example 5: Non-discrete probability space

Let $E =$ “the extraction of a number in $[0, 1]$ uniformly at random”. E is modeled with the uniform probability $\mathbb{P}$ on the space $\Omega = [0, 1]$.

$$\forall 0 \leq a \leq b \leq 1, \quad \mathbb{P}([a, b]) = b - a$$

$(\Omega, \mathbb{P})$ cannot be a discrete prob. space.

Definition 6: Probability function

We define a prob. function $\mathbb{P} : \mathcal{P}([0, 1]) \rightarrow [0, 1]$ s.t.

1. $\mathbb{P}([0, 1]) = 1$
2. $\mathbb{P}$ is σ -additive
3. $\mathbb{P}([a, b]) = b - a \quad \forall a, b$

2 Lecture 2**2.1 General Probability Spaces****Definition 1: Sigma-algebra**

Let Ω be an arbitrary set. A σ -algebra $\mathcal{A}$ is a subset of $\mathcal{P}(\Omega)$ such that:

1. $\emptyset \in \mathcal{A}$
2. If $A \in \mathcal{A}$, then $A^c \in \mathcal{A}$
3. If $(A_n)_{n \in \mathbb{N}}$ is a sequence of events in $\mathcal{A}$, then $\bigcup_{n \in \mathbb{N}} A_n$ is also in $\mathcal{A}$.

σ -algebras are subsets of $\mathcal{P}(\Omega)$ that are closed under the basic operations on sets.

Example 2: Sigma-algebras

Let Ω be an arbitrary set and $A \subseteq \Omega$. The following are σ -algebras:

- $\mathcal{P}(\Omega)$
- $\{\emptyset, A, A^c, \Omega\}$

Definition 3: General probability space

Let Ω be an arbitrary set, $\mathcal{A} \subseteq \mathcal{P}(\Omega)$ a sigma algebra and $\mathbb{P} : \mathcal{A} \rightarrow [0, 1]$. The triplet $(\Omega, \mathcal{A}, \mathbb{P})$ is called a (general) probability space if:

1. $\mathbb{P}[\Omega] = 1$
2. If $(A_n)_{n \in \mathbb{N}}$ is a sequence of disjoint events in $\mathcal{A}$, then $\mathbb{P}[\bigcup_{n \in \mathbb{N}} A_n] = \sum_{n \in \mathbb{N}} \mathbb{P}[A_n]$.

If conditions (i) and (ii) are verified, $\mathbb{P}$ is called a (general) probability.

Example 4: General probability space

If $(\Omega, \mathbb{P})$ is a discrete probability space, then $(\Omega, \mathcal{P}(\Omega), \mathbb{P})$ is a general probability space.

Proposition 5: Basic properties of probability spaces

Let $\mathbb{P} : \mathcal{P}(\Omega) \rightarrow [0, 1]$ be a probability and $A, B \subseteq \Omega$. Then:

1. Let $A_1, \dots, A_n$ be a finite collection of pairwise disjoint events. Then $\mathbb{P}[A_1 \cup \dots \cup A_n] = \sum_{j=1}^n \mathbb{P}[A_j]$. This property is called finite σ -additivity or additivity.
2. If $A \subseteq B$, $\mathbb{P}[B \setminus A] = \mathbb{P}[B] - \mathbb{P}[A]$. In particular, $\mathbb{P}[A] \leq \mathbb{P}[B]$ and $\mathbb{P}[A^c] = 1 - \mathbb{P}[A]$.
3. $\mathbb{P}[A \cup B] = \mathbb{P}[A] + \mathbb{P}[B] - \mathbb{P}[A \cap B]$. In particular, $\mathbb{P}[A \cup B] \leq \mathbb{P}[A] + \mathbb{P}[B]$.

2.2 The Borel σ -algebra

We need a σ -algebra which contains all intervals and is smaller than $\mathcal{P}(\Omega)$. The easiest choice is to take the smallest possible sigma algebra.

Definition 6: The Borel σ -algebra

Let $I \subseteq \mathbb{R}$ be an interval. The Borel sigma algebra $\mathcal{B}(I)$ is defined as:

$$\mathcal{B}(I) = \bigcap_{\{[a,b]: a,b \in I\} \subseteq \mathcal{A} \subseteq \mathcal{P}(I)} \mathcal{A}$$

To check that it is a good definition one has to verify that $\mathcal{B}(I)$ is indeed a sigma algebra.

Theorem 7: Uniform Probability Measure

There exists a general probability space $([0, 1], \mathcal{B}([0, 1]), \mathbb{P})$ such that $\mathbb{P}[[a, b]] = b - a$ for all $a, b \in \mathbb{R}$.

2.3 Random Variables on General Probability Spaces**Definition 8: Real random variable**

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space. A function $X : \Omega \rightarrow \mathbb{R}$ is called a real random variable if and only if for any $a, b \in \mathbb{R}$ we have that:

$$X^{-1}([a, b]) := \{\omega : X(\omega) \in [a, b]\} \in \mathcal{A}$$

A real random variable X defined on a general probability space $(\Omega, \mathcal{A}, \mathbb{P})$ is discrete if $X(\Omega)$ is at most countable.

Proposition 9: Discrete random variables as general r.v.

Let $(\Omega, \mathbb{P})$ be a discrete p.s. and $X : \Omega \rightarrow \mathbb{R}$ a r.v. Then X is a real random variable on $(\Omega, \mathcal{P}(\Omega), \mathbb{P})$.

Remark 10: Conventions

- If $E = \mathbb{R}$, we call X a real random variable; if $E = \mathbb{R}^n$ we call X a random vector.
- “Let X be a random variable” has to be read as “Let $(\Omega, \mathbb{P})$ be a discrete probability space and X be a random variable defined on it”.
- We may denote events without brackets, and not specify the probability space. For instance: $\mathbb{P}[\{\omega : X(\omega) \in A\}] \Leftrightarrow \mathbb{P}[X \in A]$ and $\mathbb{P}[\omega] \Leftrightarrow \mathbb{P}[\{\omega\}]$.
- We may replace the intersection with a comma. For example: $\mathbb{P}[\{X \in A\} \cap \{Y \in B\}] \Leftrightarrow \mathbb{P}[X \in A, Y \in B]$.

2.4 The Law (Distribution) of a Random Variable**Proposition 11: The law of a random variable**

Let X be a real valued r.v. The law (distribution) μ_X of X is the application $\mu_X : \mathcal{B}(\mathbb{R}) \rightarrow [0, 1]$ defined by $\mu_X(C) := \mathbb{P}[X^{-1}(C)]$. In this setting, the law $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \mu_X)$ is a general probability space.

Remark 12: Two Kinds of Random Variables

In this course, a random variable $X : \Omega \rightarrow \mathbb{R}$ will be of one of two types:

- **Discrete random variables:** it exists a countable set $E \subset \mathbb{R}$ and $p_X : E \rightarrow \mathbb{R}_+$ s.t. $\forall C \in \mathcal{B}(\mathbb{R}), \mu_X(C) = \sum_{x \in C \cap E} p_X(x)$.
- **Absolutely continuous random variables:** it exists $f_X : \mathbb{R} \rightarrow \mathbb{R}_+$ s.t. $\forall a < b \in \mathbb{R}, \mu_X([a, b]) = \int_a^b f_X(u) du$.

Fundamental intuition: $\mathbb{P}(X \in [a - \epsilon, a + \epsilon]) = \mu_X([a - \epsilon, a + \epsilon]) \approx 2\epsilon f_X(a)$. Here, p_X and f_X are respectively called the discrete density and the density of the random variable X .

2.5 Conditional Probability and Independence**Definition 13: Conditional probability**

Let $(\Omega, \mathbb{P})$ be a discrete probability space and A, B two events such that $\mathbb{P}[B] > 0$. The conditional probability of A given B is denoted $\mathbb{P}[A|B]$ and defined by:

$$\mathbb{P}[A|B] := \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]}$$

Definition 14: Independence of two events

Let $(\Omega, \mathbb{P})$ be a discrete probability space and A, B two events. We say that A, B are independent if $\mathbb{P}[A \cap B] = \mathbb{P}[A]\mathbb{P}[B]$. If $\mathbb{P}[B] > 0$, we have the equivalent definition $\mathbb{P}[A|B] = \mathbb{P}[A]$.

Definition 15: Joint and marginal law

Let $X : \Omega \rightarrow E, Y : \Omega \rightarrow F$ be two r.v. The joint law $\mu_{X,Y}$ is the law of the random vector (X, Y) . In particular, for all $C_1, C_2 \in \mathcal{B}(\mathbb{R})$:

$$\mu_{X,Y}(C_1 \times C_2) = \mathbb{P}[(X, Y) \in C_1 \times C_2] = \mathbb{P}[X \in C_1, Y \in C_2]$$

In this context μ_X, μ_Y are called marginal laws. We will write $p_{X,Y}$ and $f_{X,Y}$ for respectively the discrete joint density and the joint density according to the case we study.

Proposition 16: Marginal probabilities from joint density

Let $(\Omega, \mathbb{P})$ be a discrete probability space and $X : \Omega \rightarrow E, Y : \Omega \rightarrow F$ two random variables. Then:

$$\begin{aligned} \forall y \in F, \quad p_Y(y) &= \sum_{x \in E} p_{X,Y}(x, y) \\ \forall x \in E, \quad p_X(x) &= \sum_{y \in F} p_{X,Y}(x, y) \end{aligned}$$

Definition 17: Independence of random variables

Let $(X_1, \dots, X_N)$ be real valued random variables defined all on the same p.s. $(\Omega, \mathcal{A}, \mathbb{P})$. We say that they are independent if and only if for all possible choices of $[a_1, b_1], \dots, [a_N, b_N]$ we have:

$$\mathbb{P}[X_1 \in [a_1, b_1], \dots, X_N \in [a_N, b_N]] = \prod_{i=1}^N \mathbb{P}[X_i \in [a_i, b_i]]$$

An arbitrary family $(X_i)_{i \in I}$ of random variables defined on the same probability space is independent if and only if each finite subfamily is independent.

Proposition 18: Independence under complementation

Let $(A_i)_{i \in I}$ be independent events and $I' \subseteq I$. Then the events $(B_i)_{i \in I}$ defined by

$$B_i = \begin{cases} A_i, & i \in I' \\ A_i^c, & i \in I \setminus I' \end{cases}$$

are independent.

Proposition 19: Independence and joint density

Let $X : \Omega \rightarrow E_1, Y : \Omega \rightarrow E_2$ be two random variables defined on the same discrete probability space $(\Omega, \mathbb{P})$. Then X and Y are independent if and only if:

$$\forall x \in E_1, y \in E_2, \quad p_{X,Y}(x, y) = p_X(x)p_Y(y)$$

Proposition 20: Functions of independent random variables

Let X, Y be two independent random variables taking values in E and F respectively. Moreover, let H, K be two other sets and $f : E \rightarrow H$ and $g : F \rightarrow K$ two functions. Then the random variables $f(X)$ and $g(Y)$ are independent.

Proposition 21: The sum of independent random variables

Let X, Y be two absolutely continuous and independent random variables whose densities are f_X and f_Y . Then, the random variable $X + Y$ is absolutely continuous. Its density f_{X+Y} is given by the convolution $f_X * f_Y$ of f_X and f_Y , defined by:

$$(f_X * f_Y)(z) = \int_{-\infty}^{+\infty} f_X(x)f_Y(z-x)dx$$

It is also true in the discrete case:

$$p_{X+Y}(z) = \sum_{x \in \mathbb{R}} p_X(x)p_Y(z-x)$$

Be careful, the notation $\sum_{x \in \mathbb{R}}$ is only licit because this is the discrete case, this is highly wrong to write it in the absolutely continuous case.

3 Lecture 3**3.1 Important Distributions****Definition 1: Bernoulli Distribution**

X is a Bernoulli random variable of parameter p if $p_X(1) = p$ and $p_X(0) = (1-p)$. In this case, the distribution of X is the Bernoulli distribution of parameter p and we write $X \sim \text{Be}(p)$.

Definition 2: Geometric Distribution

X is a geometric random variable of parameter p if $p_X(0) = 0$ and $p_X(n) = p(1-p)^{n-1}$ for $n \geq 1$. In this case, the distribution of X is the geometric distribution of parameter p and we write $X \sim \text{Geo}(p)$.

Definition 3: Binomial Distribution

X is a Binomial random variable of parameters N and p if

$$p_X(k) = \binom{N}{k} p^k (1-p)^{N-k}, \quad \forall k = 0, 1, \dots, N.$$

In this case, the distribution of X is the binomial distribution of parameters N and p and we write $X \sim \text{Bin}(N, p)$.

Definition 4: Poisson Distribution

X is a Poisson random variable of parameter λ , if

$$\forall n \in \mathbb{N}, \quad p_X(n) = \exp(-\lambda) \frac{\lambda^n}{n!}$$

In this case, the distribution of X is the Poisson distribution of parameter λ and we write $X \sim \text{Poi}(\lambda)$.

3.2 Relations Between Distributions
Proposition 5: From Bernoulli to Binomial

Let $(X_i)_{i \geq 1}$ be a sequence of i.i.d. random variables such that $X_1 \sim \text{Be}(p)$. Then $\sum_{i=1}^N X_i \sim \text{Bin}(N, p)$.

PROOF. Let $X = \sum_{i=1}^N X_i$. We have:

$$\mathbb{P}(X = k) = \mathbb{P}\left(\sum_{i=1}^N X_i = k\right) = \mathbb{P}\left(\bigcup_{K \subset \{1, \dots, N\}, |K|=k} \{X_i = 1 \forall i \in K, X_i = 0 \forall i \notin K\}\right)$$

By σ -additivity, since the sets are disjoint:

$$\begin{aligned} &= \sum_{|K|=k} \prod_{i \in K} \mathbb{P}(X_i = 1) \prod_{i \notin K} \mathbb{P}(X_i = 0) \\ &= \binom{N}{k} p^k (1-p)^{N-k} \end{aligned}$$

□

Proposition 6: From Bernoulli to Geometrical

Let $(X_i)_{i \geq 1}$ be a sequence of i.i.d. random variables such that $X_1 \sim \text{Be}(p)$. Then $T = \inf\{i \geq 1 : X_i = 1\}$ is a geometric random variable of parameter p , i.e., $T \sim \text{Geo}(p)$.

PROOF. First, we find the probability that $T \geq n$:

$$\mathbb{P}(T \geq n) = \mathbb{P}(X_i = 0 \forall i \in \{1, \dots, n-1\}) = (1-p)^{n-1}$$

Then, using the difference of probabilities:

$$\begin{aligned} \mathbb{P}(T = n) &= \mathbb{P}(T \geq n) - \mathbb{P}(T \geq n+1) = (1-p)^{n-1} - (1-p)^n \\ &= (1-p)^{n-1}(1 - (1-p)) = p(1-p)^{n-1} \end{aligned}$$

□

Proposition 7: Law of Small Numbers (simplified version)

Let $\lambda > 0$ and $(X_n)_{n \geq [\lambda]}$ be a sequence of random variables such that $X_n \sim \text{Bin}(n, \lambda/n)$. In particular, this means $\mathbb{E}[X_n] = \lambda$ for all n . Then we have

$$\lim_{n \rightarrow +\infty} p_{X_n}(k) = e^{-\lambda} \frac{\lambda^k}{k!} = p_X(k), \text{ with } X \sim \text{Poi}(\lambda).$$

PROOF. Let $p = \lambda/n$. For a given r :

$$\begin{aligned} \mathbb{P}(X_n = r) &= \binom{n}{r} \left(\frac{\lambda}{n}\right)^r \left(1 - \frac{\lambda}{n}\right)^{n-r} \\ &= \frac{n(n-1)\dots(n-r+1)}{r!} \frac{\lambda^r}{n^r} \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-r} \end{aligned}$$

As $n \rightarrow \infty$, we analyze the terms:

- $\frac{n(n-1)\dots(n-r+1)}{n^r} \rightarrow 1$
- $\left(1 - \frac{\lambda}{n}\right)^n \rightarrow e^{-\lambda}$
- $\left(1 - \frac{\lambda}{n}\right)^{-r} \rightarrow 1$

Thus, $\mathbb{P}(X_n = r) \rightarrow e^{-\lambda} \frac{\lambda^r}{r!}$. □

3.3 Expectation

Definition 8: Expectation

Let X be a non negative valued random variable defined on a general probability space $(\Omega, \mathbb{P})$. We write $\mathbb{E}[X]$ the expectation of X which equals:

- $\sum_{x \geq 0} x p_X(x)$ in the discrete case,
- $\int_{\mathbb{R}} x f_X(x) dx$ in the absolute continuous case.

If X is a real valued random variable, we say that the expectation $\mathbb{E}[X]$ of X exists if one of the following conditions $\mathbb{E}[\mathbb{1}_{X \geq 0} X] < +\infty$ or $\mathbb{E}[\mathbb{1}_{X \leq 0} (-X)] < +\infty$ is verified. In that case the expectation value of X is given by

$$\mathbb{E}[X] = \mathbb{E}[\mathbb{1}_{X \geq 0} X] - \mathbb{E}[\mathbb{1}_{X \leq 0} (-X)].$$

Example 9: Expectation Examples

- **Fair dice:** If $p_X(x) = 1/6$ for $x \in \{1, 2, 3, 4, 5, 6\}$, then $\mathbb{E}[X] = 3.5$.
- **Uniform law on $[0, 1]$:** If $f_X = \mathbb{1}_{[0,1]}$, then $\mathbb{E}[X] = 0.5$.

Proposition 10: Expectation - Equivalent Definition

Let $X : \Omega \rightarrow E$ be a discrete random variable defined over a discrete probability space $(\Omega, \mathbb{P})$. We have that $\mathbb{E}[X]$ exists if and only if one of the two conditions

$$\sum_{\omega \in \Omega, X(\omega) \geq 0} X(\omega) \mathbb{P}[\{\omega\}] < +\infty, \quad \sum_{\omega \in \Omega, X(\omega) < 0} X(\omega) \mathbb{P}[\{\omega\}] > -\infty$$

is verified. In this case

$$\mathbb{E}[X] = \sum_{\omega \in \Omega} X(\omega) \mathbb{P}[\{\omega\}]$$

PROOF. Assume $X \geq 0$. Using the positive Fubini theorem:

$$\begin{aligned} \sum_{\omega \in \Omega} X(\omega) \mathbb{P}[\{\omega\}] &= \sum_{x \in X(\Omega)} \sum_{\omega \in \Omega, X(\omega)=x} X(\omega) \mathbb{P}[\{\omega\}] \\ &= \sum_{x \in X(\Omega)} x \sum_{\omega: X(\omega)=x} \mathbb{P}[\{\omega\}] = \sum_{x \in X(\Omega)} x \mathbb{P}(X = x) = \mathbb{E}[X] \end{aligned}$$

The same logic extends to $\mathbb{E}[\mathbb{1}_{X \geq 0} X]$ and $\mathbb{E}[\mathbb{1}_{X \leq 0} (-X)]$. □

Proposition 11: Expectation of a Function of a Random Variable

Let $X : \Omega \rightarrow E$ be a random variable defined over a general probability space $(\Omega, \mathbb{P})$ and $g : E \rightarrow \mathbb{R}$ any function. In the discrete case, $\mathbb{E}[g(X)]$ exists if and only if either $\sum_{x: g(x) \geq 0} g(x) p_X(x) < +\infty$ or $\sum_{x: g(x) < 0} g(x) p_X(x) > -\infty$ is verified. Then:

$$\mathbb{E}[g(X)] = \sum_{x \in X(\Omega)} g(x) p_X(x)$$

In the absolutely continuous case, if g is piecewise continuous, $\mathbb{E}[g(X)]$ exists if and only if either $\int_{\mathbb{R}} \mathbb{1}_{g(x) \geq 0} g(x) f_X(x) dx < +\infty$ or $\int_{\mathbb{R}} \mathbb{1}_{g(x) \leq 0} g(x) f_X(x) dx > -\infty$ is verified. Then:

$$\mathbb{E}[g(X)] = \int_{\mathbb{R}} g(x) f_X(x) dx$$

PROOF. For the discrete case, assuming $g \geq 0$:

$$\begin{aligned} \sum_{x \in X(\Omega)} g(x) p_X(x) &= \sum_{q \geq 0} \sum_{x: g(x)=q} q \mathbb{P}(X = x) \\ &= \sum_{q \geq 0} q \mathbb{P}(g(X) = q) = \mathbb{E}[g(X)] \end{aligned}$$

□

Proposition 12: Product of Independent Random Variables

Let X, Y be two independent random variables. If $\mathbb{E}[X], \mathbb{E}[Y] < +\infty$, then $\mathbb{E}[XY] < +\infty$ as well and

$$\mathbb{E}[XY] = \mathbb{E}[X] \mathbb{E}[Y]$$

PROOF. By Fubini positive, in the discrete case:

$$\begin{aligned}\mathbb{E}[XY] &= \sum_x \sum_y xy \mathbb{P}(X = x, Y = y) = \sum_x \sum_y xy \mathbb{P}(X = x) \mathbb{P}(Y = y) \\ &= \left(\sum_x x \mathbb{P}(X = x) \right) \left(\sum_y y \mathbb{P}(Y = y) \right) = \mathbb{E}[X] \mathbb{E}[Y]\end{aligned}$$

□

Proposition 13: Properties of Expectation

Let X, Y two real valued random variables defined on the same general probability space $(\Omega, \mathbb{P})$. The following holds:

1. **Monotonicity:** If both $\mathbb{E}[X], \mathbb{E}[Y]$ exist, and $X(\omega) \leq Y(\omega) \forall \omega \in \Omega$, then $\mathbb{E}[X] \leq \mathbb{E}[Y]$.
2. **Absolute Value:** If $\mathbb{E}[X]$ exists, then $|\mathbb{E}[X]| \leq \mathbb{E}[|X|]$.
3. **Linearity:** If X, Y have finite expectation, and $a, b \in \mathbb{R}$, then $aX + bY$ has finite expectation and $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$.

PROOF. For discrete random variables:

1. $\mathbb{E}[X] = \sum_{\omega} X(\omega) \mathbb{P}[\{\omega\}] \leq \sum_{\omega} Y(\omega) \mathbb{P}[\{\omega\}] = \mathbb{E}[Y]$.
2. $|\mathbb{E}[X]| = |\sum_{\omega} X(\omega) \mathbb{P}[\{\omega\}]| \leq \sum_{\omega} |X(\omega)| \mathbb{P}[\{\omega\}] = \mathbb{E}[|X|]$ (by triangle inequality).
3. $\mathbb{E}[aX + bY] = \sum_{\omega} (aX(\omega) + bY(\omega)) \mathbb{P}[\{\omega\}] = a\mathbb{E}[X] + b\mathbb{E}[Y]$.

□

Example 14: Uniform Random Variables Moments

If X is a uniform random variable over the interval $[a, b]$, then $f_X(x) = \frac{1}{b-a}$ for $x \in [a, b]$ and 0 otherwise.

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{b-a} \int_a^b x dx = \frac{b^2 - a^2}{2(b-a)} = \frac{a+b}{2} \\ \mathbb{E}[X^2] &= \frac{1}{b-a} \int_a^b x^2 dx = \frac{b^3 - a^3}{3(b-a)} = \frac{b^2 + ab + a^2}{3}\end{aligned}$$

3.4 Variance and Covariance

Definition 15: Variance

Let X be a real valued random variable with $\mathbb{E}[X^2] < +\infty$. The variance of X is

$$\mathbb{V}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

Definition 16: Covariance

Let X, Y be two random variables s.t. $\mathbb{E}[|X|], \mathbb{E}[|Y|] < +\infty$ and $\mathbb{E}[|XY|] < +\infty$. The covariance of X and Y is

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$$

In particular, $\mathbb{V}(X) = \text{Cov}(X, X)$.

Proposition 17: Properties of Variance and Covariance

Let X, Y be two random variables such that $\mathbb{E}[X^2], \mathbb{E}[Y^2] < +\infty$. Then we have

$$\mathbb{V}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2, \quad \text{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y].$$

and

$$\mathbb{V}(X + Y) = \mathbb{V}(X) + \mathbb{V}(Y) + 2\text{Cov}(X, Y).$$

- If $X \sim \text{Be}(p)$, then $\mathbb{V}(X) = p - p^2$.
- If X is uniformly distributed on $\{-N, -N+1, \dots, N-1, N\}$, then $\mathbb{V}(X) = \frac{N(N+1)}{3}$.