

Algebra I: Class Notes

Wilson Pan

February 8, 2026

Contents

1	Lecture 1: Group Actions & The Orbit-Stabilizer Theorem (Jan 12)	2
1.1	Permutations and Automorphisms	2
1.2	Group Actions	2
1.3	Orbits and Stabilizers	2
2	Lecture 2: Cauchy's Theorem & Conjugation (Jan 14)	4
2.1	Cauchy's Theorem	4
2.2	Conjugation	4
3	Lecture 3: Sylow Preliminaries & Isomorphism Theorems (Jan 16)	6
3.1	Product of Subgroups	6
3.2	Sylow Definitions	6
4	Lecture 4: The Sylow Theorems (Jan 21)	7
5	Lecture 5: Normal Series & Solvability (Jan 23)	10
5.1	Normal and Subnormal Series	10
5.2	Solvable Groups	10
6	Lecture 6: Solvable and Derived Series (Jan 26)	11
7	Lecture 7: Nilpotent Groups and Free Groups (Jan 28)	12
7.1	Nilpotent Groups	12
7.2	Free Groups	12
8	Lecture 8: Free Groups	14
9	Lecture 9: Generators and Relations (Feb 2)	16
10	Lecture 10: Presentation of Groups (Feb 4)	17
11	Lecture 11: Category Theory (Feb 6)	18
11.1	Categories	18

1 Lecture 1: Group Actions & The Orbit-Stabilizer Theorem (Jan 12)

1.1 Permutations and Automorphisms

Definition 1.1 (Symmetric Group). Let X be a set. The set of all permutations (bijections) of X is denoted by S_X (or sometimes Σ_X). Under function composition, S_X forms a group.

Definition 1.2 (Automorphism Group). Let (G, \cdot) be a group. An automorphism of G is a bijection $\phi : G \rightarrow G$ that is also a homomorphism, i.e.,

$$\phi(g_1 g_2) = \phi(g_1) \phi(g_2) \quad \forall g_1, g_2 \in G.$$

The set of all such automorphisms is denoted by $\text{Aut}(G)$. It forms a group under composition.

1.2 Group Actions

Definition 1.3 (Group Action). Let G be a group and X be a set. An **action** of G on X is a homomorphism $\phi : G \rightarrow S_X$.

We typically write the action as $g \cdot x := \phi(g)(x)$. This notation satisfies two axioms (equivalent to the homomorphism property):

1. **Identity:** $1 \cdot x = x$ for all $x \in X$.
2. **Compatibility:** $g_1 \cdot (g_2 \cdot x) = (g_1 g_2) \cdot x$ for all $g_1, g_2 \in G, x \in X$.

Theorem 1.4 (Equivalence Relation on X). Let G act on X . Define a relation \sim on X by:

$$x_1 \sim x_2 \iff \exists g \in G \text{ such that } g \cdot x_1 = x_2.$$

Then \sim is an equivalence relation.

Proof. We check the properties:

1. **Reflexive:** $1 \cdot x = x \implies x \sim x$.
2. **Symmetric:** If $g \cdot x_1 = x_2$, acting by g^{-1} gives $g^{-1} \cdot (g \cdot x_1) = g^{-1} \cdot x_2 \implies (g^{-1}g) \cdot x_1 = g^{-1} \cdot x_2 \implies x_1 = g^{-1} \cdot x_2$. Thus $x_2 \sim x_1$.
3. **Transitive:** If $g \cdot x_1 = x_2$ and $h \cdot x_2 = x_3$, then $h \cdot (g \cdot x_1) = x_3 \implies (hg) \cdot x_1 = x_3$. Thus $x_1 \sim x_3$.

□

1.3 Orbits and Stabilizers

The equivalence classes under the relation \sim partition the set X . These classes are called **orbits**.

Definition 1.5 (Orbit). For $x \in X$, the orbit of x is the set of all places x can be moved to by G :

$$\mathcal{O}_x = \text{Orb}(x) = \{g \cdot x : g \in G\}.$$

Definition 1.6 (Stabilizer). For $x \in X$, the stabilizer of x is the set of elements in G that fix x :

$$G_x = \text{Stab}(x) = \{g \in G : g \cdot x = x\}.$$

Note that G_x is a subgroup of G (denoted $G_x \leq G$).

Theorem 1.7 (Orbit-Stabilizer Theorem). *Let G act on X . For any $x \in X$, there is a bijection between the orbit \mathcal{O}_x and the set of left cosets G/G_x . Consequently:*

$$|\mathcal{O}_x| = [G : G_x].$$

If G is finite, $|G| = |\mathcal{O}_x| \cdot |G_x|$.

Proof. Define a map $\psi : G/G_x \rightarrow \mathcal{O}_x$ by $\psi(gG_x) = g \cdot x$.

1. **Well-defined:** Suppose $g_1G_x = g_2G_x$. Then $g_1^{-1}g_2 \in G_x$, so $(g_1^{-1}g_2) \cdot x = x$, implying $g_2 \cdot x = g_1 \cdot x$.
2. **Injectivity:** If $\psi(g_1G_x) = \psi(g_2G_x)$, then $g_1 \cdot x = g_2 \cdot x$. Multiplying by g_1^{-1} , we get $x = g_1^{-1}g_2 \cdot x$, so $g_1^{-1}g_2 \in G_x$, implying $g_1G_x = g_2G_x$.
3. **Surjectivity:** By definition, any $y \in \mathcal{O}_x$ is of the form $g \cdot x$ for some g , which is exactly $\psi(gG_x)$.

Thus, ψ is a bijection. □

2 Lecture 2: Cauchy's Theorem & Conjugation (Jan 14)

2.1 Cauchy's Theorem

Lemma 2.1. *If X is a finite set and G is a p -group (a group of order p^k) acting on X , then:*

$$|X| \equiv |X^G| \pmod{p},$$

where $X^G = \{x \in X : g \cdot x = x, \forall g \in G\}$ is the set of fixed points.

Theorem 2.2 (Cauchy's Theorem). *If G is a finite group and p is a prime dividing $|G|$, then G has an element of order p .*

Proof (McKay's Proof). Let X be the set of p -tuples of elements of G whose product is the identity:

$$X = \{(g_1, \dots, g_p) \in G^p : g_1 g_2 \cdots g_p = 1\}.$$

Notice that g_p is uniquely determined by the first $p-1$ elements ($g_p = (g_1 \cdots g_{p-1})^{-1}$), so $|X| = |G|^{p-1}$. Since $p \mid |G|$, we have $p \mid |X|$.

Let \mathbb{Z}_p (cyclic group of order p) act on X by cyclic shift:

$$\sigma \cdot (g_1, g_2, \dots, g_p) = (g_2, g_3, \dots, g_p, g_1).$$

This is a valid action because if $g_1 \cdots g_p = 1$, then $g_1(g_2 \cdots g_p) = 1 \implies g_2 \cdots g_p = g_1^{-1}$, so $(g_2 \cdots g_p)g_1 = g_1^{-1}g_1 = 1$.

By the Lemma, $|X| \equiv |X^{\mathbb{Z}_p}| \pmod{p}$. Fixed points are tuples (a, a, \dots, a) such that $a^p = 1$. Since $(1, \dots, 1) \in X^{\mathbb{Z}_p}$, the set of fixed points is non-empty. Since p divides $|X|$ and p divides the congruence $|X| - |X^{\mathbb{Z}_p}|$, p must divide $|X^{\mathbb{Z}_p}|$. Therefore, there are at least p fixed points. There must exist some $a \neq 1$ such that $a^p = 1$. \square

2.2 Conjugation

Conjugation is a specific action of G on itself.

Definition 2.3. For $g, h \in G$, the **conjugate** of g by h is $g^h := hgh^{-1}$. The map $\phi_h : G \rightarrow G$ defined by $g \mapsto hgh^{-1}$ is an automorphism (an inner automorphism).

Definition 2.4 (Classes and Centralizers). Let G act on itself by conjugation ($h \cdot g = hgh^{-1}$).

- The orbit of g is the **Conjugacy Class** of g : $\text{Cl}(g) = \{hgh^{-1} : h \in G\}$.
- The stabilizer of g is the **Centralizer** of g : $C_G(g) = \{h \in G : hg = gh\}$.

By Orbit-Stabilizer: $|\text{Cl}(g)| = [G : C_G(g)]$.

Definition 2.5 (Center of G). The **Center**, $Z(G)$, is the set of elements that commute with everything:

$$Z(G) = \{z \in G : zg = gz, \forall g \in G\} = \bigcap_{g \in G} C_G(g).$$

$Z(G)$ is the kernel of the conjugation homomorphism $G \rightarrow \text{Aut}(G)$. Thus $Z(G) \triangleleft G$.

Definition 2.6 (Normalizer). Let $H \leq G$. The **Normalizer** of H in G is:

$$N_G(H) = \{g \in G : gHg^{-1} = H\}.$$

$N_G(H)$ is the largest subgroup of G in which H is normal.

3 Lecture 3: Sylow Preliminaries & Isomorphism Theorems (Jan 16)

3.1 Product of Subgroups

Definition 3.1. Let $K, N \leq G$. Define $KN = \{kn : k \in K, n \in N\}$.

- In general, KN is not a subgroup.
- If $N \triangleleft G$ (or just $N \subseteq N_G(K)$), then KN is a subgroup.
- Size formula: $|KN| = \frac{|K||N|}{|K \cap N|}$.

Theorem 3.2 (Second Isomorphism Theorem). *Let $K \leq G$ and $N \triangleleft G$. Then $K \cap N \triangleleft K$, and*

$$\frac{KN}{N} \cong \frac{K}{K \cap N}.$$

Proof. Consider the natural projection $\pi : G \rightarrow G/N$. Restrict it to K , i.e., $\phi = \pi|_K : K \rightarrow G/N$. The image of ϕ is $\{kN : k \in K\} = KN/N$. The kernel of ϕ is $\{k \in K : kN = N\} = \{k \in K : k \in N\} = K \cap N$. By the First Isomorphism Theorem, $K/\ker(\phi) \cong \text{Im}(\phi)$. \square

3.2 Sylow Definitions

Definition 3.3 (p -group). Let p be a prime. A group G is a p -group if every element has order a power of p . For finite groups, this is equivalent to $|G| = p^k$.

Definition 3.4 (Sylow p -subgroup). Let $|G| = p^n m$ where $p \nmid m$. A subgroup $P \leq G$ is called a **Sylow p -subgroup** if $|P| = p^n$.

$$\text{Syl}_p(G) = \{P \leq G : P \text{ is a Sylow } p\text{-subgroup}\}.$$

4 Lecture 4: The Sylow Theorems (Jan 21)

Theorem 4.1 (The Sylow Theorems). Let G be a finite group of order $p^n m$ where $p \nmid m$.

1. **Existence:** $\text{Syl}_p(G) \neq \emptyset$. (There exists a subgroup of order p^n).
2. **Conjugacy:** Any two Sylow p -subgroups are conjugate in G . That is, if $P, Q \in \text{Syl}_p(G)$, $\exists g \in G$ such that $gPg^{-1} = Q$.
3. **Number:** Let $n_p = |\text{Syl}_p(G)|$. Then:
 - $n_p \equiv 1 \pmod{p}$.
 - $n_p \mid m$ (equivalently $n_p \mid |G|$).

Proof. Let p be a prime such that $p \mid |G|$. Define the set of all p -subgroups:

$$\Sigma = \{H : H \leq G, |H| = p^n \text{ for some } n > 0\}.$$

Define Ω as the set of maximal elements in Σ under inclusion:

$$\Omega = \{H : H \in \Sigma, \text{ there is no } K \in \Sigma \text{ such that } H \subsetneq K\}.$$

We let G act on the set of subgroups $\{H : H \leq G\}$ by conjugation. Since conjugation is an isomorphism ($H^g \simeq H$), it preserves the order of subgroups. Thus:

$$H \in \Sigma \iff H^g \in \Sigma \quad \text{and} \quad H \in \Omega \iff H^g \in \Omega.$$

Therefore, G acts on Ω by $g \cdot H = H^g$.

Claim 1: Let $H \in \Omega$. Consider the action of H on Ω by conjugation. Then H is the *unique* fixed point of this action.

Proof of Claim:

- **Existence:** It is trivial to see that H is a fixed point, as $H^h = hHh^{-1} = H$ for any $h \in H$.
- **Uniqueness:** Let $K \in \Omega$ be a fixed point of the action of H .

$$K \text{ is fixed by } H \iff K^h = K \quad \forall h \in H \iff H \leq N_G(K).$$

Since $K \triangleleft N_G(K)$, and $H \leq N_G(K)$, the product HK is a subgroup of $N_G(K)$, and thus $HK \leq G$. Using our counting lemma:

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

Since $H, K \in \Omega$, their orders are powers of p . Thus $|HK|$ is a power of p , meaning $HK \in \Sigma$.

We have $H \leq HK$. Since H is maximal in Σ (by definition of Ω) and $HK \in \Sigma$, it must be that $H = HK$. Similarly, $K \leq HK$ implies $K = HK$. Therefore, $H = K$.

Claim 2: Size of Ω Modulo p **Statement:** $|\Omega| \equiv 1 \pmod{p}$.*Proof of Claim:* Let $H \in \Omega$. We decompose Ω into disjoint orbits under the action of H .

$$\Omega = \{H\} \cup \bigcup_{K \neq H} \text{Orb}_H(K).$$

By Claim 1, H is the only fixed point (an orbit of size 1). For any $K \in \Omega$ with $K \neq H$, the stabilizer of K in H is a proper subgroup, so by the Orbit-Stabilizer theorem for p -groups, $|\text{Orb}_H(K)|$ is divisible by p .

$$|\Omega| = 1 + \sum (\text{multiples of } p) \implies |\Omega| \equiv 1 \pmod{p}.$$

Claim 3: Transitivity (Conjugacy of Elements in Ω)**Statement:** Any two elements of Ω are conjugate.*Proof of Claim:* Suppose for contradiction that $H, K \in \Omega$ are not conjugates. Let $\theta_1 = \text{Orb}_G(H)$ and $\theta_2 = \text{Orb}_G(K)$. Since they are distinct orbits, $\theta_1 \cap \theta_2 = \emptyset$.Consider the action of H on these sets:

- On θ_1 : $H \in \theta_1$. By Claim 1, H is the unique fixed point of H in Ω . Since $\theta_1 \subseteq \Omega$, H is the unique fixed point in θ_1 . Thus, $|\theta_1| \equiv 1 \pmod{p}$.
- On θ_2 : Since $H \in \theta_1$ and $\theta_1 \cap \theta_2 = \emptyset$, we have $H \notin \theta_2$. Therefore, θ_2 contains *no* fixed points under the action of H (because the only fixed point in all of Ω is H). Thus, every orbit of H inside θ_2 has size divisible by p . This implies $|\theta_2| \equiv 0 \pmod{p}$.

By symmetry, we can swap the roles of H and K . Running the same argument with K acting on the sets implies $|\theta_2| \equiv 1 \pmod{p}$ and $|\theta_1| \equiv 0 \pmod{p}$.

This results in a contradiction (e.g., $|\theta_1| \equiv 1$ and $|\theta_1| \equiv 0$). Thus, $\theta_1 = \theta_2$, so H and K are conjugate.

Corollary: Since there is only one orbit, $\Omega = \theta_1$. Thus $|\Omega| = |\theta_1| \equiv 1 \pmod{p}$.**Claim 4: Divisibility****Statement:** $|\Omega| \mid |G|$.*Proof of Claim:* Let $H \in \Omega$. Since Ω is exactly the orbit of H under conjugation (from Claim 3), the size of the orbit is the index of the stabilizer:

$$|\Omega| = [G : \text{Stab}_G(H)] = [G : N_G(H)] = \frac{|G|}{|N_G(H)|}.$$

Thus $|\Omega|$ divides $|G|$.**Claim 5: Identification with Sylow Subgroups****Statement:** $\Omega = \text{Syl}_p(G)$.*Proof of Claim:* We know $\text{Syl}_p(G) \subseteq \Omega$ because Sylow subgroups are maximal p -subgroups by definition. We must show the reverse: every $H \in \Omega$ is a Sylow p -subgroup.

Let $H \in \Omega$. Assume for contradiction that $H \notin \text{Syl}_p(G)$. Let $|G| = p^s m$ where $p \nmid m$. Since H is a p -group but not Sylow, $|H| = p^t$ with $t < s$.

Consider the normalizer $N_G(H)$. From Claim 2, we know $|\Omega| = [G : N_G(H)] \equiv 1 \pmod{p}$. Therefore, p does *not* divide $[G : N_G(H)]$. Since $|G| = [G : N_G(H)] \cdot |N_G(H)|$, all factors of p in $|G|$ must reside in $|N_G(H)|$. Thus, $|N_G(H)|$ is divisible by p^s .

Consequently, the index $[N_G(H) : H] = \frac{|N_G(H)|}{|H|} = \frac{(\text{multiple of } p^s)}{p^t}$ is divisible by p (since $s > t$).

By Cauchy's Theorem applied to the quotient group $N_G(H)/H$, there exists a subgroup of order p , say $\bar{K} \leq N_G(H)/H$. By the Correspondence Theorem, there exists a subgroup K such that $H \leq K \leq N_G(H)$ with $|K| = p \cdot |H| = p^{t+1}$.

Since K is a p -group, $K \in \Sigma$. However, $H \subsetneq K$, which contradicts the definition of H as a maximal element in Ω .

Therefore, H must already be of order p^s . Thus $\Omega = \text{Syl}_p(G)$. □

5 Lecture 5: Normal Series & Solvability (Jan 23)

5.1 Normal and Subnormal Series

Definition 5.1. A **subnormal series** of a group G is a chain of subgroups:

$$\{1\} = H_0 \leq H_1 \leq \cdots \leq H_k = G$$

such that $H_i \triangleleft H_{i+1}$ for all i . The quotient groups H_{i+1}/H_i are called the **factors** of the series.

Definition 5.2 (Composition Series). A subnormal series is a **composition series** if all factors H_{i+1}/H_i are **simple** groups (non-trivial groups with no normal subgroups other than $\{1\}$ and themselves).

Theorem 5.3 (Jordan-Hölder). *Any two composition series of a finite group G are equivalent. That is, they have the same length, and their factors are isomorphic (up to re-ordering).*

5.2 Solvable Groups

Definition 5.4. A group G is **solvable** if it has a subnormal series where all factors H_{i+1}/H_i are **abelian**.

Definition 5.5 (Commutator). The commutator of g, h is $[g, h] = ghg^{-1}h^{-1}$. The **Derived Subgroup** (or Commutator Subgroup) G' or $[G, G]$ is the subgroup generated by all commutators.

Proposition 5.6. G/N is abelian if and only if $[G, G] \leq N$. Thus, $[G, G]$ is the smallest normal subgroup such that the quotient is abelian.

Remark 5.7 (Derived Series). Define $G^{(0)} = G$, $G^{(1)} = [G, G]$, and $G^{(i+1)} = [G^{(i)}, G^{(i)}]$. G is solvable if and only if the derived series terminates at $\{1\}$ (i.e., $G^{(n)} = \{1\}$ for some n).

6 Lecture 6: Solvable and Derived Series (Jan 26)

Lemma 6.1. *Let $\alpha \in \text{Aut}(G)$ then $\alpha([g, h]) = [\alpha(g), \alpha(h)]$.*

Lemma 6.2. *For $N \triangleleft G$ then G/N is abelian iff $[G, G] \leq N$.*

Theorem 6.3. *If $N \triangleleft G$, $g, h \in G$ and $[gN, hN] = [g, h]N$. Additionally, gN and hN commutes iff $[g, h]N = N$ iff $[g, h] \subseteq N$.*

Definition 6.4. G is solvable iff there is a subnormal series $\langle H_i : 0 \leq i \leq t \rangle$ such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$.

Theorem 6.5. *The following are true*

1. *If G solvable and $K \leq G$ then K is solvable*
2. *If G is solvable and $N \triangleleft G$ then G/N is solvable.*
3. *For $N \triangleleft G$, G solvable iff both N and G/N are solvable*

Proof. We prove the previous theorems

1. Let $\langle H_i : 0 \leq i \leq t \rangle$ be a series in G such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$. We can let $H'_i = K \cap H_i$ for $0 \leq i \leq t$ and we need to verify $\langle H'_i : 0 \leq i \leq t \rangle$ is a solvable series in K .
2. Let $\langle H_i : 0 \leq i \leq t \rangle$ be a series in G such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$. Let $H'_i = \phi_N[H_i] = H_iN/N$. Verify it is solvable.
3. The forward direction is trivial by (1) and (2). For the reverse direction, N solvable and G/N solvable $\langle N_i : 0 \leq i \leq s \rangle$ subnormal in N and N_{i+1}/N_i abelian and similar for $\langle H_jN/N : 0 \leq j \leq t \rangle$ subnormal in G/N and $(H_{j+1}N/N)_{j+1}/(H_jN/N)$ abelian. We know $H_{j+1} \triangleleft H_j$ and $[H_{j+1}, H_{j+1}] \leq H_j$ for all $0 \leq j \leq t-1$. So $N_0 = 1 \triangleleft N_1 \cdots \triangleleft N_s = N = H_0 \triangleleft H_1 \cdots \triangleleft H_t = G$.

□

Definition 6.6. Given G , the derived series of G is given by $G'_0 = G$ and $G'_{i+1} = [G'_i, G'_i]$ for $i \geq 0$.

Theorem 6.7. *G is solvable iff there is n such that $G'_n = \{1\}$.*

Proof. For the backwards direction, by construction G'_i/G'_{i+1} is abelian. Let $H_j = G_{n-j}$. For the forward direction, let $\langle H_j : 0 \leq j \leq n \rangle$ be a subnormal series in G , H_{j+1}/H_j abelian. Show by induction that $G'_i \leq H_{n-i}$ for all i . □

Theorem 6.8. *Suppose G abelian and simple, let $a \in G$ and $a \neq 1$ then $G = \langle a \rangle$ and $|a|$ is prime.*

Theorem 6.9. *Let G be simple and let $\langle H_i : 0 \leq i \leq t \rangle$ be subnormal series in G . Then there is $n < t$ such that $H_i = 1$ for $0 \leq i \leq n$ and $H_i = G$ for $n < i \leq t$.*

7 Lecture 7: Nilpotent Groups and Free Groups (Jan 28)

7.1 Nilpotent Groups

Definition 7.1. For $H, K \leq G$, $[H, K] = \langle [h, k] : h \in H, k \in K \rangle$

Remark 7.2. G'_i is characteristic in G for all i , in particular $G'_i \triangleleft G$.

Lemma 7.3. If $N \triangleleft G$, $N \leq H \leq G$ then

$$\frac{H}{N} \leq Z\left(\frac{G}{N}\right) \iff [G, H] \leq N.$$

Definition 7.4. A normal series in G , $\langle H_i : 0 \leq i \leq t \rangle$ is a central series if $\frac{H_{i+1}}{H_i} \leq Z\left(\frac{G}{H_i}\right)$ for all $0 \leq i \leq t-1$. Equivalently, $[G, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$.

Definition 7.5. G is nilpotent iff G has a central series.

Remark 7.6. Nilpotent groups are solvable.

Definition 7.7. Let G be a group

1. The descending central series is the sequence of subgroup given by $\gamma_1(G) = G$ and $\gamma_{i+1}(G) = [G, \gamma_i(G)]$ for $i \geq 1$.
2. The ascending central series is the sequence of subgroup given by $Z_0(G) = 1$ and

$$\frac{Z_{n+1}(G)}{Z_n(G)} = Z\left(\frac{G}{Z_n(G)}\right).$$

Remark 7.8. By induction, $Z_n(G)$ and $\gamma_n(G)$ are characteristic in G . and $Z_n(G) \triangleleft G$ and $\gamma_n(G) \triangleleft G$ for all $n \geq 0$.

Remark 7.9. For any G and $N \triangleleft G$, $[G, N] \leq N$ because for any $g \in G, n \in N$ we have $[g, n] = gng^{-1}n^{-1}$ and $n^g \in N$ since $N \triangleleft G$. So $[G, N] \leq N$.

Theorem 7.10. G is nilpotent iff there is a $N \geq 0$ such that $Z_n(G) = G$ iff there is $n \geq 0$ such that $\gamma_{n+1}(G) = \{1\}$. Moreover, the least n such that $Z_n(G) = G$ is the least n such that $\gamma_{n+1}(G) = \{1\}$.

Proof. Take any central series then the decreasing and ascending central series will grow at least as fast as the arbitrary central series. \square

7.2 Free Groups

Example 7.11. Consider $\mathbb{Z} = (\mathbb{Z}, +)$, 1 is a generator $\mathbb{Z} = \langle 1 \rangle$. The universal property of \mathbb{Z} , 1. For any group G and any $g \in G$, there is a unique homomorphism $\phi : \mathbb{Z} \rightarrow G$ and $\phi(1) = g$.

Proof. If ϕ exist let $\phi(n) = g^n$ then ϕ is a homomorphism. \square

g is free because whatever 1 is mapped to in the homomorphism it won't mess you up compared to a finite group.

Definition 7.12. F is a free group on 2 element if

1. There exist $a, b \in F$, $F = \langle a, b \rangle$
2. For all G , all $g, h \in G$ there is a unique homomorphism $\phi : F \rightarrow G$ such that $\phi(a) = g$ and $\phi(b) = h$.

8 Lecture 8: Free Groups

Theorem 8.1. For any set X , there are a group F and an injective function $i : X \rightarrow F$ such that

1. For any group G and any function $f : X \rightarrow G$ there is a unique homomorphism $\phi : F \rightarrow G$ such that $\phi(i(x)) = f(x)$ for all $x \in X$.

$$\begin{array}{ccc} X & \xrightarrow{i} & F \\ & \searrow f & \downarrow \phi \\ & & G \end{array}$$

The diagram commutes: $\phi \circ i = f$. The homomorphism ϕ is unique.

2. F is generated by $i[X]$

Proof. A word (on alphabet X) is a (possibly empty) finite sequence from $X \times \mathbb{Z}$. We will write $x_1^{n_1} x_2^{n_2} \cdots x_k^{n_k}$ for the word x_1, x_2, \dots, x_k for $\langle (x_1, n_1), (x_2, n_2), \dots, (x_k, n_k) \rangle$.

Example 8.2. Let $X = \{x, y, z\}$ then $w = x^0 x^{15} y^{-23} z^{27}$ is a word.

Let w be the set of words. Given a word, we can possibly perform a reduction step

1. If word contains an entry $\langle x, 0 \rangle$, remove it.
2. If the word contains successive entries $\langle x, n \rangle$ and $\langle x, m \rangle$ then replace them with $\langle x, n + m \rangle$.

Example 8.3. $y^{15} x^3 x^{-2} x^{-1} y^2 z^5 \rightarrow y^{15} x^1 x^{-1} y^2 z^5 \rightarrow y^{15} x^0 y^2 z^5 \rightarrow y^{15} y^2 z^5 \rightarrow y^{17} z^5$.

Let R be set of all reduced words, where no more reductions can be done.

For any word w , there is a unique reduced word w' such that w can be transformed into w' by a finite set of reductions.

We will not be able to show this property using group properties but instead we'll produce a group F with the property that every word is equivalent to a unique reduced word.

Given $x \in X$, I will replace $f_x : F \rightarrow F$

1. w does not begin with some x^m then $f_x(w) = xw$
2. w is of the form $x^{-1}v$ then $f_x(w) = v$
3. w is of the form $x^n v$ for $n \neq -1$ then $f_x(w) = x^{n+1}v$

I also define $g_x : R \rightarrow R$ by

1. w does not begin with some x^n then $g_x(w) = x^{-1}w$
2. w is of the form $x^1 v$ then $g_x(w) = v$
3. w is of the form $x^n v$ for $n \neq 1$ then $g_x(w) = x^{n-1}v$

We can observe that $f_x \circ g_x = g_x \circ f_x = 1_R$.

For each $x \in X$, $f_x \in \sum_R$ (the group of permutations of R) and $g_x = f_x^{-1}$. Let F be the subgroup of \sum_R generated by $\{f_x : x \in X\}$ or $\{f_{x_1}^{n_1}, \dots, f_{x_t}^{n_t} : x_i \in X, n_i \in \mathbb{Z}\}$.

Let $i : X \rightarrow F$ with $i(x) = f_x$. For any word $w = x_1^{n_1} x_2^{n_2} \dots x_t^{n_t}$. Let $f_w = f_{x_1}^{n_1} f_{x_2}^{n_2} \dots f_{x_t}^{n_t} \in F$.

Key Facts:

1. If w is obtained from \bar{w} by a single reduction step, $f_w = f_{\bar{w}}$.
2. If w is obtained from \bar{w} by any sequence of reduction steps, then $f_w = f_{\bar{w}}$.
3. If $w \in R$ then $f_w(\langle \rangle) = w$
4. For any word w , $f_w(\langle \rangle)$ the unique reduced word to which w can be transformed by reduction step.
5. If w_1, w_2 are reduced words then $v =$ unique reduction of $w_1 w_2$, $f_v = f_{w_1} f_{w_2}$

Let F' be the group whose underlying set is R whose operation is "concatenate and reduce". Then $F \simeq F'$ by $f_v \mapsto v$. To finish, let G be a group, $h : X \rightarrow G$ function. We want to find there is a unique homomorphism ϕ with $F' \rightarrow G$ and $\phi \circ i = h$.

1. If ϕ exists then

$$\begin{aligned} \phi(x_1^{n_1} x_2^{n_2} \dots x_t^{n_t}) &= \phi(i(x_1))^{n_1} \phi(i(x_2))^{n_2} \dots \phi(i(x_t))^{n_t} \\ &= h(x_1)^{n_1} h(x_2)^{n_2} \dots h(x_t)^{n_t} \end{aligned}$$

2. Verify ϕ is a homomorphism.

□

Example 8.4. Let $X = \langle x, y \rangle$, let G be any group with 2 generators say $G = \langle a, b \rangle$. Let F be a free group on X . Let $\phi : F \rightarrow G$ be unique homomorphism such that $\phi(x) = a$ and $\phi(y) = b$. ϕ is surjective because G is generated by a and b . Thus, $G \simeq \frac{F}{\ker \phi}$.

9 Lecture 9: Generators and Relations (Feb 2)

Definition 9.1. let X be a set, let F be a free group on X . Let $A \subseteq F$ and $N =$ least normal subgroup of F containing A .

Theorem 9.2. Let $\bar{F} = F/N$ and $j : X \rightarrow \bar{F}$ by $j(x) = xN$.

$$\begin{array}{ccc} F & \xrightarrow{\phi_N} & \bar{F} \\ i \uparrow & j \nearrow & \\ X & & \end{array}$$

The following are true:

1. \bar{F} is generated by $\{j(x) : x \in X\}$. Let $\bar{x} = j(x)$

2. If $a = x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t} \in A$ then $a \in N$ and $\bar{x}_1^{n_1} \bar{x}_2^{n_2} \cdots \bar{x}_t^{n_t} = \phi_N(a) = 1_{\bar{F}}$

"Universal Property of \bar{F} ": Let G be a group and let $h : X \rightarrow G$ such that $a = x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t} \in A$ and $h(x_1)^{n_1} h(x_2)^{n_2} \cdots h(x_t)^{n_t} = 1_G$. There is an unique homomorphism $\mathcal{T} : \bar{F} \rightarrow G$ such that

$$\begin{array}{ccc} X & \xrightarrow{j} & \bar{F} \\ & h \searrow & \downarrow \mathcal{T} \\ & & G \end{array}$$

So $h = \mathcal{T} \circ j$

Proof. Since $\{\bar{x} : x \in X\}$ generates \bar{F} there is at most one τ . By universal property of F , there is unique homomorphism $\phi : F \rightarrow G$ such that

$$\phi(x^1) = h(x). \quad (\text{For all } x \in X)$$

Use property of F

$$\begin{array}{ccc} X & \xrightarrow{i} & F \\ & h \searrow & \downarrow \phi \\ & & G \end{array}$$

Claim: By hypothesis on h , $A \subseteq \ker(\phi)$.

Let $a = x_1^{t_1} x_2^{t_2} \cdots x_k^{t_k} \in A$ then $\phi(a) = \phi(x_1)^{t_1} \phi(x_2)^{t_2} \cdots \phi(x_k)^{t_k} = h(x_1)^{t_1} h(x_2)^{t_2} \cdots h(x_k)^{t_k} = 1_G$ since $h(x_1)^{t_1} h(x_2)^{t_2} \cdots h(x_k)^{t_k} = 1_G$. Thus, $a \in \ker(\phi)$.

By choice of N , $N \leq \ker(\phi)$, we'll attempt to define $\tau : \bar{F} \rightarrow G$ by $\tau(wN) = \phi(w)$ for $w \in F$.

τ is well defined because $w_1 N = w_2 N \implies w_1^{-1} w_2 \in N \implies w_1^{-1} w_2 \in \ker(\phi) \implies \phi(w_1)^{-1} \phi(w_2) = 1 \implies \phi(w_1) = \phi(w_2)$. Showing τ is a homomorphism is straightforward.

$$\begin{array}{ccccc} X & \xrightarrow{i} & F & & \\ & h \searrow & \downarrow \phi_N & \searrow \phi & \\ & & \bar{F} & \xrightarrow{\tau} & G \end{array}$$

□

10 Lecture 10: Presentation of Groups (Feb 4)

Example 10.1. Dihedral Group: D_n = symmetry group of a regular n -gon.

Let $D_n = \langle a, b \rangle$ where a = rotation by $\frac{2\pi}{n}$ and b = reflection about the vertical axis. Then D_n is generated by a and b with the relations $a^n = 1$ and $b^2 = 1$.

We have $bab^{-1}a = 1$

Let $X = \{x, y\}$ and F be free group and $A = \{x^n, y^2, yxy^{-1}x\}$ and N = least normal subgroup $\supseteq A$. Let $\bar{x} = xN$ and $\bar{y} = yN$. Then $F/N = \langle \bar{x}, \bar{y} \rangle$ and

$$\overline{yxy^{-1}x} = \bar{y}\bar{x}\bar{y}^{-1}\bar{x}^{-1} = 1_{\bar{F}}.$$

There is a unique homomorphism $\tau : F/N \rightarrow D_n$ such that $\tau(\bar{x}) = a$ and $\tau(\bar{y}) = b$. as $D_n = \langle a, b \rangle$ so τ is surjective. So $|F/N| \geq 2n$.

Claim: $|F/N| \leq 2n$ (so τ is isomorphism)

Proof: Using relations from before, we can simply write any expression in \bar{x}, \bar{y} down to $\bar{x}^s \bar{y}^t$ for $0 \leq s \leq n$ and $0 \leq t \leq 2$.

11 Lecture 11: Category Theory (Feb 6)

11.1 Categories

Example 11.1. 1. Category of sets, if we have two sets A and B then $A \rightarrow B$ is a function $f : A \rightarrow B$

2. Category of groups, if we have two groups G and H then $G \rightarrow H$ is a homomorphism $\phi : G \rightarrow H$

3. Category of topological spaces, if we have two topological spaces X and Y then $X \rightarrow Y$ is a continuous function $f : X \rightarrow Y$

Definition 11.2. A category \mathcal{C} consists of the following data:

1. A collection of objects $Ob(\mathcal{C})$
2. A collection of morphisms $Mor(\mathcal{C})$
3. For each morphism f , there are objects $dom(f)$ and $cod(f)$ such that $dom(f) \rightarrow cod(f)$ is a morphism in $Mor(\mathcal{C})$
4. A composition operation $Mor(\mathcal{C}) \times Mor(\mathcal{C}) \rightarrow Mor(\mathcal{C})$
5. An identity morphism $1_A : A \rightarrow A$ for each object $A \in Ob(\mathcal{C})$

Lemma 11.3. *The following are true:*

1. If $a \xrightarrow{f} b$ then $hom(a, b) \neq \emptyset$