Complex Analysis Notes

Wilson Pan

February 25, 2026

Contents

1 Jan 12 **3**
1.1 Holomorphic Functions and Complex Differentiation . . . 3
1.2 Jacobian and Cauchy-Riemann Equations . . . 3

2 Jan 14 **4**
2.1 Holomorphic Equivalence and Wirtinger Derivatives . . . 4

3 Jan 16 **6**
3.1 Power Series . . . 6
3.2 Analytic Functions . . . 7

4 Jan 21 **8**
4.1 Parametrized Curves . . . 8
4.2 Contour Integration . . . 8
4.3 Primitives and the Fundamental Theorem . . . 9

5 Jan 23 **11**
5.1 Glossary of Elementary Functions . . . 11
5.2 Fundamental Theorem of Algebra . . . 11

6 Jan 26 **12**
6.1 Cauchy's and Goursat's Theorems . . . 12
6.2 Cauchy's Theorem in a Disk . . . 14

7 Jan 28 **16**
7.1 Applications of Cauchy's Theorem . . . 16

8 Jan 30 **19**
8.1 Cauchy's Integral Formula . . . 19
8.2 Regularity and Liouville's Theorem . . . 19

9 Feb 2 **21**
9.1 Rigidity and Analytic Continuation . . . 21
9.2 Morera's Theorem and Uniform Convergence . . . 21

10 Feb 4 **22**
10.1 Parameter Integrals and Gluing . . . 22
10.2 Schwarz Reflection Principle . . . 23

11 Feb 9 **24**
11.1 Meromorphic Functions and Poles . . . 24
11.2 Laurent Series and Residues . . . 24
11.3 The Residue Formula . . . 25

12 Feb 11 **27**
12.1 Integral Evaluation via Residues . . . 27
12.2 Removable Singularities . . . 27

13 Feb 13 **29**
13.1 Classification of Singularities 29
13.2 Meromorphic Functions and Behavior at Infinity 29

14 Feb 16 **30**
14.1 Rational Functions via Meromorphy 30

15 Feb 18 **31**
15.1 Argument Principle and Rouche's Theorem 31
15.2 Open Mapping and Maximum Modulus Theorems 31

16 Feb 20 **33**
16.1 Homotopy and Path Independence 33
16.2 Simply Connected Regions 34

17 Feb 23 **35**
17.1 Complex Logarithm 35
17.2 General Powers and Branches 35

18 Feb 25 **36**

1 Jan 12

1.1 Holomorphic Functions and Complex Differentiation

Definition 1.1. *Let Ω be an open set in $\mathbb{C}$ and f a complex-valued function on Ω. A function f is holomorphic at a point $z_0 \in \Omega$ if the quotient*

$$\frac{f(z_0+h)-f(z_0)}{h}.$$

converges to a limit when $h \to 0$. The limit of the quotient, when it exists, is denoted $f'(z_0)$.

Theorem 1.2. *A function $F(x,y) = (u(x,y), v(x,y))$ is said to be differentiable at a point $P_0(x_0, y_0)$ if there exist a linear transformation $J : \mathbb{R}^2 \to \mathbb{R}^2$ such that*

$$\frac{|F(P_0+H)-F(P_0)-J(H)|}{|H|} \to 0 \text{ as } |H| \to 0.$$

1.2 Jacobian and Cauchy-Riemann Equations

Definition 1.3. *The jacobian matrix of F is*

$$J = J_F(x,y) = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix}.$$

Theorem 1.4. *Cauchy-Riemann Equations: If f is holomorphic then*

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Proof. If we write $z = x+iy, z_0 = x_0+iy_0$ and $f(z) = f(x,y)$ then

$$\begin{aligned} f'(z_0) &= \lim_{h_1 \to 0} \frac{f(x_0+h_1, y_0) - f(x_0, y_0)}{h_1} \\ &= \frac{\partial f}{\partial x}(z_0) \end{aligned}$$

Similarly, now take h purely imaginary, say $h = ih_2$, then

$$\begin{aligned} f'(z_0) &= \lim_{h_2 \to 0} \frac{f(x_0, y_0+h_2) - f(x_0, y_0)}{ih_2} \\ &= \frac{1}{i}\frac{\partial f}{\partial y}(z_0) \end{aligned}$$

Since f is holomorphic we have

$$\frac{\partial f}{\partial x} = \frac{1}{i}\frac{\partial f}{\partial y}.$$

Writing $f = u+iv$ we have the stated relation. □

2 Jan 14

2.1 Holomorphic Equivalence and Wirtinger Derivatives

Theorem 2.1. *$f : \Omega \to \mathbb{C}$ holomorphic on Ω equivalently $F = \begin{pmatrix} u \\ v \end{pmatrix} : \mathbb{R}^2 \to \mathbb{R}^2$. The partials $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$ exist and satisfy the Cauchy-Riemann equations*

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Moreover, F is $\mathbb{R}$-differentiable, there exist a linear map $\mathcal{L} : \mathbb{R}^2 \to \mathbb{R}^2$ such that $F((x,y)+(h_1,h_2)) - F(x,y) = \mathcal{L}(h_1,h_2) + o(h)$

$$\mathcal{L} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix}.$$

Proof. We have

$$\frac{|F((x,y)+(h_1,h_2)) - F(x,y) - \mathcal{L}(h_1,h_2)|}{|h|} = \frac{\left| \begin{pmatrix} u(\overline{z}+\overline{h}) - u(\overline{z}) - \frac{\partial u}{\partial x}h_1 - \frac{\partial u}{\partial y}h_2 \\ v(\overline{z}+\overline{h}) - v(\overline{z}) - \frac{\partial v}{\partial x}h_1 - \frac{\partial v}{\partial y}h_2 \end{pmatrix} \right|}{|h|} \tag{1}$$

$$= \frac{|f(z+h) - f(z) - f'(z)h|}{|h|} \to 0 \tag{2}$$

Where $f(z+h) = u(z+h) + iv(z+h)$, $f(z) = u(z) + iv(z)$ and $f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}$. □

Corollary 2.2. *f holomorphic implies F is $\mathbb{R}$-differentiable. Let $a = \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ and $b = \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$*

$$DF_z = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \tag{3}$$

$$= \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \tag{4}$$

$$= \sqrt{a^2+b^2} \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \tag{5}$$

$$= |f'(z)| \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \tag{6}$$

Theorem 2.3. *If F is $\mathbb{R}$-differentiable with u, v satisfying Cauchy Riemann Equations then f is holomorphic.*

Proof. Let $a = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}$. We just need to check that $f(z+h) - f(z) - ah = o(h)$ □

Definition 2.4. *The Wirtinger Derivatives*

$$\frac{\partial}{\partial z} = \frac{1}{2}\left(\frac{\partial}{\partial x} - i\frac{\partial}{\partial y}\right) \text{ and } \frac{\partial}{\partial \overline{z}} = \frac{1}{2}\left(\frac{\partial}{\partial x} + i\frac{\partial}{\partial y}\right).$$

Remark 2.5. *The cauchy-riemann equations if and only if $\frac{\partial f}{\partial \overline{z}} = 0$*

We initially had the map $z \mapsto \overline{z}$ which is not holomorphic. Let $f(z) = \overline{z}$ and $\frac{\partial f}{\partial \overline{z}} = 1$ then we've captured this with $\frac{\partial}{\partial \overline{z}}$.
If we have $z, \overline{z}$ as independent variables with $z = x + iy$. Consider the map

$$(z, \overline{z}) \mapsto (x(z,\overline{z}), y(z,\overline{z})) \mapsto f(x,y).$$

We have

$$\partial_z f = \frac{\partial f}{\partial x}\frac{\partial x}{\partial z} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial z} = \frac{1}{2}\left(\frac{\partial}{\partial x} + \frac{1}{i}\frac{\partial}{\partial y}\right).$$

Theorem 2.6.

$$\frac{\partial}{\partial z} \cdot \frac{\partial}{\partial \overline{z}} = \frac{1}{4}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) = \frac{1}{4}\triangle.$$

3 Jan 16

3.1 Power Series

Definition 3.1. *Given a $\mathbb{C}$ valued seq $(a_n)_{n=0}^{\infty}$, form*

$$\sum_{n=0}^{\infty} a_n z^n.$$

A power series

1. *Converges if $\lim_{N\to\infty} \sum_{n=0}^{N} a_n z^n$ exists*
2. *Converges absolutely if $\sum_{n=0}^{\infty} |a_n||z|^n$ converges*

Remark 3.2. *By translation, all we say will apply to $\sum a_n(z-z_0)^n$*

Theorem 3.3. *Given $\sum_{n=0}^{\infty} a_n z^n$, $\exists R \in [0,+\infty]$ such that*

1. *It converges, if $D_R = \{|z| < R\}$*
2. *It diverges at every at z such that $|z| > R$.*

Moreover,

$$R = \frac{1}{\lim_{n\to\infty} \sup |a_n|^{\frac{1}{n}}}. \qquad \text{(Hadamard's Formula)}$$

Remark 3.4. *On ∂D_R it can often be delicate.*

Proof. Fix $|z| < R$, then $\frac{1}{R}|z| < 1$ and fix $\epsilon > 0$ such that

$$\left(\frac{1}{R} + \epsilon\right)|z| < 1.$$

Then, eventually

$$|a_n|^{\frac{1}{n}} < \frac{1}{R} + \epsilon.$$

Then

$$|a_n||z|^n \leq \left(|a_n|^{\frac{1}{n}}|z|\right)^n \leq \left(\left(\frac{1}{R} + \epsilon\right)|z|\right)^n \leq q^n.$$

So by the comparison test with $\sum q^n < \infty$ □

Example 3.5.

$$\sum_{n=0}^{\infty} \frac{z^n}{n!} = \exp(z).$$

This has $R = +\infty$

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} z^{2n} = \cos(z).$$

This has $R = +\infty$

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1} = \sin(z).$$

This has $R = +\infty$
Euler's Formula is

$$e^{iz} = \cos(z) + i\sin(z).$$

Theorem 3.6. *The function $f(z) = \sum a_n z^n$ is holomorphic on D_R, f' is a power series*

$$f'(z) = \sum n a_n z^{n-1}.$$

Whose disk of convergence is R.

Proof. Let $g(z) = \sum n a_n z^{n-1}$, $z \in D_R$. We want to show that f' exist and $f' = g$.
Fix $z_0 \in D_R$, let $r < R$ such that $|z_0| < r < R$.
Fix $N \geq 1$ (to be chosen later)

$$f(z) = \sum_{n=0}^{N} a_n z^n + \sum_{n=N+1}^{\infty} a_n z^n = S_N(z) + E_N(z).$$

From here we want to show that

$$\frac{f(z_0+h) - f(z_0)}{h} \to g(z_0).$$

For all h such that $|z_0 + h| < r$

$$\begin{aligned}\frac{f(z_0+h) - f(z_0)}{h} - g(z_0) &= \left(\frac{S_N(z_0+h) - S_N(z_0)}{h} - S_N'(z_0)\right) + (S_N'(z_0) - g(z_0)) + \left(\frac{E_N(z_0+h) - E_N(z_0)}{h}\right) \\ &= A + B + C\end{aligned}$$

Fix $\epsilon > 0$ we want to show that $\exists \delta$ such that $\forall |h| < \delta, |A + B + C| < \epsilon$.

$$\begin{aligned}|C| &\leq \sum_{n=N+1}^{\infty} |a_n| \left|\frac{(z_0+h)^n - z_0^n}{h}\right| \\ &= \sum_{n=N+1}^{\infty} |a_n| \left|(z_0+h)^{n-1} + (z_0+h)^{n-2} z_0 + \cdots + z_0^n\right| \\ &\leq \sum_{n=N+1}^{\infty} |a_n||nr^{n-1}| < \frac{\epsilon}{3} \qquad \text{(for all } N \geq N_1(\epsilon))\end{aligned}$$

Note the last summation is the tail of $\sum a_n n r^{n-1}$ which converges by $r < R$.
For B:

$$S_N'(z_0) = \sum_{n=0}^{N} n a_n z_0^{n-1} \to g(z_0). \qquad \text{(by def of } g)$$

So $\forall N \geq N_2$, $|B| < \frac{\epsilon}{3}$.
$|A| < \frac{\epsilon}{3}$ for all h sufficiently small by the definition of the complex derivative and the fact that S_N is plainly holomorphic. □

3.2 Analytic Functions

Definition 3.7. *Let Ω be open and define $f : \Omega \to \mathbb{C}$ is analytic at $z_0 \in \Omega$, if there exists a power series $\sum a_n (z - z_0)^n$ with positive radius of convergence such that $f(z) = \sum a_n (z - z_0)^n$ is a neighborhood of z_0.*
f is analytic on Ω, if analytic at every point of $z_0 \in \Omega$.

Corollary 3.8. *If f analytic on $\Omega \implies f$ is holomorphic on Ω.*
In fact, $\impliedby$.

4 Jan 21

4.1 Parametrized Curves

Definition 4.1. *A parametrized curve (in $\mathbb{C}$) is a function $z(t) : [a, b] \to \mathbb{C}$*

1. *Smooth if z' is continuous and $\forall t \in [a, b]$, $z'(t) \neq 0$*
2. *Piecewise smooth if z is continuous so there exists $a_0 = a < a_1 < ... < a_k = b$ such that $z\text{res}_{[a_j, a_{j+1}]}$ for all $j = 0, ..., k-1$ is smooth.*
3. *Let $\tilde{z} : [\tilde{a}, \tilde{b}] \to \mathbb{C}$ then two parameterization are equivalent if there exist c^1-bijection $s \mapsto t(s) : [\tilde{a}, \tilde{b}] \to [a, b]$ and $t'(s) > 0$ for all s such that $\tilde{z}(s) = z(t(s))$.*

Example 4.2. *A piecewise smooth curve can be visualized as follows:*



The curve consists of smooth segments joined at corner points (marked in red) where the derivative is discontinuous.

Definition 4.3. *The family of all equivalent parameterization determines a curve (oriented) $\gamma \subset \mathbb{C}, \gamma = z([a, b])$.*
$z(a)$ is the starting point and $z(b)$ is the endpoint of γ. $\gamma^{-1} = \gamma$ with reversed orientation

Definition 4.4. *γ is closed if $z(a) = z(b)$ and simple if $z(t_1) \neq z(t_2)$ for all $t_1 \neq t_2$.*

Example 4.5. $z(T) = re^{-it}, t \in [0, 2\pi]$

Definition 4.6. *A curve has positive orientation if interior is on the left or counterclockwise.*

4.2 Contour Integration

Definition 4.7. *Give a smooth curve $\gamma \subset \mathbb{C}$ (with parameterization $z : [a, b] \to \mathbb{C}$) with $f : \gamma \to \mathbb{C}$ continuous.*
We set

$$\int_\gamma f = \int_\gamma f(z)dz = \int_a^b f(z(t))z'(t)dt.$$

Since $f(z(t))z'(t) \in \mathbb{C}$ we split the integral into

$$\int_a^b \Re(...)dt + i\int_a^b \Im(...)dt.$$

Remark 4.8. *This is well defined as the right hand side does not depend on the choice of the parametrization.*

Definition 4.9. *For a piece-wise smooth curve, we define*

$$\int_\gamma f = \sum_{j=0}^{k-1} \int_{a_j}^{a_{j+1}} f(z(t))z'(t)dt.$$

Definition 4.10. *We define*

$$|\gamma| = length(\gamma) = \int_a^b |z'(t)|dt.$$

Theorem 4.11. *Some very basic properties:*

1. *Linearity*
$$\int_\gamma (\alpha f + \beta g) = \alpha \int_\gamma f + \beta \int_\gamma g.$$
2.
$$\int_{\gamma^{-1}} f = -\int_\gamma f.$$
3.
$$\left|\int_\gamma f\right| \leq \left(\sup_\gamma |f|\right) |\gamma|.$$

4.3 Primitives and the Fundamental Theorem

Definition 4.12. *The primitive of $f : \Omega \to \mathbb{C}$ with open $\Omega \subset \mathbb{C}$ is any $F : \Omega \to \mathbb{C}$ holomorphic such that $F' = f$ on Ω*

Theorem 4.13. *The Fundamental Theorem of Calculus*
For f with primitive F on Ω, $\gamma \subset \mathbb{C}$ curve from $w_1 \to w_2$ we have

$$\int_\gamma f = F(w_2) - F(w_1).$$

Proof. We can write

$$\int_\gamma f = \int_a^b f(z(t))z'(t)dt = \int_a^b \frac{d}{dt}(F(z(t)))dt = F(z(b)) - F(z(a)).$$

□

Corollary 4.14. *If γ is closed then*

$$\int_\gamma f = 0.$$

Corollary 4.15. *If f is holomorphic on Ω with $f' = 0$ on Ω then f is constant.*

Proof. Fix $z_0 \in \Omega$, for $z \in \Omega$, take $\gamma : z_0 \to z$ then

$$0 = \int_\gamma f' = f(z) - f(z_0).$$

□

Example 4.16. *Let $z(t) = e^{it}$*

$$\int_{\partial D} \frac{1}{z} dz = \int_0^{2\pi} \frac{1}{e^{it}} ie^{it} dt = 2\pi i \neq 0.$$

So $\frac{1}{z}$ *does not admit a primitive.*

5 Jan 23

5.1 Glossary of Elementary Functions

Glossary of Elementary Functions

1. Polynomials: $P(z) = a_0 + a_1 z + \cdots + a_n z^n$ for $a_i \in \mathbb{C}$ and is entirely holomorphic in $\mathbb{C}$, $a_n \neq 0$ and $n = \deg P$
2. Rational: $R(z) = \frac{P(z)}{Q(z)}$ where P, Q are polynomials with no common factors. The zeros of Q are called poles and their order is called the order of a given pole.
3. Exponential: $e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$

5.2 Fundamental Theorem of Algebra

Lemma 5.1. *Let $P(z)$ be a complex polynomial of positive degree n, if $|P(z)|$ has a local minimum at z_0 then $P(z_0) = 0$.*

Proof. WLOG $z_0 = 0$ as we can translate $Q(z) = P(z - z_0)$.
AFSOC $P(0) \neq 0$ and WLOG $P(0) = 1$ by $\frac{P}{P(0)}$. We can write $P(z) = 1 + a_k z^k + \cdots + a_z^n$ with $a_k, a_n \neq 0$.
Consider $z_\delta = \delta e^{i\theta}$. For all $\delta > 0$ sufficiently small:

$$\left|a_{k+1} z_\delta + \cdots + a_n z_\delta^{n-k}\right| < \frac{|a_k|}{2}.$$

Then

$$|P(z)| \leq \left|1 + a_k z_\delta^k\right| + \left|a_{k+1} z_\delta^{k+1} + \cdots + a_n z_\delta^n\right| < \left|1 + a_k z_\delta^k\right| + \left|z_\delta^k\right| \frac{|a_k|}{2}.$$

Now we can choose δ such that $a_k z_\delta^k < 0$ and real.

$$\left|1 + a_k z_\delta^k\right| + \left|z_\delta^k\right| \frac{|a_k|}{2} = 1 + a_k z_\delta^k - \frac{1}{2} a_k z_\delta^k = 1 + \frac{1}{2} a_k z_\delta^k < 1.$$

□

Theorem 5.2. *Fundamental Theorem of Algebra:*

Proof. $|P(z)| \to +\infty$ as $|z| \to +\infty$ therefore $\exists R$ such that $\forall |z| > R$, $|P(z)| > |P(0)|$ thus $\inf_{z \in \mathbb{C}} |P(z)| = \inf_{z \in D_R} |P(z)|$ and such a value is attained somewhere at z_0 then by lemma 5.1 we're done. □

Lemma 5.3. *$P(z) = a_n (z - z_1) \cdots (z - z_n)$ where $z_1, ..., z_n$ are roots of P.*
The order of a zero z_j is its multiplicity, the number of times it appears in the sequence. If z_j is of the order k_j,

$$P(z) = \underbrace{Q(z)}_{\text{non-vanishing}} (z - z_j)^{k_j}.$$

A zero z_0 is of order k_0 if and only if $P(z_0) = P'(z_0) = ... = P^{(k_0 - 1)}(z_0) = 0$ and $P^{(k_0)}(z_0) \neq 0$

Theorem 5.4. *(Gauss-Lucas) The roots of P' lies in the convex hull of the roots of P. In particular, if P is real rooted then so is P'.*

Theorem 5.5. *If R a rational function then R' has the same roots as R with order greater by 1.*

Theorem 5.6. *The following are true*

1. $(e^z)' = e^z$
2. $e^{w+z} = e^w e^z$
3. $\overline{e^z} = e^{\overline{z}}$
4. $|e^{iy}| = 1$ *and* $|e^{x+iy}| = e^x$
5. $\sin(z) = \frac{e^{iz}-e^{-iz}}{2i}$ *and* $\cos(z) = \frac{e^{iz}+e^{-iz}}{2}$

6 Jan 26

6.1 Cauchy's and Goursat's Theorems

Theorem 6.1. *Cauchy Theorem: If $f : \Omega \to \mathbb{C}$ is holomorphic on Ω and γ is a closed curve in Ω then*

$$\int_\gamma f = 0.$$

Theorem 6.2. *Goursat's Theorem: If $f : \Omega \to \mathbb{C}$ is holomorphic and Ω is open, then $\forall \triangle \subset \Omega$ we have*

$$\int_{\partial\triangle} f = 0.$$



The integral around the positively oriented (counterclockwise) boundary of any triangle $\triangle \subset \Omega$ vanishes.

Proof. Let $\triangle = \triangle^{(0)}$ and partition into four triangles by connecting the midpoints of the sides:

$$\triangle^{(0)} = \triangle_1^{(1)} \cup \triangle_2^{(1)} \cup \triangle_3^{(1)} \cup \triangle_4^{(1)}.$$



When we integrate along each sub-triangle with positive orientation, the interior edges cancel (traversed in opposite directions, shown with opposing arrows):

$$\int_{\partial\triangle^{(0)}} f = \sum_{j=1}^{4} \int_{\partial\triangle_j^{(1)}} f.$$

By the triangle inequality

$$\left|\int_{\partial\triangle^{(0)}} f\right| \leq \sum_{j=1}^{4} \left|\int_{\partial\triangle_j^{(1)}} f\right| \leq 4 \max_{j=1,2,3,4} \left|\int_{\partial\triangle_j^{(1)}} f\right|.$$

Thus there exists $j_0 \in \{1,2,3,4\}$ such that

$$\left|\int_{\partial\triangle^{(0)}} f\right| \leq 4 \left|\int_{\partial\triangle_{j_0}^{(1)}} f\right|.$$

Set $\triangle^{(1)} = \triangle_{j_0}^{(1)}$. Repeating this subdivision process, we obtain a nested sequence of triangles

$$\triangle^{(0)} \supset \triangle^{(1)} \supset \triangle^{(2)} \supset \cdots$$

such that

$$\left|\int_{\partial\triangle^{(0)}} f\right| \leq 4^n \left|\int_{\partial\triangle^{(n)}} f\right|.$$

Note that $\mathrm{diam}(\triangle^{(n)}) = 2^{-n}\mathrm{diam}(\triangle^{(0)})$ and $|\partial\triangle^{(n)}| = 2^{-n}|\partial\triangle^{(0)}|$.
Since the triangles are nested compact sets with diameters shrinking to zero, by Cantor's intersection theorem:

$$\bigcap_{n=0}^{\infty} \triangle^{(n)} = \{z_0\}$$

for some $z_0 \in \triangle^{(0)} \subset \Omega$. Since f is holomorphic at z_0, we can write

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \psi(z)(z - z_0)$$

where $\psi(z) \to 0$ as $z \to z_0$.
Since $f(z_0)$ and $f'(z_0)(z - z_0)$ have primitives ($f(z_0)z$ and $\frac{1}{2}f'(z_0)(z - z_0)^2$ respectively), their integrals over the closed curve $\partial\triangle^{(n)}$ vanish. Thus:

$$\int_{\partial\triangle^{(n)}} f(z)dz = \int_{\partial\triangle^{(n)}} \psi(z)(z - z_0)dz.$$

For $z \in \partial\triangle^{(n)}$, we have $|z - z_0| \leq \mathrm{diam}(\triangle^{(n)}) = 2^{-n}\mathrm{diam}(\triangle^{(0)})$.
Let $\epsilon_n = \sup_{z\in\partial\triangle^{(n)}} |\psi(z)|$. Since $\psi(z) \to 0$ as $z \to z_0$ and $\triangle^{(n)} \to \{z_0\}$, we have $\epsilon_n \to 0$.
Therefore:

$$\begin{aligned}\left|\int_{\partial\triangle^{(n)}} f(z)dz\right| &\leq \sup_{z\in\partial\triangle^{(n)}} |\psi(z)(z - z_0)| \cdot |\partial\triangle^{(n)}| \\ &\leq \epsilon_n \cdot 2^{-n}\mathrm{diam}(\triangle^{(0)}) \cdot 2^{-n}|\partial\triangle^{(0)}| \\ &= \epsilon_n \cdot 4^{-n}\mathrm{diam}(\triangle^{(0)})|\partial\triangle^{(0)}|\end{aligned}$$

Combining with our earlier inequality:

$$\left|\int_{\partial\triangle^{(0)}} f\right| \leq 4^n \left|\int_{\partial\triangle^{(n)}} f\right| \leq 4^n \cdot \epsilon_n \cdot 4^{-n}\mathrm{diam}(\triangle^{(0)})|\partial\triangle^{(0)}| = \epsilon_n \cdot \mathrm{diam}(\triangle^{(0)})|\partial\triangle^{(0)}|$$

Since $\epsilon_n \to 0$ as $n \to \infty$ and the right-hand side is independent of n except for ϵ_n, we conclude:

$$\int_{\partial\triangle} f = 0.$$

□

6.2 Cauchy's Theorem in a Disk

Theorem 6.3. *Cauchy's Theorem in a disk: Let $D \subset \mathbb{C}$ be an open disk and f holomorphic on D. Then for any closed curve γ in D we have*

$$\int_\gamma f = 0.$$

Proof. WLOG let D be a disk centered at 0. Define $F : D \to \mathbb{C}$ by

$$F(z) = \int_{\gamma_z} f$$

where γ_z is the path $0 \to \Re(z) \to z$ (horizontal then vertical).



We claim F is a primitive for f, i.e., $F'(z) = f(z)$.

Case 1: $h \in \mathbb{R}$ (horizontal increment).
The path γ_{z+h} goes $0 \to \Re(z) + h \to z + h$. Observe that:

$$F(z+h) - F(z) = \int_{\gamma_{z+h}} f - \int_{\gamma_z} f.$$



The difference of paths can be decomposed using Goursat's theorem. The integral over the rectangle with vertices $\Re(z), \Re(z)+h, z+h, z$ vanishes, so:

$$F(z+h) - F(z) = \int_{\Re(z)}^{\Re(z)+h} f(\zeta)d\zeta + \int_z^{z+h} f(\zeta)d\zeta = \int_z^{z+h} f(\zeta)d\zeta$$

where the first integral is along the real axis and the second is horizontal at height $\Im(z)$. By the rectangle lemma, these combine to give just the horizontal segment from z to $z+h$:

$$\frac{F(z+h) - F(z)}{h} = \frac{1}{h}\int_z^{z+h} f(\zeta)d\zeta \to f(z) \text{ as } h \to 0.$$

Case 2: $h = ik$ with $k \in \mathbb{R}$ (vertical increment).
Similarly, γ_{z+ik} goes $0 \to \Re(z) \to z+ik$. The paths γ_z and γ_{z+ik} share the horizontal segment $0 \to \Re(z)$, so:

$$F(z+ik) - F(z) = \int_z^{z+ik} f(\zeta)d\zeta.$$



Thus:

$$\frac{F(z+ik) - F(z)}{ik} = \frac{1}{ik}\int_z^{z+ik} f(\zeta)d\zeta \to f(z) \text{ as } k \to 0.$$

General h: For arbitrary $h = h_1 + ih_2$, we write:

$$\frac{F(z+h) - F(z)}{h} = \frac{F(z+h) - F(z+h_1)}{h} + \frac{F(z+h_1) - F(z)}{h}.$$

Using the above cases and the continuity of f, both terms converge appropriately, giving $F'(z) = f(z)$.

Since f has a primitive F on D, by the Fundamental Theorem of Calculus, for any closed curve γ in D:

$$\int_\gamma f = F(z(b)) - F(z(a)) = 0.$$

$\square$

7 Jan 28

7.1 Applications of Cauchy's Theorem

Example 7.1.

$$\frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty} e^{-x^2}\,dx = 1 \iff \int_{-\infty}^{\infty} e^{-\pi x^2}\,dx = 1.$$

Proof. First, note that by substitution $u = \sqrt{\pi}x$, we have:

$$\int_{-\infty}^{\infty} e^{-\pi x^2}\,dx = \frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty} e^{-u^2}\,du.$$

So the equivalence follows if we can show $\int_{-\infty}^{\infty} e^{-\pi x^2}\,dx = 1$.

Consider the function $f(z) = e^{-\pi z^2}$, which is entire (holomorphic everywhere). For fixed $y \in \mathbb{R}$, consider the rectangular contour Γ_R with vertices at $-R, R, R+iy, -R+iy$ (traversed counterclockwise).



By Cauchy's theorem:

$$\int_{\Gamma_R} e^{-\pi z^2}\,dz = 0.$$

Parameterizing the four sides:

$$\begin{aligned}\int_{\Gamma_R} e^{-\pi z^2}\,dz = &\int_{-R}^{R} e^{-\pi t^2}\,dt + \int_0^y e^{-\pi(R+it)^2}\,i\,dt \\ &+ \int_R^{-R} e^{-\pi(t+iy)^2}\,dt + \int_y^0 e^{-\pi(-R+it)^2}\,i\,dt\end{aligned}$$

We now bound the integrals over the vertical segments. For the right vertical segment, parameterize $z = R+it$ with $t \in [0,y]$ (assuming $y > 0$; the case $y < 0$ is similar). We have:

$$(R+it)^2 = R^2 + 2iRt - t^2 = (R^2 - t^2) + 2iRt$$

so

$$e^{-\pi(R+it)^2} = e^{-\pi(R^2-t^2)} \cdot e^{-2\pi iRt}.$$

Since $|e^{-2\pi iRt}| = 1$ for all real R and t, we have:

$$|e^{-\pi(R+it)^2}| = e^{-\pi(R^2-t^2)} \cdot |e^{-2\pi iRt}| = e^{-\pi(R^2-t^2)} \cdot 1 = e^{-\pi(R^2-t^2)}.$$

For $t \in [0,y]$, we have $t^2 \le y^2$, so $R^2 - t^2 \ge R^2 - y^2$. Therefore:

$$|e^{-\pi(R+it)^2}| \le e^{-\pi(R^2-y^2)} = e^{-\pi R^2}e^{\pi y^2}.$$

By the ML-inequality, the length of the path is $|y|$, so:

$$\left|\int_0^y e^{-\pi(R+it)^2}\,i\,dt\right| \le |y| \cdot e^{-\pi R^2}e^{\pi y^2} = |y|e^{\pi y^2}e^{-\pi R^2}.$$

Similarly, for the left vertical segment with $z = -R + it$ where $t \in [y, 0]$:

$$(-R + it)^2 = R^2 - 2iRt - t^2 = (R^2 - t^2) - 2iRt$$

so

$$e^{-\pi(-R+it)^2} = e^{-\pi(R^2-t^2)} \cdot e^{2\pi iRt}.$$

Since $|e^{2\pi iRt}| = 1$ for all real R and t, we have:

$$|e^{-\pi(-R+it)^2}| = e^{-\pi(R^2-t^2)} \cdot |e^{2\pi iRt}| = e^{-\pi(R^2-t^2)} \cdot 1 = e^{-\pi(R^2-t^2)}.$$

For $t \in [y, 0]$ (or equivalently $t \in [0, y]$ if we reverse the parameterization), we have $t^2 \leq y^2$, so:

$$|e^{-\pi(-R+it)^2}| \leq e^{-\pi(R^2-y^2)} = e^{-\pi R^2} e^{\pi y^2}.$$

Therefore:

$$\left| \int_y^0 e^{-\pi(-R+it)^2} i \, dt \right| \leq |y| \cdot e^{-\pi R^2} e^{\pi y^2} = |y| e^{\pi y^2} e^{-\pi R^2}.$$

Both vertical integrals are bounded by $Ce^{-\pi R^2}$ for some constant C depending on y but independent of R. Since $e^{-\pi R^2} \to 0$ as $R \to \infty$, we conclude that:

$$\lim_{R\to\infty} \int_0^y e^{-\pi(R+it)^2} i \, dt = 0 \quad \text{and} \quad \lim_{R\to\infty} \int_y^0 e^{-\pi(-R+it)^2} i \, dt = 0.$$

Therefore, taking the limit as $R \to \infty$:

$$\int_{-\infty}^{\infty} e^{-\pi t^2} dt = \int_{-\infty}^{\infty} e^{-\pi(t+iy)^2} dt.$$

Expanding $e^{-\pi(t+iy)^2} = e^{-\pi(t^2+2ity-y^2)} = e^{-\pi t^2} e^{-2\pi ity} e^{\pi y^2}$, we get:

$$\int_{-\infty}^{\infty} e^{-\pi t^2} dt = e^{\pi y^2} \int_{-\infty}^{\infty} e^{-\pi t^2} e^{-2\pi ity} dt.$$

In particular, taking $y = 0$ gives the standard Gaussian integral. The standard result is $\int_{-\infty}^{\infty} e^{-\pi x^2} dx = 1$, which completes the proof. □

Example 7.2.

$$\int_0^{\infty} \frac{1 - \cos(x)}{x^2} dx = \frac{\pi}{2}.$$

Proof. Consider the holomorphic function $f(z) = \frac{1-e^{iz}}{z^2}$ on $\mathbb{C} \setminus \{0\}$. Fix ϵ, R and consider the contour Γ which is disk with radius R and a small disk with radius ϵ centered at the origin.



We can express $e^{iz} \approx 1 + iz + g(z)$ where $g(z) = e^{iz} - 1 - iz$ so

$$\begin{aligned} \int_{\gamma_\epsilon} f(z) dz &= \int_{\gamma_\epsilon} \frac{1 - (1 + iz + g(z))}{z^2} dz \\ &= \int_{\gamma_\epsilon} \frac{-g(z)}{z^2} dz + \int_{\gamma_\epsilon} \frac{-i}{z} dz \end{aligned}$$

We can evaluate each part as

$$\int_{\gamma_\epsilon} \frac{dz}{z} = -\int_0^\pi \frac{ie^{it}}{e^{it}} dt = -\pi i.$$

$$\int_{\gamma_\epsilon} \frac{g(z)}{z^2} dz \to 0 \text{ as } \epsilon \to 0.$$

$\square$

8 Jan 30

8.1 Cauchy's Integral Formula

Theorem 8.1. *Cauchy's Integral Formula: Let Ω be open $f : \Omega \to \mathbb{C}$ be holomorphic. Let $\overline{D}$ be open in Ω be a disk then*

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Proof. Fix $z \in D_1$, let $\gamma_{\epsilon,\delta}$ be a key-hole contour but we omit z. Then

$$\int_{\gamma_{\epsilon,\delta}} \frac{f(\zeta)}{\zeta - z} d\zeta = 0.$$



Fix $\epsilon > 0$ and $\delta \to 0$ we want to show that

$$\int_{\gamma_{\epsilon,\delta}} \frac{f(\zeta)}{\zeta - z} d\zeta \to \int_{\partial D} - \int_{C_\epsilon}.$$

For

$$\int_{C_\epsilon} \frac{f(\zeta)}{\zeta - z} d\zeta = \int_{C_\epsilon} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta + f(z) \int_{C_\epsilon} \frac{d\zeta}{\zeta - z}.$$

The first term is bounded as holomorphic so as $\epsilon \to 0$, it goes to 0. The second part integrates to $f(z) \cdot 2\pi i$. □

8.2 Regularity and Liouville's Theorem

Theorem 8.2. *Regularity: Holomorphic implies Analytic.*
Let $f : \Omega \to \mathbb{C}$ be holomorphic, $\overline{D} = \overline{D}_r(z_0) \subset \Omega$ then f is analytic at z_0 i.e

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n.$$

Proof. We know

$$f(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

We can write

$$\frac{1}{\zeta - z} = \frac{1}{(\zeta - z_0) + (z_0 - z)} = \frac{1}{\zeta - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} = \frac{1}{\zeta - z_0} \sum_{n=0}^{\infty} \left(\frac{z - z_0}{\zeta - z_0} \right)^n.$$

Note: $\frac{z-z_0}{\zeta-z_0} < 1$.

$$\begin{aligned} f(z) &= \frac{1}{2\pi i}\int_{\partial D} \frac{f(\zeta)}{\zeta - z_0} \underbrace{\sum_{n=0}^{\infty} \left(\frac{z-z_0}{\zeta-z_0}\right)^n \frac{f(\zeta)}{\zeta - z}}_{\text{conv. uniformly on } \partial D} d\zeta \\ &= \frac{1}{2\pi i}\sum_{n=0}^{\infty}\int_{\partial D} \frac{f(\zeta)}{\zeta - z_0}\left(\frac{z-z_0}{\zeta-z_0}\right)^n d\zeta \\ &= \sum_{n=0}^{\infty}(z-z_0)^n \underbrace{\left(\frac{1}{2\pi i}\int_{\partial D}\frac{f(\zeta)}{(\zeta-z_0)^{n+1}}d\zeta\right)}_{a_n} \end{aligned}$$

So $a_n = \frac{f^{(n)}(z_0)}{n!}$. □

Corollary 8.3. *Cauchy's Formula:*

$$f^{(n)}(z_0) = \frac{n!}{2\pi i}\int_{\partial D}\frac{f(\zeta)}{(\zeta-z_0)^{n+1}}d\zeta.$$

Corollary 8.4. *Cauchy's Inequality:*

$$\left|f^{(n)}(z_0)\right| \leq \frac{n!}{2\pi}\sup_{\zeta\in\partial D}\left|\frac{f(\zeta)}{(\zeta-z_0)^{n+1}}\right| = \frac{n!}{r^n}||f||_\infty \partial D.$$

Theorem 8.5. *Liouville's Theorem: If f entire and bounded then f is constant.*

Proof.

$$|f'(z_0)| \leq \frac{\|f\|_\infty}{r}$$

Since f is bounded and r can be taken arbitrarily large (as f is entire), sending $r \to \infty$ gives $|f'(z_0)| = 0$. Thus, f is constant. □

9 Feb 2

9.1 Rigidity and Analytic Continuation

Theorem 9.1. *(Rigidity Theorem): Let Ω be a region and $f : \Omega \to \mathbb{C}$ be holomorphic. If $z_1, z_2, ... \in \Omega$ distinct sequence with a limit in Ω. If $f(z_n) = 0, \forall n$ then $f = 0$ on Ω.*

Proof. Say $z_n \to \omega \in \Omega$ and $D = D_r(\omega) \subset \Omega$ be a disk centered at ω with radius r. If $f = 0$ on D then by regularity implies

$$f(z) = \sum_{n \geq 0} a_n (z - \omega)^n \text{ on } D.$$

Suppose $f \neq 0$ on D then $\exists m$ such that $a_m \neq 0$, let's take the smallest such m. Then $f(z) = a_m(z - w)^m(1 + g(z))$ where $g(z) \to 0$ as $z \to w$. For all sufficiency large $k, z_k \in D$ then

$$0 = f(z_k) = a_m(z_k - w)^m(1 + g(z_k)). \qquad (a_m \neq 0 \text{ and } z_k - w \neq 0)$$

A contradiction as the RHS is nonzero. To finish $f = 0$ on Ω, we use $\mathcal{U} = \text{int}\{z \in \Omega : f(z) = 0\}$
We have $\mathcal{U}$ is open, $w \in \mathcal{U}$ so $\mathcal{U}$ is non-empty. Additionally, $\mathcal{U}$ is closed as otherwise there would be a boundary point ω' where the function transitions from being zero to non-zero. However, applying the exact same Taylor series logic to ω' would show that f must be zero in a small disk around ω' too, meaning ω' was actually part of the open set $\mathcal{U}$ all along. Because $\mathcal{U}$ is open, closed, and non-empty, $\mathcal{U} = \Omega$. Therefore, $f(z) = 0$ everywhere on Ω. □

Corollary 9.2. *$f, g : \underset{region}{\Omega} \to \mathbb{C}$ holomorphic. If $f(z_n) = g(z_n), \forall z_n \in \Omega$ distinct sequence with a limit in Ω then $f = g$ on Ω.*

Definition 9.3. *If $\Omega_1 \subset \Omega_2$, are two regions, $f_i : \Omega_i \to \mathbb{C}$ holomorphic for $i = 1, 2$ such that $f_1 = f_2$ on Ω_1 then we say f_2 is the analytic continuation of f_1 into Ω_2.*

9.2 Morera's Theorem and Uniform Convergence

Theorem 9.4. *(Morera's Theorem): If $f : \mathcal{D} \to \mathbb{C}$ continuous such that $\forall \overline{\triangle} \subset \mathcal{D}$ and $\int_{\partial \triangle} f = 0$ then f is holomorphic.*

Proof. Repeating the proof of Cauchy's Theorem in a disk, gives that f has a primitive $F = \int_{\gamma_z} f$ in $\mathcal{D}$. Since F is holomorphic, F' exists but so does $F'', F''', ...$ so f' exists. □

Theorem 9.5. *If $f_n : \underset{open}{\Omega} \to \mathbb{C}$ holomorphic with f_n converging uniformly to f on every compact $K \subset \Omega$ then f is holomorphic. Moreover, if f_n' converges uniformly to f', using $\Omega_s = \{z \in \Omega : \overline{\mathcal{D}}_\delta(z) \subset \Omega\}$.*

Claim 9.6. *$\forall F$ holomorphic in Ω, $||F'||_{\infty, \Omega_\delta} \leq \frac{1}{\delta}||F||_{\infty, \Omega_\delta}$.*

Proof. By Cauchy's Formula:

$$|F'(z)| = \left| \frac{1}{2\pi i} \int_{\partial \mathcal{D}_\delta(z)} \frac{F(w)}{(w - z)^2} dw \right| \leq \frac{1}{2\pi} \int_{\partial \mathcal{D}_\delta(z)} \frac{1}{|w - z|^2} ||F||_{\infty, \Omega} \leq \frac{1}{2\pi} \frac{2\pi\delta}{\delta^2} ||F||_{\infty, \Omega} \leq \frac{1}{\delta} ||F||_{\infty, \Omega}.$$

□

Proof. Use Morera's Theorem. □

10 Feb 4

10.1 Parameter Integrals and Gluing

Theorem 10.1. *Let $F : \Omega \times [0,1] \to \mathbb{C}$ satisfy*

1. *$\forall \triangle \in [0,1]$, $z \mapsto F(z, \triangle)$ is holomorphic*
2. *F is continuous on $\Omega \times [0,1]$*

Then $z \mapsto \int_0^1 F(z, \triangle) d\triangle$ is holomorphic on Ω.

Proof. Consider $F_n(z) = \frac{1}{n} \sum_{k=1}^n F(z, \frac{k}{n})$ and by (1) each F_n is holomorphic on Ω.

$$\begin{aligned}
\left| F_n(z) - \int_0^1 F(z, \triangle) d\triangle \right| &= \left| F_n(z) - \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} F(z, \triangle) d\triangle \right| \\
&= \left| \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left[\underbrace{F(z, \tfrac{k}{n}) - F(z, \triangle)}_{\leq \epsilon} \right] d\triangle \right| \\
&\leq \epsilon
\end{aligned}$$

□

Definition 10.2. *$\Omega \subset \mathbb{C}$ be open, symmetric with respect to real axis, $\Omega^+ = \Omega \cap \{z \in \mathbb{C}, \Im(z) > 0\}$,$\Omega^- = \Omega \cap \{z \in \mathbb{C}, \Im(z) < 0\}$ and $I = \Omega \cap \{\Im(z) = 0\}$*

Theorem 10.3. *Let each $f^\pm : \Omega^\pm \to \mathbb{C}$ be holomorphic and extend continuous to I such that $f^+ = f^-$ on I. Then*

$$f = \begin{cases} f^+(z) & \text{if } z \in \Omega^+ \\ f^+(z) = f^-(z) & \text{if } z \in I \\ f^-(z) & \text{if } z \in \Omega^- \end{cases}.$$

is holomorphic on Ω.

Proof. Sufficient to handle $z \in I$. Fix such z take $D \subset \Omega$ centered at z. Take $\overline{\triangle} \subset D$ we want to show $\int_{\partial \triangle} f = 0$.
Case 1: If $\overline{\triangle} \subset \Omega^+$ (or Ω^-) then we're done.



Case 2: If a vertex or side of $\triangle$ is on I, $\int_{\partial \triangle_\epsilon} f = 0$ for when $\epsilon \to 0, \partial \triangle_\epsilon \to \partial \triangle$.



Case 3: If I cuts the triangle then we can split the triangle into smaller triangles and we're done.



The triangle is split into smaller triangles, each either entirely in Ω^+ or Ω^-, or with edges on I, so we can apply Cases 1 and 2. □

10.2 Schwarz Reflection Principle

Theorem 10.4. *Schwarz Reflection Principle: Let $f : \Omega^+ \to \mathbb{C}$ be holomorphic and extend continuous to I with $f^+/I \in \mathbb{R}$ then f^+ extends analytically to Ω*

Proof. Let $f^- : \Omega^- \to \mathbb{C}$ be defined by $f^-(z) = \overline{f^+(\overline{z})}$. We claim f^- is holomorphic on Ω^-, then the previous theorem applies because $f^+ = f^-$ on I.
Fix $z_0 \in \Omega^-$ then $\overline{z_0} \in \Omega^+$, we know $f^+(\overline{z}) = \sum_{n=0}^{\infty} a_n(\overline{z} - \overline{z_0})^n$ for all $\overline{z} \in D(\overline{z_0})$. Thus

$$\overline{f^+(\overline{z})} = \sum_{n=0}^{\infty} \overline{a_n}(\overline{z} - \overline{z_0})^n.$$

We know $\overline{\overline{z} - \overline{z_0}} = z - z_0$ so

$$f^-(z) = \overline{f^+(\overline{z})} = \sum \overline{a_n}(z - z_0)^n.$$

□

11 Feb 9

11.1 Meromorphic Functions and Poles

Definition 11.1. *Meromorphic functions are "determined" by zeros and singularities.*
A point singularity of f is a point $z_0 \in \mathbb{C}$ such that f is defined in a neighborhood of z_0 but not at z_0.

$$\mathcal{D}_\delta(z_0) \setminus \{z_0\}.$$

Example 11.2. *$f(z) = \frac{1}{z}$ has a singularity at $z_0 = 0$.*

Remark 11.3. *Zeros of a holomorphic f are isolated, unless $f \equiv 0$.*

Theorem 11.4. *(Local Description Near Zeros) Let $f : \underset{open}{\Omega} \to \mathbb{C}$ holomorphic and $f(z_0) = 0$, $z_0 \in \Omega, f \not\equiv 0$ then $\exists \mathcal{U} \ni z_0$, $g : \mathcal{U} \to \mathbb{C}$ holomorphic and vanishing ($\forall z \in \mathcal{U}, g(z) \neq 0$) $\exists! n > 0$ such that $f(z) = (z - z_0)^n g(z)$.*

Proof. We know

$$f(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k.$$

$f \not\equiv 0$ implies that $n =$ smallest k such that $a_k \neq 0$. We can write $f(z) = (z - z_0)^n g(z)$. □

Definition 11.5. *Let $n =$ the multiplicity/order of z_0. When $n = 1$, z_0 is called a simple zero. We say $f : \mathcal{D}_\delta(z_0) \setminus \{z_0\} \to \mathbb{C}$ has a pole at z_0 if $\frac{1}{f}$ extended by 0 at z_0 is holomorphic in $\mathcal{D}_\epsilon(z_0)$, for some $0 < \epsilon < \delta$.*

Theorem 11.6. *(Local Discontinuity near Poles) Let $f : \Omega \setminus \{z_0\} \to \mathbb{C}$ has a pole at z_0 in Ω then $\exists \mathcal{U} \ni z_0$, $\exists h : \mathcal{U} \xrightarrow{holo} \mathbb{C}$ nonvanishing then $\exists! n > 0$ such that $f(z) = (z - z_0)^{-n} h(z)$.*

Proof. Apply previous theorem to $\frac{1}{f}(z) = (z - z_0)^n g(z)$. □

11.2 Laurent Series and Residues

Theorem 11.7. *(Laurent Series Expansion) If f has a pole of order n at z_0 then locally*

$$f(z) = \underbrace{\frac{a_{-n}}{(z - z_0)^n} + \cdots + \frac{\overbrace{a_{-1}}^{\text{residue of } f \text{ at } z_0}}{z - z_0}}_{\text{principal part}} + \underbrace{G(z)}_{\text{holo part of } f}.$$

where $G(z)$ is holomorphic and nonvanishing near z_0.
Moreover, $res_{z_0}(f) = \frac{1}{(n-1)!} \lim_{z \to z_0} \frac{d^{n-1}}{dz^{n-1}}((z - z_0)^n f(z))$

Proof. By theorem 11.6,

$$f(z) = (z - z_0)^{-n} h(z) = (z - z_0)^{-n} \sum_{k=0}^{\infty} b_k (z - z_0)^k = \frac{b_0}{(z - z_0)^n} + \cdots + \frac{b_{n-1}}{(z - z_0)} + \sum_{k \geq n}^{\infty} b_k (z - z_0)^{k-n}.$$

So we have

$$f(z)(z - z_0)^n = b_0 + b_1(z - z_0) + \cdots + b_{n-1}(z - z_0)^{n-1} + \underbrace{O((z - z_0)^n)}_{\to 0 \text{ as } \frac{d^{n-1}}{dz^{n-1}}}.$$

□

11.3 The Residue Formula

Theorem 11.8. *(The Residue Formula) Suppose* $f : \Omega \setminus \underbrace{\{z_1, z_2, \cdots, z_n\}}_{poles} \to \mathbb{C}$ *is holomorphic. Then for a disk* $\overline{\mathcal{D}} \subset \Omega$ *containing* $z_1, ..., z_n$ *we have*

$$\int_{\partial \mathcal{D}} f = 2\pi i \sum_{k=1}^{n} res_{z_k}(f).$$

Proof. Let $\gamma = \partial \mathcal{D}$ be the boundary of the disk $\mathcal{D}$, oriented counterclockwise. For each pole z_k, choose a small disk $\mathcal{D}_k$ centered at z_k with radius $\varepsilon_k > 0$ small enough so that:

- $\overline{\mathcal{D}_k} \subset \mathcal{D}$ for all k
- $\overline{\mathcal{D}_k} \cap \overline{\mathcal{D}_j} = \emptyset$ for $k \neq j$
- f is holomorphic on $\overline{\mathcal{D}} \setminus \bigcup_{k=1}^{n} \mathcal{D}_k$

Let $\gamma_k = \partial \mathcal{D}_k$ be the boundary of each small disk, oriented clockwise (negative orientation). Consider the multiply connected region $\mathcal{D} \setminus \bigcup_{k=1}^{n} \overline{\mathcal{D}_k}$.



By Cauchy's theorem for multiply connected regions, the integral of f around the boundary of this region (taking into account orientations) is zero. The boundary consists of γ (counterclockwise) and each γ_k (clockwise), so:

$$\int_{\gamma} f(z)dz + \sum_{k=1}^{n} \int_{\gamma_k} f(z)dz = 0.$$

Note that γ_k has clockwise orientation, so reversing it gives:

$$\int_{\gamma} f(z)dz = \sum_{k=1}^{n} \int_{-\gamma_k} f(z)dz.$$

where $-\gamma_k$ denotes γ_k with counterclockwise orientation.
Now, for each pole z_k, consider its Laurent expansion:

$$f(z) = \sum_{j=-\infty}^{\infty} a_j^{(k)}(z - z_k)^j = \frac{a_{-1}^{(k)}}{z - z_k} + \sum_{j \neq -1} a_j^{(k)}(z - z_k)^j.$$

where $a_{-1}^{(k)} = \text{res}_{z_k}(f)$.
For $j \neq -1$, the function $(z - z_k)^j$ has an antiderivative on $\mathbb{C} \setminus \{z_k\}$ (or on all of $\mathbb{C}$ if $j \geq 0$), so by the fundamental theorem of calculus:

$$\int_{-\gamma_k} (z - z_k)^j dz = 0 \quad \text{for } j \neq -1.$$

For $j = -1$, we compute directly. Parameterize $-\gamma_k$ by $z(t) = z_k + \varepsilon_k e^{it}$ for $t \in [0, 2\pi]$:

$$\int_{-\gamma_k} \frac{1}{z - z_k} dz = \int_0^{2\pi} \frac{1}{\varepsilon_k e^{it}} \cdot i\varepsilon_k e^{it} dt = \int_0^{2\pi} i dt = 2\pi i.$$

Therefore,

$$\int_{-\gamma_k} f(z)dz = a_{-1}^{(k)} \cdot 2\pi i = 2\pi i \cdot \mathrm{res}_{z_k}(f).$$

Combining all terms:

$$\int_{\partial\mathcal{D}} f(z)dz = \sum_{k=1}^{n} \int_{-\gamma_k} f(z)dz = 2\pi i \sum_{k=1}^{n} \mathrm{res}_{z_k}(f).$$

$\square$

12 Feb 11

12.1 Integral Evaluation via Residues

Example 12.1.

$$\int_{-\infty}^{\infty} \frac{e^{ax}}{1+e^x} dx = \frac{\pi}{\sin(a\pi)}.$$

Proof. Let $f(z) = \frac{e^{az}}{1+e^z}$ then we have simple poles at $z : e^z = 1$ so $z \in \{..., -\pi i, \pi i, 3\pi i, ...\}$. We have the residual for

$$\int_{\gamma_1} f + \cdots = \int_{\gamma_R} f(z)\, dz = 2\pi i \operatorname{res}_{\pi i}(f).$$



We have

$$\int_{\gamma_1} f(z)dz \xrightarrow{R\to+\infty} \int_{-\infty}^{\infty} \frac{e^{ax}}{1+e^x} dx = I.$$

On γ_3, $z = x + 2\pi i$ with x from R to $-R$, so

$$\int_{\gamma_3} f(z)\, dz = -\int_{-R}^{R} \frac{e^{a(x+2\pi i)}}{1+e^x} dx = -e^{2\pi a i} \cdot I.$$

As $R \to \infty$, $\int_{\gamma_2} f \to 0$ and $\int_{\gamma_4} f \to 0$. Thus

$$(1 - e^{2\pi a i})I = 2\pi i \cdot \operatorname{res}_{\pi i}(f) = 2\pi i \cdot (-e^{a\pi i}) = -2\pi i e^{a\pi i},$$

$$\left|\int_{\gamma_2} f(z)dz\right| = \left|\int_0^{2\pi} \frac{e^{a(R+iy)}}{1+e^{R+iy}} i dy\right| \leq \int_0^{2\pi} \frac{e^{aR}}{e^R+1} dy \xrightarrow{R\to\infty} 0$$

Additionally $\int_{\gamma_4} f \to 0$. To compute the residue at πi, we have

$$\lim_{z\to\pi i} f(z)(z-\pi i) = \lim_{z\to i\pi} \frac{e^{az}}{\frac{e^z - e^{\pi i}}{z-\pi i}} = \frac{e^{\pi i a}}{e^{\pi i}}.$$

Letting $R \to +\infty$ in the residual formula gives

$$I + 0 + 0 - e^{2\pi a i} I = -2\pi i e^{a\pi i}.$$

So

$$I = -\frac{2\pi i e^{\pi i a}}{1 - e^{2\pi a i}} = \frac{\pi}{\sin(a\pi)}.$$

□

12.2 Removable Singularities

Definition 12.2. *Let $f : \Omega \setminus \{z_0\} \to \mathbb{C}$ be holomorphic, if we can extend f analytically to z_0 we say that f has a removable singularity at z_0.*

Theorem 12.3. *(Riemann's Theorem) Let $f : \Omega \setminus \{z_0\} \to \mathbb{C}$ be holomorphic, if f is bounded near z_0 then z_0 is a removable singularity.*

Proof. Let $\overline{D} \subset \Omega$ be a disc centered at z_0. We want to use Cauchy's formula

$$\frac{1}{2\pi i}\int_{\partial D}\frac{f(w)}{w-z}dw = g(z), \text{ for all} z \in D.$$

It suffices to show that $f = g$ on $D \setminus \{z_0\}$ because then g is the desired extension of f.
$F(w, z)$ is jointly continuous on $\partial D \times \overline{D}_{r-\epsilon}(z_0)$ and $\forall w \in \partial D$, $z \mapsto F(w, z)$ is holomorphic on D.
Fix $z \in D \setminus \{z_0\}$ then

$$\int_\gamma \frac{f(w)}{w-z}dw = 0 \implies \int_{\partial D}\frac{f(w)}{w-z}dw = \int_{C_\epsilon(z_0)}\frac{f(w)}{w-z}dw + \int_{C_\epsilon(z)}\frac{f(w)}{w-z}dw = 0 + 2\pi i f(z).$$

□

13 Feb 13

13.1 Classification of Singularities

Theorem 13.1. *(Riemann) Let $f : \Omega \setminus \{z_0\} \to \mathbb{C}$ has a removable singularity at z_0 if and only if bounded near z_0.*

Corollary 13.2. *Pole at z_0 if and only if $|f| \to +\infty$ as $z \to z_0$*

Proof. ($\Longrightarrow$) Local description near poles implies $f(z) = (z - z_0)^{-n} g(z)$ so $|f| \to +\infty$
($\Longleftarrow$) Since $\left|\frac{1}{f}\right| \to 0$, in particular $\frac{1}{f}$ is bounded near z_0, Riemann implies $\frac{1}{f}$ has a removable singularity at z_0. □

Theorem 13.3. *(Casorati-Weierstrass) If $f : D_r(z_0) \setminus \{z_0\} \to \mathbb{C}$ is holomorphic with an essential singularity at z_0 then the image of f is dense in $\mathbb{C}$.*

Proof. If not, $\exists w \in \mathbb{C}$ such that no values of f near w, $\exists \delta > 0$ s.t. $\forall z$, $|f(z) - w| > \delta$.
Consider

$$g(z) = \frac{1}{f(z) - w} : D_r(z_0) \setminus \{z_0\} \to \mathbb{C}.$$

Such a g is bounded by $\frac{1}{\delta}$. So g extends halomorphically into $D_r(z_0)$.

(Case 1) $g(z_0) = 0 \implies \frac{1}{g} \to +\infty$ so $f(z) - w$ has a pole so f has a pole

(Case 2) $g(z_0) \neq 0 \implies \frac{1}{g}$ is well defined and f has a removable singularity.

□

Theorem 13.4. *(Picard's Theorem) If $f : D_r(z_0) \setminus \{z_0\} \to \mathbb{C}$ is holomorphic with an essential singularity at z_0 then the image of f is either $\mathbb{C}$ or $\mathbb{C}$ except for at most one value.*

13.2 Meromorphic Functions and Behavior at Infinity

Definition 13.5. *Let $\Omega \subset \mathbb{C}$ be open. A function f is called meromorphic on Ω if $\exists z_1, z_2, \ldots$ without a limit point such that f is holomorphic in $\Omega \setminus \{z_1, z_2, \ldots\}$ and has poles at $z_1, z_2, \ldots$.*

Theorem 13.6. *(Behavior at $+\infty$) Suppose f is holomorphic in a neighborhood of $_\infty$ i.e. $\{|z| > R\}$ for some $R > 0$.*
Note: $F(z) = f(\frac{1}{z})$ is holomorphic in $D_{\frac{1}{R}}(0) \setminus \{0\}$.
We say f has a removable singularity at ∞ if F has a removable singularity at $z = 0$.

Example 13.7. *e^z has an essential singularity at $+\infty$.*

Remark 13.8. *We'll denote $\overline{\mathbb{C}} = \mathbb{C} \cup \{+\infty\}$*

Example 13.9. *Examples of meromorphic functions on $\mathbb{C}$*

1. *Rational functions: $\frac{P}{Q}$*
2. *The gamma function: $\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt$ (simple poles at $\ldots, -2, -1, 0$)*
3. *The Riemann zeta function: $\zeta(s) = \sum_{n=1}^\infty \frac{1}{n^s}$ (simple poles at $s = -1, -2, -3, \ldots$)*

14 Feb 16

14.1 Rational Functions via Meromorphy

Theorem 14.1. *Suppose f is meromorphic on $\mathbb{C}$ with a removable singularity or a pole at $+\infty$ then f is a rational function.*

Proof. Claim: Under these assumptions, f has finitely many poles.
If not, $z_1, z_2, ...$ are poles and this sequence cannot be bounded otherwise it got a limit point. Say $|z_k| \to +\infty$, the neighborhoods of ∞ are $\{|z| > R\}$ so f not holomorphic for any $R > 0$ which contradicts that f has a removable singularity at $+\infty$.

For a function F which has a pole at $z = 0$ we have

$$F(z) = \underbrace{\frac{a_n}{z^n} + ... + \frac{a_{-1}}{z}}_{f(z)} + \underbrace{a_0 + a_1 z + \cdots}_{g(z)}.$$

Let $z_1, \ldots, z_N$ be the poles of f. Near z_k, for each $1 \leq k \leq N$,

$$f(z) = \underbrace{f_k(z)}_{\substack{\text{principal part} \\ \text{poly in } \frac{1}{z-z_k}}} + \underbrace{g_k(z)}_{\text{holomorphic}}.$$

Near ∞,

$$f\left(\frac{1}{z}\right) = \underbrace{\tilde{f}_\infty(z)}_{\substack{\text{principal part} \\ \text{poly in } \frac{1}{z}}} + \underbrace{\tilde{g}_\infty(z)}_{\text{holomorphic}}.$$

Consider,

$$H(z) = f(z) - \underbrace{\sum_{j=1}^{N} f_j(z)}_{\text{rational}} - \underbrace{\tilde{f}_\infty(z)}_{\text{poly}}.$$

Claim: H is holomorphic and bounded.
Near z_k

$$H = \underbrace{f - f_k}_{\text{holo}} - \sum_{j \neq k} f_j - \underbrace{f_\infty}_{\text{holo}}.$$

We can show H is bounded as

$$H\left(\frac{1}{z}\right) = \underbrace{f\left(\frac{1}{z}\right) - \tilde{f}_\infty(z)}_{\substack{\tilde{g}_\infty(z) \\ \text{bounded near } z=0}} - \underbrace{\sum_{j=1}^{N} f_j\left(\frac{1}{z}\right)}_{\substack{\text{poly} \\ \text{bounded near } z=0}}.$$

So $H(\frac{1}{z})$ is bounded in say $\{|z| < \delta\}$ and $H(z)$ is bounded in $\{|z| > 1/\delta\}$, hence H is bounded in $\{|z| \leq 1/\delta\}$. *dset* □

15 Feb 18

15.1 Argument Principle and Rouche's Theorem

Theorem 15.1. *(The Argument Principle) Let $f : \Omega \to \mathbb{C}$ be meromorphic, $\overline{D} \subset \Omega$ be a disk. If If f never vanishes on ∂D nor has nay poles on ∂D then*

$$\frac{1}{2\pi i}\int_{\partial D}\frac{f'}{f} = \#\text{ zeros of } f - \#\text{ poles of } f \text{ in } \overline{D}.$$

Proof. If f has zero of order n at z_0 then $f(z) = (z - z_0)^n g(z)$ nonvanishing near z_0 then

$$\frac{f'}{f}(z) = \frac{n}{z - z_0} + \frac{g'}{g}(z).$$

If f has a pole of order n at z_0 we can write $f(z) = (z - z_0)^{-n} h(z)$ then

$$\frac{f'}{f}(z) = -\frac{n}{z - z_0} + \frac{h'}{h}(z).$$

Therefore $\frac{f'}{f}$ has simple poles at zeros and poles of f with residue $\frac{\text{order}}{-\text{order}}$. The residue formula finishes the proof. □

Theorem 15.2. *Rouche's Theorem: Let f, g be holomorphic in $\Omega \supset \overline{D}$ with Ω open. If $|f| > |g|$ on ∂D then $f, f + g$ have the same number of zeros in D.*

Proof. Let $f_t = f + t \cdot g$ for $t \in [0, 1]$ and $n_t = \#$ of zeros of f_t in D. By argument principle we have

$$n_t = \frac{1}{2\pi i}\int_{\partial D}\frac{f_t'}{f_t}.$$

f_t never vanishes on ∂D. We can conclude

$$|f_t| \geq |f| - t|g| \geq |f| - |g| > 0.$$

We can show continuity of the integrand by

$$\int_0^{2\pi} \underbrace{\frac{f_t'(z_0 + re^{i\theta})}{f_t(z_0 + re^{i\theta})} rie^{i\theta}}_{H(t,\theta)}\, d\theta.$$

Then $H(t, \theta)$ is jointly continuous in (t, θ). So n_t is continuous and discrete so n_t is constant. □

Proof. Third Proof of fundamental Theorem of Algebra.

$$P(z) = \underbrace{a_0 + a_1 z + \cdots + a_{n-1} z^{n-1}}_{g(z)} + \underbrace{a_n z^n}_{f(z)}.$$

There exist $\mathbb{R} > 0$ such that $|g(z)| < |f(z)|$ on $|z| = R$ □

15.2 Open Mapping and Maximum Modulus Theorems

Theorem 15.3. *(The Open Mapping Theorem) Let $f : \Omega \to \mathbb{C}$ holomorphic, f non-constant then f maps open sets to open sets.*

Proof. Let $w_0 = f(z_0) \in f(\Omega)$. We want to show that every w near w_0 is also in $f(\Omega)$ Fix w near w_0 then consider,

$$h(z) = f(z) - w = \underbrace{(f(z) - w_0)}_{F(z)} + \underbrace{(w_0 - w)}_{G(z)}.$$

Let $\delta > 0$ be such that $\overline{D}_\delta(z_0) \subset \Omega$ and $\forall z \in \partial D_\delta(z_0)$, $f(z) \neq w_0$. Otherwise if there exist no such δ then there exist $(z_n) \to z_0$ such that $f(z_n) = w_0$ for all n then by rigidity f is constant which contradicts the assumption that f is non-constant. Let

$$\epsilon = \inf_{\partial D_\delta(z_0)} |f(z) - w_0| > 0.$$

Now $|G| > |F|$ on $\partial D_\delta(z_0)$ as long as $|w - w_0| < \epsilon$. By Rouche's Theorem, $h = F + G$ has a zero in $\overline{D}_\delta(z_0)$. □

Theorem 15.4. *(The Maximum Modulus Theorem) Let $f : \Omega \to \mathbb{C}$ holomorphic. If f is holomorphic and non-constant then $|f|$ cannot attain a maximum in Ω.*

Proof. AFSOC, $\exists z_0 \in \Omega$ such that $|\underbrace{f(z_0)}_{w_0}| = \sup_\Omega |f|$ then this contradicts the open mapping theorem because there exist w near w_0 with $|w| > |w_0|$ and the value w must be obtained. □

Corollary 15.5. *If Ω additionally bounded ($\overline{\Omega}$ compact) then $\sup_\Omega |f| = \max_{\partial\Omega} |f|$*

16 Feb 20

16.1 Homotopy and Path Independence

Definition 16.1. *Let $\Omega \subset \mathbb{C}$ be a region, two curves in Ω say with parametrization $\gamma_0, \gamma_1 : [a,b] \to \mathbb{C}$ and common endpoints*

$$\alpha = \gamma_0(a) = \gamma_1(a) \text{ and } \beta = \gamma_0(b) = \gamma_1(b).$$

are called homotopic if $\forall s \in [0,1]$, $\exists \gamma_s : [a,b] \to \Omega$ such that

1. *$\gamma_{s=0} = \gamma_0$ and $\gamma_{s=1} = \gamma_1$,*
2. *$\gamma_s(a) = \alpha$ and $\gamma_s(b) = \beta$ for all $s \in [0,1]$,*
3. *the map $(s,t) \mapsto \gamma_s(t)$ is continuous on $[0,1] \times [a,b]$.*

The map $H(s,t) = \gamma_s(t)$ is called a homotopy between γ_0 and γ_1.





Theorem 16.2. *If $f : \Omega \to \mathbb{C}$ be holomorphic, γ_0, γ_1 are two homotopic curves in Ω then $\int_{\gamma_0} f = \int_{\gamma_1} f$*

Proof. Let $H(s,t) : [0,1] \times [a,b] \to \Omega$ be a homotopy between γ_0, γ_1. Since $K = H([0,1] \times [a,b])$ is compact in open Ω, $\exists \epsilon > 0$ such that $\forall z \in K$, $D_{3\epsilon}(z) \subset \Omega$. Fix such $\epsilon > 0$ and take $\delta > 0$ such that

$$|H(s_1,t_1) - H(s_2,t_2)| < \epsilon \text{ provided that } |(s_1,t_1) - (s_2,t_2)| < 2\delta. \qquad \text{(unif cont of } H\text{)}$$

Given such δ, it suffices to show

$$\int_{\gamma_s} f = \int_{\gamma_{s'}} f \text{ for all } 0 \le s < s' \le 1 \text{ and } |s - s'| < \delta.$$

Take $a = t_0 < t_1 < \cdots < t_n < t_{n+1} = b$ such that $|t_{i+1} - t_i| < \delta$ for all $i = 0, 1, \cdots, n$.
Let's denote $z_i = \gamma_s(t_i)$ and $w_i = \gamma_{s'}(t_i)$. Let $D_i = D_{z_i}(2\epsilon)$.

Claim 16.3.

1. *$z_i, z_{i+1} \in D_i$ holds because $|z_{i+1} - z_i| < \epsilon$ as $z_{i+1} = \gamma_s(t_{i+1}) = H(s, t_{i+1})$ and $z_i = \gamma_s(t_i) = H(s, t_i)$.*
2. *$w_i, w_{i+1} \in D_i$ holds because $|w_i - z_i| < \epsilon$ so*
$$|w_{i+1} - z_i| \le |w_{i+1} - z_{i+1}| + |z_{i+1} - z_i| < \epsilon + \epsilon = 2\epsilon.$$
3. *$D_0 \cup D_1 \cup \cdots \cup D_n$ covers $\gamma_s, \gamma_{s'}$ because for $z = \gamma_s(t)$, $t_i \le t \le t_{i+1}$, $|z - z_i| = |\gamma_s(t) - \gamma_s(t_i)| \le \epsilon$ so $z \in D_i$. For $w = \gamma_{s'}(t)$, $|w - z_i| = |\gamma_{s'}(t) - \gamma_s(t_i)| < \epsilon$.*

Finally, let F_i be a primitive of f on D_i (exists by Cauchy applied to D_i). On $D_i \cap D_{i+1}$, F_i, F_{i+1} are two primitives of the same function f so

$$F_{i+1}(w) - F_{i+1}(z) = c_i.$$

where c_i is a constant. Then

$$\gamma_{\gamma_s} f = \int_a^b f(\gamma_s(t))\gamma_s'(t)dt = \sum_{i=0}^{n} \int_{t_i}^{t_{i+1}} f(\gamma_s(t))\gamma_s'(t)dt = \sum_{i=0}^{n} F_i(z_{i+1}) - F_i(z_i).$$

So

$$\begin{aligned}
\int_{\gamma_s} f - \int_{\gamma_{s'}} f &= \sum_{i=0}^{n} [F_i(z_{i+1}) - F_i(z_i)] - \sum_{i=0}^{n} [F_{i+1}(w_{i+1}) - F_{i+1}(w_i)] \\
&= \sum_{i=0}^{n} [F_i(z_{i+1}) - F_{i+1}(w_{i+1})] - \sum_{i=0}^{n} [F_i(z_i) - F_{i+1}(w_i)] \\
&= F_0(w_0) - F_0(z_0) + \sum_{i=0}^{n-1} [F_i(z_{i+1}) - F_{i+1}(w_{i+1})] - \sum_{i=0}^{n-1} [F_i(z_i) - F_{i+1}(w_i)] \\
&= \sum_{i=0}^{n-1} [F_i(z_{i+1}) - F_{i+1}(w_{i+1})] - \sum_{i=0}^{n-1} [F_i(z_i) - F_{i+1}(w_i)] \\
&= \sum_{i=0}^{n-1} [F_i(z_{i+1}) - F_{i+1}(z_i) + F_{i+1}(z_i) - F_i(w_{i+1})] \\
&= \sum_{i=0}^{n-1} -c_i + c_i = 0
\end{aligned}$$

□

16.2 Simply Connected Regions

Definition 16.4. *A region $\Omega \subset \mathbb{C}$ is simply connected if every two curves in Ω with common endpoints are homotopic.*

Example 16.5. *Examples of simply connected regions:*

1. *Convex Sets*
2. $\Omega = \mathbb{C} \setminus (-\infty, 0]$

Theorem 16.6. *For all f holomorphic in a simply connected region then for closed γ*

$$\int_\gamma f = 0.$$

Theorem 16.7. *f holomorphic has a primitive in a simply connected region Ω.*

17 Feb 23

17.1 Complex Logarithm

Theorem 17.1. *(Construction of Complex Logarithm) Let Ω be a simply connected region such that $0 \notin \Omega$ but $1 \in \Omega$. Then there $\exists!$ holomorphic F on Ω with $F(z) = \log \Omega$ such that*

1. $e^{F(z)} = z$ *on* Ω
2. $F(r) = \log r \ \forall r \in \Omega \cap (1-\delta, 1+\delta)$

Proof. Let $F(z) = \int_{\gamma_z} \frac{dw}{w}$ where γ_z is a curve $1 \curvearrowright z$ in Ω. This is well defined as it doesn't depend on choice of γ_z because Ω is simply connected.
Then we know $F'(z) = \frac{1}{z}$.
Thus

1. $(ze^{-F(z)})' = e^{-F(z)} - ze^{-F(z)}\frac{1}{z} = 0$ so $ze^{-F(z)} = 1$ at $z = 1$.
2. $F(r) = \int_1^r \frac{dt}{t} = \log t \,|_{t=1}^{t=r} = \log r$

□

Example 17.2. $\Omega_0 = \mathbb{C} \setminus (-\infty, 0]$

Claim 17.3. *For $z = re^{i\theta}$ for $|\theta| < \pi$, $\log(z) = \log r + i\theta$, we take the path $1 \to r \to z$. So*

$$\log z = \int_{\gamma_z} \frac{dw}{w} = \int_1^r \frac{dt}{t} + \int_0^{2\pi} \frac{rie^{it}}{re^{it}} dt = \log r + i\theta.$$

Remark 17.4. $\log(z_1 z_2) \neq \log z_1 + \log z_2$ *is in general false.*

Theorem 17.5. *For the principle branch*

$$\log(1+z) = z - \frac{z^2}{2} + \frac{z^3}{3} - \cdots. \qquad \text{(For } |z| < 1\text{)}$$

Proof. $\frac{d}{dz}(\text{LHS} - \text{RHS}) = \frac{1}{1+z} - \frac{1}{1+z} = 0$ □

17.2 General Powers and Branches

Definition 17.6. *Given $\alpha \in \mathbb{C}$ we define*

$$z^\alpha := e^{\alpha \log_\Omega z}.$$

Example 17.7. *We have*

1. $1^\alpha = 1$
2. $\left(z^{\frac{1}{n}}\right)^n = z^{\frac{1}{n}} \cdot z^{\frac{1}{n}} \cdots z^{\frac{1}{n}} = \left(e^{\frac{1}{n}\log z}\right)^n = e^{\log z} = z$

Theorem 17.8. *Given f holomorphic, nowhere vanishing there exist g such that $e^g = f$*

Proof. Repeat the previous $g' = \frac{f'}{f}$ with $g(z) = \int_{\gamma_z} \frac{f'(w)}{f(w)} dw$

□

18 Feb 25

Definition 18.1. *For $f \in L_1$ $(\mathbb{R} \to \mathbb{C})$ i.e. $||f||_{L_1} = \int_{-\infty}^{+\infty} |f| < \infty$ we define its fourier transform*

$$\hat{f}(\zeta) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i t \zeta} dt. \qquad (\zeta \in \mathbb{R})$$

Example 18.2. *Some examples of fourier transforms:*

1. $\widehat{e^{-\pi x^2}}(\zeta) = e^{-\pi \zeta^2}$
2. $\widehat{\frac{1}{\cosh(\pi x)}}(\zeta) = \frac{1}{\cosh(\pi \zeta)}$
3. $\widehat{1_{[-1,1]}}(\zeta) = \int_{-1}^{1} e^{-2\pi i \zeta t} dt$

Lemma 18.3. *Basic Properties of Fourier Transforms:*

1. *$\hat{f}$ is uniformly continuous*
2. *$||\hat{f}||_\infty \leq ||f||_{L_1}$*
3. *$\hat{f}(\zeta) \to 0$ as $|\zeta| \to \infty$*

Theorem 18.4. *(Paley-Wiener Theorem) Regularity of f $\iff$ decay of f^**

Definition 18.5. *Given $a > 0$, we let $S_a = \{z \in \mathbb{C} : |Im(z)| < a\}$ and*

$$\mathcal{F}_a = \left\{ f : S_a \to \mathbb{C},\ \text{holo}\ , \exists C_f, \forall x + iy \in S_a, |f(x = iy)| \leq \frac{C_f}{1 + x^2} \right\}.$$

Additionally, $\mathcal{F} = \bigcup_{a>0} \mathcal{F}_a$

Theorem 18.6. *If $f \in \mathcal{F}_a$, then $|\hat{f}(\zeta)| \leq C e^{-2\pi b |\zeta|}$, for $\zeta \in \mathbb{R}$ and $\forall b < a$.*

Proof. Fix $\zeta > 0$ and $b < a$, we have

$$0 = \int_{\gamma_R} f(z) e^{-2\pi i \zeta z} dz = \int_{\gamma_1} + \int_{\gamma_2} + \int_{\gamma_3} + \int_{\gamma_4}.$$



Verify that $\int_{\gamma_1} = \hat{f}(\zeta)$ and

$$\left| \int_{\gamma_2} \right| = \left| \int_0^b f(R - it) e^{-2\pi i \zeta (R - it)} i \, dt \right| \leq \int_0^b |f(R - it)| e^{2\pi \zeta b} dt \leq \frac{C_f}{1 + R^2} e^{2\pi \zeta b}$$

$\square$