# Algebra I: Class Notes

Wilson Pan

March 27, 2026

## Contents

# 1 Lecture 1: Group Actions & The Orbit-Stabilizer Theorem (Jan 12)

## 1.1 Permutations and Automorphisms

**Definition 1.1** (Symmetric Group)**.** *Let $X$ be a set. The set of all permutations (bijections) of $X$ is denoted by $\mathrm{Sym}_X$ (or sometimes $\Sigma_X$). Under function composition, $\mathrm{Sym}_X$ forms a group.*

**Definition 1.2** (Automorphism Group)**.** *Let $(G, \cdot)$ be a group. An automorphism of $G$ is a bijection $\phi : G \to G$ that is also a homomorphism, i.e.,*

$$\phi(g_1 g_2) = \phi(g_1)\phi(g_2) \quad \forall g_1, g_2 \in G.$$

*The set of all such automorphisms is denoted by $\mathrm{Aut}(G)$. It forms a group under composition.*

## 1.2 Group Actions

**Definition 1.3** (Group Action)**.** *Let $G$ be a group and $X$ be a set. An **action** of $G$ on $X$ is a homomorphism $\phi : G \to \mathrm{Sym}_X$.*
*We typically write the action as $g \cdot x := \phi(g)(x)$. This notation satisfies two axioms (equivalent to the homomorphism property):*

1. ***Identity:*** *$1 \cdot x = x$ for all $x \in X$.*
2. ***Compatibility:*** *$g_1 \cdot (g_2 \cdot x) = (g_1 g_2) \cdot x$ for all $g_1, g_2 \in G, x \in X$.*

**Theorem 1.4** (Equivalence Relation on $X$)**.** *Let $G$ act on $X$. Define a relation $\sim$ on $X$ by:*

$$x_1 \sim x_2 \iff \exists g \in G \text{ such that } g \cdot x_1 = x_2.$$

*Then $\sim$ is an equivalence relation.*

*Proof.* We check the properties:

1. **Reflexive:** $1 \cdot x = x \implies x \sim x$.
2. **Symmetric:** If $g \cdot x_1 = x_2$, acting by $g^{-1}$ gives $g^{-1} \cdot (g \cdot x_1) = g^{-1} \cdot x_2 \implies (g^{-1}g) \cdot x_1 = g^{-1} \cdot x_2 \implies x_1 = g^{-1} \cdot x_2$. Thus $x_2 \sim x_1$.
3. **Transitive:** If $g \cdot x_1 = x_2$ and $h \cdot x_2 = x_3$, then $h \cdot (g \cdot x_1) = x_3 \implies (hg) \cdot x_1 = x_3$. Thus $x_1 \sim x_3$.

□

## 1.3 Orbits and Stabilizers

The equivalence classes under the relation $\sim$ partition the set $X$. These classes are called **orbits**.

**Definition 1.5** (Orbit)**.** *For $x \in X$, the orbit of $x$ is the set of all places $x$ can be moved to by $G$:*

$$\mathcal{O}_x = Orb(x) = \{g \cdot x : g \in G\}.$$

**Definition 1.6** (Stabilizer)**.** *For $x \in X$, the stabilizer of $x$ is the set of elements in $G$ that fix $x$:*

$$G_x = \mathrm{Stab}(x) = \{g \in G : g \cdot x = x\}.$$

*Note that $G_x$ is a subgroup of $G$ (denoted $G_x \leq G$).*

**Theorem 1.7** (**Orbit-Stabilizer Theorem**)**.** *Let $G$ act on $X$. For any $x \in X$, there is a bijection between the orbit $\mathcal{O}_x$ and the set of left cosets $G/G_x$. Consequently:*

$$|\mathcal{O}_x| = [G : G_x].$$

*If $G$ is finite, $|G| = |\mathcal{O}_x| \cdot |G_x|$.*

*Proof.* Define a map $\psi : G/G_x \to \mathcal{O}_x$ by $\psi(gG_x) = g \cdot x$.

1. **Well-defined:** Suppose $g_1G_x = g_2G_x$. Then $g_1^{-1}g_2 \in G_x$, so $(g_1^{-1}g_2) \cdot x = x$, implying $g_2 \cdot x = g_1 \cdot x$.
2. **Injectivity:** If $\psi(g_1G_x) = \psi(g_2G_x)$, then $g_1 \cdot x = g_2 \cdot x$. Multiplying by $g_1^{-1}$, we get $x = g_1^{-1}g_2 \cdot x$, so $g_1^{-1}g_2 \in G_x$, implying $g_1G_x = g_2G_x$.
3. **Surjectivity:** By definition, any $y \in \mathcal{O}_x$ is of the form $g \cdot x$ for some $g$, which is exactly $\psi(gG_x)$.

Thus, $\psi$ is a bijection. $\square$

# 2 Lecture 2: Cauchy's Theorem & Conjugation (Jan 14)

## 2.1 Cauchy's Theorem

**Lemma 2.1.** *If $X$ is a finite set and $G$ is a $p$-group (a group of order $p^k$) acting on $X$, then:*

$$|X| \equiv |X^G| \pmod{p},$$

*where $X^G = \{x \in X : g \cdot x = x, \forall g \in G\}$ is the set of fixed points.*

**Theorem 2.2** (Cauchy's Theorem)**.** *If $G$ is a finite group and $p$ is a prime dividing $|G|$, then $G$ has an element of order $p$.*

*Proof (McKay's Proof).* Let $X$ be the set of $p$-tuples of elements of $G$ whose product is the identity:

$$X = \{(g_1, \ldots, g_p) \in G^p : g_1 g_2 \cdots g_p = 1\}.$$

Notice that $g_p$ is uniquely determined by the first $p-1$ elements ($g_p = (g_1 \cdots g_{p-1})^{-1}$), so $|X| = |G|^{p-1}$. Since $p \mid |G|$, we have $p \mid |X|$.
Let $\mathbb{Z}_p$ (cyclic group of order $p$) act on $X$ by cyclic shift:

$$\sigma \cdot (g_1, g_2, \ldots, g_p) = (g_2, g_3, \ldots, g_p, g_1).$$

This is a valid action because if $g_1 \cdots g_p = 1$, then $g_1(g_2 \cdots g_p) = 1 \implies g_2 \cdots g_p = g_1^{-1}$, so $(g_2 \cdots g_p)g_1 = g_1^{-1} g_1 = 1$.
By the Lemma, $|X| \equiv |X^{\mathbb{Z}_p}| \pmod{p}$. Fixed points are tuples $(a, a, \ldots, a)$ such that $a^p = 1$. Since $(1, \ldots, 1) \in X^{\mathbb{Z}_p}$, the set of fixed points is non-empty. Since $p$ divides $|X|$ and $p$ divides the congruence $|X| - |X^{\mathbb{Z}_p}|$, $p$ must divide $|X^{\mathbb{Z}_p}|$. Therefore, there are at least $p$ fixed points. There must exist some $a \neq 1$ such that $a^p = 1$. □

## 2.2 Conjugation

Conjugation is a specific action of $G$ on itself.

**Definition 2.3.** *For $g, h \in G$, the **conjugate** of $g$ by $h$ is $g^h := hgh^{-1}$. The map $\phi_h : G \to G$ defined by $g \mapsto hgh^{-1}$ is an automorphism (an inner automorphism).*

**Definition 2.4** (Classes and Centralizers)**.** *Let $G$ act on itself by conjugation ($h \cdot g = hgh^{-1}$).*

- *The orbit of $g$ is the **Conjugacy Class** of $g$: $Cl(g) = \{hgh^{-1} : h \in G\}$.*
- *The stabilizer of $g$ is the **Centralizer** of $g$: $C_G(g) = \{h \in G : hg = gh\}$.*

*By Orbit-Stabilizer: $|Cl(g)| = [G : C_G(g)]$.*

**Definition 2.5** (Center of $G$)**.** *The* ***Center****, $Z(G)$, is the set of elements that commute with everything:*

$$Z(G) = \{z \in G : zg = gz, \forall g \in G\} = \bigcap_{g \in G} C_G(g).$$

*$Z(G)$ is the kernel of the conjugation homomorphism $G \to \mathrm{Aut}(G)$. Thus $Z(G) \trianglelefteq G$.*

**Definition 2.6** (Normalizer)**.** *Let $H \leq G$. The* ***Normalizer*** *of $H$ in $G$ is:*

$$N_G(H) = \{g \in G : gHg^{-1} = H\}.$$

*$N_G(H)$ is the largest subgroup of $G$ in which $H$ is normal.*

# 3 Lecture 3: Sylow Preliminaries & Isomorphism Theorems (Jan 16)

## 3.1 Product of Subgroups

**Definition 3.1.** *Let $K, N \leq G$. Define $KN = \{kn : k \in K, n \in N\}$.*

- *In general, $KN$ is not a subgroup.*
- *If $N \trianglelefteq G$ (or just $N \subseteq N_G(K)$), then $KN$ is a subgroup.*
- *Size formula: $|KN| = \frac{|K||N|}{|K \cap N|}$.*

**Theorem 3.2** (Second Isomorphism Theorem)**.** *Let $K \leq G$ and $N \trianglelefteq G$. Then $K \cap N \trianglelefteq K$, and*

$$\frac{KN}{N} \cong \frac{K}{K \cap N}.$$

*Proof.* Consider the natural projection $\pi : G \to G/N$. Restrict it to $K$, i.e., $\phi = \pi \mathrm{res}_K : K \to G/N$. The image of $\phi$ is $\{kN : k \in K\} = KN/N$. The kernel of $\phi$ is $\{k \in K : kN = N\} = \{k \in K : k \in N\} = K \cap N$. By the First Isomorphism Theorem, $K/\ker(\phi) \cong \mathrm{Im}(\phi)$. □

## 3.2 Sylow Definitions

**Definition 3.3** ($p$-group)**.** *Let $p$ be a prime. A group $G$ is a $p$-group if every element has order a power of $p$. For finite groups, this is equivalent to $|G| = p^k$.*

**Definition 3.4** (Sylow $p$-subgroup)**.** *Let $|G| = p^n m$ where $p \nmid m$. A subgroup $P \leq G$ is called a **Sylow $p$-subgroup** if $|P| = p^n$.*

$$Syl_p(G) = \{P \leq G : P \text{ is a Sylow } p\text{-subgroup}\}.$$

# 4 Lecture 4: The Sylow Theorems (Jan 21)

**Theorem 4.1** (**The Sylow Theorems**). *Let $G$ be a finite group of order $p^n m$ where $p \nmid m$.*

1. ***Existence:*** *$Syl_p(G) \neq \emptyset$. (There exists a subgroup of order $p^n$).*
2. ***Conjugacy:*** *Any two Sylow $p$-subgroups are conjugate in $G$. That is, if $P, Q \in Syl_p(G)$, $\exists g \in G$ such that $gPg^{-1} = Q$.*
3. ***Number:*** *Let $n_p = |Syl_p(G)|$. Then:*
   - $n_p \equiv 1 \pmod{p}$.
   - $n_p \mid m$ *(equivalently $n_p \mid |G|$).*

*Proof.* Let $p$ be a prime such that $p \,\Big|\, |G|$. Define the set of all $p$-subgroups:

$$\Sigma = \{H : H \leq G, |H| = p^n \text{ for some } n > 0\}.$$

Define $\Omega$ as the set of maximal elements in $\Sigma$ under inclusion:

$$\Omega = \{H : H \in \Sigma, \text{ there is no } K \in \Sigma \text{ such that } H \subsetneq K\}.$$

We let $G$ act on the set of subgroups $\{H : H \leq G\}$ by conjugation. Since conjugation is an isomorphism ($H^g \simeq H$), it preserves the order of subgroups. Thus:

$$H \in \Sigma \iff H^g \in \Sigma \quad \text{and} \quad H \in \Omega \iff H^g \in \Omega.$$

Therefore, $G$ acts on $\Omega$ by $g \cdot H = H^g$.

**Claim 1: Let $H \in \Omega$. Consider the action of $H$ on $\Omega$ by conjugation. Then $H$ is the *unique* fixed point of this action.**

*Proof of Claim:*

- **Existence:** It is trivial to see that $H$ is a fixed point, as $H^h = hHh^{-1} = H$ for any $h \in H$.
- **Uniqueness:** Let $K \in \Omega$ be a fixed point of the action of $H$.

  $$K \text{ is fixed by } H \iff K^h = K \quad \forall h \in H \iff H \leq N_G(K).$$

  Since $K \trianglelefteq N_G(K)$, and $H \leq N_G(K)$, the product $HK$ is a subgroup of $N_G(K)$, and thus $HK \leq G$. Using our counting lemma:

  $$|HK| = \frac{|H||K|}{|H \cap K|}.$$

  Since $H, K \in \Omega$, their orders are powers of $p$. Thus $|HK|$ is a power of $p$, meaning $HK \in \Sigma$.

  We have $H \leq HK$. Since $H$ is maximal in $\Sigma$ (by definition of $\Omega$) and $HK \in \Sigma$, it must be that $H = HK$. Similarly, $K \leq HK$ implies $K = HK$. Therefore, $H = K$.

**Claim 2: Size of $\Omega$ Modulo $p$**

**Statement:** $|\Omega| \equiv 1 \pmod{p}$.

*Proof of Claim:* Let $H \in \Omega$. We decompose $\Omega$ into disjoint orbits under the action of $H$.

$$\Omega = \{H\} \cup \bigcup_{K \neq H} \mathrm{Orb}_H(K).$$

By Claim 1, $H$ is the only fixed point (an orbit of size 1). For any $K \in \Omega$ with $K \neq H$, the stabilizer of $K$ in $H$ is a proper subgroup, so by the Orbit-Stabilizer theorem for $p$-groups, $|\mathrm{Orb}_H(K)|$ is divisible by $p$.

$$|\Omega| = 1 + \sum(\text{multiples of } p) \implies |\Omega| \equiv 1 \pmod{p}.$$

**Claim 3: Transitivity (Conjugacy of Elements in $\Omega$)**

**Statement:** Any two elements of $\Omega$ are conjugate.
*Proof of Claim:* Suppose for contradiction that $H, K \in \Omega$ are not conjugates. Let $\theta_1 = \mathrm{Orb}_G(H)$ and $\theta_2 = \mathrm{Orb}_G(K)$. Since they are distinct orbits, $\theta_1 \cap \theta_2 = \emptyset$.

Consider the action of $H$ on these sets:

- On $\theta_1$: $H \in \theta_1$. By Claim 1, $H$ is the unique fixed point of $H$ in $\Omega$. Since $\theta_1 \subseteq \Omega$, $H$ is the unique fixed point in $\theta_1$. Thus, $|\theta_1| \equiv 1 \pmod{p}$.
- On $\theta_2$: Since $H \in \theta_1$ and $\theta_1 \cap \theta_2 = \emptyset$, we have $H \notin \theta_2$. Therefore, $\theta_2$ contains *no* fixed points under the action of $H$ (because the only fixed point in all of $\Omega$ is $H$). Thus, every orbit of $H$ inside $\theta_2$ has size divisible by $p$. This implies $|\theta_2| \equiv 0 \pmod{p}$.

By symmetry, we can swap the roles of $H$ and $K$. Running the same argument with $K$ acting on the sets implies $|\theta_2| \equiv 1 \pmod{p}$ and $|\theta_1| \equiv 0 \pmod{p}$.

This results in a contradiction (e.g., $|\theta_1| \equiv 1$ and $|\theta_1| \equiv 0$). Thus, $\theta_1 = \theta_2$, so $H$ and $K$ are conjugate.

*Corollary:* Since there is only one orbit, $\Omega = \theta_1$. Thus $|\Omega| = |\theta_1| \equiv 1 \pmod{p}$.

**Claim 4: Divisibility**

**Statement:** $|\Omega| \,\Big|\, |G|$.
*Proof of Claim:* Let $H \in \Omega$. Since $\Omega$ is exactly the orbit of $H$ under conjugation (from Claim 3), the size of the orbit is the index of the stabilizer:

$$|\Omega| = [G : \mathrm{Stab}_G(H)] = [G : N_G(H)] = \frac{|G|}{|N_G(H)|}.$$

Thus $|\Omega|$ divides $|G|$.

**Claim 5: Identification with Sylow Subgroups**

**Statement:** $\Omega = Syl_p(G)$.
*Proof of Claim:* We know $Syl_p(G) \subseteq \Omega$ because Sylow subgroups are maximal $p$-subgroups by definition. We must show the reverse: every $H \in \Omega$ is a Sylow $p$-subgroup. Let $H \in \Omega$. Assume for contradiction that $H \notin Syl_p(G)$. Let $|G| = p^s m$ where $p \nmid m$. Since $H$ is a $p$-group but not Sylow, $|H| = p^t$ with $t < s$.
Consider the normalizer $N_G(H)$. From Claim 2, we know $|\Omega| = [G : N_G(H)] \equiv 1 \pmod{p}$. Therefore, $p$ does *not* divide $[G : N_G(H)]$. Since $|G| = [G : N_G(H)] \cdot |N_G(H)|$, all factors of $p$ in $|G|$ must reside in $|N_G(H)|$. Thus, $|N_G(H)|$ is divisible by $p^s$.
Consequently, the index $[N_G(H) : H] = \frac{|N_G(H)|}{|H|} = \frac{(\text{multiple of } p^s)}{p^t}$ is divisible by $p$ (since $s > t$).
By Cauchy's Theorem applied to the quotient group $N_G(H)/H$, there exists a subgroup of order $p$, say $\bar{K} \leq N_G(H)/H$. By the Correspondence Theorem, there exists a subgroup $K$ such that $H \leq K \leq N_G(H)$ with $|K| = p \cdot |H| = p^{t+1}$.
Since $K$ is a $p$-group, $K \in \Sigma$. However, $H \subsetneq K$, which contradicts the definition of $H$ as a maximal element in $\Omega$.
Therefore, $H$ must already be of order $p^s$. Thus $\Omega = Syl_p(G)$. $\square$

# 5 Lecture 5: Normal Series & Solvability (Jan 23)

## 5.1 Normal and Subnormal Series

**Definition 5.1.** *A **subnormal series** of a group $G$ is a chain of subgroups:*

$$\{1\} = H_0 \leq H_1 \leq \cdots \leq H_k = G$$

*such that $H_i \trianglelefteq H_{i+1}$ for all $i$. The quotient groups $H_{i+1}/H_i$ are called the **factors** of the series.*

**Definition 5.2** (Composition Series)**.** *A subnormal series is a **composition series** if all factors $H_{i+1}/H_i$ are **simple** groups (non-trivial groups with no normal subgroups other than $\{1\}$ and themselves).*

**Theorem 5.3** (Jordan-Hölder)**.** *Any two composition series of a finite group $G$ are equivalent. That is, they have the same length, and their factors are isomorphic (up to reordering).*

## 5.2 Solvable Groups

**Definition 5.4.** *A group $G$ is **solvable** if it has a subnormal series where all factors $H_{i+1}/H_i$ are **abelian**.*

**Definition 5.5** (Commutator)**.** *The commutator of $g, h$ is $[g, h] = ghg^{-1}h^{-1}$. The **Derived Subgroup** (or Commutator Subgroup) $G'$ or $[G, G]$ is the subgroup generated by all commutators.*

**Proposition 5.6.** *$G/N$ is abelian if and only if $[G, G] \leq N$. Thus, $[G, G]$ is the smallest normal subgroup such that the quotient is abelian.*

**Remark 5.7** (Derived Series)**.** *Define $G^{(0)} = G$, $G^{(1)} = [G, G]$, and $G^{(i+1)} = [G^{(i)}, G^{(i)}]$. $G$ is solvable if and only if the derived series terminates at $\{1\}$ (i.e., $G^{(n)} = \{1\}$ for some $n$).*

# 6 Lecture 6: Solvable and Derived Series (Jan 26)

**Lemma 6.1.** *Let $\alpha \in Aut(G)$ then $\alpha([g,h]) = [\alpha(g), \alpha(h)]$.*

**Lemma 6.2.** *For $N \trianglelefteq G$ then $G/N$ is abelian iff $[G,G] \leq N$.*

**Theorem 6.3.** *If $N \trianglelefteq G$, $g, h \in G$ and $[gN, hN] = [g,h]N$. Additionally, $gN$ and $hN$ commutes iff $[g,h]N = N$ iff $[g,h] \subseteq N$.*

**Definition 6.4.** *$G$ is solvable iff there is a subnormal series $\langle H_i : 0 \leq i \leq t \rangle$ such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$.*

**Theorem 6.5.** *The following are true*
1. *If $G$ solvable and $K \leq G$ then $K$ is solvable*
2. *If $G$ is solvable and $N \trianglelefteq G$ then $G/N$ is solvable.*
3. *For $N \trianglelefteq G$, $G$ solvable iff both $N$ and $G/N$ are solvable*

*Proof.* We prove the previous theorems

1. Let $\langle H_i : 0 \leq i \leq t \rangle$ be a series in $G$ such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$. We can let $H_i' = K \cap H_i$ for $0 \leq i \leq t$ and we need to verify $\langle H_i' : 0 \leq i \leq t \rangle$ is a solvable series in $K$.

2. Let $\langle H_i : 0 \leq i \leq t \rangle$ be a series in $G$ such that $[H_{i+1}, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$. Let $H_i' = \phi_N[H_i] = H_iN/N$. Verify it is solvable.

3. The forward direction is trivial by (1) and (2). For the reverse direction, $N$ solvable and $G/N$ solvable $\langle N_i : 0 \leq i \leq s \rangle$ subnormal in $N$ and $N_{i+1}/N_i$ abelian and similar for $\langle H_jN/N : 0 \leq j \leq t \rangle$ subnormal in $G/N$ and $(H_{j+1}N/N)_{j+1}/(H_jN/N)$ abelian. We know $H_{j+1} \trianglelefteq H_j$ and $[H_{j+1}, H_{j+1}] \leq H_j$ for all $0 \leq j \leq t-1$. So $N_0 = 1 \trianglelefteq N_1 \cdots \trianglelefteq N_s = N = H_0 \trianglelefteq H_1 \cdots \trianglelefteq H_t = G$.

□

**Definition 6.6.** *Given $G$, the derived series of $G$ is given by $G_0' = G$ and $G_{i+1}' = [G_i', G_i']$ for $i \geq 0$.*

**Theorem 6.7.** *$G$ is solvable iff there is $n$ such that $G_n' = \{1\}$.*

*Proof.* For the backwards direction, by construction $G_i'/G_{i+1}'$ is abelian. Let $H_j = G_{n-j}$. For the forward direction, let $\langle H_j : 0 \leq j \leq n \rangle$ be a subnormal series in $G$, $H_{j+1}/H_j$ abelian. Show by induction that $G_i' \leq H_{n-i}$ for all $i$. □

**Theorem 6.8.** *Suppose $G$ abelian and simple, let $a \in G$ and $a \neq 1$ then $G = \langle a \rangle$ and $|a|$ is prime.*

**Theorem 6.9.** *Let $G$ be simple and let $\langle H_i : 0 \leq i \leq t \rangle$ be subnormal series in $G$. Then there is $n < t$ such that $H_i = 1$ for $0 \leq i \leq n$ and $H_i = G$ for $n < i \leq t$.*

# 7 Lecture 7: Nilpotent Groups and Free Groups (Jan 28)

## 7.1 Nilpotent Groups

**Definition 7.1.** *For $H, K \leq G$, $[H, K] = \langle [h, k] : h \in H, k \in K \rangle$*

**Remark 7.2.** *$G_i'$ is characteristic in $G$ for all $i$, in particular $G_i' \trianglelefteq G$.*

**Lemma 7.3.** *If $N \trianglelefteq G, N \leq H \leq G$ then*

$$\frac{H}{N} \leq Z\left(\frac{G}{N}\right) \iff [G, H] \leq N.$$

**Definition 7.4.** *A normal series in $G$, $\langle H_i : 0 \leq i \leq t \rangle$ is a central series if $\frac{H_{i+1}}{H_i} \leq Z\left(\frac{G}{H_i}\right)$ for all $0 \leq i \leq t-1$. Equivalently, $[G, H_{i+1}] \leq H_i$ for all $0 \leq i \leq t-1$.*

**Definition 7.5.** *$G$ is nilpotent iff $G$ has a central series.*

**Remark 7.6.** *Nilpotent groups are solvable.*

**Definition 7.7.** *Let $G$ be a group*

1. *The descending central series is the sequence of subgroup given by $\gamma_1(G) = G$ and $\gamma_{i+1}(G) = [G, \gamma_i(G)]$ for $i \geq 1$.*
2. *The ascending central series is the sequence of subgroup given by $Z_0(G) = 1$ and*

$$\frac{Z_{n+1}(G)}{Z_n(G)} = Z\left(\frac{G}{Z_n(G)}\right).$$

**Remark 7.8.** *By induction, $Z_n(G)$ and $\gamma_n(G)$ are characteristic in $G$. and $Z_n(G) \trianglelefteq G$ and $\gamma_n(G) \trianglelefteq G$ for all $n \geq 0$.*

**Remark 7.9.** *For any $G$ and $N \trianglelefteq G, [G, N] \leq N$ because for any $g \in G, n \in N$ we have $[g, n] = gng^{-1}n^{-1}$ and $n^g \in N$ since $N \trianglelefteq G$. So $[G, N] \leq N$.*

**Theorem 7.10.** *$G$ is nilpotent iff there is a $n \geq 0$ such that $Z_n(G) = G$ iff there is $n \geq 0$ such that $\gamma_{n+1}(G) = \{1\}$. Moreover, the least $n$ such that $Z_n(G) = G$ is the least $n$ such that $\gamma_{n+1}(G) = \{1\}$.*

*Proof.* Take any central series then the decreasing and ascending central series will grow at least as fast as the arbitrary central series. □

## 7.2 Free Groups

**Example 7.11.** *Consider $\mathbb{Z} = (\mathbb{Z}, +)$, 1 is a generator $\mathbb{Z} = \langle 1 \rangle$. The universal property of $\mathbb{Z}$, 1. For any group $G$ and any $g \in G$, there is a unique homomorphism $\phi : \mathbb{Z} \to G$ and $\phi(1) = g$.*

*Proof.* If $\phi$ exist let $\phi(n) = g^n$ then $\phi$ is a homomorphism. □

*$g$ is free because whatever 1 is mapped to in the homomorphism it won't mess you up compared to a finite group.*

**Definition 7.12.** *$F$ is a free group on 2 element if*

1. *There exist $a, b \in F$, $F = \langle a, b \rangle$*
2. *For all $G$, all $g, h \in G$ there is a unique homomorphism $\phi : F \to G$ such that $\phi(a) = g$ and $\phi(b) = h$.*

# 8 Lecture 8: Free Groups

**Theorem 8.1.** *For any set $X$, there are a group $F$ and an injective function $i : X \to F$ such that*

1. *For any group $G$ and any function $f : X \to G$ there is a unique homomorphism $\phi : F \to G$ such that $\phi(i(x)) = f(x)$ for all $x \in X$.*

$$\begin{tikzcd} X \arrow[r, hook, "i"] \arrow[dr, "f"'] & F \arrow[d, dashed, "\phi"] \\ & G \end{tikzcd}$$

   *The diagram commutes: $\phi \circ i = f$. The homomorphism $\phi$ is unique.*

2. *$F$ is generated by $i[X]$*

*Proof.* A word (on alphabet $X$) is a (possibly empty) finite sequence from $X \times \mathbb{Z}$. We will write $x_1^{n_1} x_2^{n_2} \cdots x_k^{n_k}$ for the word $x_1, x_2, \ldots, x_k$ for $\langle (x_1, n_1), (x_2, n_2), \ldots, (x_k, n_k) \rangle$.

**Example 8.2.** *Let $X = \{x, y, z\}$ then $w = x^0 x^{15} y^{-23} z^{27}$ is a word.*

Let $w$ be the set of words. Given a word, we can possibly perform a reduction step

1. If word contains an entry $\langle x, 0 \rangle$, remove it.
2. If the word contains successive entries $\langle x, n \rangle$ and $\langle x, m \rangle$ then replace them with $\langle x, n + m \rangle$.

**Example 8.3.** $y^{15} x^3 x^{-2} x^{-1} y^2 z^5 \to y^{15} x^1 x^{-1} y^2 z^5 \to y^{15} x^0 y^2 z^5 \to y^{15} y^2 z^5 \to y^{17} z^5$.

Let $R$ be set of all reduced words, where no more reductions can be done.
For any word $w$, there is a unique reduced word $w'$ such that $w$ can be transformed into $w'$ by a finite set of reductions.
We will not be bale to show this property using group properties but instead we'll produce a group $F$ with the property that every word is equivalent to a unique reduced word.

Given $x \in X$, I will replace $f_x : F \to F$

1. $w$ does not begin with some $x^m$ then $f_x(w) = xw$
2. $w$ is of the form $x^{-1}v$ then $f_x(w) = v$
3. $w$ is of the form $x^n v$ for $n \neq -1$ then $f_x(w) = x^{n+1} v$

I also define $g_x : R \to R$ by

1. $w$ does not begin with some $x^n$ then $g_x(w) = x^{-1} w$
2. $w$ is of the form $x^1 v$ then $g_x(w) = v$
3. $w$ is of the form $x^n v$ for $n \neq 1$ then $g_x(w) = x^{n-1} v$

We can observe that $f_x \circ g_x = g_x \circ f_x = 1_R$.
For each $x \in X$, $f_x \in \sum_R$ (the group of permutations of $R$) and $g_x = f_x^{-1}$. Let $F$ be the subgroup of $\sum_R$ generated by $\{f_x : x \in X\}$ or $\{f_{x_1}^{n_1}, ..., f_{x_t}^{n_t} : x_i \in X, n_i \in \mathbb{Z}\}$.

Let $i : X \to F$ with $i(x) = f_x$. For any word $w = x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t}$. Let $f_w = f_{x_1}^{n_1} f_{x_2}^{n_2} \cdots f_{x_t}^{n_t} \in F$.
Key Facts:

1. If $w$ is obtained from $\overline{w}$ by a single reduction step, $f_w = f_{\overline{w}}$.
2. If $w$ is obtained from $\overline{w}$ by any sequence of reduction steps, then $f_w = f_{\overline{w}}$.
3. If $w \in R$ then $f_w(\langle\rangle) = w$
4. For any word $w$, $f_w(\langle\rangle)$ the unique reduced word to which $w$ can be transformed by reduction step.
5. If $w_1, w_2$ are reduced words then $v =$ unique reduction of $w_1 w_2$, $f_v = f_{w_1} f_{w_2}$

Let $F'$ be the group whose underlying set is $R$ whose operation is "concatenate and reduce". Then $F \simeq F'$ by $f_v \mapsto v$. To finish, let $G$ be a group, $h : X \to G$ function. We want to find there is a unique homomorphism $\phi$ with $F' \to G$ and $\phi \circ i = h$.

1. If $\phi$ exists then

$$\begin{aligned}\phi(x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t}) &= \phi(i(x_1))^{n_1} \phi(i(x_2))^{n_2} \cdots \phi(i(x_t))^{n_t} \\ &= h(x_1)^{n_1} h(x_2)^{n_2} \cdots h(x_t)^{n_t}\end{aligned}$$

2. Verify $\phi$ is a homomorphism.

$\square$

**Example 8.4.** *Let $X = \langle x, y\rangle$, let $G$ be any group with 2 generators say $G = \langle a, b\rangle$. Let $F$ be a free group on $X$. Let $\phi : F \to G$ be unique homomorphism such that $\phi(x) = a$ and $\phi(y) = b$. $\phi$ is surjective because $G$ is generated by $a$ and $b$. Thus, $G \simeq \frac{F}{\ker \phi}$.*

# 9 Lecture 9: Generators and Relations (Feb 2)

**Definition 9.1.** *let $X$ be a set, let $F$ be a free group on $X$. Let $A \subseteq F$ and $N =$ least normal subgroup of $F$ containing $A$.*

**Theorem 9.2.** *Let $\overline{F} = F/N$ and $j : X \to \overline{F}$ by $j(x) = xN$.*

$$\begin{tikzcd} F \arrow[r, "\phi_N"] & \overline{F} \\ X \arrow[u, "i"] \arrow[ur, "j"] & \end{tikzcd}$$

*The following are true:*

1. *$\overline{F}$ is generated by $\{j(x) : x \in X\}$. Let $\overline{x} = j(x)$*
2. *If $a = x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t} \in A$ then $a \in N$ and $\overline{x_1}^{n_1} \overline{x_2}^{n_2} \cdots \overline{x_t}^{n_t} = \phi_N(a) = 1_{\overline{F}}$*

*"Universal Property of $\overline{F}$": Let $G$ be a group and let $h : X \to G$ such that $a = x_1^{n_1} x_2^{n_2} \cdots x_t^{n_t} \in A$ and $h(x_1)^{n_1} h(x_2)^{n_2} \cdots h(x_t)^{n_t} = 1_G$. There is an unique homomorphism $\mathcal{T} : \overline{F} \to G$ such that*

$$\begin{tikzcd} X \arrow[r, "j"] \arrow[dr, "h"'] & \overline{F} \arrow[d, dashed, "\mathcal{T}"] \\ & G \end{tikzcd}$$

*So $h = \mathcal{T} \circ j$*

*Proof.* Since $\{\overline{x} : x \in X\}$ generates $\overline{F}$ there is at most one $\tau$. By universal property of $F$, there is unique homomorphism $\phi : F \to G$ such that

$$\phi(x^1) = h(x). \qquad \text{(For all } x \in X\text{)}$$

Use property of $F$

$$\begin{tikzcd} X \arrow[r, "i"] \arrow[dr, "h"'] & F \arrow[d, dashed, "\phi"] \\ & G \end{tikzcd}$$

Claim: By hypothesis on $h$, $A \subseteq \ker(\phi)$.
Let $a = x_1^{t_1} x_2^{t_2} \cdots x_k^{t_k} \in A$ then $\phi(a) = \phi(x_1)^{t_1} \phi(x_2)^{t_2} \cdots \phi(x_k)^{t_k} = h(x_1)^{t_1} h(x_2)^{t_2} \cdots h(x_k)^{t_k} = 1_G$ since $h(x_1)^{t_1} h(x_2)^{t_2} \cdots h(x_k)^{t_k} = 1_G$. Thus, $a \in \ker(\phi)$.

By choice of $N$, $N \leq \ker(\phi)$, we'll attempt to define $\tau : \overline{F} \to G$ by $\tau(wN) = \phi(w)$ for $w \in F$.
$\tau$ is well defined because $w_1 N = w_2 N \implies w_1^{-1} w_2 \in N \implies w_1^{-1} w_2 \in \ker(\phi) \implies \phi(w_1)^{-1} \phi(w_2) = 1 \implies \phi(w_1) = \phi(w_2)$. Showing $\tau$ is a homomorphism is straightforward.

$$\begin{tikzcd} X \arrow[r, "i"] \arrow[dr, "h"'] & F \arrow[d, "\phi_N"] \arrow[dr, "\phi"] & \\ & \overline{F} \arrow[r, dashed, "\mathcal{T}"] & G \end{tikzcd}$$

$\square$

# 10 Lecture 10: Presentation of Groups (Feb 4)

**Example 10.1.** *Dihedral Group: $D_n$ = symmetry group of a regular $n$-gon.*
*Let $D_n = \langle a, b\rangle$ where $a$ = rotation by $\frac{2\pi}{n}$ and $b$ = reflection about the vertical axis. Then $D_n$ is generated by $a$ and $b$ with the relations $a^n = 1$ and $b^2 = 1$.*
*We have $bab^{-1}a = 1$*
*Let $X = \{x, y\}$ and $F$ be free group and $A = \{x^n, y^2, yxy^{-1}x\}$ and $N$ = least normal subgroup $\supseteq A$. Let $\overline{x} = xN$ and $\overline{y} = yN$. Then $F/N = \langle \overline{x}, \overline{y}\rangle$ and*

$$\overline{yxy^{-1}x} = \overline{yxy}^{-1}\overline{x}^{-1} = 1_{\overline{F}}.$$

*There is a unique homomorphism $\tau : F/N \to D_n$ such that $\tau(\overline{x}) = a$ and $\tau(\overline{y}) = b$ as $D_n = \langle a, b\rangle$ so $\tau$ is surjective. So $|F/N| \geq 2n$.*
*Claim: $|F/N| \leq 2n$ (so $\tau$ is isomorphism)*
*Proof: Using relations from before, we can simply write any expression in $\overline{x}, \overline{y}$ down to $\overline{x}^s\overline{y}^t$ for $0 \leq s \leq n$ and $0 \leq t \leq 2$.*

# 11 Lecture 11: Category Theory (Feb 6)

## 11.1 Categories

**Example 11.1.** *1. Category of sets, if we have two sets $A$ and $B$ then $A \to B$ is a function $f : A \to B$*
*2. Category of groups, if we have two groups $G$ and $H$ then $G \to H$ is a homomorphism $\phi : G \to H$*
*3. Category of topological spaces, if we have two topological spaces $X$ and $Y$ then $X \to Y$ is a continuous function $f : X \to Y$*

**Definition 11.2.** *A category $\mathcal{C}$ consists of the following data:*
1. *A collection of objects $Ob(\mathcal{C})$*
2. *A collection of morphisms $Mor(\mathcal{C})$*
3. *For each morphism $f$, there are objects $\mathrm{dom}(f)$ and $\mathrm{cod}(f)$ such that $dom(f) \to cod(f)$ is a morphism in $Mor(\mathcal{C})$*
4. *A composition operation $Mor(\mathcal{C}) \times Mor(\mathcal{C}) \to Mor(\mathcal{C})$*
5. *An identity morphism $1_A : A \to A$ for each object $A \in Ob(\mathcal{C})$*

**Lemma 11.3.** *The following are true:*
1. *If $a \xrightarrow{f} b$ then $f1_a = f$ and $1_b f = f$.*
2. *If $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} d$ then $h(gf) = (hg)f$*

**Example 11.4.** *Posets: $\mathbb{P}$ is a poset with $\leq$ binary relation such that*
1. *$a \leq a$ for all $a \in \mathbb{P}$*
2. *$a \leq b$ and $b \leq a \implies a = b$*
3. *$a \leq b$ and $b \leq c \implies a \leq c$*

*Given poset $\mathbb{P}$, make "poset category" with objects the elements of $\mathbb{P}$. There is only one morphism $a \to b$ if $a \leq b$ and no morphism if $a \not\leq b$.*

**Definition 11.5.** *Let $\mathcal{C}$ be a category. An object is an initial if for all object $b$ there is a exactly one arrow $a \to b$.*

**Definition 11.6.** *An arrow $a \xrightarrow{f} b$ in $\mathcal{C}$ is an isomorphism iff there is another $b \xrightarrow{f} a$ such that $gf = 1_b$ and $fg = 1_a$.*

**Lemma 11.7.** *If $a, b$ are initial objects in $\mathcal{C}$ they are uniquely isomorphic.*

*Proof.* Let $a \xrightarrow{f} b$ and $b \xrightarrow{g} a$ be unique arrows from $a$ to $b$ and $b$ to $a$ respectively. Then we have $a \xrightarrow{gf} a$ and $b \xrightarrow{fg}$ so $gf = 1_a$ and $fg = 1_b$. □

**Definition 11.8.** *Object $b$ is terminal iff for all $a$ there is a unique $a \to b$ iff $b$ is initial in $\mathcal{C}^{op}$*

**Lemma 11.9.** *Two terminal objects are uniquely isomorphic.*

**Definition 11.10.** *Let $\mathcal{C}$ and $\mathcal{D}$ be categories. A functor from $\mathcal{C}$ to $\mathcal{D}$ is $\mathcal{C} \xrightarrow{F} \mathcal{D}$. Assign to each object $a$ of $\mathcal{C}$ an object $F(a)$ of $\mathcal{D}$. For each arrow $a \xrightarrow{f} b$ there is an arrow $F(a) \xrightarrow{F(f)} F(b)$ with $f(1_a) = 1_{F(a)}$. Additionally, $F(gf) = F(g)F(f)$*

# 12 Lecture 12: Ring Theory (Feb 8)

**Definition 12.1.** *A ring is a set $R$ with two binary operations $+$ and $\times$ such that*
1. *$(R, +)$ is an abelian group*
2. *$(R, \times)$ is associative*
3. *$a \times (b + c) = a \times b + a \times c$ and $(b + c) \times a = b \times a + c \times a$ for all $a, b, c \in R$*

**Example 12.2.** *The following are rings*
1. $\mathbb{Z}, 2\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$
2. $M_2(\mathbb{Z})$
3. $\{f : \mathbb{R} \to \mathbb{R}$ *continuous*$\}$
4. $\{f : \mathbb{R} \to \mathbb{R}$ *differentiable*$\}$

**Definition 12.3.** *$R$ is commutative iff $rs = sr$ for all $rs \in R$*

**Definition 12.4.** *$R$ is unital iff there exists $1_R \in R$ such that $1_R r = r 1_R = r$ for all $r \in R$*

**Definition 12.5.** *Let $R, S$ be rings. A ring homomorphism from $R$ to $S$ if $\phi : R \to S$ satisfy:*
1. *$\phi(r + s) = \phi(r) + \phi(s)$ for all $r, s \in R$*
2. *$\phi(rs) = \phi(r)\phi(s)$ for all $r, s \in R$*
3. *$\phi(1_R) = 1_S$*

**Definition 12.6.** *Let $S$ be a subring of $R$ if $S$ is a ring and the inclusion map $i : S \to R$ is a ring homomorphism.*

**Definition 12.7.** *Let $\phi : R \to S$ be a ring homomorphism then*
1. *$\ker(\phi) = \{r \in R : \phi(r) = 0\}$ is a subring of $R$*
2. *$\phi(R) = \{s \in S : \exists r \in R, \phi(r) = s\}$ is a subring of $S$*

**Remark 12.8.** *In general, not the case that $\ker(\phi)$ is a subring of $R$.*

**Definition 12.9.** *Let $R$ be a ring. An ideal of $R$ is a subring $I$ of $R$ such that*
1. *$I \leq (R, +)$*
2. *For all $a \in R$ and $b \in I$, $ab \in I$.*

**Definition 12.10.** *Let $I$ be an ideal of $R$ then we define $r + I = \{r + i : i \in I\}$*

# 13 Lecture 13: Ideals and Quotient Rings (Feb 10)

**Theorem 13.1.** *First Isomorphism Theorem of Rings: Let $R, S$ be rings and $\phi : R \to S$ be a homomorphism with $\ker(\phi)$ an ideal of $R$ and $\mathrm{im}(\phi)$ a subring of $S$. Then we get isomorphism*

$$\tau : \frac{R}{\ker(\phi)} \xrightarrow{\cong} \mathrm{im}(\phi) \text{ and } \tau : r + \ker(\phi) \mapsto \phi(r).$$

**Theorem 13.2.** *The following are true for any ideals $I, J$ of $R$:*

1. *$I + J =$ smallest ideal containing $I$ and $J$*
2. *$I \cap J =$ largest ideal contained in $I$ and $J$*
3. *$IJ = \{\sum_{i=1}^{n} r_i s_i | n \in \mathbb{N}, r_i \in I, s_i \in J\} \subseteq I \cap J$*
4. *Ideals of $R/I$ are in bijection with ideals of $R$ that contain $I$*

**Definition 13.3.** *$R$ is a zero ring if $R = \langle 0_R \rangle$*
*Note: If $1_R = 0_R$ then $r \cdot 1 = r \cdot 0 = 0$ and $R$ is a zero ring.*

**Definition 13.4.** *$R$ is an integral domain iff*

1. *$1_R \neq 0_R$ ($R$ is not a zero ring)*
2. *For all $a, b \in R$*

$$ab = 0 \implies a = 0 \text{ or } b = 0.$$

**Definition 13.5.** *$R$ is a field iff*

1. *$1_R \neq 0_R$*
2. *Every element of $R \setminus \{0_R\}$ has a multiplicative inverse*

**Exercise 13.6.** *If $R$ is a field then $R$ is an integral domain.*

**Definition 13.7.** *Let $R$ be a ring. Then $u \in R$ is a unit iff $u$ has a multiplicative inverse.*
*We let*

$$U(R) = \{u \in R : u \text{ is a unit}\}.$$

*$U(R)$ is a group under multiplication.*

**Definition 13.8.** *An ideal $I$ of $R$ is principal iff $I = aR$*

**Lemma 13.9.** *Let $R$ be any ring then $R$ is the largest ideal and $\{0_R\}$ is the smallest ideal.*

**Remark 13.10.** *We have $R/R = \{0_R\}$ and $R/\{0_R\} \simeq R$*

**Theorem 13.11.** *Let $R$ be a field and let $I$ be an ideal then $I = 0$ or $I = R$*

*Proof.* Assume $I \neq 0$, let $a \in I$ and $a \neq 0$. There exist $b \in R$ such that $ab = 1 \in I$. In

any ring $1 \in I$ iff $I = R$.

$\square$

**Theorem 13.12.** *If $R \neq 0$ are the only two ideals of $R$ then $R$ is a field.*

*Proof.* Let $a \in R, a \neq 0$ then $(a) \neq \emptyset$ as $a \in (a)$ so $(a) = R$ so $1 \in (a)$ and $1 = ab$ for some $b$. $\square$

**Definition 13.13.** *An ideal $I$ of $R$ is maximal iff*

1. *$I \neq R$*
2. *For all ideals $J \supseteq I$, $J = I$ or $J = R$*

*$I$ is maximal in poset $\{J : J \text{ ideal}, J \neq R\}$ ordered by $\subseteq$*

**Lemma 13.14.** *If $I$ is an ideal of $R$,*

$$I \text{ maximal} \iff R/I \text{ is a field.}$$

*Proof.* $R \neq I \iff 1_{R/I} \neq 0_{R/I}$. By $I$ maximal iff there are only two ideals of $R/I$ which are 0 and $R/I$. $\square$

**Definition 13.15.** *Ideal $I$ is prime iff*

1. *$I \neq R$*
2. *For all $a, b \in R$, $ab \in I \implies a \in I$ or $b \in I$*

**Lemma 13.16.** *$R/I$ is an integral domain iff $I$ is a prime ideal.*

**Definition 13.17.** *$Spec(R) = \{I : I \text{ is a prime ideal of } R\}$*

**Theorem 13.18.** *Let $R$ be a ring. $I$ is an ideal of $R$, $I \neq R$. Then there is a maximal ideal $J$ such that $I \subseteq J$.*

# 14 Lecture 14: Zorn's Lemma and Modules (Feb 13)

**Lemma 14.1.** *Zorn's Lemma: If $\mathbb{P}$ is a poset such that every chain in $\mathbb{P}$ has an upper bound, then for every element $p$ of $\mathbb{P}$ there is a $q \geq p$ with $q$ maximal.*

**Example 14.2.** *Claim: $\mathbb{Q}$ satisfies the hypothesis of Zorn's Lemma.*

*Proof.* Let $(I_a : a \in A)$ be a chain in $\mathbb{Q}$. That is, $I_a$ is ideal with $I_a \neq R$ and for $a, b \in A, I_a \subseteq I_b$ or $I_b \subseteq I_a$. We need upper bound, let

$$J = \bigcup_{a \in A} I_a = \{r : \text{there is } a \in A, r \in I_a\}.$$

Verify $J$ is an ideal and for $r, s \in J$ then $r \in I_a$ and $s \in I_b$. WLOG let $I_a \subseteq I_b$ so $r, s \in I_b$ so $r + s \in I_b \subseteq J$.
Verify $J \neq R$, easy as $1 \notin I_a$ all $a$ so $1 \notin J$. □

## 14.1 Modules

**Definition 14.3.** *Let $R$ be a ring. A $R$-module is an abelian group $(M, +)$ equipped with a function $R \times M \to M$ by $(r, m) \mapsto rm$ such that*
1. *$0_R m = 0_M, 1m = m$ for all $m \in M$*
2. *$(r_1 + r_2) = r_1 m + r_2 m$ and $m(r_1 + r_2) = mr_1 + mr_2$*
3. *$r(sm) = (rs)m$ for $r, s \in R$ and $m \in M$*

**Definition 14.4.** *If $M$ is a $R$-module, we can form "linear combinations"*

$$\sum_{i=1}^{n} r_i m_i \text{ for } r_i \in R \text{ and } m_i \in M.$$

**Example 14.5.** *Let $R = \mathbb{Z}$ and $(G, +)$ be an abelian group. Then $G$ is a $\mathbb{Z}$-module.*

**Definition 14.6.** *If $M$ is a $R$-module, a submodule of $M$ is $N$ such that $N \leq (M, +)$, $rn \in N$ for $r \in R$ and $n \in N$.*

**Remark 14.7.** *If $R$ is a ring, we can view $R$ as a $R$-module. The ideals of $R$ are the submodules of $R$ (viewed as a $R$-module).*

**Definition 14.8.** *Let $M, N$ be $R$-modules. A $R$-module homomorphism is a function $\phi : M \to N$ such that*
1. *$\phi$ is a group homomorphism for $(M, +)$ to $(N, +)$*
2. *$\phi(rm) = r\phi(m)$ for $r \in R$ and $m \in M$*

*Note: This is like linear transformation in linear algebra.*

# 15 Lecture 15: Modules (Feb 16)

**Definition 15.1.** *Let $M$ and $N$ be $R$-modules then we say $\phi : M \to N$ is $R$-linear if*

$$\phi\left(\sum_{i=1}^{n} r_i m_i\right) = \sum_{i=1}^{n} r_i \phi(m_i) \text{ for } r_i \in R \text{ and } m_i \in M.$$

**Definition 15.2.** *Let $M \leq N$ ($M$ is submodule of $N$), $N/M = \{n + M : n \in N\}$. We define*

1. $(n_1 + M) + (n_2 + M) = (n_1 + n_2) + M$
2. $r(n + M) = (rn) + M$

*We can define the $R$-linear quotient map $\phi_M : N \to N/M$ by $\phi : n \mapsto n + M$ with $\ker(\phi_M) = M$.*

**Theorem 15.3.** *First Isomorphism Theorem for Modules: Let $\phi : M_1 \to M_2$ linear, then*

$$\mathrm{im}(\phi) \simeq \frac{M_1}{\ker(\phi)}.$$

*via $m + \ker(\phi) \mapsto \phi(m)$*

**Definition 15.4.** *Given $R$-module $M$ and set $X \subseteq M$, the least submodule containing $X$ is*

$$\mathrm{Span}(X) = \left\{\sum_{i=1}^{n} r_i x_i | n \in \mathbb{N}, r_i \in R, x_i \in X\right\}.$$

*We specifically define* $\mathrm{Span}(\emptyset) = \{0_M\}$

**Definition 15.5.** *If $R$ is a ring, $X \subseteq R$ we write $\langle X \rangle$ for the least ideal containing $X$*

$$\langle X \rangle = \left\{\sum_{i=1}^{n} r_i x_i | n \in \mathbb{N}, r_i \in R, x_i \in X\right\}.$$

**Definition 15.6.** *A ring $R$ is a principal ideal ring iff all ideals of $R$ are principal (a principal ideal is of form $Ra$ for some $a \in R$).*

**Definition 15.7.** *$R$ is a principal ideal domain iff $R$ is integral domain and principal ideal ring.*

**Definition 15.8.** *A ring $R$ is noetherian iff every ideal of $R$ is finitely generated. (An ideal is finitely generated if $I = Ra_1 + \cdots + Ra_n$ for some $a_1, \ldots, a_n \in R$.)*

**Theorem 15.9.** *If $I$ is an ideal of $R$ the ideals $R/I$ are in bijection with $\{J : J \text{ ideal of } R, J \supseteq I\}$.*
*Also prime ideals of $R/I$ correspond to prime ideals of $R$ that contain $I$.*

**Example 15.10.** *Common PID's*
1. $\mathbb{Z}$
2. $k[x]$ *for* $k$ *a field*

**Definition 15.11.** *$R$ is a Euclidean Domain iff*
1. *$R$ is integral domain*
2. *There is a function $\phi : R \setminus \{0\} \to \mathbb{N}$ such that for all $a, b \in R$, for $b \neq 0$, there exist $q, r$ such that $a = bq + r$ either $r = 0$ or $\phi(r) < \phi(b)$*

**Example 15.12.** *If $R = \mathbb{Z}$ then $\phi(n) = |n|$ is a Euclidean Domain.*

**Example 15.13.** *If $R = k[x]$ for $k$ a field then $\phi(f) = \deg(f)$ is a Euclidean function.*

**Theorem 15.14.** *If $R$ is a Euclidean Domain then $R$ is a PID.*

*Proof.* Let $R$ be euclidean domain and let $I$ be the ideal of $R$. If $I = 0$ then we're done. If $I \neq 0$ consider $\{\phi(r) | r \in I, r \neq 0\} \subseteq \mathbb{N}$. Let $n$ be the minimal element and let $a \in I \setminus \{0\}$ such that $\phi(a) = n$. As $a \in I, Ra \subseteq I$. Let $b \in I$, find $q, r$ such that $b = aq + r$ either $r = 0$ or $\phi(r) < n$. We have $r = b - aq \in I$. So $r = 0$ and $b \in Ra$. □

**Lemma 15.15.** *Facts about Integral Domains: Let $R$ be an integral domain.*
1. *If $a \neq 0$ and $ab = ac$ then $a(b - c) = 0$*
2. *$Ra = Rb$ iff $a$ and $b$ are associates*

**Lemma 15.16.** *Let $R$ be any ring, the following are true about units*
1. *$u$ unit iff $u$ has a multiplicative inverse*
2. *Units of $R$ form a group under multiplication*
3. *$a, b \in R$ are called associates iff there is an unit $u$ such that $au = b$.*

**Example 15.17.** *$\mathbb{Z}[x]$ is not a PID. Consider the ideal $\langle 2, x \rangle = \{f :$ constant term of $f$ is even$\}$.*

**Theorem 15.18.** *$R$ Noetherian iff $R[x]$ is Noetherian.*

# 16 Lecture 16: Integral Domains

**Definition 16.1.** *Let $p \in R$*

1. *$p$ is prime if and only if $p$ is a nonzero non-unit and for all $r, s \in R$, $p|rs$ implies $p|r$ or $p|s$.*
2. *$p$ is irreducible if and only if $p$ is a nonzero non-unit and for all $r, s \in R$ and $p = rs$ then one of $r$ or $s$ is a unit.*

**Lemma 16.2.** *The following are true*

1. *As $R$ is an ID, for any $a \in R$, $a$ nonzero non-unit iff $(a) \notin \{0, R\}$*
2. *If $p$ is prime, then $(p)$ is a prime ideal*
3. *$p$ prime implies $p$ is irreducible.*
4. *If $R$ PID then prime = irreducible.*
5. *In any ID, $(0)$ is a prime ideal.*

**Definition 16.3.** *$R$ is a unique factorization domain $(UFD)$ iff*

1. *$R$ is an integral domain*
2. *Every non-zero nonunit in $R$ is a product $u_1, ..., u_n$, $n > 0, u_i$ irreducible.*
3. *If $m, n > 0$ then $u_1...u_n = v_1...v_m$ and $u_i, v_j$ irreducible, then $m = n$ and $u_i$ are associates and permutations of $v_j$*

**Recall 16.4.** *An $R$-module $M$ is finitely generated $(fg)$ if there are $m_1, ..., m_n \in M$ such that $M = \mathrm{Span}(m_1, ..., m_n)$.*

**Definition 16.5.** *An $R$-module $N$ is Noetherian iff every submodule of $N$ is finitely generated.*
*Equivalently, Ring $R$ is Noetherian iff $R$ is Noetherian as an $R$-module that is all ideals are finitely generated.*

**Example 16.6.** *Non-Noetherian Rings:*

$$R = \bigcup_{n=1}^{\infty} \mathbb{Z}[x_1, ..., x_n].$$

*Consider the ideal*

$$I = \langle x_1, x_2, ... \rangle = \{f \in R : f \text{ has no constant term}\} = \{f : f(0, ...) = 0\}.$$

*Proof.* If $I$ is finitely generated there is $h_1, ..., h_n \in I$ such that

$$I = Rh_1 + \cdots + Rh_n = (h_1) + \cdots + (h_n).$$

Each $h_i$ has finite $x_j$'s in it so find $N$ so large that $x_N$ does not appear in any $h_i$.
As $x_N \in I$ there are $f_1, ..., f_n \in R$ such that $x_N = \sum_{i=1}^{n} f_i h_i$. Substitute

$$x_j = \begin{cases} 0 & \text{if } j < N \\ 1 & \text{if } j \geq N \end{cases}.$$

Then we conclude $1 = 0$ which is a contradiction. $\square$

**Theorem 16.7.** *Let $M$ be a $R$-module then the following are equivalent*

1. *$M$ is Noetherian*
2. *For any increasing change of submodules $N_0 \subseteq N_1 \subseteq N_2 \subseteq \cdots$ of $M$, the chain is eventually constant.*
3. *For any nonempty set $X$ of submodules of $M$, there is some $N \in X$ which is maximal under inclusion.*

*Proof.* (1) $\implies$ (2): Consider $N_\infty = \bigcup_{i=0}^{\infty} N_i = \{m : \exists i \in \mathbb{N}, m \in N_i\}$.
Verify that $N_\infty$ is a submodule of $M$, as $M$ is noetherian, $N_\infty$ is finitely generated say $N_\infty = \operatorname{Span}(X)$ with $X \subseteq N_\infty$ finite. Find $i$ such that $X \subseteq N_i$ then $N_\infty = \operatorname{Span}(X) \subseteq N_i \subseteq N_j \subseteq N_\infty$ for $j \geq i$.

(2) $\implies$ (1): AFSOC there is $N \leq M$ with $N$ not finitely generated. Take $n_0 \in N$ and $n_0 \neq 0$. Choose $n_1 \in N \setminus Rn_0$ and generally

$$n_k \subseteq N \setminus \left( \sum_{i<k} Rn_i \right).$$

In the end, let $N_i = \sum_{j<i} Rn_j$ then $N_0 < N_1 < N_2 < \cdots$, a contradiction. $\square$

# 17 Lecture 17: PID (Feb 19)

**Lemma 17.1.** *If $R$ PID then*
1. *$R$ is Noetherian*
2. *$0$ is a prime ideal and $0$ is maximal iff $R$ is a field*
3. *For $a \neq 0, (a)$ prime iff $a$ prime iff $(a)$ maximal iff $a$ irreducible*

**Theorem 17.2.** *If $R$ is a PID then $R$ UFD*

**Claim 17.3.** *Every nonzero non-unit $a \in R$ is a finite product of irreducibles.*

*Proof.* If not, choose $a$ that is not a finite product of irreducibles such that $(a)$ is maximal under inclusion, $a$ not finite product of irreducibles so $a$ is not irreducible.
So $a = bc$ where $b, c$ not units. So $(a) \subsetneq (b)$ and $(a) \subsetneq (c)$ so $b, c$ are finite products of irreducibles as $(a)$ is maximal under inclusion. Contradiction so $a$ is a finite product of irreducibles. □

**Claim 17.4.** *If $u_1...u_n = v_1...v_n$ with $u_i, v_j$ irreducible then $m = n$ and $u_i, v_j$ are the same up to permutations and associates.*

*Proof.* $u_1$ divides $v_1...v_m$. As $R$ PID, $u_1$ is prime so $u_1|v_j$ for some $j$. As $v_j$ is irreducible, $u_i$ either unit or an associate of $v_j$. As $u_i$ irreducible, $u_i$ not a unit. So $u_1$ and $v_j$ are associates. Say $u_1 = x \cdot v_j$ for $x$ unit. Then

$$xu_2...u_n = v_1...v_{j-1}v_{j+1}...v_m.$$

Finish by induction. □

**Corollary 17.5.** *If $k$ field, $k[x]$ is a UFD.*

**Lemma 17.6.** *If $R$ PID and $p \in R$ irreducible then $R/(p)$ is a field.*

**Theorem 17.7.** *(Hilbert's Basissatz Theorem) If $R$ Noetherian ring then $R[x]$ is Noetherian ring.*

*Proof.* Let $I$ be ideal of $R[x]$. For each $n$ let

$$J_n = \left\{ r \in R : \exists c_0, ..., c_{n-1} \in R, rx^n + \sum_{i=0}^{n-1} c_i x^i \in I \right\}.$$

$J_n$ is an ideal of $R$.
Claim: $J_n \subseteq J_{n+1}$
Let $r \in J_n$ say $rx^n + \sum_{i=0}^{n-1} c_i x^i \in I$, as $I$ ideal, $x(rx^n + ...) \in I$ so $r \in J_{n+1}$.

As $R$ is noetherian, there is $N$ such that $J_n = J_N$ for all $n \geq N$.
For each $n \leq N$, as $R$ noetherian, $J_n$ is finitely generated ideal of $R$. For each $n \leq N$,

choose a finite set $F_n \subseteq I$ such that if $\deg(f) = n$ then $f \in F_n$ and $\{$leading coeff of $f$ : $f \in F_n\}$ generates $J_n$.
Let $F = \bigcup_{n \leq N} F_n$ and claim that $F$ generates $I$ (as an ideal of $R[x]$). Show by induction on $\deg(h)$ that every $h \in I$ is an $R[x]$-linear combination of elements of $F$.
(Base Case) $h = 0$, $\deg(h) \leq N$ and $h = rx^n + \sum_{in} c_i x^i$ where $r \in J_n$. SO find $\lambda_1, ..., \lambda_t \in R$ and $g_1, ..., g_t \in F_n$ such that $r =$ leading coefficient of $\sum_{i=1}^{t} \lambda_i g_i$. Look at $h - \sum_{i=1}^{t} \lambda_i g_i \in I$ has lower degree.
(Induction Step) Let $h \in I$ with $\deg(h) = n > N$, $h = rx^n + \sum_{i<n} c_i x^i$ with $r \in J_n = J_N$. Find $g'_1, ..., g'_s \in F_N, \lambda_1, ..., \lambda_s \in R$ such that $r =$ leading coefficients of $\sum_{j=1}^{s} \lambda_j g'_j$ with leading term in $rx^N$. So now look at $h - \sum_{j=1}^{s} \lambda_j x^{n-N} g'_j$. This has degree $< n$, so done by induction. $\square$

# 18 Lecture 18: Unique Factorization Domains (Feb 23)

**Theorem 18.1.** *$R$ UFD $\implies$ $R[x]$ UFD*

**Definition 18.2.** *Field of Fractions: Given any integral domain $R$, constructing its "field of fractions" as $X = \{(a,b) \in R \times R : b \neq 0\}$.*
*Define binary relation $\sim$ on $X$ by*

$$(a_1, b_1) \sim (a_2, b_2) \iff a_1 b_2 = a_2 b_1.$$

**Definition 18.3.** *Define $\frac{a}{b}$ to be the $\sim$-equivalence class of $(a,b)$.*
*The fractions form a ring $F$ with $1_F = \frac{1}{1}$ and $0_F = \frac{0}{1}$. Every element has an inverse, so $F$ is a field.*

**Remark 18.4.** *If $R$ is integral domain and $F$ is a field of fractions then $F[x]$ is a PID.*

**Definition 18.5.** *Let $R$ be an integral domain and let $a_1, ..., a_n \in R, a_i \neq 0$. A greatest common divisor (gcd) for $a_1, ..., a_n$ is $a \in R$ such that*
1. *$a | a_i$ for $i = 1, ..., n$*
2. *If $b | a_i$ for $i = 1, ..., n$ then $b | a$*

**Remark 18.6.** *If $\gcd(a_1, ..., a_n)$ exists, it is unique up to associates.*

**Lemma 18.7.** *If $R$ UFD, then gcd's exist.*

**Lemma 18.8.** *If $R$ PID and $(a_1, ..., a_n) = Ra_1 + \cdots + Ra_n$ then*

$$(a_1, ..., a_n) = (d) \iff d = \gcd(a_1, ..., a_n).$$

**Definition 18.9.** *Let $R, S$ be rings, $\phi : R \to S$ a ring homomorphism.*
1. *If $I$ is an ideal of $R$, $I^e$ (extension of $I$) = ideal generated by $\phi[I]$ in $S$.*
2. *If $J$ ideal of $S$, $J^c$ (contraction) of $J$ is $\phi^{-1}[J] = \{r \in R : \phi(r) \in J\}$.*

**Lemma 18.10.** *If we define $\psi : R/J^c \to S/J$ by $\psi : r + J^c \mapsto \phi(r) + J$, $\psi$ is well defined injective homomorphism.*

**Lemma 18.11.** *If $J$ is a prime ideal of $S$, then $J^c$ is a prime ideal of $R$.*

*Proof.* $S/J$ is ID, so all subrings of $S/J$ are ID's, so $R/J^c$ is ID, so $J^c$ is prime ideal. $\square$

**Recall 18.12.** *If $R$ a ring, a subset $S \subseteq R$ is multiplicatively closed iff $1 \in S$ and for all $s, t \in S$, $st \in S$.*
*Define a "ring of fractions" $S^{-1}R$.*

**Theorem 18.13.** *Let $X = \{(a, b) : a \in R, b \in S\}$ then*

$$(a_1, b_1) \sim (a_2, b_2) \iff \exists s \in S, s(a_1 b_2 - a_2 b_1) = 0.$$

**Theorem 18.14.** *Let the set be $S^{-1}R$. Define the map $\phi : R \to S^{-1}R$ by $\phi : r \mapsto \frac{r}{1}$. Then $\phi$ is a ring homomorphism but it may not be injective.*

**Theorem 18.15.** *Universal Property: For all rings $A$ and all homomorphism $\psi : R \to A$ such that $\phi(s)$ is a unit in $A$ for all $s \in S$. There is unique $X : S^{-1}R \to A$ such that $\psi = X \circ \phi$.*

$$\begin{tikzcd} R \arrow[r, "\phi"] \arrow[dr, "\psi"'] & S^{-1}R \arrow[d, dashed, "X"] \\ & A \end{tikzcd}$$

*Proof.* $X\left(\frac{x}{s}\right) = \psi(r)\psi(s)^{-1}$ □

# 19 Lecture 19: (Feb 25)

**Theorem 19.1.** *$R$ UFD $\implies$ $R[x]$ UFD*

**Remark 19.2.** *As $R$ ID for nonzero, $f, g \in R[x]$, $\deg(fg) = \deg(f) + \deg(g)$*

**Definition 19.3.** *Let $f \in R[x]$ with $f = 0$, $f$ is primitive iff there is no prime $p \in R$ such that $p|f$ in $R[x]$*

**Remark 19.4.** *Informally, the* gcd *of the coefficients of $f$ is 1.*

**Lemma 19.5.** *(Gauss's Lemma) If $f, g \in R[x]$ both primitive then $fg$ is primitive.*

*Proof.* Suppose not, and let $p$ be prime in $R$ such that $p|fg$. As $p$ prime, $P = (p)$ is prime ideal so $R/P$ is ID. Let $\phi : R \to R/P$ quotient map.
Define $\phi^* : R[x] \to R/P[x]$ by

$$\phi^*\left(\sum_{i=0}^{n} a_i x^i\right) = \sum_{i=0}^{n} \phi(a_i) x^i.$$

$\phi^*$ is homomorphism. So $\phi^*(fg) = \phi(f)\phi(g)$ but $\phi^*(fg) = 0$ as $p|fg$.
As $R/P$ ID, $\frac{R}{P}[x]$ is ID so $\phi^*(f) = 0$ or $\phi^*(g) = 0$. So $p|f$ or $p|g$ so a contradiction. □

**Lemma 19.6.** *Let $f \in R[x]$, $f \neq 0$. Then there are $C \in R$, $g \in R[x]$, $g$ primitive and $f = Cg$*

*Proof.* $C = \gcd(a_0, ..., a_n)$ if $f(x) = a_0 + a_1 x + \cdots + a_n x^n$. □

**Lemma 19.7.** *Let $h \in F[x]$, then there is a nonzero $a \in F$ such that $ah \in R[x]$ and $ah$ primitive.*

**Lemma 19.8.** *Let $f \in R[x]$ be irreducible with $\deg f > 0$. Then $f$ irreducible in $F[x]$*

*Proof.* Proof of Theorem 19.1:
To show existence of factorization: Let $f \in R[x]$ nonzero and nonunit. Write $f = cg$ where $c \in R$ and $g \in R[x]$ primitive. If $C$ nonunit (in $R$), use $R$ being UFD to write $c = r_1 ... r_n$ with $r_i$ irreducible in $R$.
Note: $r_i$ still irreducible in $R[x]$.
We have $g \in R[x] \implies g \in F[x]$. If $\deg(g) > 0$ then $g$ is not a unit and using $F[x]$ is UFD to write

$$g = h_1 ... h_n \text{ where } h_i \in F[x], h_i \text{ irreducible in } F[x].$$

Find $a_i \neq 0$ in $F$ such that $H_i = a_i h_i \in R[x]$ primitive.
Note: $H_i$ irreducible in $F[x]$.

So we can write

$$\prod_{i=1}^{n} H_i = \left(\prod_{i=1}^{n} a_i\right)\left(\prod_{i=1}^{n} h_i\right) = \left(\prod_{i=1}^{n} a_i\right) g.$$

Let $\prod a_i = \frac{A}{B}$ for $A, B \in R$ and $A, B \neq 0$. Then $B(\prod H_i) = Ag$. Argue that $A, B$ must be associates so $\prod a_i$ unit of $R$ and WLOG $\prod a_i = 1$.

To show uniqueness of factorization:
Suppose $f = r_1 ... r_n g_1 ... g_s = r'_1 ... r'_m g'_1 ... g'_s$ for $r_i, r'_i \in R$ irreducible and $g_i, g'_i \in R[x]$ irreducible and $\deg(g_i), \deg(g'_i) > 0$ for all $i$.
As $g_i, g'_i$ are primitive, $g_1 ... g_s$ and $g'_1 ... g'_s$ are primitive.
So $v_1 ... v_n, v'_1, ..., v'_m$ are associates in $R$ and using $R$ being UFD. By lemma 19.8, $g, g'$ are irreducible in $F[x]$. $\square$

# 20 Lecture 20: (Mar 2)

**Definition 20.1.** *Let $\mathcal{C}$ be a category and $c, d$ objects then a cone over $c, d$ is a diagram*

$$\begin{tikzcd} & e \arrow[dl, "\alpha"'] \arrow[dr, "\beta"] & \\ c & & d \end{tikzcd}$$

*A cone over $c, d$ is a product of $c, d$ iff for all cones*

$$\begin{tikzcd} & e' \arrow[dl, "\alpha'"'] \arrow[dr, "\beta'"] & \\ c & & d \end{tikzcd}$$

*there is a unique $\phi : e' \to e$ such that the diagram*

$$\begin{tikzcd} & e' \arrow[dl, "\alpha'"'] \arrow[d, dashed, "\phi"] \arrow[dr, "\beta'"] & \\ c & e \arrow[l, "\alpha"] \arrow[r, "\beta"'] & d \end{tikzcd}$$

*commutes (i.e. $\alpha \circ \phi = \alpha'$ and $\beta \circ \phi = \beta'$).*

**Example 20.2.** *Groups: Let $G_1, G_2, H$ be groups and $\phi : H \to G_1$, $\psi : H \to G_2$ homomorphisms. The product cone is $(G_1 \times G_2, \pi_1, \pi_2)$ where $\pi_1(g_1, g_2) = g_1$ and $\pi_2(g_1, g_2) = g_2$. The unique $\rho : H \to G_1 \times G_2$ with $\rho(h) = (\phi(h), \psi(h))$ makes the diagram*

$$\begin{tikzcd} & H \arrow[dl, "\phi"'] \arrow[d, dashed, "\rho"] \arrow[dr, "\psi"] & \\ G_1 & G_1 \times G_2 \arrow[l, "\pi_1"] \arrow[r, "\pi_2"'] & G_2 \end{tikzcd}$$

*commute ($\pi_1 \circ \rho = \phi$ and $\pi_2 \circ \rho = \psi$).*

**Recall 20.3.** *For any category $\mathcal{D}$, a terminal object is $d$ object such that all objects $\overline{d}$ there is exactly one $\overline{d} \to d$.*

**Lemma 20.4.** *Given $\mathcal{C}$ and objects $c, d$ make a new category whose objects are the cones over $c, d$.*

***Cone 1***

$$\begin{tikzcd} & e \arrow[dl, "\alpha"'] \arrow[dr, "\beta"] & \\ c & & d \end{tikzcd}$$

***Cone 2***

$$\begin{tikzcd} & e' \arrow[dl, "\alpha'"'] \arrow[dr, "\beta'"] & \\ c & & d \end{tikzcd}$$

*Drawing the arrow from cone 1 to cone 2 we have*

$$\begin{tikzcd} & e \arrow[dl, "\alpha"'] \arrow[d, dashed, "\phi"] \arrow[dr, "\beta"] & \\ c & e' \arrow[l, "\alpha'"] \arrow[r, "\beta'"'] & d \end{tikzcd}$$

*such that $\alpha \circ \phi = \alpha'$ and $\beta \circ \phi = \beta'$.*
*Being a product of $c, d$ is equivalent to being a terminal object in category of cones over $c, d$*

**Definition 20.5.** *A co-cone over $c, d$ is a diagram*

$$\begin{tikzcd} c \arrow[dr, "\alpha"'] & & d \arrow[dl, "\beta"] \\ & e & \end{tikzcd}$$

*is a coproduct of $c, d$ iff for all cocones*

$$\begin{tikzcd} c \arrow[dr, "\alpha'"'] & & d \arrow[dl, "\beta'"] \\ & e' & \end{tikzcd}$$

*there is a unique $\phi : e \to e'$ such that the diagram*

$$\begin{tikzcd} c \arrow[dr, "\alpha"] \arrow[ddr, "\alpha'"'] & & d \arrow[dl, "\beta"'] \arrow[ddl, "\beta'"] \\ & e \arrow[d, dashed, "\phi"] & \\ & e' & \end{tikzcd}$$

*commutes (i.e. $\alpha' \circ \phi = \alpha$ and $\beta' \circ \phi = \beta$).*

**Example 20.6.** *Sets: Let $X, Y, Z$ be sets then define $X \cup Y = (X \times \{0\}) \cup (Y \times \{1\})$. Define $i : x \mapsto (x, 0)$ and $j : y \mapsto (y, 1)$. Then the coproduct is $h(x, 0) = f(x)$ and $h(y, 1) = g(y)$.*

$$\begin{tikzcd} X \arrow[d, "i"] \arrow[dr, "g"] & Y \arrow[d, "f"] \arrow[dl, "j"'] \\ X \cup Y \arrow[r, dashed, "h"] & Z \end{tikzcd}$$

*such that $h(i(x)) = f(x)$ and $h(j(y)) = g(y)$.*

**Definition 20.7.** *Let $M_1, M_2$ be $R$-modules. Define*

$$M_1 \oplus M_2 = \{(m_1, m_2) : m_i \in M_i\}.$$

*Define $i : M_1 \to M_1 \oplus M_2$ by $i(m_1) = (m_1, 0)$ and $j : M_2 \to M_1 \oplus M_2$ by $j(m_2) = (0, m_2)$. For any $R$-module $N$ and homomorphisms $\psi_1 : M_1 \to N$, $\psi_2 : M_2 \to N$, there is a unique $\rho : M_1 \oplus M_2 \to N$ such that the diagram*

$$\begin{tikzcd}
M_1 \arrow[d, "i"] \arrow[dr, "\psi_1"] & M_2 \arrow[d, "j"] \arrow[dl, "\psi_2"'] \\
M_1 \oplus M_2 \arrow[r, dashed, "\rho"] & N
\end{tikzcd}$$

*commutes ($\rho \circ i = \psi_1$ and $\rho \circ j = \psi_2$). We have $\rho(m_1, m_2) = \rho((m_1, 0) + (0, m_2)) = \psi_1(m_1) + \psi_2(m_2)$.*

*For the coproducts, $\pi_1 : M_1 \oplus M_2 \to M_1$ by $\pi_1(m_1, m_2) = m_1$ and $\pi_2 : M_1 \oplus M_2 \to M_2$ by $\pi_2(m_1, m_2) = m_2$. For any $R$-module $N$ and homomorphisms $\psi_1 : N \to M_1$, $\psi_2 : N \to M_2$, there is a unique $\rho : N \to M_1 \oplus M_2$ such that the diagram*

$$\begin{tikzcd}
N \arrow[d, "\psi_1"] \arrow[dr, "\psi_2"'] & M_1 \arrow[d, "\pi_1"] \arrow[dl, "\pi_2"] \\
M_1 \oplus M_2 \arrow[r, dashed, "\rho"] & N
\end{tikzcd}$$

*commutes ($\rho \circ \pi_1 = \psi_1$ and $\rho \circ \pi_2 = \psi_2$). We have $\rho(n) = \rho(\psi_1(n), \psi_2(n)) = (\psi_1(n), 0) + (0, \psi_2(n))$.*

# 21 March 11

**Lemma 21.1.** *Fix $R$-module, $M$ and $N$ consider $Hom_R(M, N) = \{\phi : M \to N : \phi R - linear\}$. This forms an $R$-module.*

**Theorem 21.2.** *Given ring $R$ and set $X$, to construct a $R$-module on $X$. Let $M = \{g : X \to R, \{x \in X : g(x) \neq 0\}$ is finite$\}$. To make $M$ an $R$-module, $(g + h)(x) = g(x) + h(x)$ and $(rg)(x) = r(g(x))$.*
*We'll write elements of $M$ as $\lambda_1 x_1 + \cdots + \lambda_n x_n$ where $\lambda_i \in R$ and $x_i \in X$*

**Claim 21.3.** *For any $R$-module $N$, for any function $f : X \to N$, there is a unique $R$-linear $\psi : M \to N$ such that $\psi(1x) = f(x)$ for all $x \in X$.*

*Proof.*

$$\psi\left(\sum_{i=1}^{n} \lambda_i x_i\right) = \sum_{i=1}^{m} \lambda_i f(x_i).$$

□

**Theorem 21.4.** *Any module is isomorphic to a quotient of a free module.*

*Proof.* Let $N$ be arbitrary, let $X =$ underlying set of $N$, $f : X \to N$ where $f(n) = n$. $\psi : M \to N$ with $M = Free_R(X)$, $\psi$ is surjective. □

**Remark 21.5.** *When I constructed $Free_R(X)$, every element of $Free_R(X)$ is a unique $R$-linear combination of elements of elements from the set $\{1x : x \in X\}$.*

**Definition 21.6.** *$R$-module $M$ is free iff $M$ has a linearly independent generating set.*

**Lemma 21.7.** *If $B$ is such a set, $M \simeq Free_R(B)$*

**Definition 21.8.** *Let $M, N, P$ be $R$-modules. A function $\phi : M \times N \to P$ is $R$-bilinear if*

1. *For all $n$, $m \mapsto \phi(m, n)$ is $R$-linear*
2. *For all $m$, $n \mapsto \phi(m, n)$ is $R$-linear*

**Remark 21.9.** *Bilinear functions from $M \times N$ to $P$ are very different from linear functions from $M \oplus N$ to $P$.*

**Example 21.10.** *$R = \mathbb{R}$ and $M = N = \mathbb{R}'$:*

1. *$R' \times \mathbb{R}' \to \mathbb{R}' : (x, y) \mapsto xy$ is bilinear, not linear*
2. *$R' \oplus \mathbb{R}' = \mathbb{R}' : (x, y) \mapsto x + y$ is linear, not bilinear*

**Definition 21.11.** *Given $M, N$ $R$-modules we will construct an $R$-module $M \otimes_R N$ (tensor product of $M$ and $N$) together with a bilinear map $\otimes : M \times N \to M \otimes_R N$ by $(m, n) \mapsto m \otimes n$ such that for all $R$-module $P$ and all bilinear $\phi : M \times N \to P$ there is unique $R$-linear $\rho : M \otimes_R N \to P$ such that $\phi = \rho \circ \otimes$*

$$\begin{tikzcd} M \times N \arrow[r, "\otimes"] \arrow[dr, "\phi"'] & M \otimes N \arrow[d, dashed, "\rho"] \\ & P \end{tikzcd}$$

*commutes.*
*"$\otimes$ is initial in the category of bilinear maps with domain $M \times N$"*

**Theorem 21.12.** *$Hom_R(M, Hom_R(N, P)) \simeq$ bilinear maps from $M \times N \to P \simeq Hom_R(M \otimes N, P)$*

# 22 March 11

## 22.1 Goal and Preliminaries

**Remark 22.1** (Note on Bilinearity)**.** *Let $\phi : M \times N \to P$ be a bilinear map. If $\beta : P \to Q$ is a linear map, then the composition $\beta \circ \phi$ is bilinear.*

**Definition 22.2.** *(Universal Property of Tensor Products)*
***Goal:*** *Construct a module $M \otimes N$ and a bilinear map $\otimes : M \times N \to M \otimes N$ such that for any module $Q$ and any bilinear map $\psi : M \times N \to Q$, there exists a unique linear map $\beta : M \otimes N \to Q$ that makes the following diagram commute:*

$$\begin{tikzcd} M \times N \arrow[r, "\otimes"] \arrow[dr, "\psi"'] & M \otimes N \arrow[d, dashed, "\exists!\beta"] \\ & Q \end{tikzcd}$$

*In other words, $\psi = \beta \circ \otimes$.*

**Lemma 22.3** (Easy Fact on Quotients)**.** *Let $A \leq B$ be submodules. Then:*

1. *The natural projection $\phi_A : B \to B/A$ is a homomorphism, and it is $0$ on $A$.*
2. *For any module $C$ and homomorphism $\gamma : B \to C$ such that $\gamma(A) = 0$, there is a unique linear map $\delta : B/A \to C$ such that the diagram commutes:*

$$\begin{tikzcd} B \arrow[r, "\phi_A"] \arrow[dr, "\gamma"'] & B/A \arrow[d, dashed, "\exists!\delta"] \\ & C \end{tikzcd}$$

*Specifically, $\delta$ is defined by mapping $b + A \mapsto \gamma(b)$.*

## 22.2 Construction of the Tensor Product

**Definition 22.4** (Constructing $M \otimes N$ and $\otimes$)**.** *Let $X = M \times N$ be the Cartesian product. Construct the free $R$-module on $X$, denoted $Fr(X)$, whose elements are finite sums of the form:*

$$\sum \lambda_i(m_i, n_i) \quad \text{where } \lambda_i \in R.$$

*Consider the map $\sigma : M \times N \to Fr(X)$ defined by $(m, n) \mapsto 1(m, n)$.*
*For $\sigma$ to be bilinear, we would need equations like $(m_1 + m_2, n) - (m_1, n) - (m_2, n) = 0$ and $(\lambda m, n) - \lambda(m, n) = 0$ to hold. However, in $Fr(X)$, these are* not *true!*
*To fix this, let $B$ be the submodule of $Fr(X)$ generated by all elements of the form that "need to be $0$" to satisfy bilinearity. Define the tensor product as the quotient module:*

$$M \otimes N = Fr(X)/B$$

*and define the map $\otimes : M \times N \to M \otimes N$ by:*

$$\otimes : (m, n) \mapsto 1(m, n) + B = m \otimes n.$$

**Claim 22.5.** *The map $\otimes$ is bilinear.*

*Proof.* We check one of the bilinearity conditions (the others follow similarly).

$$\begin{aligned}\otimes(\lambda m, n) - \lambda \otimes (m, n) &= 1(\lambda m, n) + B - \lambda(m, n) + B \\ &= [1(\lambda m, n) - \lambda(m, n)] + B \\ &= 0 + B\end{aligned}$$

This evaluates to $0 + B$ because the term in the brackets is one of the generators of the submodule $B$, meaning it lives inside $B$. The same logic applies for the other three bilinearity conditions. □

**Claim 22.6** (Fulfilling the Goal)**.** *The constructed $M \otimes N$ satisfies the universal property. Let $\psi : M \times N \to Q$ be bilinear. There exists a unique linear $\beta : M \otimes N \to Q$.*

*Proof.* Let $\phi : Fr(X) \to Q$ be the $R$-module homomorphism defined on the basis by the unique $R$-linear extension:

$$\phi\left(\sum \lambda_i(m_i, n_i)\right) = \sum \lambda_i \psi(m_i, n_i) \implies \phi(1(m, n)) = \psi(m, n).$$

We claim $\phi(b) = 0$ for all $b \in B$. Since $B$ is generated by the bilinearity relations, we evaluate $\phi$ on those generators. For instance:

$$\phi((\lambda m, n) - \lambda(m, n)) = \psi(\lambda m, n) - \lambda\psi(m, n) = 0$$

because $\psi$ is explicitly given as bilinear. Since $\phi$ maps all 4 types of generators of $B$ to 0, $\phi(B) = 0$.
By our "Easy Fact", there exists a unique linear map $\beta : \frac{Fr(X)}{B} \to Q$ such that the diagram commutes:

$$\begin{tikzcd}
M \times N \arrow[r, "\sigma"] \arrow[dr, "\otimes"'] \arrow[ddr, bend right, "\psi"'] & Fr(X) \arrow[d, "\text{quotient}"] \arrow[dd, bend left, "\phi"] \\
& M \otimes N \arrow[d, dashed, "\beta"] \\
& Q
\end{tikzcd}$$

For elements in $M \otimes N$, we have $m \otimes n = (m, n) + B \implies \beta(m \otimes n) = \psi(m, n)$. □

**Remark 22.7** (Uniqueness of $M \otimes N$)**.** *Suppose $\otimes' : M \times N \to M \otimes' N$ is another bilinear map satisfying the universal property. Then, any bilinear map is of the form $\delta \circ \otimes'$ for a linear map $\delta$ on $M \otimes' N$. By universal properties, there is a unique isomorphism $I : M \otimes N \to M \otimes' N$ such that $I(m \otimes n) = m \otimes' n$ for all $m, n$.*

## 22.3 Tensor Product with the Base Ring

**Theorem 22.8.** *Let $M$ be an $R$-module. Then $R \otimes_R M \cong M$.*

*Proof.* Consider the scalar multiplication map $t : R \times M \to M$ defined by $(r, m) \mapsto rm$. For any bilinear $\psi : R \times M \to Q$, we want to show there exists a unique linear map $\beta : M \to Q$ such that $\psi = \beta \circ t$.

$$\begin{tikzcd} R \times M \arrow[r, "t"] \arrow[dr, "\psi"'] & M \arrow[d, dashed, "\exists!\beta"] \\ & Q \end{tikzcd}$$

If $\beta$ exists, we must have:

$$\beta(m) = \beta(1 \cdot m) = \beta(t(1, m)) = \psi(1, m).$$

This defines $\beta$ uniquely. We verify two claims:

1. **Claim 1:** $\beta$ is linear because $\psi$ is bilinear.
2. **Claim 2:** $\beta \circ t = \psi$.
   *Proof:* $\psi(r, m) = r\psi(1, m) = \psi(1, rm) = \beta(rm) = \beta(t(r, m))$.

By the uniqueness property of the tensor product, we conclude:

1. $R \otimes M \cong M$.
2. There exists a unique isomorphism $I : R \otimes M \to M$ mapping $r \otimes m \mapsto rm$.

$\square$

# 23 March 16

**Definition 23.1.** *An $R$-module $M$ is cyclic iff there exist $m$ such that $M = Rm$*

**Definition 23.2.** *$M$ is an $R$-module, $m \in M$, the annihilator of $m$ is $Ann(m) = \{r : rm = 0\}$*

**Lemma 23.3.** *For any $M$ $R$-module any $m \in M$, have $R$-module homomorhpism $\phi : R \to M$, $\phi : r \mapsto rm$.. Then we have $Im(\phi) = Rm$ and $\ker(\phi) = Ann(m)$ so*

$$Rm \simeq \frac{R}{Ann(m)}.$$

**Remark 23.4.** *If $M$ cyclic, $M = Rm$, $M \simeq \frac{R}{Ann(m)}$*

**Remark 23.5.** *If we have $I$ ideal of $R$ and view $R$-module, then*

$$R/I = R(1 + I) \text{ and } Ann(1 + I) = I.$$

**Remark 23.6.** *Let $R$ be PID, cyclic $R$-module $\simeq R/(c)$ for $c \in R$.*

**Definition 23.7.** *Let $M$ be $R$-module, $P, Q$ submodules of $M$ are complementary if $M = P + Q$ and $P \cap Q = 0$*

**Definition 23.8.** *If $P$, a submodule of $M$, $P$ is a direct summand of $M$ if there is a complementary $Q$.*

**Example 23.9.** *The following are examples of direct summand*

1. $R = \mathbb{R}, M = \mathbb{R}^2$
2. $R = \mathbb{Z}, M = \mathbb{Z}/4\mathbb{Z}$, $P = \{0, 2\}$ *then $P$ has no complement*
3. $M_i$ *$R$-module,* $i = 1, ..., n$, $M_1 \oplus \cdots \oplus M_n = \{(m_1, ..., m_n) : m_i \in M_i\}$ *with obvious operations*
4. *If $P_1, ..., P_n$ submodules of $N$ and $N = P_1 + P_2 + \cdots + P_n$ and $\sum_{i=1}^n p_i = 0$ ($p_i \in P_i$) then all $p_i = 0, N \simeq P_1 \oplus \cdots \oplus P_n$*

**Lemma 23.10.** *A free module on a finite set $X = \{x_1, ..., x_n\}$ with $|X| = n$ then $Fr_R(X) \simeq R^n$ by*

$$\sum_{i=1}^n \lambda_i x_i \mapsto (\lambda_1, ..., \lambda_n).$$

**Definition 23.11.** *Let $R$ be ID, $M$ $R$-module, $m \in M$ is a torsion element if there is $r \neq 0$ such that $rm = 0$.*

**Definition 23.12.** *$R$ ID, $M$ is $R$-module, $M$ is torsion free if $0$ is the only torsion element.*

**Definition 23.13.** *If $M$ is torsion free $R$-module, $R$ ID, then $m_1, ..., m_n$ are linearly independent if they are distinct and*

$$\sum_{i=1}^{n} r_i m_i = 0.$$

**Definition 23.14.** *$R$ ID, $M$ torsion free $R$-module. $M$ has rank $n$ iff $n = \max\{|I| : I \subseteq M, I$ independent$\}$.*

**Theorem 23.15.** *Let $R$ be PID and let $M$ be a finitely generated $R$-module. Then $M$ is isomorphic to*

$$\frac{R}{(c_1)} \oplus \cdots \oplus \frac{R}{(c_n)}.$$

*for $c_i$'s are nonzero nonunit elements of $R$ and $c_i$ divides $c_{i+1}$ Moreover:*

1. *$t$ unique*
2. *$n$ unique*
3. *ideals $(c_i)$ are unique equivalently, $c_i$'s are unique up to associates.*

**Lemma 23.16.** *If $R$ ID, $R^n$ has rank $n$.*

*Proof.* Let $e_i = (0, ..., 1, ..., 0)$ where 1 is in the $i$-th position. We have $e_1, ..., e_n$ are independent. Let $I \subseteq R^n$ be independent. Let $k$ be the field of fraction of ID $R$. $I \subseteq R^n \subseteq k^n$.
Claim: $I$ is an independent subset of the $k$-vs $k^n$. Prove this by clearing fraction.
Then by linear algebra $|I| \leq \dim_k(k^n) = n$.
The result will follow from the following theorem

**Theorem 23.17.** *If $R$ PID, $N$ free of rank $n$ and $M \leq N$ then $M$ is free of some rank $m \leq n$.*

There is a basis $z_1, ..., z_n$ of $N$ and there are nonzero nonunit $c_1, ..., c_m$ such that $c_i$ divides $c_{i+1}$ and $c_1 z_1, ..., c_m z_m$ forms a basis of $M$. □

# 24 March 18

**Theorem 24.1.** *Let $R$ be a principal ideal domain (PID) and $M$ be a finitely generated $R$-module. Then*

$$M \simeq \frac{R}{(c_1)} \oplus \cdots \oplus \frac{R}{(c_n)} \oplus R^t$$

*where $t \geq 0$, $c_i \neq 0$, $c_i \mid c_{i+1}$ for all $1 \leq i < n$, and the elements $c_i$ are non-units.*

**Lemma 24.2.** *Let $R$ be a ring with $1 \neq 0$, and let $R^n$ be a free $R$-module of rank $n$ with standard basis $e_i = (0, \ldots, 1, \ldots, 0)$. Then $Hom_R(R^n, R) \simeq R^n$, and it is generated by the dual basis $e_j^*$, defined by*

$$e_j^* : (\lambda_1, \ldots, \lambda_n) \mapsto \lambda_j.$$

**Lemma 24.3.** *Let $R$ be a PID, $N$ be a free $R$-module of rank $n$, and $M \leq N$ be a submodule. Then:*

1. *$M$ is a free $R$-module of rank $m$, where $m \leq n$.*
2. *There exists a basis $z_1, \ldots, z_n$ of $N$ and non-zero elements $d_1, \ldots, d_m \in R$ such that $d_i \mid d_{i+1}$ for all $1 \leq i < m$, and $d_1 z_1, \ldots, d_m z_m$ forms a basis for $M$.*

*Proof.* **Step 1: Construction of direct summands.**
We aim to find an element $z \in N$ ($z \neq 0$) and $d \neq 0$ in $R$ such that $dz \in M$, $Rz$ is a direct summand of $N$, and $R(dz)$ is a direct summand of $M$.

First, we claim that if $M \neq 0$, there exists $\psi \in \mathrm{Hom}_R(N, R)$ such that $\psi(M) \neq 0$. To show this, without loss of generality, assume $N = R^n$. Let $m = (a_1, \ldots, a_n) \in M$ be a non-zero element. Then $a_j \neq 0$ for some $j$. Using the dual basis functional, $e_j^*(m) = a_j \neq 0$, which implies $e_j^*(M) \neq 0$.

For any $\phi \in \mathrm{Hom}_R(N, R)$, the image $\phi(M)$ is an ideal of the PID $R$. Consider the set of ideals:

$$\Sigma = \{\phi(M) : \phi \in \mathrm{Hom}_R(N, R)\}.$$

Since $M \neq 0$, $\Sigma$ is a non-empty set containing at least one non-zero ideal. Because $R$ is a PID, it is Noetherian, so $\Sigma$ possesses a maximal element with respect to inclusion. Choose $\psi \in \mathrm{Hom}_R(N, R)$ such that $\psi(M) = (d)$ is maximal in $\Sigma$. Choose $y \in M$ such that $\psi(y) = d$.

We claim that $d \mid \phi(y)$ for all $\phi \in \mathrm{Hom}_R(N, R)$. To show this, let $d' = \phi(y)$ and let $e = \gcd(d, d')$. Since $R$ is a PID, we can write $e = \lambda d + \mu d'$ for some $\lambda, \mu \in R$. Then:

$$e = \lambda\psi(y) + \mu\phi(y) = (\lambda\psi + \mu\phi)(y) \in (\lambda\psi + \mu\phi)(M).$$

This implies that:

$$(d) \subseteq (e) \subseteq (\lambda\psi + \mu\phi)(M).$$

By the maximality of $(d) = \psi(M)$ in $\Sigma$, we must have $(d) = (e)$. Thus, $d$ and $e$ are associates, which means $d \mid d'$, and consequently $d \mid \phi(y)$.

Now, let $w_1, \ldots, w_n$ be a basis of $N$, and let $w_j^* \in \mathrm{Hom}_R(N, R)$ be the corresponding dual basis functionals. We can express $y = \sum_{j=1}^n \mu_j w_j$, meaning $w_j^*(y) = \mu_j$. By our previous

claim, $d \mid w_j^*(y)$, so $d \mid \mu_j$ for all $j$. Let:

$$z = \sum_{j=1}^{n} \left(\frac{\mu_j}{d}\right) w_j \in N.$$

Notice that $dz = y$. Furthermore, since $\psi(y) = d$, we have $\psi(dz) = d$. By linearity, $d\psi(z) = d$. Since $R$ is an integral domain and $d \neq 0$, we conclude that $\psi(z) = 1$.

We now claim:

(a) $N = \ker(\psi) \oplus Rz$

(b) $M = (\ker(\psi) \cap M) \oplus Ry$

To show (a), note that if $x \in \ker(\psi) \cap Rz$, then $x = rz$ for some $r \in R$, and $0 = \psi(x) = r\psi(z) = r(1) = r$, so $x = 0$. For any $n \in N$, let $\lambda = \psi(n)$. We can write $n = (n - \lambda z) + \lambda z$. Observe that $\psi(n - \lambda z) = \lambda - \lambda\psi(z) = 0$, so $(n - \lambda z) \in \ker(\psi)$ and $\lambda z \in Rz$. Thus $N = \ker(\psi) \oplus Rz$. The proof of (b) is identical, substituting $M$ for $N$ and $y$ for $z$.

**Step 2: Proof of Part (1).**
We proceed by induction on the rank of $M$ (with $N$ fixed). If the rank of $M$ is 0, $M = 0$, which is trivially free. Assume the result holds for submodules of smaller rank. Using our construction, $M = Ry \oplus (\ker(\psi) \cap M)$. Because $Ry$ is generated by a single non-torsion element, it has rank 1. Thus, the rank of $\ker(\psi) \cap M$ is strictly less than the rank of $M$. By the induction hypothesis, $\ker(\psi) \cap M$ is a free $R$-module. Adding $y$ to a basis of $\ker(\psi) \cap M$ yields a free basis for $M$. Thus $M$ is free of rank $m \leq n$.

**Step 3: Proof of Part (2).**
We proceed by induction on the rank $n$ of $N$, proving the statement for all submodules $M \leq N$ simultaneously. If $n = 0$, it is trivial. Suppose it holds for free modules of rank $n - 1$. From Step 1, $N = \ker(\psi) \oplus Rz$. Since $N$ has rank $n$ and $Rz$ has rank 1, $\ker(\psi)$ is a free $R$-module of rank $n - 1$. By the induction hypothesis applied to the submodule $\ker(\psi) \cap M \leq \ker(\psi)$, there exists a basis $z_2, \ldots, z_n$ of $\ker(\psi)$ and elements $d_2, \ldots, d_m \in R$ with $d_2 \mid d_3 \mid \cdots \mid d_m$ such that $d_2 z_2, \ldots, d_m z_m$ is a basis for $\ker(\psi) \cap M$.

Let $z_1 = z$ and $d_1 = d$. Then $z_1, z_2, \ldots, z_n$ is a basis for $N$, and $d_1 z_1, d_2 z_2, \ldots, d_m z_m$ is a basis for $M$.

It remains only to show that $d_1 \mid d_2$. Define a homomorphism $f : N \to R$ by mapping $z_1 \mapsto 1$, $z_2 \mapsto 1$, and $z_i \mapsto 0$ for $i > 2$. We evaluate $f(M)$: $f(d_1 z_1) = d_1$ and $f(d_2 z_2) = d_2$, so the ideal $f(M)$ contains both $d_1$ and $d_2$, meaning $f(M) = (d_1, d_2)$. Because $(d_1) = \psi(M)$ is maximal among such ideals, and $(d_1) \subseteq (d_1, d_2) = f(M)$, we must have $(d_1) = (d_1, d_2)$. This implies $d_2 \in (d_1)$, so $d_1 \mid d_2$. This completes the proof. $\square$

# 25 March 20

**Theorem 25.1.** *Let $F$ be a finitely genereated module over PID $R$ say $F = Rf_1 + \cdots + Rf_t$, $f_i \in F$. Define $\pi : R^t \to F$ by $\pi : (\lambda_1, ..., \lambda_t) \mapsto \sum_{i=1}^t \lambda_i t_i$. $\pi$ is a surjective homomorphism so*

$$F \simeq \frac{N}{\ker(\pi)}.$$

*We have $N, M$ having rank $t$ and $s$.*

**Theorem 25.2.** *Let $R$ be a PID, $M$ be a finitely generated $R$-module. Then $M$ is isomorphic to*

$$\frac{R}{(c_1)} \oplus \cdots \oplus \frac{R}{(c_n)} \oplus R^t$$

*where $t \geq 0$, $c_i \neq 0$, $c_i \mid c_{i+1}$ for all $1 \leq i < n$, and the elements $c_i$ are non-units.*

**Lemma 25.3.** *By previous result, we get a basis $n_1, ..., n_t$ for $N$ and nonzero constants $r_1, ..., r_s \in R$ such that $r_i | r_{i+1}$ for $1 \leq i < s$ and $r_1 n_1, ..., r_s n_s$ is a basis for $M$.*

**Exercise 25.4.**

$$\frac{N}{M} \simeq \frac{R}{(r_1)} \oplus \cdots \oplus \frac{R}{(r_s)} \oplus R^{t-s}.$$

**Lemma 25.5.** *If $r$ nonzero nonunit, $r = u\pi_1^{k_1} \cdots \pi_n^{k_n}$ for $u$ unit, $\pi$ non associate and irreducibles.*

**Theorem 25.6** (Chinese Remainder Theorem)**.** *If $(r) = (\pi_1)^{k_1} \cdots (\pi_n)^{k_n}$ then*

$$\frac{R}{(r)} \simeq \frac{R}{(\pi_1^{k_1})} \oplus \cdots \oplus \frac{R}{(\pi_n^{k_n})}.$$

## 25.1 Adjoint Functors

**Definition 25.7.** *Let $\mathcal{U}$ be "forgetful" functor from group to set.*

$$\mathcal{U}(G) = \text{underlying set of } G.$$

*If $\phi : G \to H$, $\mathcal{U}(\phi)$ is $\phi$ viewed as a function.*

**Lemma 25.8.** *We have for any group $G$ and any set $X$, there is a bijection between $Hom_{Gr}(Fr(X), G)$ and $Hom_{Set}(X, \mathcal{U}(G))$*

**Definition 25.9** (Natural Transformation)**.** *Let $F_1, F_2 : \mathcal{C} \to \mathcal{D}$. A natural transformation from $F_1 \to F_2$ is a family $v = (v_c)$, for $c$ object of $\mathcal{C}$, $F_1 c \xrightarrow{v_c} F_2 c$ in $\mathcal{D}$ such that for all $c \xrightarrow{\alpha} c'$*

$$\begin{tikzcd} F_1 c \arrow[r, "v_c"] \arrow[d, "F_1\alpha"'] & F_2 c \arrow[d, "F_2\alpha"] \\ F_1 c' \arrow[r, "v_{c'}"'] & F_2 c' \end{tikzcd}$$

*commutes.*

**Definition 25.10** (Opposite Category)**.** *$\mathcal{C}$ category then $\mathcal{C}^{op}$ is the category with the same objects but with the arrows reversed.*

# 26 Mar 23

**Remark 26.1.** *If $Fc \xrightarrow{\alpha} d$ (in $\mathcal{D}$) write $\alpha^*$ for $v_{cd}(\alpha)$ and if $c \xrightarrow{\beta} Gd$ (in $\mathcal{C}$) write $\beta_*$ for $v_{cd}^{-1}(\beta)$.*

**Definition 26.2.** *Let $F : \mathcal{C} \to \mathcal{D}$ and $G : \mathcal{D} \to \mathcal{C}$ with $\mathcal{C}, \mathcal{D}$ categories and $F, G$ functors. An adjunction between $F$ and $G$ is a family of bijections $v_{cd} : Hom_{\mathcal{D}}(Fc, d) \simeq Hom_{\mathcal{C}}(c, Gd)$ for objects $c$ of $\mathcal{C}, d$ of $\mathcal{D}$ such that*

1. *Naturality in $c$: If $y = \delta(F\alpha)$ commutes then $y^* = \delta^*\alpha$ commutes.*

$$\begin{tikzcd} Fc_1 \arrow[r, "F\alpha"] \arrow[dr, "y"'] & Fc_2 \arrow[d, "\delta"] \\ & d \end{tikzcd} \implies \begin{tikzcd} c_1 \arrow[r, "\alpha"] \arrow[dr, "y^*"'] & c_2 \arrow[d, "\delta^*"] \\ & Gd \end{tikzcd}$$

2. *Naturality in $d$: Let $d_1 \xrightarrow{\beta} d_2$*

**Recall 26.3.** *Universal Property of Products:*

**Theorem 26.4.** *If $G$ is a right adjoint of some $F$ then $G$ preserves products. If $d \xrightarrow{\pi_1} d_1$ and $d \xrightarrow{\pi_2} d_2$ exhibits $d = d_1 \times d_2$ in $\mathcal{D}$, then*

$$\begin{tikzcd} & d \arrow[dl, "\pi_1"'] \arrow[dr, "\pi_2"] & \\ d_1 & & d_2 \end{tikzcd} \implies \begin{tikzcd} & Gd \arrow[dl, "G\pi_1"'] \arrow[dr, "G\pi_2"] & \\ Gd_1 & & Gd_2 \end{tikzcd}$$

*and $Gd$ is a product of $Gd_1$ and $Gd_2$ in $\mathcal{C}$ (so $Gd \simeq Gd_1 \times Gd_2$).*

**Remark 26.5.** *Let $R$ be a ring, $M, N, P$ be $R$-modules. We can view $\phi : M \times N \to P$ bilinear as an element of $Hom(M, Hom(N, P))$. This is from $\phi(m, -)$ is a linear map from $N$ to $P$ and $m \mapsto \phi(m, -)$ is a linear map from $M$ to $Hom(N, P)$.*

**Lemma 26.6.** *$Hom(M \otimes N, P) \simeq Hom(M, Hom(N, P)) \simeq$ bilinear maps from $M \times N \to P$*

**Lemma 26.7.** *Fixing $N$, this gives adjunction between $- \otimes N$ and $Hom(N, -)$*

# 27 Mar 25

**Recall 27.1.** *If $R$ PID and $M$ finitely generated $R$-module then*

$$M \simeq \frac{R}{(c_1)} \oplus \cdots \oplus \frac{R}{(c_n)} \oplus R^t.$$

*where $c_i$ nonzero nonunit and $c_i | c_{i+1}$.*

**Corollary 27.2.** *If $R$ PID and $M$ finitely generated torsion-free then $M$ is free.*

**Recall 27.3.** *An $R$-module $M$ is noetherian if it satisfies any of the equivalent condition*

1. *Every $N \leq M$ is finitely generated*
2. *Every chain $N_0 \subseteq N_1 \subseteq \cdots$ of submodules is eventually constant.*
3. *Any nonempty set of submodules of $M$ has a maximal element*

**Lemma 27.4.** *If $M$ noetherian and $N \leq M$ then*

1. *$N$ noetherian*
2. *$M/N$ noetherian (submodules of $M/N \simeq$ submodules $M$ that contain $N$)*

**Lemma 27.5.** *If $N \leq M$ and both $N$ and $M/N$ are noetherian then $M$ noetherian.*

*Proof.* Let $(K_i)_i$ with $K_0 \subseteq K_1 \subseteq \cdots$ and $K_i$ submodules of $M$. Then $(K_i \cap N)_i$ stabilizes. Consider $(\phi_N[K_i])_i$ with $\phi_N : M \to M/N$ stablizes. Let $j$ be so large that $i \geq j$, $K_i \cap N = K_j \cap N$ and $\frac{K_i+N}{N} = \frac{K_j \cap N}{N}$.
Let $i \geq j$, we've already shown $K_j \subseteq K_i$ so we just have to show that $K_i \subseteq K_j$. Let $k \in K_i$ then $k + N \in \frac{K_i+N}{N} = \frac{K_j+N}{N}$. So there is $k' \in K_j$ such that $k + N = k' + N$ and so $k - k' \in N$. As $k, k' \in K_i$ we also have $k - k' \in N \cap K_i = N \cap K_j$. □

**Lemma 27.6.** *If $M_1, M_2$ are noetherian $R$-modules then $M_1 \oplus M_2$ is noetherian.*

*Proof.* If $M_1 \simeq M_1' = M_1 \otimes 0$ then $\frac{M_1 \oplus M_2}{M_1'} \simeq M_2$ and we're done by the second lemma. □

**Lemma 27.7.** *A finite direct sum of noetherian $R$-modules is noetherian.*

*Proof.* By induction. □

**Lemma 27.8.** *If $R$ is noetherian ring then for all $t$, $R^t$ is a noetherian $R$-module.*

*Proof.* By the previous lemma □

**Lemma 27.9.** *If $S \subseteq R$ MC and $R$ noetherian ring then $S^{-1}R$ is a noetherian ring.*

**Lemma 27.10.** *If $R$ is noetherian then all finitely generated $R$-modules are noetherian.*

*Proof.* Let $M$ be $Rm_1 + \cdots + Rm_t$. Let $\phi : R^t \to M$ by $\phi(\lambda_1, ..., \lambda_t) = \sum_{i=1}^t \lambda_i m_i$. We have $\phi$ is surjective so $M \simeq \frac{R^t}{\ker(\phi)}$. By previous theorem, $M$ is noetherian. □

**Definition 27.11.** *An $R$-module $M$ is faithful if for all $r \in R$, if $rm = 0$ for all $m \in M$ then $r = 0$*

**Remark 27.12.** *If $R \leq S$ then*
1. *$S$ can be viewed as an $R$-module*
2. *$S$ is faithful as $rs = 0$ for all $s \in S$ implies $r_1 = 0$ so $r = 0$*

**Lemma 27.13.** *Suppose $R \leq S$ and $a \in S$, the least subring of $S \supseteq R \cup \{a\} = \{f(a) : f \in R[x]\}$ (write $R[a]$ for this)*

**Theorem 27.14.** *If $\phi$ HM and $im(\phi) = R[a]$ so*

$$R[a] \simeq \frac{R[x]}{(\ker(\phi))}.$$

**Definition 27.15.** *Let $R \leq S$, let $a \in S$ with $a$ integral over $R$ if there is monic $f \in R[x], f(a) = 0$.*

**Lemma 27.16.** *The following are equivalent where $R \leq S, s \in S$*
1. *$s$ integral over $R$*
2. *$R[S]$ is finitely generated as $R$-module.*
3. *There is a faithful $R[S]$-module which is finitely generated as an $R$-module.*

# 28 Mar 27

**Recall 28.1.** *for*