

t-SNE Problem Boot Camp

Yi Gu

December 1, 2024

1 Setup

This document is dedicated to the reconstruction of [1] with low-dimension kernel g replaced by a more general kernel. This would serve as a great way to understand the material and help with the completion of the original project. For the sake of simplicity, we will use the exact same notation as in [1] and skip all the definitions. Recall in [1], g is defined as

$$g(y, y') = \frac{1}{1 + \|y - y'\|^2} \quad (1.1)$$

With all these being said, we will first start by giving our general low dimension kernel g some general assumptions, and modify it as we move on. We first require that $g(y, y') = k(\|y - y'\|)$ such that:

1. $k : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is decreasing and bounded.
2. k satisfies

$$\iint_{(0, \infty)^2} k(x, x') dx dx' < \infty. \quad (1.2)$$

We will kick off this warmup by listing all the important lemmas and results in order, and prove our own versions when the need arises. The final goal is to prove a same set of results as in [1] with the y kernel replaced. Recall that

$$\begin{aligned} F_{\rho, \mu}(x, \sigma) = & - \int \frac{\|x - x'\|^2}{2\sigma^2} \frac{\exp(-\|x - x'\|^2 / 2\sigma^2)}{\int \exp(-\|x - x'\|^2 / 2\sigma^2) d\mu(x')} d\mu(x') \\ & - \log \left(\int \exp(-\|x - x'\|^2 / 2\sigma^2) \right) + \log \rho. \end{aligned} \quad (1.3)$$

Proposition 1. *If μ has a C^1 density f such that both f and $\|\nabla f\|$ are bounded, then the equation $F_{\rho, \mu}(x, \sigma) = 0$ has a unique solution $\sigma_{\rho, \mu}^*(x)$ for all $\rho \in (0, 1)$ and $x \in \mathbb{R}^d$. Moreover, $x \mapsto \sigma_{\rho, \mu}^*(x)$ is smooth.*

Proof. This proposition is about the kernel of x . The proof stays unchanged from [1]. □

The next couple of lemmas are general results in section 3 of [1]. We will omit the proof because they are not directly discussing the quantities in the t-sne algorithm. Recall that X or the corresponding probability measure μ is called sub-Gaussian if $\mathbb{P}(X > t) \leq 2 \exp(-Ct^2)$ for some positive constant C .

Lemma 1. Let ν be sub-Gaussian with second moment equal to 1 and suppose $X \in \mathbb{R}^d$ has i.i.d. entries drawn from ν . Then there exists constant C, c_1 such that

$$\mathbb{P}(\|X\| \geq \sqrt{d} + t) \leq Ce^{-c_1 t^2}. \quad (1.4)$$

Lemma 2. Let μ be sub-Gaussian and c_1 be the constant from lemma 1. Let $X_1, \dots, X_n \in \mathbb{R}^d$ have i.i.d. entries drawn from μ . Then there exists a constant C such that

$$\mathbb{P}\left(\max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\frac{\log n}{c_1}} + \sqrt{d} + t\right) \leq Ce^{-c_1 t^2}. \quad (1.5)$$

Lemma 3. Let $\delta > 0$ be constant, $K_n = [\delta, \sqrt{n}]$ and $h : \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$. Suppose the following hold:

1. μ is sub-Gaussian;
2. $h(x, x', \sigma)$ and $\nabla_x h(x, x', \sigma)$ are bounded for all x, x' and all $\delta \in K_n$;
3. $h(x, x', \sigma) \rightarrow 0$ as $\|x - x'\|/\sigma \rightarrow \infty$.

Let $X_1, \dots, X_n$ be i.i.d. random variables drawn from μ . Let $\mu_n = (1/n) \sum_i \delta_{X_i}$. Then for any $\epsilon > 0$ with probability at least $1 - e^{-c\epsilon^2 n}$ we have

$$\sup_{\sigma \in K_n, x} \left| \int h(x, x', \sigma) d\mu_n(x) - \int h(x, x', \sigma) d\mu(x') \right| \leq \epsilon + O\left(\sqrt{\frac{\log n}{n}}\right). \quad (1.6)$$

Define

$$f_\sigma(x, x') = \exp(-\|x - x'\|^2 / 2\sigma^2), \quad g_\sigma = \frac{\|x - x'\|^2}{2\sigma^2} \exp(-\|x - x'\|^2 / 2\sigma^2). \quad (1.7)$$

The next two lemmas use similar techniques to prove.

Lemma 4. Let c_1 be the decay constant in lemma 1 and let $\delta > 0$. Then almost surely in $X_1, \dots, X_n$,

$$\sup_{\substack{\sigma \in [\delta, \sqrt{n}] \\ i=1, \dots, n}} \left| \frac{\int g_\sigma(X_i, x') d\mu_n(x')}{\int f_\sigma(X_i, x') d\mu_n(x')} - \frac{\int g_\sigma(X_i, x') d\mu(x')}{\int f_\sigma(X_i, x') d\mu(x')} \right| = O(n^{8/c_1 \delta^2 - 1/2} \sqrt{\log n}). \quad (1.8)$$

Lemma 5. Let c_1 be the decay constant in lemma 1 and let $\delta > 0$ be such that $c_1 \delta^2 > 8$. Then almost surely in $X_1, \dots, X_n$,

$$\sup_{\substack{\sigma \in [\delta, \sqrt{n}] \\ i=1, \dots, n}} \left| \log \left(\int f_\sigma(X_i, x') d\mu_n(x') \right) - \log \left(\int f_\sigma(X_i, x') d\mu(x') \right) \right| = O(n^{4/c_1 \delta^2 - 1/2} \sqrt{\log n}) \quad (1.9)$$

For this pair of lemmas, [1] provides the proof to the first lemma. We will provide the second for the sake of exercise.

Proof of lemma 5. First, notice that for arbitrary $x > 0$, the following elementary inequality holds:

$$\log(x) \leq x - 1. \quad (1.10)$$

Therefore, the lemma can be proved if we are able to show that

$$\begin{aligned} \left| \frac{\int f_\sigma(X_i, x') d\mu_n(x')}{\int f_\sigma(X_i, x') d\mu(x')} - 1 \right| &= \left| \frac{\int f_\sigma(X_i, x') d\mu_n(x') - \int f_\sigma(X_i, x') d\mu(x')}{\int f_\sigma(X_i, x') d\mu(x')} \right| \\ &= O(n^{4/c_1 \delta^2 - 1/2} \sqrt{\log n}). \end{aligned} \quad (1.11)$$

We can achieve that by bounding the numerator and the denominator of the previous formula. On one hand, consider set

$$E'_1 = \left\{ \max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\alpha + \frac{1}{c_1}} \sqrt{\log n} + \sqrt{d} \right\} \quad (1.12)$$

Setting $t = \sqrt{\alpha \log n}$ in lemma 2, we have $\mathbb{P}(E'_1) \leq Cn^{-c_1\alpha}$. To absorb the $\sqrt{d}$ term, we can find $\epsilon_n \rightarrow 0$ such that

$$E_1 = \left\{ \max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\alpha + \frac{1}{c_1} + \epsilon_n} \sqrt{\log n} \right\} \quad (1.13)$$

with $\mathbb{P}(E_1) \leq Cn^{-c_1\alpha}$. Because the maximum of norm of each X_i is restricted, on E_1^c we have for all i, j , that

$$\|X_i - X_j\|^2 \leq 4 \left(\alpha + \frac{1}{c_1} + \epsilon_n \right) \log n =: 4\gamma \log n. \quad (1.14)$$

Then for $\sigma \in [\delta, \sqrt{n}]$, on E_1^c for all i we have

$$\int f_\sigma(X_i, x') d\mu(x') \geq (1 - Cn^{-c_1\alpha}) \exp\left(-\frac{2\gamma \log n}{\delta^2}\right) \geq \frac{1}{2} \exp\left(-\frac{2\gamma \log n}{\delta^2}\right) = \frac{1}{2} n^{-2\gamma/\delta^2}. \quad (1.15)$$

This helps us bound the denominator. On the other hand, by lemma 3, we are able to find set E_2 with $\mathbb{P}(E_2) \leq e^{-c\epsilon^2 n}$, such that, on E_2^c ,

$$\sup_{\substack{\sigma \in [\delta, \sqrt{n}] \\ i=1, \dots, n}} \left| \int f_\sigma(X_i, x') d\mu_n(x') - \int f_\sigma(X_i, x') d\mu(x') \right| \leq \epsilon + O\left(\frac{\sqrt{\log n}}{\sqrt{n}}\right). \quad (1.16)$$

Therefore, we can combine (1.15) and (1.16) and conclude that on $E_1^c \cap E_2^c$,

$$\left| \frac{\int f_\sigma(X_i, x') d\mu_n(x') - \int f_\sigma(X_i, x') d\mu(x')}{\int f_\sigma(X_i, x') d\mu(x')} \right| \leq C\epsilon n^{2\gamma/\delta^2} + O(\sqrt{\log n} \cdot n^{-1/2+2\gamma/\delta^2}). \quad (1.17)$$

In order for the RHS to converge to zero, we need

$$\epsilon n^{2\gamma/\delta^2} \rightarrow 0, \quad 1/2 - 2\gamma/\delta^2 > 0. \quad (1.18)$$

Moreover, we have $\mathbb{P}(E_1 \cup E_2) \leq Cn^{-c_1\alpha} + e^{-c\epsilon^2 n}$. Set $\epsilon = n^{-\eta}$ for some $\eta > 0$. In order to get almost sure convergence using Borel-Cantelli, we also need

$$c_1\alpha > 1, \quad 1 - 2\eta > 0. \quad (1.19)$$

Hence for the entire lemma to work we need the following four inequalities to hold:

$$\frac{2\gamma}{\delta^2} - \eta < 0, \quad \frac{2\gamma}{\delta^2} - 1/2 < 0, \quad c_1\alpha > 1, \quad 1 - 2\eta > 0. \quad (1.20)$$

The last inequality implies that $\eta < 1/2$ so the first inequality implies the second. Now, recall that

$$\gamma := \alpha + \frac{1}{c_1} + \epsilon_n. \quad (1.21)$$

The collection of inequalities is equivalent to

$$2 \left(\alpha + \frac{1}{c_1} + \epsilon_n \right) < \eta \delta^2, \quad c_1\alpha > 1, \quad 0 < \eta < 1/2. \quad (1.22)$$

Here, we are free to choose α and η , but c_1 and δ are fixed. Now let

$$\alpha = \frac{1}{c_1} + \epsilon, \quad \eta = \frac{1}{2} - \epsilon. \quad (1.23)$$

Then the second and third inequalities in (1.22) are satisfied. The first is equivalent to

$$\frac{4}{c_1 \delta^2} + \epsilon < \frac{1}{2} - \epsilon \quad (1.24)$$

as $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$. In addition, since ϵ is arbitrary, we have again, equivalently,

$$\frac{4}{c_1 \delta^2} - \frac{1}{2} < 0. \quad (1.25)$$

When this is satisfied, we have:

$$\sup_{\substack{\sigma \in [\delta, \sqrt{n}] \\ i=1, \dots, n}} \left| \int f_\sigma(X_i, x') d\mu_n(x') - \int f_\sigma(X_i, x') d\mu(x') \right| = O(n^{4/c_1 \delta^2 - 1/2} \sqrt{\log n}). \quad (1.26)$$

The proof is thus finished. $\square$

Then combining the previous result we have an important convergence:

Proposition 2. *Let μ be sub-Gaussian and c_1 be the constant from lemma 1. $X_1, \dots, X_n$ are i.i.d. random variables with probability measure μ and $\mu_n = \frac{1}{n} \sum \delta_{X_i}$ is the empirical measure. Additionally, set $\delta > 0$ to be a constant such that $c_1 \delta^2 > 16$. Then, there exists $\epsilon > 0$, such that almost surely in $X_1, \dots, X_n$:*

$$\sup_{\substack{\sigma \in [\delta, \sqrt{n}] \\ i=1, \dots, n}} |F_{\mu_n}(X_i, \sigma) - F_\mu(X_i, \sigma)| = O(n^{-\epsilon}). \quad (1.27)$$

Let us summarize the important results upon this point, which will be used later:

There exists $\gamma = 2/c_1 + \epsilon$ such that the following holds almost surely

- (1) $\max_{i=1, \dots, n} \|X_i\| \leq \sqrt{\gamma \log n}$;
- (2) $\max_{i,j} \|X_i - X_j\|^2 \leq 4\gamma \log n$;
- (3) For $\sigma \in [\delta, \sqrt{n}]$,

$$\int \exp(-\|X_i - x'\|^2 / 2\sigma^2) d\mu_n(x), \int \exp(-\|X_i - x'\|^2 / 2\sigma^2) d\mu(x') \geq C n^{-2\gamma/\sigma^2}. \quad (1.28)$$

Next couple of lemmas in [1] are about lower bounds on $\sigma_{\rho, \mu}^*$. Recall that $\sigma_{\rho, \mu}^*$ is the unique solution of the equation $F_{\rho, \mu}(x, \sigma_{\rho, \mu}^*) = 0$. For simplicity of notation, we write $\sigma^*(x) = \sigma_{\rho, \mu}^*(x)$ and $\sigma_n^*(x) = \sigma_{\rho, \mu_n}^*(x)$.

Lemma 6. *If μ satisfies the condition in lemma 1 of [1], then $\sigma^*(x)$ and $\sigma_n^*(x)$ are uniformly bounded from below in both i and n almost surely.*

Proposition 3. *Suppose μ has compact support and C^1 density. Then almost surely, as $n \rightarrow \infty$,*

$$\sup_{i=1, \dots, n} |\sigma_n^*(X_i) - \sigma^*(X_i)| \rightarrow 0. \quad (1.29)$$

Proof. By [1], we can quickly reach to the stage where

$$|\sigma_n^*(x) - \sigma^*(x)| = O(|F_{\mu_n}(x, \sigma_n^*(x)) - F_{\mu}(x, \sigma_n^*(x))|) \rightarrow 0. \quad (1.30)$$

Moreover, we know that μ has compact support and $x \mapsto \sigma^*(x)$ is smooth, $\sigma^*(X_i)$ is bounded above and below, meaning that $\sigma_n^*(X_i)$ is bounded uniformly from above and below as well. This completes the proof in the original paper. $\square$

2 More Lemmas

Starting this section we are gradually approaching our point of work. We need to list a few more lemmas before that. To make notations cleaner, [1] introduced some abbreviation in section 4.2. We will use all of them by default.

Lemma 7. *Any sequence in $\tilde{\mathcal{P}}_X$ is tight.*

Recall Prokhorov's theorem:

Theorem (Prokhorov's Theorem). *If μ_n is a tight sequence of measures in Euclidean space. There exists a subsequence μ_{n_k} and a measure μ such that μ_{n_k} converges to μ weakly.*

We immediately get the following lemma:

Lemma 8. *There exists μ^* such that*

$$I_\rho(\mu^*) = \inf_{\mu \in \tilde{\mathcal{P}}_X} I_\rho(\mu) \quad (2.1)$$

Recall that $I_\rho(\mu)$ is the continuous version of $L_{n,\rho}(X, Y)$. It only depends on the measure μ and the perplexity constant ρ . We have not reached the part that requires we have a common perplexity, but from now on we should pay close attention as it is one of the core setup of the original paper.

Lemma 9. *As $n \rightarrow \infty$, for some $\epsilon > 0$, we have*

$$|f_n(X_i, X_j) - f(X_i, X_j)| = O(n^{-\epsilon}). \quad (2.2)$$

Proof. There are typos in the original proof so we prove this lemmas again despite it being very easy and elementary. We compute directly

$$\begin{aligned} & |f_n(x, x') - f(x, x')| = |\exp(-\|x - x'\|^2 / 2\sigma_n(x)^2) - \exp(-\|x - x'\|^2 / 2\sigma(x)^2)| \\ &= \left| \exp\left(\frac{|\sigma_n(x)^2 - \sigma(x)^2| \|x - x'\|^2}{2\sigma_n(x)^2 \sigma(x)^2}\right) - 1 \right| \cdot \exp(-\|x - x'\|^2 / 2\sigma(x)^2) \\ &\leq \left| \exp\left(\frac{|\sigma_n(x)^2 - \sigma(x)^2| \|x - x'\|^2}{2\sigma_n(x)^2 \sigma(x)^2}\right) - 1 \right|. \end{aligned} \quad (2.3)$$

Recall that the sum of two sub-Gaussian random variables is still sub-Gaussian. If μ_X is sub-Gaussian, let C be the constant of decay for $X_i - X_j$ and we have

$$\mathbb{P}(|X_i - X_j|^2 \geq \log n) \leq 2 \exp(-\log n / C^2) = 2 \exp\left(-\frac{1}{C^2}\right) \cdot \frac{1}{n} \rightarrow 0. \quad (2.4)$$

This means $|X_i - X_j|^2 = O(\log n)$ almost surely. On the other hand, if μ_X has compact support then $|X_i - X_j|$ will be bounded, even easier. Moreover, since σ_n and σ are both uniformly bounded below by lemma (6), by setting $x = X_i$ and $x' = X_j$ we get

$$\begin{aligned}
& \left| \exp \left(\frac{|\sigma_n(x)^2 - \sigma(x)^2| \|x - x'\|^2}{2\sigma_n(x)^2 \sigma(x)^2} \right) \right| \\
& \leq \frac{1}{\sigma_n(x) \sigma(x)} \left(\frac{1}{\sigma_n(x)} + \frac{1}{\sigma(x)} \right) |\sigma_n(x) - \sigma(x)| \|x - x'\|^2 \\
& = O(|\sigma_n(x) - \sigma(x)| \|x - x'\|^2) \\
& = O(|\sigma_n(x) - \sigma(x)|) O(\log n) \\
& = O(n^{-\epsilon}) O(\log n) \\
& = O(n^{-\epsilon}),
\end{aligned} \tag{2.5}$$

by proposition (3). This finishes the proof. $\square$

One thing we should highlight here is that we are abusing the notation of σ and σ_n . They are defined as the unique solution to $F = 0$. However, the definition of σ_i is not abused and we will later see why such abuse makes sense.

Lemma 10. *If $c_1 \delta > 32$, then there exists $\epsilon > 0$ such that as $n \rightarrow \infty$, we have*

$$\begin{aligned}
\sup_{i,j} |p'_n(X_i, X_j) - p'(X_i, X_j)| &= O(n^{-\epsilon}), \\
\sup_{i,j} |p_n(X_i, X_j) - p(X_i, X_j)| &= O(n^{-\epsilon}).
\end{aligned} \tag{2.6}$$

Supplementary Proof. The overall methodology of the original proof in [1] is natural and clear. However, it can still be slightly confusing as it is missing the actual values we assign to ϵ . In the original paper, we can take, for some constant $k > 0$, that

$$n^{-\epsilon} = n^{8/c_1 \delta^2 - 1/2} (\log n)^k. \tag{2.7}$$

This comes from the detailed computations in the original proof and the quantity above is the slowest decaying speed, the 'worst case scenario' given all the conditions. Plug this into the computations and everything is clear. The proof uses a little bit extra than just the bounds in section 3.1. $\square$

Lemma 11. *If $c_1 \delta > 40$, then there exists $\epsilon > 0$ such that as $n \rightarrow \infty$, we have*

$$\sup_{i,j} |p_n(X_i, X_j) \log p_n(X_i, X_j) - p(X_i, X_j) \log p(X_i, X_j)| = O(n^{-\epsilon}). \tag{2.8}$$

Lemma 12. *If $c_1 \delta^2 > 8$, then as $n \rightarrow \infty$ we have*

$$\sum_j p_{j|i} \log p_{j|i} + \log(\rho n) = o(1). \tag{2.9}$$

[Yi: TODO: Ask Tuca about the last step of the proof in [1].]

3 The Actual Computation

From this point onward, we are starting to have lemmas and results related to the output kernel g and this is where the trail begins. Along the way we will see if there is a need to add more conditions to g and if we do, every time we add more constraints we will reiterate the entire settings for g for clarity. To refresh our memory, here is the initial assumptions we have for g : $g(y, y') = k(\|y - y'\|)$ such that:

1. $k : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is decreasing and bounded.
2. k satisfies

$$\iint_{(0, \infty)^2} k(x, x') dx dx' < \infty. \quad (3.1)$$

Lemma 13. *There exists a set of points $y^*(X)$ such that*

$$\inf_{Y \in (\mathbb{R}^s)^n} L_{n, \rho}(X, Y) = L_{n, \rho}(X, y^*). \quad (3.2)$$

[Yi: TODO: Ask Tuca about the $1/n^2$ in the original proof. The scaling seems to not play a role in the proof.]

Proof. Fortunately, the first proof is almost identical to the original one. Let $k \in \{1, \dots, n\}$. We are going to show that

$$\lim_{|y_k| \rightarrow \infty} L_{n, \rho}(X, y) = \infty. \quad (3.3)$$

By definition, L is the relative entropy and

$$L_{n, \rho}(X, y) = \sum_{i \neq j} p_{ij} \log \left(\frac{p_{ij}}{q_{ij}} \right) = \sum_{i \neq j} p_{ij} \log p_{ij} - \sum_{i \neq j} p_{ij} \log q_{ij}. \quad (3.4)$$

For n given and fixed, it suffices to show that

$$\sum_{i \neq j} p_{ij} \log(q_{ij}) = \sum_{i \neq j} p_{ij} \log \left(\frac{g(y_i, y_j)}{\sum_{m \neq l} g(y_m, y_l)} \right) \rightarrow -\infty. \quad (3.5)$$

Recall that $g(y, y')$ is bounded. We have

$$\begin{aligned} & - \sum_{i \neq j} p_{ij} \log \left(\sum_{m \neq l} g(y_m, y_l) \right) = - \log \left(\sum_{m \neq l} g(y_m, y_l) \right) \\ & \leq - \sum_{i \neq j} p_{ij} \log \left(\sum_{m \neq k, l \neq k, m \neq l} g(y_m, y_l) \right) < \infty. \end{aligned} \quad (3.6)$$

On the other hand, we have $g(y_i, y_k) = k(\|y_i - y_k\|) \rightarrow 0$ as $|y_k| \rightarrow \infty$ because $\|y_i - y_k\| \rightarrow \infty$ and k satisfies

$$\iint_{(0, \infty)^2} k(x, x') dx dx' < \infty. \quad (3.7)$$

This means

$$\lim_{|y_k| \rightarrow \infty} \sum_{i \neq j} p_{ij} \log(g(y_i, y_j)) \rightarrow -\infty. \quad (3.8)$$

Hence, the minimum of the $L_{n, \rho}(X, y)$ is achieved on a bounded set, the lemma follows by the continuity of $L_{n, \rho}(X, y)$.

□

Lemma 14. *As $n \rightarrow \infty$, we have*

$$\sum_{i \neq j} p_{ij} \log p_{ij} = \iint p(x, x') \log p(x, x') d\mu d\mu - 2 \log n + o(1). \quad (3.9)$$

Proof. The original paper does provide a detailed proof, so it is our turn to exercise. First recall that $p_{ij} = p_n(X_i, X_j)/n^2$. Therefore we have

$$\begin{aligned} \sum_{i \neq j} p_{ij} \log p_{ij} &= \sum_{i \neq j} \frac{1}{n^2} p_n(X_i, X_j) \log \left(\frac{1}{n^2} p_n(X_i, X_j) \right) \\ &= \frac{1}{n^2} \sum_{i \neq j} p_n(X_i, X_j) \log(p_n(X_i, X_j)) - 2 \log n \sum_{i \neq j} \frac{1}{n^2} p_n(X_i, X_j) \\ &= \frac{1}{n^2} \sum_{i \neq j} p_n(X_i, X_j) \log(p_n(X_i, X_j)) - 2 \log n, \end{aligned} \quad (3.10)$$

because the second summation is just the sum of all probability mass. Now we have the $2 \log n$ term and we are left to deal with the first summation. Inspired by the proofs in [1], we begin by showing that the diagonal terms do not matter in our estimation. consider

$$\begin{aligned} &\frac{1}{n^2} \sum_{i \neq j} p_n(X_i, X_j) \log(p_n(X_i, X_j)) - \iint p_n(x, x') \log p_n(x, x') d\mu_n(x) d\mu_n(x') \\ &= -\frac{1}{n^2} \sum_{t=1}^n p'_n(X_t, X_t) \\ &= -\frac{1}{n^2} \sum_{t=1}^n \frac{1}{\int f(X_t, x') d\mu_n(x') - \frac{1}{n}} \log \left(\frac{1}{\int f(X_t, x') d\mu_n(x') - \frac{1}{n}} \right). \end{aligned} \quad (3.11)$$

If we refer to the proof of lemma 14 in [1], we can see that the difference of diagonal term vanishes and it does not depend on the specific X_i . Therefore in our proof, when we sum over all the X_i and normalize by dividing n^2 , we reach the exact same conclusion. The diagonal term is $O(n^{4/c_1 \delta^2 - 1} \log n)$ which is $o(1)$ depending on our choice of c_1 .

Then we apply lemma 11 and since $d\mu_n \times d\mu_n$ is a probability measure, we have

$$\iint p_n(x, x') \log p_n(x, x') d\mu_n(x) d\mu_n(x') = \iint p(x, x') \log p(x, x') d\mu_n(x) d\mu_n(x') + o(1), \quad (3.12)$$

which immediately implies

$$\sum_{i \neq j} p_{ij} \log p_{ij} = \iint p(x, x') \log p(x, x') d\mu_n(x) d\mu_n(x') - 2 \log n + o(1). \quad (3.13)$$

The proof will be finished if we are able to replace the measure in the last integral with $d\mu$. We verify that

$$h(x, x', \sigma) := p(x, x') \log p(x, x') \quad (3.14)$$

satisfies all the requirements in lemma 3. First μ is sub-Gaussian by construction. Then is easy to verify that h is C^1 for all x, x' and σ . Moreover, by calculus we have $p(x, x') \in (0, 1]$ and $\lim_{a \rightarrow 0} a \log a = 0$. and $p \rightarrow 0$ as $\|x - x'\|/\sigma \rightarrow \infty$, it is obvious that h is bounded and $h \rightarrow 0$ as $\|x - x'\|/\sigma \rightarrow \infty$. The rest of the conditions are also very easily to prove. We can finally conclude that

$$\iint p(x, x') \log p(x, x') d\mu_n(x) d\mu_n(x') = \iint p(x, x') \log p(x, x') d\mu(x) d\mu(x') + o(1). \quad (3.15)$$

Combine this estimation with (3.13) and we can conclude

$$\sum_{i \neq j} p_{ij} \log ij = \iint p(x, x') \log p(x, x') d\mu(x) d\mu(x') - 2 \log n + o(1). \quad (3.16)$$

The proof is thus finished. □

References

- [1] Antonio Auffinger and Daniel Fletcher. Equilibrium distributions for t-distributed stochastic neighbour embedding, 2023, 2304.03727. URL <https://arxiv.org/abs/2304.03727>.
- [2] Aditya Ravuri and Neil D. Lawrence. Towards one model for classical dimensionality reduction: A probabilistic perspective on umap and t-sne, 2024, 2405.17412. URL <https://arxiv.org/abs/2405.17412>.
- [3] Pavlin G. Poličar and Blaž Zupan. Visualizing high-dimensional temporal data using direction-aware t-sne, 2024, 2403.19040. URL <https://arxiv.org/abs/2403.19040>.
- [4] Laurens van der Maaten and Geoffrey Hinton. Visualizing data using t-sne. *Journal of Machine Learning Research*, 9(2579-2605), 2008.
- [5] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.