

Final Endeavor for the Generalized t -SNE Problem

Yi Gu

January 1, 2025

1 Progress So Far and Setup

OK this is it. We have reached this step and the end is within our grasp. The stake is high and I can literally feel the adrenaline coursing my vein, along with my fear. So far we have finished two crucial steps. First, we have finished the proof of convergence for the generalized output kernel. Denote $\mathbf{Ker}_\mathbf{O}$ the output kernel and $\mathbf{Ker}_\mathbf{I}$ the input kernel. The main theorems in [1] will hold if the following is true:

Condition 1 (Output Kernel Conditions). *The output kernel $\mathbf{Ker}_\mathbf{O}(y, y') = k(\|y - y'\|)$ depends only on the distance between y and y' and that:*

1. $k : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is decreasing and bounded;
2. g satisfies

$$\iint_{(0, \infty)^2} g(y, y') dy dy' < \infty; \quad (1.1)$$

3. k has bounded derivative;
4. $k'(0) = 0$.

Moreover, the lemmas that deal with the output does not interfere with those related to the input kernel. The initial exploration we did around the output kernel not only provides us a good opportunity to understand the problem and the literature, but also serves as an integral part of the final paper. If we are able to tackle the lemmas related to the input kernel we will be able to join the results seamlessly.

On the other hand, we have established the form of the generalized input kernel, set a basis for us to work on, and prove the existence of $\sigma_{\rho, \mu}^*(x)$, the unique solution to $F_{\rho, \mu}(x, \sigma) = 0$. For this to happen, the generalized input kernel should satisfy the following conditions:

Condition 2 (Initial Input Kernel Conditions). *Let $w : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be some fixed function. The input kernel has the form of $\mathbf{Ker}_\mathbf{I}(x_i, x_j, \sigma_i) = \exp(-w(\sigma_i \|x_j - x_i\|^\theta))$. Here, σ_i is used to control entropy (thus perplexity) and $\theta > 0$ is a fixed constant. (We can replace σ_i with $\psi : \mathbb{R}^d \rightarrow \mathbb{R}^+$ in the continuous version of the kernel.) w satisfies:*

1. $w'(t) > 0$ for all $t \geq 0$ with $\lim_{t \rightarrow \infty} w(t) = \infty$;
2. $w'(t) + tw''(t) \geq 0$ for all $t \geq 0$;
3. The following two integrals converge:

$$\int_0^\infty t^{d-1} \exp(-w(t^\theta)) dt < \infty, \quad \int_0^\infty t^{d-1} w(t^\theta) \exp(-w(t^\theta)) dt < \infty. \quad (1.2)$$

Since the probability masses are all normalized, we can assume without loss of generality that $w(0) = 0$. Note that condition 2 is only enough to prove the existence and uniqueness of σ^* . We may add more assumptions later on if necessary. Before we march towards the final battle, let us remind ourselves some important definitions and notations, as well as some of the lemmas from [1] that we can borrow. Recall that X or the corresponding probability measure μ is called sub-Gaussian if $\mathbb{P}(X > t) \leq 2 \exp(-Ct^2)$ for some positive constant C .

Lemma 1. *Let ν be sub-Gaussian with second moment equal to 1 and suppose $X \in \mathbb{R}^d$ has i.i.d. entries drawn from ν . Then there exists constant C, c_1 such that*

$$\mathbb{P}(\|X\| \geq \sqrt{d} + t) \leq Ce^{-c_1 t^2}. \quad (1.3)$$

Lemma 2. *Let μ be sub-Gaussian and c_1 be the constant from lemma 1. Let $X_1, \dots, X_n \in \mathbb{R}^d$ have i.i.d. entries drawn from μ . Then there exists a constant C such that*

$$\mathbb{P}\left(\max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\frac{\log n}{c_1}} + \sqrt{d} + t\right) \leq Ce^{-c_1 t^2}. \quad (1.4)$$

Lemma 3. *Let $\delta > 0$ be constant, $K_n = [\delta, \sqrt{n}]$ and $h : \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$. Suppose the following hold:*

1. μ is sub-Gaussian;
2. $h(x, x', \sigma)$ and $\nabla_x h(x, x', \sigma)$ are bounded for all x, x' and all $\delta \in K_n$;
3. $h(x, x', \sigma) \rightarrow 0$ as $\sigma \|x - x'\| \rightarrow \infty$.

Let $X_1, \dots, X_n$ be i.i.d. random variables drawn from μ . Let $\mu_n = (1/n) \sum_i \delta_{X_i}$. Then for any $\epsilon > 0$ with probability at least $1 - e^{-c\epsilon^2 n}$ we have

$$\sup_{\sigma^{-1} \in K_n, x} \left| \int h(x, x', \sigma) d\mu_n(x) - \int h(x, x', \sigma) d\mu(x') \right| \leq \epsilon + O\left(\sqrt{\frac{\log n}{n}}\right). \quad (1.5)$$

Finally, let us also recall that

$$\begin{aligned} & F_{\rho, \mu}(x, \sigma) \\ &= \int \frac{\exp(-w(\sigma \|x - x'\|^\theta))}{\int \exp(-w(\sigma \|x - x'\|^\theta)) d\mu(x')} \log \left(\frac{\exp(-w(\sigma \|x - x'\|^\theta))}{\int \exp(-w(\sigma \|x - x'\|^\theta)) d\mu(x')} \right) d\mu(x') + \log \rho \\ &= - \int \frac{w(\sigma \|x - x'\|^\theta) \exp(-w(\sigma \|x - x'\|^\theta))}{\int \exp(-w(\sigma \|x - x'\|^\theta)) d\mu(x')} d\mu(x') \\ &\quad - \log \left(\int \exp(-w(\sigma \|x - x'\|^\theta)) d\mu(x') \right) + \log \rho. \end{aligned} \quad (1.6)$$

The two terms in the formula of F are the pillars of this manuscript. Their convergence will lead us to prove the convergence of F , and most importantly, the uniform convergence of $\sigma(x)$. **[Yi: As for the question Tuca asked me during our discussion, yes we can define $\sigma(x)$ to be the unique solution to $F(x, \sigma) = 3$ or whatever number we peek as long as the number is greatly than zero (to ensure the existence of uniqueness of the solution). If we move the constant to the left it is equivalent to just pick a different perplexity. The convergence of F works for any σ and that is the key. As long as control the entropy, we are expecting convergence as long as we set up the continuous version to have the same perplexity.]**

2 The Two Most Important Lemmas

The biggest obstacles left are to proof two lemmas similar to lemma 6 and lemma 7 in [1]. These two lemmas will give us the convergence of F and thus the convergence of $\sigma(x)$. We

will try to tackle this part using similar methods from the original paper. Write

$$\begin{aligned} f_\sigma(x, x') &= \exp(-w(\sigma \|x - x'\|^\theta)), \\ g_\sigma(x, x') &= w(\sigma \|x - x'\|^\theta) \exp(-w(\sigma \|x - x'\|^\theta)). \end{aligned} \quad (2.1)$$

Note that for any real numbers A, B, C, D , we have

$$\left| \frac{A}{B} - \frac{C}{D} \right| = \left| \frac{AD - BC}{BD} \right| = \left| \frac{D(A - C) + C(D - B)}{BD} \right| \leq \left| \frac{A - C}{B} \right| + \left| \frac{C(D - B)}{BD} \right|. \quad (2.2)$$

This means

$$\begin{aligned} \left| \frac{\int g_\sigma(x, x') d\mu_n(x')}{\int f_\sigma(x, x') d\mu_n(x')} - \frac{\int g_\sigma(x, x') d\mu(x')}{\int f_\sigma(x, x') d\mu(x')} \right| &\leq \left| \frac{\int g_\sigma(x, x') d\mu_n(x') - \int g_\sigma(x, x') d\mu(x')}{\int f_\sigma(x, x') d\mu_n(x')} \right| \\ &+ \left| \frac{\int g_\sigma(x, x') d\mu(x') (\int f_\sigma(x, x') d\mu_n(x') - \int f_\sigma(x, x') d\mu(x'))}{\int f_\sigma(x, x') d\mu(x') \int f_\sigma(x, x') d\mu_n(x')} \right|. \end{aligned} \quad (2.3)$$

Let $\alpha > 0$. Define

$$E'_1 = \left\{ \max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\left(\alpha + \frac{1}{c_1}\right) \sqrt{\log n}} + \sqrt{d} \right\}. \quad (2.4)$$

Setting $t = \sqrt{\alpha \log n}$ in lemma 2 we have $\mathbb{P}(E'_1) \leq Cn^{-c_1\alpha}$. To absorb the $\sqrt{d}$ term in E'_1 , we can find $\epsilon_n \rightarrow 0$ such that

$$E_1 = \left\{ \max_{i=1, \dots, n} \|X_i\| \geq \sqrt{\left(\alpha + \frac{1}{c_1} + \epsilon_n\right) \sqrt{\log n}} \right\} \quad (2.5)$$

with $\mathbb{P}(E_1) \leq Cn^{-c_1\alpha}$. On the other hand, on E_1^c for all i, j ,

$$\|X_i - X_j\| \leq 2\sqrt{\left(\alpha + \frac{1}{c_1} + \epsilon_n\right) \sqrt{\log n}}. \quad (2.6)$$

3 The Rest of the Proof

4 Wrap Up and Conclusion

References

- [1] Antonio Auffinger and Daniel Fletcher. Equilibrium distributions for t-distributed stochastic neighbour embedding, 2023, 2304.03727. URL <https://arxiv.org/abs/2304.03727>.
- [2] Aditya Ravuri and Neil D. Lawrence. Towards one model for classical dimensionality reduction: A probabilistic perspective on umap and t-sne, 2024, 2405.17412. URL <https://arxiv.org/abs/2405.17412>.
- [3] Pavlin G. Poličar and Blaž Zupan. Visualizing high-dimensional temporal data using direction-aware t-sne, 2024, 2403.19040. URL <https://arxiv.org/abs/2403.19040>.
- [4] Laurens van der Maaten and Geoffrey Hinton. Visualizing data using t-sne. *Journal of Machine Learning Research*, 9(2579-2605), 2008.
- [5] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.
- [6] Ben Weinkove. Dimension reduction and the gradient flow of relative entropy, 2024, 2409.16963. URL <https://arxiv.org/abs/2409.16963>.